

Nurse-led interventions among older adults affected by cancer: an integrative review.

ALEMANIA, E., SAMARA, J., RALPH, N. and PATERSON, C.


2023

© 2023 The Author(s). Published by Elsevier Inc. on behalf of Asian Oncology Nursing Society.

Journal Pre-proof

Nurse-led interventions among older adults affected by cancer: An integrative review

E. Alemania, GradCert Renal Nursing, BN, A. Hind, RN, MGer, J. Samara, MNNP, GradCert Cancer & Haematology Nursing, BN, M. Turner, BAppSci, MBA, N. Ralph, PhD, BN, MCLinPrac, GCertClinTrialsRes, C. Paterson, PhD, BA, MSc, PgCert LTA, FHEA, RAN



PII: S2347-5625(23)00107-5

DOI: <https://doi.org/10.1016/j.apjon.2023.100289>

Reference: APJON 100289

To appear in: *Asia-Pacific Journal of Oncology Nursing*

Received Date: 1 June 2023

Revised Date: 26 July 2023

Accepted Date: 31 July 2023

Please cite this article as: Alemania E, Hind A, Samara J, Turner M, Ralph N, Paterson C, Nurse-led interventions among older adults affected by cancer: An integrative review, *Asia-Pacific Journal of Oncology Nursing*, <https://doi.org/10.1016/j.apjon.2023.100289>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Inc. on behalf of Asian Oncology Nursing Society.

Title: Nurse-led interventions among older adults affected by cancer: An integrative review

Authors: Alemania, E., (GradCert Renal Nursing, BN)^a, Hind, A., (RN, MGer)^a, Samara, J., (MNNP, GradCert Cancer & Haematology Nursing, BN)^f, Turner, M., (BAppSci, MBA)^{ab}, Ralph, N., (PhD, BN, MClinPrac, GCertClinTrialsRes)^{g,h}, Paterson, C (PhD, BA, MSc, PgCert LTA, FHEA, RAN)^{a,b,c,d,e}

a Faculty of Health, University of Canberra, Bruce ACT, Australia

b Prehabilitation, Activity, Cancer, Exercise and Survivorship (PACES) Research Group, University of Canberra, Bruce ACT, Australia

c School of Nursing, Midwifery and Public Health, University of Canberra, Bruce ACT, Australia

d Canberra Health Services & ACT Health, SYNERGY Nursing & Midwifery Research Centre, ACT Health Directorate Level 3, Building 6, Canberra Hospital, Australia

e Robert Gordon University, Aberdeen, Scotland, UK

f Calvary Public Hospital Bruce, Clare Holland House Specialist Palliative Aged Care, Barton ACT, Australia

g School of Nursing & Midwifery, University of Southern Queensland, Ipswich, QLD, Australia

h Centre for Health Research, University of Southern Queensland, Springfield, QLD, Australia

Corresponding Author:

Professor Catherine Paterson

Clinical Professor of Nursing

Faculty of Health, University of Canberra, Bruce ACT, Australia

Email: catherine.paterson@canberra.edu.au

Declarations

Funding: none

Conflicts of interest/Competing interests: none to declare

Ethics approval: not applicable

Consent to participate: not applicable

Consent for publication: not applicable

Availability of data and material: not applicable

Code availability: not applicable

Authors' contributions: CRediT author statement

Elizabeth Alemania: Screening, data extraction, reviewing, writing, reviewing final draft.

Alicia Hind: Screening, data extraction, reviewing.

Juliana Samara: Reviewing, editing draft, writing, reviewing final draft.

Murray Turner: Database search strategy design and execution, data extraction, writing, reviewing, editing.

Nick Ralph: Writing, reviewing

Catherine Paterson: Conceptualization, methodology, validation, screening, data extraction, formal analysis, interpretation, writing and reviewing & editing, supervision.

Acknowledgments:

We would like to extend our thanks to the NSW and ACT Australia and New Zealand Urological Nurses Society for providing a research fellowship for this work.

Journal Pre-proof

Title: Nurse-led interventions among older adults affected by cancer: An integrative review

Objective: Aging can introduce significant changes in health, cognition, function, social, and emotional status among older adults affected by cancer. Little is known about how existing nurse-led interventions address the needs of older adults. The objective was to identify existing nurse-led interventions among older adults to optimize recovery and survivorship needs.

Methods: A integrative systematic review was reported conducted according to the PRISMA 2020 Guidelines. Electronic databases (APA PsycINFO, CINAHL, MEDLINE, Scopus, and Google Scholar databases) were searched using key search terms. Articles were assessed for inclusion according to a pre-determined eligibility criterion. Data extraction and quality appraisal were conducted. Findings were integrated in a narrative synthesis.

Results: 21 studies were included and a total of 4,253 participants represented. There were a range of study designs: quantitative (n=10), randomised controlled trials (n=6), mixed methods studies (n=3), qualitative (n=1) and a non-randomised controlled study (n=1). Most participants had prostate cancer with some representation in colorectal, lung, head and neck, renal, oesophageal, and mixed cancer patient populations.

Conclusion: This review shows a lack of evidence on the inclusion of geriatric assessments for older people with cancer within existing nurse-led interventions. Further research is needed to test nurse-led interventions with the inclusion of geriatric assessments, and their contribution in the multidisciplinary team across the cancer care continuum for various cancer patient populations.

Key words: older people; medical oncology; oncology nursing; integrative oncology; geriatric assessment

1. Introduction

As the population ages, significant numbers of older adults will be diagnosed with cancer.^{1,2} The care of older adults with cancer is an increasing challenge for multidisciplinary oncology healthcare teams globally.³ Treatment of older patients with cancer will be an increasing challenge as the population ages because cancer is primarily a disease of older people.⁴ Specifically, the 2015 World Report on Ageing and Health underscores that the number of people older than 60 years will double by 2050.² There is a need to develop new initiatives to improve the quality of care for older adults with cancer, and to translate them into broader standards of person-centred care.⁵

Aging can introduce significant changes in health, cognition, function, social, and emotional status. Therefore, addressing the supportive care needs of the older adult affected by cancer is complex^{6,7}, underscoring the importance of comprehensive geriatric assessment and care to improve quality of life, reduce decrements in health, avoid complications, and reduce the risk of hospitalization or prolonged hospital stay. It is essential to integrate geriatric principles of care into oncology⁸, which includes the recognition of: (1) multiple chronic conditions, (2) polypharmacy, (3) social determinants of health, (4) screening for geriatric syndromes, and (5) incorporating functional assessments in practice across the cancer care continuum.⁹ As ageing is related to an alteration in physiologic functions, individualised treatment options in older patients with cancer needs to be evaluated.¹⁰ Complex health care conditions in the elderly are multifactorial and compounded by geriatric syndromes (e.g., falls, nutritional deficits, sensory loss, cognitive impairment, frailty, multiple chronic conditions, and functional status) largely not addressed in oncology care.^{6,7,11-13}

Identifying appropriate cancer treatments in older people can be complicated due to the presence of chronic health conditions and different health priorities.¹⁴ Many older people

with cancer do however, tolerate cancer treatments¹² but many will not have access to tailored treatment options sensitive to the complex issues compounded by geriatric health, largely due to a lack of representation in clinical trials.¹⁵ Geriatric conditions and frailty can lead to loss of independence, transition to a higher level of care, longer hospital stays, and higher mortality rates, all of which can negatively impact cancer survivorship and recovery. Furthermore, the majority of older adults living with cancer are likely to have two or more long-term conditions.¹⁶

Older adults affected by cancer are more likely to have an increased prevalence of multimorbidity compared to those without cancer demonstrated in age-matched control group research.¹⁷ The practicalities of implementing comprehensive geriatric oncology models of care have been acknowledged as a stumbling point in clinical services.¹⁸ Challenges often relate to a lack of time and limited access to specialised healthcare professionals who can perform a comprehensive assessment in practice. Oncology nurses are central to the multidisciplinary team (MDT) to promote and optimise patient-centred care for all older adults with cancer, and are necessary to achieve optimum care.¹⁹ Internationally, there has been a development of various nurse-led geriatric oncology models of care.²⁰ However, little is known about oncology nurse-led interventions in older adults and whether comprehensive geriatric assessments are included, and whether nursing assessments and interventions are sensitive to the unique needs of this older patient group. Therefore, this integrative systematic review set out to identify existing nurse-led interventions in older adults affected by cancer to advance future directions for practice and research.²¹

2. Methods

Design: This integrative systematic review was conducted and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines.

²² This review also followed a systematic review protocol available on request.

2.1 Eligibility Criteria

Types of studies

This review included all qualitative, quantitative, and mixed methods studies irrespective of research design, with the only limiter of being published in the English language. All commentaries, editorials, and studies where nurse-led interventions for patients affected by cancer with a mean age of <65 years were excluded.

Types of participants

All older participants (where studies reported a mean study age of ≥ 65 years) diagnosed with cancer (irrespective of cancer stage, treatment, or time since diagnosis) and received a nurse-led intervention were included. All other long-term conditions, younger study samples <65 years, and non-nurse led interventions were excluded.

Types of outcomes

The primary outcome of this review was related to supportive care needs provided by nurse-led interventions for older adults in study samples with a mean age of ≥ 65 years. The Supportive Care Needs Framework ⁶ guided the classification of supportive care provided by nurse-led interventions through the identification of inclusion of comprehensive geriatric assessments.

2.2. Literature search

The APA PsycINFO, CINAHL, MEDLINE, Scopus, and Google Scholar databases were searched from inception to September 2022 for all relevant studies published. To capture as many studies as possible, the database search architecture utilised a wide range of key words (nurse-led OR nurse-managed) (care OR model* OR program* OR intervention*) AND (cancer* OR oncolog* OR neoplasm*) designed and conducted by an expert systematic review librarian, see Supplementary Table 1. Limiters were placed on all searches for studies published in the English language, no other limiters were set to ensure all nurse-led interventions were identified to be as inclusive and sensitive as possible. All articles were assessed independently by two reviewers to identified studies in older people where studies reported a mean study age of ≥ 65 years) diagnosed with cancer (irrespective of cancer stage, treatment, or time since diagnosis) and received a nurse-led intervention. The reference lists of all the full-text articles included were checked to identify any other relevant studies. Citations were managed with EndNote 20 and imported into Covidence systematic review software to facilitate the systematic review process.

2.3 Selection of studies

Duplication of publications were removed in Covidence. All titles and abstracts were independently screened by three reviewers for eligibility, and any disagreements were resolved by discussion. Full text publications were then retrieved, independently screened by reviewers, and linked multiple records of the same study together. Any disagreements were resolved by discussion and reasons for exclusion documented.

2.4 Data extraction and management

Data extraction was performed on the retained full-text studies meeting the inclusion criteria. The data was extracted by three reviewers, and independently quality checked among all

reviewers for accuracy and quality assurance in the process. The data extraction tables were developed and tested on a small sample of studies and then further refined through discussion among the reviewers. The first table of data extraction included information on the purpose, setting, country, sample size, participant characteristics, sampling used, response rate, attrition, design, time points, and data collection tools. The second data extraction table related to the nurse-led interventions and the supportive care needs outcome data (see **Table 1**)²³.

2.5 Assessment of risk of bias in included studies

The included full-text studies all underwent a methodological quality assessment. None of the studies were excluded based upon their methodological quality score to enable a comprehensive overview of the current state of the evidence. The methodological quality assessment was conducted using the Mixed Methods Assessment Tool (MMAT).²⁴ The MMAT tool was selected because it enabled a plethora of study designs to be critically appraised. This assessment tool enabled critical appraisal of all qualitative, quantitative, and mixed methods studies. Each domain of assessment was rated against, “no”, “yes” and “unclear”. Methodological quality assessment was performed by one reviewer and quality checked by a second reviewer until consensus reached.

2.6 Data synthesis

This integrative systematic review used a narrative synthesis.²⁵ The steps in the narrative synthesis involved: 1) data reduction by tabulation, 2) data comparison between studies, and finally, 3) drawing conclusions. This process involved reading and re-reading full text publications linking together similarities and differences between the studies, and quality checking with the primary sources. The data comparison phase involved the reviewers' identifying patterns and themes through counting, clustering, and making comparisons across

the study findings in tabulated format grouped together by cancer type. The data synthesis was conducted by two authors and checked by a third author.

3. Results

The search identified a total of 1244 publications, see **Figure 1**. A total of 106 full-text articles were assessed and 85 were excluded with reasons (see **Figure 1**). The included studies were conducted in a range of countries including United Kingdom (n=13), United States of America (n=3), Italy (n=1), Netherlands (n=2), Australia (n=1), and Sweden (n=1), see **Table 2** for an overview of the included studies. Across the included studies the sample sizes ranged from 13 to 815, with a total of 4,253 participants represented in this review. There were a range of study designs: quantitative (n=10)²⁶⁻³⁵, randomised controlled trials (n=6)³⁶⁻⁴¹, mixed methods studies (n=3)⁴²⁻⁴⁴, qualitative (n=1)⁴⁵ and a quantitative non-randomised controlled study (n=1)⁴⁶. The majority of the participants included had prostate^{27,30,33,34,39,40,44,45}, colorectal^{32,42}, lung^{41,43}, head and neck⁴⁶, renal³⁵, oesophageal³⁷ and mixed cancer.^{26,28,29,31,36,38} The methodological quality of the included studies was overall good, but some caution should be taken in the interpretation of the results due to non-response bias, and a lack of acknowledgement of the divergence between the qualitative and quantitative data in the mixed methods studies, see **Table 3**.

3.1 Types of Nurse-Led Interventions

The nurse-led interventions were cancer specific and included: prostate^{27,30,33,34,39,40,44,45}, colorectal^{32,42}, lung^{41,43}, head and neck⁴⁶, renal³⁵, radiotherapy³⁶, palliative care^{28,38}, virtual-telephone²⁶ and only one geriatric-oncology³¹ intervention, see **Table 4**.

Heterogeneity existed in the nurse-led interventions for patients ≥ 65 years in terms of the duration, composition, mode of delivery, and the outcomes measured to quantify their impact. Given that the types of nurse-led interventions were cancer specific the findings of this

review have been synthesised according to cancer type, with the underpinning clinical rationale that each type of cancer has its own unique implications and requirements for supportive care ²³.

Prostate Cancer Nurse-Led Interventions

It is unsurprising that most of the included nurse-led interventions were focussed on the clinical management of prostate cancer, given that this is largely a disease among older men. Nurse-led interventions were focussed on mixed treatment groups and stages ^{27,30,39,40,44}, active surveillance ³³, men on androgen deprivation therapy for metastatic disease ⁴⁵, and the stage and treatments were not reported in one study. ³⁴ For most of the studies it was unclear how the nursing process was implemented to assess, plan, implement interventions and evaluate patient care outcomes within the reported interventions. The inherent lack of information about the nursing process is an important omission in the included studies. Consequently, little is known about the cycle of decision-making that nurses used to capture (assess) patient information, how they created care plans and implemented interventions, and what strategies were used to evaluate whether care episodes were effective or not. Five studies used Patient Reported Outcome Measures (PROMs) to collect information to assess lower urinary track symptoms (International Prostate Symptom Score [IPSS]) ²⁷ or used a comprehensive holistic assessment tool. ^{30,39,40,45} Most of the prostate cancer nurse-led interventions did not report how they performed their clinical assessments to determine the patients' baselines or deterioration in symptoms, including timely identification of risk factors such as nutritional deficits, falls, cognitive impairment, frailty, multiple chronic conditions, and functional status. Two of the included studies ^{30,45} delivered an educational program which focussed on treatment, side-effects, and self-management strategies. One study delivered a nurse-led telephone service ²⁷ and the majority of the participants reported that they found the intervention to be convenient, informative, reassuring and beneficial in

terms of cost savings due to travel. Despite a clear lack of assessment of geriatric oncology risk factors among men who received nurse-led interventions for prostate cancer, men articulated value because they were provided with self-management support to reduce distress and recover from treatment side-effects including both physical and psychological difficulties.^{30,33,39} Commonly, patients reported that they were treated with dignity and respect, they were listened to and had time to ask questions^{34,45}, and that they were happy to see the specialist nurse for prostate cancer follow-up care compared to consultants.^{33,45} Studies demonstrated cost-effectiveness in nurse-led interventions of prostate cancer compared to consultant led care.^{36,38}

Nurse-Led Models of Colorectal Cancer Care

There were only two studies^{32,42} which explored nurse-led interventions of care among people affected by colorectal cancer. Both studies did not include nurse-led geriatric assessments, but one study used the Supportive Care Needs Survey (SCNS)⁴² to tailor consultations and improve supported self-management tailored to areas of patient distress or concern. All patients were satisfied with the nurse-led intervention, with reported reductions in physical and psychological symptoms, improved access to timely informational support, and patients valued using the SCNS to bring their needs and concerns to the forefront of the consultation.^{32,42} Cost saving reported from the UK-based studies were estimated to save £28,030 to the National Health Service (NHS).³²

Lung Cancer Nurse-Led Interventions

Two studies explored nurse-led interventions among people living with lung cancer and both did not include comprehensive geriatric assessments.^{41,43} One study explored the routine use of PROMS (Sheffield Profile for Assessment and Referral to Care [SPARC]) in identifying unmet supportive care needs to tailor individualised support.⁴³ Patients reported improved

symptom management and reduced distress caused by breathlessness, with a sense of improved control and empowerment as active partners in their own care.⁴³ Patients viewed that the time with the specialist nurse enabled them to build rapport and have the one-to-one time to explore sensitive issues such as death and dying, family, and sexuality related issues. Whereas the second study explored a palliative care nurse-led educational intervention to optimise symptom management in people diagnosed with advanced lung cancer compared to standard care (albeit standard care was not described in the context of this study).⁴¹ This study⁴¹ did not report any statistically significant differences in quality of life or satisfaction with care between both study arms. No cost effectiveness outcome data was included in either study.

Head and Neck Cancer Nurse-Led Interventions

This intervention was tested among 48 participants diagnosed with head and neck cancer and treated by radiotherapy.⁴⁶ The intervention includes a structured consultation with a specialist nurse and covered a variety of self-care topics related to psycho-social concerns, wound care, work, and financial support. There were no PROMS to structure the consultation or evidence of consideration to include comprehensive geriatric assessments in this older patient population.

Renal Cancer Nurse-led Interventions

A single study provided insights into a nurse-led intervention for renal cancer surveillance for patients treated by either partial or total nephrectomy.³⁵ There were no details on the nurse-led intervention in terms of the nursing process of care, no considerations of comprehensive geriatric assessments or documented survivorship care plans. The model of care was largely focus on cancer surveillance only and lacked information on the contribution of nursing support to address survivorship issues in this patient group.

Palliative Care Nurse-led Intervention

Two studies^{28,38} delivered nurse-led palliative interventions among patients with different cancers. Both nurse-led interventions involved educational and informational support to optimise symptom control and quality of life, nurse-patient-family engagement in advanced care planning, and a daily multidisciplinary staff briefing about holistic person-centred care with clear goals, and preferences of care documented. Of note, both studies did not include comprehensive geriatric assessments in this patient population of a mean age of 74 years²⁸, and 69 years.³⁸ Both studies identified improvement in physical and psychological wellbeing in favour of the nurse-led interventions.

Nurse-led Interventions Among Mixed Cancer Groups

Only two studies^{26,36} explored nurse-led interventions care among patients diagnosed with different cancers, as opposed to single tumour specialised nursing interventions. One study²⁶ explored a virtual intervention during the COVID-19 pandemic. Patients were diagnosed with various types of cancer, the most common were breast, gastrointestinal and haematological. The nurse-led consultation was conducted via telephone or video calls to provide timely support and assess practical daily living needs. Patients reported that they received emotional support, COVID-19 education and nutritional advice, medication support and signposting for financial assistance.²⁶ The second study³⁶ provided a nurse-led radiotherapy intervention for mixed cancer groups. The intervention provided education and informational support and practical advice to optimise self-management throughout radiotherapy treatment.³⁶ Physical and psychological wellbeing among the participants was in favour of the intervention arm compared to standard treatment. Participants articulated that they felt that their needs and concerns were taken seriously and that they valued the

experience of continuity of care. No comprehensive geriatric assessments were conducted in either of these studies.

Geriatric Oncology Nurse-led Intervention

Only one study³¹ explored nurse-led geriatric oncology intervention in 197 participants with a mean age of 78 years. This study represented mixed cancer patient groups and the geriatric assessment involved somatic, social, psychological, and functional assessments.

Polypharmacy considerations were considered for participants taking five or more medication prescription drugs. Frailty was assessed using both the Groningen Frailty Indicator and The Letter Fluency Test for a measure cognition. Other assessments included mobility, activities of daily living, and co-morbidity. The central premise of this nurse-led intervention was to embed comprehensive geriatric assessments and the patients' preferences within an MDT oncogeriatric approach to tailor cancer treatment recommendations. Over half (52.3%) of the patients were frail, at risk of polypharmacy (52.7%) and experiencing cognitive decline. Of note, 27% of the cancer treatment recommendations for the patients within the nurse-led oncogeriatrics MDT, differed from the cancer tumour board MDT treatment recommendations, and the modifications were largely related to less intensive or invasive treatments.

4. Discussion

This integrative systematic review set out to identify existing nurse-led interventions among older adults across all different cancer types. Of the 21 included studies within this review, only one study³¹ incorporated comprehensive geriatric nurse-led assessments underscoring fundamental shortcomings within existing nurse-led interventions for the older person with cancer.⁴⁷ This review has identified that nurse-led interventions are highly specialised by

cancer type only and lacked the integration of the inclusion of geriatric assessments within the multidisciplinary team.

Several possible reasons exist for a lack of nurse-led geriatric assessment among older people with cancer which may include: a lack of funding and resources; reduced capacity to meet clinical demand; time; and poor communication processes for referrals.⁴⁸ Historically, these barriers have led to variation and poor documentation around the development and implementation of nurse-led interventions in cancer care.⁴⁹ The findings of this review reflect similar shortfalls in practice around the integration of geriatric assessment with the exception of Festen et.al³¹ who did include a geriatric oncology intervention. However, Festen et al.³⁰ reported that the reason for including geriatric assessment in their model was driven by the ‘accumulating evidence’ on the value of their predictive validity in determine clinical outcomes. While this may be true, four newer studies included in the review did not acknowledge this accumulating evidence and suggests an evidence-practice gap in the process of assessing and implementing appropriate care interventions and treatment for older people with cancer.

Addressing this evidence-practice gap is a priority for improving the assessment and treatment of older people living with cancer as clinical assessments can influence different cancer treatment regimens for older patients diagnosed with cancer. Consideration of age and tumour characteristics alone is insufficient to determine fitness for treatment for people >65 years. Comprehensive geriatric assessments may trigger the use of less aggressive or more aggressive treatments which can affect quality of life, implications for supportive care, and overall survival.^{31,50} Validated and practical geriatric assessment tools to assess function, falls, comorbidity, cognition, depression and nutrition can be used to predict adverse outcomes in patients aged 65 years and older receiving treatment for cancer.⁵¹ The use of validated geriatric assessment tools can assist with developing goal-directed interventions and

guide management in older people with cancer.⁵² The transfer of information between health professionals using standardised instruments enhances timely communication exchange and teamwork to accurately document decision-making and goals of care to optimise quality and quantity of life over time.

One approach in optimising physical and psychological health of older people with cancer is through the implementation of prehabilitation models of care for the older person with cancer. Prehabilitation is a model of cancer care implemented between the time of a cancer diagnosis and the beginning of acute treatment. A recent systematic review⁵³ in older men affected by prostate cancer identified that multi-component nurse-led prehabilitation interventions of supportive care may provide benefit in optimising physical and psychological reserve before treatments and reduce treatment-related deconditioning. Prehabilitation includes physical and psychological assessments⁵⁴, including comprehensive geriatric assessments, that establish baseline functioning and identifies impairments that can impact cancer treatment-related morbidity, as well as provide targeted interventions to optimize overall well-being prior to treatment.

This systematic review highlighted the need for further research to explore the impact of nurse-led geriatric oncology interventions which are safely embedded in the MDT. Despite the growing need for appropriate models of cancer care for older people⁵¹ this review has identified that there is a dearth of inclusion of geriatric assessment to identify age-related vulnerabilities such as functional, medical, cognitive, psychosocial, and nutritional needs in existing nurse-led interventions. Quantity of life versus quality of life in older people with cancer must be carefully considered in cancer MDTs, to ensure informed consent and shared treatment decision-making processes, however, these important considerations were seldom discussed in any of the studies included. A recent systematic review⁵⁵ identified that only a small percentage of patients diagnosed with cancer will ever be discussed in a cancer MDT

meeting which therefore, further compounds the challenge of integrating timely and effective MDT geriatric oncology considerations in patient care. Importantly, this review has underscored the need for further research to test nurse-led interventions in geriatric oncology and future research directions in prehabilitation for the older person with cancer.

4.1 Limitations

Despite this review following a clear, rigorous, and transparent review process there are some important limitations to point out. This review included studies which were published in the English language only, and as such may have excluded publications in other languages which might have omitted important information. However, the review did represent evidence from a range of international countries. One of the major challenges of this review was synthesising evidence from heterogeneous study designs and methodologies, and our findings are constrained due to the methodological limitations of the primary studies included. This review has enabled a broad summary of the evidence in relation to nurse-led interventions for older person with cancer which has provided some clinical practice insights gaps and facilitated refinement of future research directions. We included studies where the mean age of the patient sample was ≥ 65 years. As a result, we identified studies with nurse-led interventions among people with a mean age of 65 years, not studies specifically designed for people ≥ 65 years. Therefore, the included studies underscore that existing nurse-led interventions included some older people which are largely not addressing or including comprehensive geriatric assessments and remains a significant gap in practice and research.

5. Conclusion

The findings of this review highlight a concerning lack of evidence on integrating geriatric assessments into nurse-led interventions. With most developed countries reporting an aging population and a general trend of cancer survivors living longer following diagnosis, the need

to incorporate geriatric assessments into routine care are a priority for optimising the health of older people living with cancer. Since few approaches are reported in the literature, adopting innovative strategies such as the use of prehabilitation and involving cancer MDTs to facilitate geriatric assessment should be explored further.

Journal Pre-proof

References

1. United Nations Department of Economic Social Affairs Population Division. *World population ageing: 2017 highlights*. UN; 2017.
2. World Health Organization. *World report on ageing and health*. World Health Organization; 2015.
3. Morris L, Turner S, Thiruthaneeswaran N, Agar M. Improving the Education of Radiation Oncology Professionals in Geriatric Oncology: Where Are We and Where Should We Be? *Seminars*

- in Radiation Oncology*. 2022/04/01/ 2022;32(2):109-114.
doi:<https://doi.org/10.1016/j.semradonc.2021.11.008>
4. Pilleron S, Soto-Perez-de-Celis E, Vignat J, et al. Estimated global cancer incidence in the oldest adults in 2018 and projections to 2050. *International journal of cancer*. 2021;148(3):601-608.
 5. Extermann M, Brain E, Canin B, et al. Priorities for the global advancement of care for older adults with cancer: an update of the International Society of Geriatric Oncology Priorities Initiative. *The Lancet Oncology*. 2021/01/01/ 2021;22(1):e29-e36. doi:[https://doi.org/10.1016/S1470-2045\(20\)30473-3](https://doi.org/10.1016/S1470-2045(20)30473-3)
 6. O'Dea A, Gedye C, Jago B, Paterson C. Identifying the unmet supportive care needs of people affected by kidney cancer: a systematic review. *J Cancer Surviv*. Oct 1 2021;doi:10.1007/s11764-021-01113-8
 7. Paterson C, Primeau C, Howard N, Xiberras P, Pillay B, Crowe H. Experiences of Unmet Need and Access to Supportive Care for Men Receiving Androgen Deprivation Therapy for Prostate Cancer: A Bi-national Study. *Semin Oncol Nurs*. Aug 2020;36(4):151049. doi:10.1016/j.soncn.2020.151049
 8. Gray-Miceli D, Wilson LD, Stanley J, et al. Improving the Quality of Geriatric Nursing Care: Enduring Outcomes From the Geriatric Nursing Education Consortium. *Journal of Professional Nursing*. 2014/11/01/ 2014;30(6):447-455. doi:<https://doi.org/10.1016/j.profnurs.2014.05.001>
 9. Leff B, Kao H, Ritchie C. How the principles of geriatric care can be used to improve care for Medicare patients. *Generations*. 2015;39(2):99-105.
 10. Presley CJ, Krok-Schoen JL, Wall SA, et al. Implementing a multidisciplinary approach for older adults with Cancer: geriatric oncology in practice. *BMC Geriatr*. Jul 6 2020;20(1):231. doi:10.1186/s12877-020-01625-5
 11. Paterson C, Jensen BT, Jensen JB, Nabi G. Unmet informational and supportive care needs of patients with muscle invasive bladder cancer: A systematic review of the evidence. *Eur J Oncol Nurs*. Aug 2018;35:92-101. doi:10.1016/j.ejon.2018.05.006
 12. Paterson C, Alashkham A, Lang S, Nabi G. Early oncological and functional outcomes following radical treatment of high-risk prostate cancer in men older than 70 years: A prospective longitudinal study. *Urol Oncol*. Aug 2016;34(8):335.e1-7. doi:10.1016/j.urolonc.2016.03.002
 13. Paterson C, Alashkham A, Windsor P, Nabi G. Management and treatment of men affected by metastatic prostate cancer: evidence-based recommendations for practice. <https://doi.org/10.1111/ijun.12093>. *International Journal of Urological Nursing*. 2016/03/01 2016;10(1):44-55. doi:<https://doi.org/10.1111/ijun.12093>
 14. Puts MT, Tapscott B, Fitch M, et al. A systematic review of factors influencing older adults' decision to accept or decline cancer treatment. *Cancer treatment reviews*. 2015;41(2):197-215.
 15. Seghers PA, Wiersma A, Festen S, et al. Patient Preferences for Treatment Outcomes in Oncology with a Focus on the Older Patient—A Systematic Review. *Cancers*. 2022;14(5):1147.
 16. Corbett T, Bridges J. Multimorbidity in older adults living with and beyond cancer. *Curr Opin Support Palliat Care*. Sep 2019;13(3):220-224. doi:10.1097/spc.0000000000000439
 17. Pergolotti M, Deal AM, Williams GR, et al. Activities, function, and health-related quality of life (HRQOL) of older adults with cancer. *J Geriatr Oncol*. Jul 2017;8(4):249-254. doi:10.1016/j.jgo.2017.02.009
 18. Ramjaun A, Nassif MO, Krotneva S, Huang AR, Meguerditchian AN. Improved targeting of cancer care for older patients: a systematic review of the utility of comprehensive geriatric assessment. *J Geriatr Oncol*. Jul 2013;4(3):271-81. doi:10.1016/j.jgo.2013.04.002
 19. Morgan B, Tarbi E. The Role of the Advanced Practice Nurse in Geriatric Oncology Care. *Seminars in Oncology Nursing*. 2016/02/01/ 2016;32(1):33-43. doi:<https://doi.org/10.1016/j.soncn.2015.11.005>
 20. Burhenn PS, Perrin S, McCarthy AL. Models of care in geriatric oncology nursing. Elsevier; 2016:24-32.
 21. Van Cleave JH, Kenis C, Sattar S, Jabloo VG, Ayala AP, Puts M. A research agenda for geriatric oncology nursing. Elsevier; 2016:55-64.
 22. Page MJ, McKenzie JE, Bossuyt PM, et al. Updating guidance for reporting systematic reviews: development of the PRISMA 2020 statement. *Journal of clinical epidemiology*. 2021;134:103-112.

23. Paterson C, Toohey K, Bacon R, Kavanagh PS, Roberts C. What are the unmet supportive care needs of people affected by cancer: an umbrella systematic review. Elsevier; 2022:151353.
24. Pace R, Pluye P, Bartlett G, et al. Testing the reliability and efficiency of the pilot Mixed Methods Appraisal Tool (MMAT) for systematic mixed studies review. *International journal of nursing studies*. 2012;49(1):47-53.
25. Lisy K, Porritt K. Narrative synthesis: considerations and challenges. *JBI Evidence Implementation*. 2016;14(4):201.
26. Bigelow SM, Hart E, Shaban T, et al. A new proactive virtual resource center navigation model identifies patient risk factors to reduce barriers to cancer care during the COVID-19 pandemic. *Support Care Cancer*. Oct 2021;29(10):6069-6077. doi:10.1007/s00520-021-06147-3
27. Casey RG, Powell L, Braithwaite M, Booth CM, Sizer B, Corr JG. Nurse-Led Phone Call Follow-Up Clinics Are Effective for Patients With Prostate Cancer. *Journal of Patient Experience*. 2017/09/01 2017;4(3):114-120. doi:10.1177/2374373517706613
28. Catania G, Zanini M, Signori A, et al. Providing a nurse-led complex nursing INtervention FOCused on quality of life assessment on advanced cancer patients: The INFO-QoL pilot trial. *Eur J Oncol Nurs*. Jun 2021;52:101961. doi:10.1016/j.ejon.2021.101961
29. Craven O, Hughes CA, Burton A, Saunders MP, Molassiotis A. Is a nurse-led telephone intervention a viable alternative to nurse-led home care and standard care for patients receiving oral capecitabine? Results from a large prospective audit in patients with colorectal cancer. *Eur J Cancer Care (Engl)*. May 2013;22(3):413-9. doi:10.1111/ecc.12047
30. Ferguson J, Aning J. Prostate cancer survivorship: a nurse-led service model. *Br J Nurs*. Oct 8-21 2015;24(18):S14-21. doi:10.12968/bjon.2015.24.Sup18.S14
31. Festen S, Kok M, Hopstaken JS, et al. How to incorporate geriatric assessment in clinical decision-making for older patients with cancer. An implementation study. *Journal of Geriatric Oncology*. 2019/11/01/ 2019;10(6):951-959. doi:https://doi.org/10.1016/j.jgo.2019.04.006
32. Knowles G, Sherwood L, Dunlop MG, et al. Developing and piloting a nurse-led model of follow-up in the multidisciplinary management of colorectal cancer. *Eur J Oncol Nurs*. Jul 2007;11(3):212-23; discussion 224-7. doi:10.1016/j.ejon.2006.10.007
33. Martin E, Persaud S, Corr J, Casey R, Pillai R. Nurse-led active surveillance for prostate cancer is safe, effective and associated with high rates of patient satisfaction-results of an audit in the East of England. *Ecancermedicalscience*. 2018;12:854. doi:10.3332/ecancer.2018.854
34. McGlynn B, White L, Smith K, et al. A service evaluation describing a nurse-led prostate cancer service in NHS, Ayrshire and Arran. *International Journal of Urological Nursing*. 2014;8(3):166-180. doi:https://doi.org/10.1111/ijun.12049
35. Sibbons A, Pillai R, Corr J, Persaud S. Nurse-led renal cancer follow-up is safe and associated with high patient satisfaction-an audit from the East of England. *Ecancermedicalscience*. 2019;13:955. doi:10.3332/ecancer.2019.955
36. Faithfull S, Corner J, Meyer L, Huddart R, Dearnaley D. Evaluation of nurse-led follow up for patients undergoing pelvic radiotherapy. *Br J Cancer*. Dec 14 2001;85(12):1853-64. doi:10.1054/bjoc.2001.2173
37. Malmström M, Ivarsson B, Klefsgård R, Persson K, Jakobsson U, Johansson J. The effect of a nurse led telephone supportive care programme on patients' quality of life, received information and health care contacts after oesophageal cancer surgery-A six month RCT-follow-up study. *Int J Nurs Stud*. Dec 2016;64:86-95. doi:10.1016/j.ijnurstu.2016.09.009
38. Schenker Y, Althouse AD, Rosenzweig M, et al. Effect of an Oncology Nurse-Led Primary Palliative Care Intervention on Patients With Advanced Cancer: The CONNECT Cluster Randomized Clinical Trial. *JAMA Intern Med*. Nov 1 2021;181(11):1451-1460. doi:10.1001/jamainternmed.2021.5185
39. Schofield P, Gough K, Lotfi-Jam K, et al. Nurse-led group consultation intervention reduces depressive symptoms in men with localised prostate cancer: a cluster randomised controlled trial. *BMC Cancer*. Aug 16 2016;16:637. doi:10.1186/s12885-016-2687-1
40. Stanciu MA, Morris C, Makin M, et al. Trial of personalised care after treatment-Prostate cancer: A randomised feasibility trial of a nurse-led psycho-educational intervention. *Eur J Cancer Care (Engl)*. Mar 2019;28(2):e12966. doi:10.1111/ecc.12966

41. Reinke LF, Sullivan DR, Slatore C, et al. A Randomized Trial of a Nurse-Led Palliative Care Intervention for Patients with Newly Diagnosed Lung Cancer. *J Palliat Med.* Jun 1 2022;doi:10.1089/jpm.2022.0008
42. Kotronoulas G, Papadopoulou C, MacNicol L, Simpson M, Maguire R. Feasibility and acceptability of the use of patient-reported outcome measures (PROMs) in the delivery of nurse-led supportive care to people with colorectal cancer. *European Journal of Oncology Nursing.* 2017/08/01/ 2017;29:115-124. doi:https://doi.org/10.1016/j.ejon.2017.06.002
43. Kotronoulas G, Papadopoulou C, Simpson MF, McPhelim J, Mack L, Maguire R. Using patient-reported outcome measures to deliver enhanced supportive care to people with lung cancer: feasibility and acceptability of a nurse-led consultation model. *Supportive Care in Cancer.* 2018/11/01 2018;26(11):3729-3737. doi:10.1007/s00520-018-4234-x
44. Ream E, Wilson-Barnett J, Faithfull S, Fincham L, Khoo V, Richardson A. Working patterns and perceived contribution of prostate cancer clinical nurse specialists: a mixed method investigation. *Int J Nurs Stud.* Oct 2009;46(10):1345-54. doi:10.1016/j.ijnurstu.2009.03.006
45. Primeau C, Paterson C, Nabi G. A Qualitative Study Exploring Models of Supportive Care in Men and Their Partners/Caregivers Affected by Metastatic Prostate Cancer. *Oncol Nurs Forum.* Nov 1 2017;44(6):E241-e249. doi:10.1188/17.Onf.E241-e249
46. van der Meulen IC, de Leeuw JR, Gamel CJ, Hafsteinsdóttir TB. Educational intervention for patients with head and neck cancer in the discharge phase. *Eur J Oncol Nurs.* Apr 2013;17(2):220-7. doi:10.1016/j.ejon.2012.05.007
47. Dufton P, Tarasenko E, Mellerick A, Yates P, Lee K, Parakh S. Implementation of a Nurse-led Geriatric Oncology Assessment Model in the Lung Cancer Care Pathway. *Journal of Thoracic Oncology.* 2022;17(9):S24-S25.
48. Douglas C, Schmalkuche D, Nizette D, Yates P, Bonner A. Nurse-led services in Queensland: A scoping study. *Collegian.* 2018;25(4):363-370.
49. Monterosso L, Platt V, Bulsara M, Berg M. Systematic review and meta-analysis of patient reported outcomes for nurse-led models of survivorship care for adult cancer patients. *Cancer treatment reviews.* 2019;73:62-72.
50. Decoster L, Kenis C, Van Puyvelde K, et al. The influence of clinical assessment (including age) and geriatric assessment on treatment decisions in older patients with cancer. *Journal of geriatric oncology.* 2013;4(3):235-241.
51. Mohile SG, Dale W, Somerfield MR, et al. Practical assessment and management of vulnerabilities in older patients receiving chemotherapy: ASCO guideline for geriatric oncology. *Journal of Clinical Oncology.* 2018;36(22):2326.
52. Magnuson A, Allore H, Cohen HJ, et al. Geriatric assessment with management in cancer care: Current evidence and potential mechanisms for future research. *Journal of geriatric oncology.* 2016;7(4):242-248.
53. Paterson C, Roberts C, Kozlovskaja M, et al. The Effects of Multimodal Prehabilitation Interventions in Men Affected by Prostate Cancer on Physical, Clinical and Patient Reported Outcome Measures: A Systematic Review. *Seminars in Oncology Nursing.* 2022/08/20/ 2022:151333. doi:https://doi.org/10.1016/j.soncn.2022.151333
54. Paterson C, Roberts C, Toohey K, McKie A. Prostate Cancer Prehabilitation and the Importance of Multimodal Interventions for Person-centred Care and Recovery. *Semin Oncol Nurs.* Aug 2020;36(4):151048. doi:10.1016/j.soncn.2020.151048
55. Askelin B, Hind A, Paterson C. Exploring the impact of uro-oncology multidisciplinary team meetings on patient outcomes: A systematic review. *European Journal of Oncology Nursing.* 2021;54:102032.

Table 1. Classification of Supportive Care Needs












Domain of need	Description
 Physical needs	Experience of symptoms such as fatigue, pain, etc. co-morbidities, nutritional deficits, frailty, functional status
 Psychological/emotional needs	Experience of depression, anxiety, sadness, fear, distress, etc.
 Cognitive needs	Individual experience of cognitive impairment or decline, memory problems, sensory loss, etc.
 Patient-clinician communication needs	Quality of communication and co-ordination between patients and health care professionals, shared decision making, etc.
 Health system/information needs	Information needs, uncertainty of follow-up, lack of information about diagnosis and treatment, etc.
 Spiritual needs	Fear of death and dying, fears regarding the afterlife, etc.
 Daily living needs	Experience of restrictions to daily living, exercise, housekeeping, etc
 Interpersonal/intimacy needs	Experience of difficulties with body image, masculinity, sexual dysfunction, compromised intimacy with partner, etc.
 Practical needs	Related to daily task restrictions, employment, accessing benefits, life insurance, etc.
 Family related needs	Experience of fears/concerns of the family, dysfunctional relationships, etc
 Social needs	Experience of reduced social support, social isolation, loneliness, lack of peer support, etc

Table 2. Overview of the Included Studies.

Author and Year	Purpose	Sample and age (mean)	Participants (cancer, stage, treatment)	Design	Data Collection tools Comprehensive geriatric assessment (CGA)
Bigelow et al. 2021 ²⁵ USA	To describe the implementation, associated interventions, and outcomes of the PFRC's proactive virtual resource centre navigation model.	Sample size: 586 65 years	Cancer: Most common were haematological, breast and gastrointestinal Cancer stage: Not reported Treatment: Surgery, radiation, chemotherapy, or other systemic treatments	Quality Improvement Study	Referral information, patient demographics, risk characteristics, visit data, interventions, and outcomes CGA: not included.
Casey et al. 2017 ²⁶ UK	To evaluate patient satisfaction with a nurse-led phone call follow up clinic for patients with prostate cancer.	Sample size: 815 75 years	Cancer: Prostate Cancer Cancer stage: Mixed Treatment: Surgery, radiotherapy, combined surgery and radiotherapy, brachytherapy, hormone manipulation	Satisfaction survey	Satisfaction survey CGA: not included.
Catania et al. 2021 ²⁷ Italy	To pilot a nurse-led complex intervention focused on QoL assessment in advanced- disease cancer patients.	Sample size: 187 74 years	Cancer: Mixed Cancer stage: Advanced disease/last phase of life Treatment: Not reported	Quasi-Experimental design	Integrated Palliative Care Outcome Scale (I-POS) CGA: not included. Cognitively impaired patients were excluded
Craven et al. 2012 ²⁸ UK	A prospective audit exploring the usefulness of a nurse-led telephone intervention for supporting cancer patients treated with Capecitabine.	Sample size: 462 65 years	Cancer: Colorectal and breast Cancer stage: Not reported. Treatment: Capecitabine	Longitudinal prospective evaluation	National Cancer Institute Common Toxicity Criteria (NCI-CTC), patient satisfaction questionnaire was completed. CGA: not included.
Faithfull et al. 2001 ³⁵ UK	To compare outcomes in terms of toxicity, symptoms experienced, quality of life, satisfaction with care and health care costs between those receiving nurse-led care vs. group receiving standard care.	Sample size: 115 70 years	Cancer: Prostate and bladder Cancer stage: Mixed Treatment: Radiotherapy	RCT	Observer- rated RTOG Toxicity scores, EORTC QLQ C30, Satisfaction Questionnaire, Economic Appraisal Information CGA: not included.
Ferguson and Aning 2015 ²⁹ UK	To describe the implementation of a nurse-led survivorship programme for men with Prostate Cancer.	Sample size: 76 65 years	Cancer: Prostate Cancer stage: Not reported Treatment: Robotic Radical Prostatectomy, Laparoscopic Radical Prostatectomy, Radiotherapy, ADT, Active surveillance, chemotherapy	Service Evaluation	Clinical and demographic data CGA: not included.
Festen et al. 2019 ³⁰ Netherlands	To evaluate nurse-led geriatric assessment and assessment of patient preferences for oncological treatment decisions for older patients with solid malignancy.	Sample size: 197 78 years	Cancer: Mixed Cancer stage: Mixed Treatment: Curative Intent: 159 (80.7%), Palliative Intent: 38 (19.3%)	Prospective Cohort Study	CGA: Yes. GA involved an evaluation on four domains: somatic, social, psychological and functional. Polypharmacy was defined as taking ≥5 prescription drugs. Groningen Frailty Indicator (GFI), The Letter Fluency Test (LFT) was used as a measure of cognition, The Timed Up and Go (TUG) test was

					used as a measure of mobility, The Groningen Activity Restriction Scale (GARS) is a combined Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (iADL), Outcome Prioritization Tool (OPT), Charlson Comorbidity Index (CCI)
Knowles et al. 2007 ³¹ UK	To assess the feasibility of a follow up programme led by nurse specialist for patients with colorectal cancer.	Sample size: 60 67 years	Cancer: Colorectal Cancer Stage: Mixed Treatment: Short course Radiotherapy, Colectomy, Resection, Chemotherapy and Radiation	Pilot Study	QLQ - C30, QLQ- CR38, Satisfaction Questionnaire CGA: not included.
Kotronoulas et al. 2017 ⁴¹ UK	To explore the feasibility and acceptability of PROMs-driven, CNS-led consultations to enhance delivery of supportive care to people with CRC completing adjuvant chemotherapy.	Sample size: 13 65 years	Cancer: Colorectal Cancer stage: Mixed Treatment: Surgery, Chemotherapy, radiotherapy	Systematic literature review, focus groups and repeated measure exploratory study	Supportive Care Needs Survey e Short Form 34 (SCNS-SF34), Twelve patients initially consented to end-of-study interviews CGA: not included.
Kotronoulas et al. 2018 ⁴² UK	To examine whether a nurse-led PRO measure driven is feasible and acceptable in identifying unmet needs in patients with lung cancer.	Sample size: 20 67 years	Cancer: Lung Cancer stage: Mixed Treatment: Not reported	Mixed Methods	Nine patients with lung cancer (6 men, 3 women) took part in interviews. Sheffield Profile for Assessment and Referral to Care (SPARC) CGA: not included.
Malmstrom et al. 2016 ⁵⁵ Sweden	To evaluate the effect of a nurse-led telephone supportive care program on QoL compared to conventional care on patients following oesophageal resection for cancer.	Sample size: 82 66.4 years	Cancer: Oesophageal cancer Cancer stage: Not reported Treatment: Oesophagectomy or oesophago-gastrectomy	RCT	QLQ-C30, QLQ-0625, QLQ-INFO25. CGA: not included.
Martin et al 2018 ³² UK	To evaluate a nurse-led service for men affected by PC on AS.	Sample size: 104 patients 66 years	Cancer: Prostate Cancer stage: Localised prostate cancer Treatment: Active surveillance	Retrospective audit, patient satisfaction survey and staff satisfaction survey	National Cancer Patient Experience Survey. CGA: not included.
McGlynn et al. 2014 ³³ UK	A local evaluation of the nurse-led collaborative care model for the management of patients with prostate cancer.	Sample size: 71 No mean age provided. participants aged approximately between 63-82 years	Cancer: Prostate Cancer stage: Not reported. Treatment: Not reported	Retrospective audit, patient satisfaction survey and staff satisfaction survey	Patient satisfaction questionnaire. CGA: not included.
Primeau et al. 2017 ⁴⁴	To explore the experience of patients and their partner/caregiver as well as MDT members of a nurse-led	Sample size: 19 patients, 7 partners/caregivers, 7 MDT members Range 67-84 years	Cancer: Prostate Cancer stage: Metastatic Treatment: Androgen deprivation therapy	Qualitative Study	Semi-structured interviews. CGA: not included.

UK	multimodality supportive care intervention in men with metastatic prostate cancer as well as standard care.				
Ream et al. 2009 ⁴³ UK	To investigate the role of Prostate Cancer Clinical Nurse Specialists and determine their targeted services, work practises and perceived contribution.	Sample size: 4 PCNS, 19 clinical colleagues, 40 patients 67 years	Cancer: Prostate Cancer stage: Not reported Treatment: prostatectomy, radiotherapy, hormone therapy, active monitoring, brachytherapy, cryotherapy, chemotherapy	Mixed method design	Nurse diary based on prototype developed by Macmillan's and interviews. CGA: not included.
Reinke et al. 2022 ⁴⁰ USA	To assess the effect of a nurse-led telephone-based primary palliative care intervention for patient with lung cancer.	Sample size: 151 70 years	Cancer: Lung Cancer stage: 64% III-IV Treatment: Mixed	RCT	FACT-L, Satisfaction of care was measured using the FAMCARE-P13 Patient Scale. CGA: not included.
Schenker et al. 2021 ³⁷ USA	To assess the effect of CONNECT (Care Management by Oncology Nurses to Address Supportive Care Needs).	Sample size: 672 69 years	Cancer: The most common cancers lung and gastrointestinal Cancer stage: Not reported Treatment: Chemotherapy, radiotherapy, hormonal therapy, immunotherapy	RCT	Functional Assessment of Chronic Illness Therapy-Palliative care, Edmonton Symptom Assessment Scale, Hospital Anxiety and Depression Scale, ECOG Performance Status score CGA: not included.
Schofield et al. 2016 ³⁸ Australia	To investigate the benefits of a group nurse-led intervention in men receiving radiotherapy for Prostate Cancer.	Sample size: 331 67.2 years	Cancer: Prostate Cancer stage: Not reported Treatment: Surgery, radiotherapy, hormonal therapy	RCT	HADS, CaTS, SCNS-SF 34-F, EPIC-26, DT CGA: not included.
Sibbons et al. 2019 ³⁴ UK	To evaluate a nurse-led service for patients affected by renal cancer	Sample size: 89 67 years	Cancer: Renal Cancer stage: Not reported Treatment: Partial nephrectomy and radical nephrectomies	Clinical audit and service evaluation	Retrospective clinical audit of medical records and patient satisfaction survey CGA: not included.
Stanciu et al. 2018 ³⁹ UK	To evaluate a nurse-led model of personalised care after prostate cancer treatment	Sample size: 45 (intervention), 47 (control group) range 66-94 years old	Cancer: Prostate Cancer stage: Mixed Treatment: surgery, radiotherapy, hormone therapy, or deemed unlikely to receive further treatment (watchful waiting)	RCT	Clinical and demographic data, EPIC-26, HADS, SCNS-SF34, EQ-5D-5L, CSRI, Confidence Managing Own Health, Satisfaction with Healthcare Survey CGA: not included.
van der Meulen et al. 2013 ⁴⁵ Netherlands	To test a nurse-led educational intervention for patients with head and neck cancer	Sample size: 48 65 years	Cancer: Head and neck Cancer stage: Not reported Treatment: Radiotherapy	Quasi-Experimental study	PINQ, SCIP CGA: not included.

Urinary Tract Symptoms; Nodal (N), extent of the tumour; PCNS, Prostate Cancer Nurse Specialist; PFRC NN, Patient and Family Resource Centre Nurse Navigator; PINQ, Patient Information Need Questionnaire; PRO, Patient Reported Outcome; PSA, Prostate Specific Antigen; QLQ-INFO25, Perceived level of information; QLQ-0625, European Quality of Life in Cancer of Oesophagus, Oesophago-gastric junction or Stomach; QLQ-C30, European Organisation for Research and Treatment of Cancer Quality of Life with Cancer; QoL, Quality of Life; RCT, Randomised Controlled Trial; RTOG, Radiation Therapy Oncology Group; SCIP, Satisfaction with Cancer Information Profile; SCNS-SF34-R, Supportive Care Needs Short Form Revised; SCNS-SF36, Supportive Care Needs Survey Short-Form; SPARC, Sheffield Profile For Assessment and Referral to Care; SRM, Standardised Response Mean; TURP, Transurethral Resection of the Prostate

Journal Pre-proof

Table 3. Results of Quality Assessment.

Qualitative Study	Item number of checklist						
	S1	S2	1.1	1.2	1.3	1.4	1.5
Primeau et al. 2017 ⁴⁴	Y	Y	Y	Y	Y	Y	Y
Item number checklist key: S1. Are there clear research questions, S2. Do the collected data allow to address the research questions, 1.1. Is the qualitative approach appropriate to answer the research question, 1.2. Are the qualitative data collection methods adequate to address the research question, 1.3. Are the findings adequately derived from the data, 1.4. Is the interpretation of results sufficiently substantiated by data, 1.5. Is there coherence between qualitative data sources, collection, analysis and interpretation.							
Quantitative Randomised Controlled Trials	Item number of checklist						
	S1	S2	2.1	2.2	2.3	2.4	2.5
Faithfull et al. 2001 ³⁵	Y	Y	Y	Y	Y	U	Y
Malmstrom et al. 2016 ⁵⁵	Y	Y	Y	Y	Y	Y	Y
Reinke et al. 2022 ⁴⁰	Y	Y	U	Y	Y	U	U
Schofield et al. 2016 ³⁸	Y	Y	Y	Y	Y	Y	U
Schenker et al. 2021 ³⁷	Y	Y	Y	Y	Y	Y	U
Stanciu et al. 2018 ³⁹	Y	Y	Y	Y	Y	Y	Y
Item number check list key: S1. Are there clear research questions, S2. Do the collected data allow to address the research questions, 2.1. Is randomisation appropriately performed, 2.2. Are the groups comparable at baseline, 2.3. Are there complete outcome data, 2.4. Are outcome assessors blinded to the intervention provided, 2.5. Did the participants adhere to the assigned intervention							
Quantitative Non- Randomised Controlled Trials	Item number of checklist						
	S1	S2	3.1	3.2	3.3	3.4	3.5
Van der Meulen et al. 2013 ⁴⁵	Y	Y	Y	Y	Y	U	Y

S1. Are there clear research questions, S2. Do the collected data allow to address the research questions, 3.1. Are the participants representative of the target population, 3.2. Are measurements appropriate regarding both the outcome and intervention (or exposure), 3.3. Are there complete outcome data, 3.4. Are the confounders accounted for in the design and analysis, 3.5. During the study period, is the intervention administered (or exposure occurred) as intended							
Quantitative Descriptive Studies	Item number of checklist						
	S1	S2	4.1	4.2	4.3	4.4	4.5
Bigelow et al. 2021 ²⁵	Y	Y	Y	Y	Y	Y	Y
Casey et al. 2017 ²⁶	Y	Y	Y	U	Y	U	Y
Catania et al. 2021 ²⁷	Y	Y	Y	U	Y	U	Y
Craven et al. 2012 ²⁸	Y	Y	Y	U	Y	U	Y
Festen et al. 2019 ³⁰	Y	Y	Y	Y	U	U	Y
Ferguson and Aning 2015 ²⁹	Y	Y	Y	Y	U	U	Y
Knowles et al. 2007 ³¹	Y	Y	U	Y	Y	Y	Y
Martin et al. 2018 ³²	Y	Y	Y	Y	U	U	Y
McGlynn et al. 2014 ³³	Y	Y	Y	U	U	U	U
Sibbons et al. 2019 ³⁴	Y	Y	Y	U	U	U	Y
Item number check list key: S1. Are there clear research questions, S2. Do the collected data allow to address the research questions, 4.1. Is the sampling strategy relevant to address the research question, 4.2. Is the sample representative of the target population, 4.3. Are the measurements appropriate, 4.4. Is the risk of non-response bias low, 4.5. Is the statistical analysis appropriate to answer the research question							
Mixed Methods	Item number of checklist						
	S1	S2	5.1	5.2	5.3	5.4	5.5
Kotronoulas et al. 2017 ⁴¹	Y	Y	Y	Y	Y	U	Y
Kotronoulas et al. 2018 ⁴²	Y	Y	Y	Y	Y	U	Y

Ream et al. 2009 ⁴³	Y	Y	Y	Y	Y	U	Y
Item number check list key: S1. Are there clear research questions, S2. Do the collected data allow to address the research questions, 5.1. Is there an adequate rationale for using a mixed methods design to address the research question, 5.2. Are the different components of the study effectively integrated to answer the research question, 5.3. Are the outputs of the integration of qualitative and quantitative components adequately interpreted, 5.4. Are divergences and inconsistencies between quantitative and qualitative results adequately addressed, 5.5. Do the different components of the study adhere to the quality criteria of each tradition of the methods involved							

Three levels of assessment quality scores

Yes (Y)
Unclear (U)
No (N)

Journal Pre-proof

Table 4. Overview Nurse-Led Interventions in Older Adults affected by Cancer

Author and Year	Purpose	Intervention
Bigelow et al. 2021 ²⁵	To describe the implementation, associated interventions, and outcomes of the PFRC's proactive virtual resource centre navigation model.	This virtual nurse-led interventions to reduce barriers to care during the pandemic. The nurse navigator determines the complexity of needs and risk factors then triaged to the appropriate team member. Visits were completed by phone or video. When unmet need is identified, respective interventions internal or external such as financial relief, food pantry resources, external referrals, transport, and educational support are provided.
Casey et al. 2017 ²⁶	To evaluate patient satisfaction with a nurse-led phone call follow up clinic for patients with prostate cancer.	A nurse-led telephone follow up service for patients with stable prostate cancer. A satisfaction survey was used to assess patient's satisfaction with the service. All patients were assessed every 6 months by phone at a pre-agreed time and date. This consisted of a recent PSA test and assessment of IPSS score, ECOG status, and side effects from treatment or any new symptoms consistent with local or metastatic disease progression. Triggers to discuss patients with their consultant included any change in symptoms or rise in PSA nadir.
Craven et al. 2012 ²⁸	To explore the usefulness of a nurse-led telephone intervention for supporting cancer patients treated with Capecitabine.	A nurse-led telephone follow-up service. Patients were assessed at baseline and thereafter had two phone calls during cycle 1 (Day 3 & 10) and one phone call during cycle 2 (Day 10). Call duration was 5–10 min (10–15 min for new patients). Chemotherapy booklet and written information about capecitabine and related toxicities. Patients were given the 24-h 'hotline' phone number of the cancer centre.
Catania et al. 2021 ²⁷	To pilot a nurse-led complex intervention focused on QoL assessment in advanced- disease cancer patients	The INFO-QoL intervention aimed at improving patients' outcomes and overall QoL in advanced-disease cancer patients with palliative care needs. Three main components: 1) a small group interdisciplinary team educational program focusing on QoL issues and interventions that promote better outcomes in advanced-disease cancer care; 2) nurse-patient and nurse-family face-to-face interaction to educate patients and their families on QoL issues; 3) patients' outcomes and QoL assessment and appointing a nurse in charge of the process. The care plan was developed during the daily multidisciplinary staff briefing and included changing treatments/route of administration, monitoring vital signs, providing emotional support, educating patients and their families about the illness and options for care based on their goals and preferences, and initiating decision-making conversations.
Faithfull et al. 2013 ³⁵	To compare outcomes in terms of toxicity, symptoms experienced, quality of life, satisfaction with care and health care costs between those receiving nurse-led care vs. groups receiving standard care.	This nurse-led intervention approach exploring patients' understanding of cancer diagnosis, symptoms, providing information and practical advice, leaflets on healthy eating and radiotherapy in outpatient appointments for 20 mins. Telephone contact was also maintained between clinic appointments to assess health status. Contact was established at the start of radiotherapy and continued throughout treatment until 12 weeks. The provision of information and practical advice on how to recognise early symptoms, what to expect from treatment and how to manage existing problems were considered. A protocol of medication and management for symptoms was agreed with the responsible consultants.

Ferguson and Aning 2015 ²⁹	To describe the implementation of a nurse-led survivorship programme for men with Prostate Cancer	This nurse-led intervention to deliver a survivorship program. Patients were given overview of the role of the survivorship nurse specialist, then were invited to attend face to face appointments for 45 minutes at 10 weeks post treatment, 6 months and 1 year. All men were also invited to attend a 6-week course of "Living with and Beyond" education. The educational programme was developed in collaboration with staff at the Maggie's Centre (prostate cancer overview, radiotherapy overview, side-effects of hormone therapy, post-prostatectomy continence, psychosexual implications of treatment).
Festen et al. 2019 ³⁰	To evaluate nurse-led geriatric assessment and assessment of patient preferences for oncological treatment decisions for older patients with solid malignancy	This nurse-led intervention included the integration of geriatric assessment and assessment of patient's preferences in a multidisciplinary approach to reach tailored treatment advice. Inclusion of the study took place in the outpatient clinic wherein patients will be discussed in the conventional tumour board as well as in Onco-Geriatric MDT where nurses had an active role to compare recommendations and implementation purposes.
Knowles et al. 2007 ³¹	To assess the feasibility of a follow up programme led by nurse specialist for patients with colorectal cancer	This nurse-led intervention included placement of CNS in the multidisciplinary team to coordinate follow up programs with adherence to follow up protocols at each clinic visit. A baseline QoL is measured pre-operatively then in each 4 visits. A questionnaire which is a self-rated tool is then used at 12 month follow up. Clinician satisfaction is also used at the completion of study.
Kotronoulas et al 2017 ⁴¹	To explore the impact of nurse-led PROMS consultation with patients.	This nurse-led intervention included a pre-consultation PROM (SCNS) data were collected during three consecutive, monthly consultations, and used by the CNS to enable delivery of personalised supportive care.
Kotronoulas et al. 2018 ⁴²	To examine whether a nurse-led PRO measure driven is feasible and acceptable in identifying unmet needs in patients with Lung Cancer.	This nurse-led intervention included three consecutive monthly consultations with patients using PRO measure. Subsequently, the lung CNS met with the patient and used PRO data to identify unmet needs, direct discussions, and intervene accordingly. The lung CNS documented any identified needs and clinical interventions/advice.
Malmstrom et al. 2016 ⁵⁵	To evaluate the effect of a nurse-led telephone supportive care program on QoL compared to conventional care on patients following esophageal resection for cancer.	This nurse-led intervention included a meeting before discharge where the patients had the opportunity to ask questions, discuss their concerns and received both oral and written information focusing on life after surgery, self-care, plans for the future, and where to turn to for help if needed. After discharge, the follow-up by the nurse was proactive and focused on the patients individual needs of support as well as areas known to be problematic for patients after this type of surgery e.g. nutrition, elimination, pain and psychological issues aiming to detect possible problems in an early stage and to help patients to manage them.
Martin et al. 2018 ³²	To explore the impact of nurse-led service among men undergoing AS for prostate cancer.	The nurse-led intervention included nurse assessment of current health status (and sexual function/ability, where appropriate) and LUTS symptoms, review of PSA and other biochemical results, DRE at 6/12 intervals or if LUTS deteriorated, tailored discussion of all findings with patient ± spouse/partner/carers and offer copy of clinic letter, arrange 2 yearly transrectal ultrasound/template biopsies or sooner if PSA/DRE deteriorates ± MRI, arrange follow-up and arrange PSA before next visit. Clear rationale and trigger for safety net to refer to consultants.

McGlynn et al. 2014 ³³	A local evaluation of the innovative nurse-led collaborative care model for the management of patients with prostate cancer.	A nurse led collaborative care model for the management of patients with prostate cancer. The nurse makes a full assessment and advises/plans further management appropriately, with advice as required either from the Nurse Consultant or consultant/medical team. All patients have contact details for the urology oncology nursing team and are encouraged to be in touch as required.
Primeau et al. 2017 ⁴⁴	To explore the experience of patients and their partner/caregiver as well as MDT members of a nurse-led multimodality supportive care intervention in men with metastatic prostate cancer as well as standard care.	This nurse-led intervention included patients and their partners/caregivers by completing a holistic needs assessment prior to routine 3 monthly follow up. A clinical review is then conducted by the PCNS that lasts for 40-90 minutes. Information collected used to identify supportive care needs and tailor self-management support through ThriveCare intervention.
Ream et al. 2009 ⁴³	To investigate the roles of Prostate Cancer Clinical Nurse Specialists and determine their targeted services, work practises and perceived contribution.	This nurse-led intervention related to caring activities related to care co-ordination, emotional care, treatment advice, symptom assessment and management, giving results, treatment administration, preoperative preparation, and monitoring at risk patients.
Reinke et al. 2022 ⁴⁰	To assess the effect of a nurse-led telephone-based primary palliative care intervention in patients with lung cancer.	This nurse-led intervention included a one-day End of Life Nursing Education Course for Veterans diagnosed with lung cancer and an eight-hour online communication course, and nurse support and education on lung cancer symptom management.
Schenker et al. 2021 ³⁷	To evaluate palliative nurse-led model.	The CONNECT intervention included 3 monthly visits with an existing infusion room nurse who was trained to address symptoms, provide emotional support, engage in advance care planning, and coordinate care. Conceptually grounded in the chronic care model described by Wagner and colleagues CONNECT used an oncology nurse-led care management approach to improve the provision of primary palliative care within outpatient oncology practices
Schofield et al. 2016 ³⁸	To investigate the benefits of a group nurse-led intervention in men receiving radiotherapy for prostate cancer.	The intervention package was designed to: 1) systematically assess patient needs and values to direct the content of consultations; 2) provide timely information on basic prostate anatomy, side effects, treatment, and survivorship issues at critical points in the treatment trajectory; 3) coach men in evidence-based self-care and communication strategies with their treatment team to assist them to achieve optimal health status; and 4) offer a forum for psychosocial peer support and information exchange. It consists of four group consultations and one individual consultation.
Sibbons et al. 2019 ³⁴	To explore the impact of nurse-led service among patients affected with renal cancer.	This nurse-led intervention included patients for follow-up appointments after either radical or partial nephrectomy surgery for histologically proven renal cell carcinomas at either 3, 6 or 12 monthly intervals depending upon their stage, grade and original diagnosis. The clinic is run by two clinical nurse specialists on a weekly basis and consists on average of six patients per clinic, utilising 30-minute slots. No further details reported about the nursing process of care.

Stanciu et al. 2018 ³⁹	To evaluate a nurse-led model of personalised care after prostate cancer treatment	This nurse-led intervention included the use of a comprehensive holistic needs assessment tool and care plan, specifically exploring the physical, emotional, spiritual, lifestyle and family aspects of cancer survivorship, together with an additional bespoke instrument developed in secondary care to monitor physical symptoms. Following the assessment, the nurse will provide individualised information, advice and support tailored to each patient, to help men, improve their symptoms or cope better with symptoms they cannot improve. Patient referral to GP or secondary care and signposting to community or third sector support services was made as appropriate.
Van der Meulen et al. 2013 ⁴⁵	To develop a nurse-led educational intervention to provide information during a discharge interview and to investigate the effects of the intervention on information needs and satisfaction with information in head and neck cancer.	This nurse-led intervention provided educational intervention in a 30-45 minute structured conversation about general information, wound care, physical-social problems, work and finances.




Figure. PRISMA Diagram

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof

Supplementary Table 1. Exemplar of Database searches

Database	Date	Search Terms	Limiters	# Results
CINAHL	14/09/2021 UPDATED 20/09/2022	((nurse-led OR nurse-managed) N3 (care OR model* OR program* OR intervention*)) AND (cancer* OR oncolog* OR neoplasm*)	Date: 1990- Language: English	292 49
Google Scholar	14/09/2021 UPDATED 20/09/2022	nurse-led nurse-managed care model program intervention cancer oncology neoplasm	Date 1990-	34 2
MEDLINE	14/09/2021 UPDATED 20/09/2022	((nurse-led OR nurse-managed) N3 (care OR model* OR program* OR intervention*)) AND (cancer* OR oncolog* OR neoplasm*)	Date: 1990- Language: English	322 64
PsycINFO	14/09/2021 UPDATED 20/09/2022	((nurse-led OR nurse-managed) N3 (care OR model* OR program* OR intervention*)) AND (cancer* OR oncolog* OR neoplasm*)	Date: 1990- Language: English	90 3
Scopus	14/09/2021 UPDATED 20/09/2022	((nurse-led OR nurse-managed) W/3 (care OR model* OR program* OR intervention*)) AND (cancer* OR oncolog* OR neoplasm*)	Date: 1990- Language: English	328 60
Total:				1066 1244