

MaineHealth

MaineHealth Knowledge Connection

Costas T. Lambrew Research Retreat 2023

Costas T. Lambrew Research Retreat

5-2023

The Use of the iPACK Block with the Adductor Canal Block (ACB) Decreases 48-hour Narcotic Usage and Postoperative Pain following Total Knee Arthroplasty

Bailey Shevenell

Johanna Mackenzie

Matthew Hersch

Michael Becker

Brian J. McGrory

See next page for additional authors

Follow this and additional works at: <https://knowledgeconnection.mainehealth.org/lambrew-retreat-2023>



Part of the [Surgery Commons](#)

Authors

Bailey Shevenell, Johanna Mackenzie, Matthew Hersch, Michael Becker, Brian J. McGrory, Nicholas Colacchio, Adam Rana, Ryan Mountjoy, and Rob Hubbs

Adductor Canal and iPACK Blocks reduce pain levels and narcotic consumption following Total Knee Arthroplasty

Adductor Canal and iPACK Blocks reduce the length of stay and Narcotic consumption after Total Knee Arthroplasty

Bailey Shevenell, BA; Johanna Mackenzie, MPH; Ryan Mountjoy, MD; Matthew Hersch; Brian McGrory, MD; Peter R Hubbs, MD; Michael Becker, MD; Adam Rana, MD

Introduction

- Multimodal analgesic pain protocols, such as the use of nerve block techniques, have been implemented in TKA procedures to improve recovery and limit the use of prescribed narcotics.
- Traditionally, the adductor canal block (ACB) has utilized local anesthetics for pain management, however recent data shows that pain management is improved when administering liposomal bupivacaine into the ACB and using additional nerve blocks such as the iPACK (interspace between the popliteal artery and capsule of the posterior knee) block.
- This study compares postoperative outcomes of patients that received either the ACB block alone or in addition to the iPACK block.

Methods

1. Retrospective cohort study of all primary, unilateral TKA patients between 2015-2021.
2. Patient demographics, preoperative, perioperative, and postoperative data was collected from EPIC.
3. Statistical analysis was performed in Excel.

Results

- There were 2,321 patients identified for this study.
- The ACB and iPACK group (n = 577) showed a decreased MME consumption during initial hospital stay ($p < 0.001$) and had lower pain levels at 6-weeks postoperatively ($p = 0.003$) than the ACB group (n = 1,744).
- The ACB and iPACK group also had a shorter length of stay ($p < 0.001$) and were more likely to be discharged home ($p < 0.001$).

Discussion

- When used in conjunction, the ACB and iPACK nerve block techniques are associated with a decrease in analgesic narcotics, length of stay, and postoperative pain scores than the use of ACB alone.
- Our results suggest a decreased period of recovery following TKA surgeries, leading to better outcomes and higher patient satisfaction with their surgery.

Short-term Postoperative MME Consumption after Total Knee Arthroplasty





Figure 1. Average MME consumption by postoperative time period and total length of acute hospital stay between ACB and ACB + iPACK block.



Take a picture to
download the full paper

Maine Medical

PARTNERS

A department of Maine Medical Center

Maine Medical Center
MaineHealth