

Grassland dynamics of soil moisture and temperature

Original

Grassland dynamics of soil moisture and temperature / Ferraris, Stefano; N'Sassila, Mesmer; Gentile, Alessio; Galvagno, Marta; Stevenin, Herve; Canone, Davide; Previati, Maurizio; Bevilacqua, Ivan; Gisolo, Davide; Painter, KEVIN JOHN. - ELETTRONICO. - (2021). (Intervento presentato al convegno EGU General Assembly 2021 tenutosi a online nel 19-30 Aprile 2021) [10.5194/egusphere-egu21-15800].

Availability:

This version is available at: 11583/2977678 since: 2023-07-12T11:31:11Z

Publisher:

EGU

Published

DOI:10.5194/egusphere-egu21-15800

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

EGU21-15800, updated on 28 Apr 2021
<https://doi.org/10.5194/egusphere-egu21-15800>
EGU General Assembly 2021
© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Grassland dynamics of soil moisture and temperature

Stefano Ferraris¹, Mesmer N'Sassila¹, Alessio Gentile¹, Marta Galvagno², Herve Stevenin³, Davide Canone¹, Maurizio Previati¹, Ivan Bevilacqua¹, Davide Gisolo¹, and Kevin Painter¹

¹Politecnico and Università of Torino, DIST, Torino, Italy (stefano.ferraris@unito.it)

²ARPA Valle d'Aosta, Italy

³Centro Funzionale Regione Valle d'Aosta, Italy

Alpine ecosystems are vulnerable to climate and land use changes. Measurement sites at different altitude and aspect can provide precious information on them. Also, vadose rootzone plays a major role in partitioning fluxes. In this work field data of soil water content, matric potential and soil temperature in some mountain grassland sites are compared with simulations results of the CLM model (The Community Land Model, NCAR, US). These are key state variables regulating the physical processes that determine the flows of two main greenhouse gases, water vapour and carbon dioxide, to the atmosphere in the presence of vegetation.

Some transients show significant differences between data and CLM simulation results and further analyses are performed using the HYDRUS model from the US Salinity Laboratory, in order to better explore the soil, grass, and atmosphere roles in the dynamics of those state variables. Some insight is finally provided about the effects on water vapour and carbon dioxide fluxes.