

Actual evapotranspiration of abandoned grassland on a slope in the Western Italian Alps: Impact of shrub encroachment

*Original*

Actual evapotranspiration of abandoned grassland on a slope in the Western Italian Alps: Impact of shrub encroachment / Gisolo, Davide; Bevilacqua, Ivan; van Ramshorst, Justus; Knohl, Alexander; Siebicke, Lukas; Gentile, Alessio; Previati, Maurizio; Canone, Davide; Ferraris, Stefano. - (2022). (Intervento presentato al convegno EGU General Assembly 2022 tenutosi a Vienna nel 23-27 Maggio 2022) [10.5194/egusphere-egu22-7320].

*Availability:*

This version is available at: 11583/2977684 since: 2023-03-31T13:49:28Z

*Publisher:*

EGU

*Published*

DOI:10.5194/egusphere-egu22-7320

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)



## Actual evapotranspiration of abandoned grassland on a slope in the Western Italian Alps: Impact of shrub encroachment

**Davide Gisolo**<sup>1</sup>, Ivan Bevilacqua<sup>1</sup>, Justus van Ramshorst<sup>2</sup>, Alexander Knohl<sup>2,3</sup>, Lukas Siebicke<sup>2</sup>, Alessio Gentile<sup>1</sup>, Maurizio Previati<sup>1</sup>, Davide Canone<sup>1</sup>, and Stefano Ferraris<sup>1</sup>

<sup>1</sup>Interuniversity Dept. of Regional and Urban Studies and Planning (DIST), Polytechnic Univ. of Turin, Italy

<sup>2</sup>Bioclimatology, Faculty of Forest Sciences and Forest Ecology, University of Göttingen, Germany

<sup>3</sup>Centre of Biodiversity and Sustainable Land Use (CBL), University of Göttingen, Göttingen, Germany

Land cover changes affect the local hydrological cycle, including actual evapotranspiration (ET<sub>a</sub>). Encroachment by shrubs on abandoned grasslands is an increasing phenomenon in the Alps, a region already suffering climate change effects. In addition, shrub encroachment is thought to occur faster on steep slopes. Unfortunately, steep mountain slopes are rarely studied because of complex morphologies, despite a need for data to better understand these changing ecosystems.

Four growing seasons (two wet – 2014 and 2015 and two dry – 2016 and 2017) of eddy covariance, meteorological, hydrological, and soil data were collected at an abandoned grassland on a slope encroached by shrubs in the Italian Western Alps. The objectives were to: 1) study the ET<sub>a</sub> differences between two land cover types, grassland and shrubland, based on Hydrus 1D model simulations. 2) Compare the simulated ET<sub>a</sub> from the two land covers (ET<sub>a,Sim grass</sub> and ET<sub>a,Sim shrub</sub>) with the observed eddy covariance-derived evapotranspiration (ET<sub>a,Obs</sub>).

The simulated ET<sub>a</sub> from shrubland showed a better agreement with the observed ET<sub>a</sub> ( $R^2=0.4$  to  $0.5$ , slope= $0.8$  to  $1.3$ ). The simulated ET<sub>a</sub> from shrubland (ET<sub>a,Sim shrub</sub>) was higher compared to the simulated ET<sub>a</sub> from grassland (ET<sub>a,Sim grass</sub>) with the observations (ET<sub>a,Obs</sub>) in between, confirming that ET<sub>a,Obs</sub> represents a mixture of shrubland and grassland contributions. The relative contribution was different for each year due to meteorological conditions. On average across all years, a 51:49% contribution from respectively grassland and shrubland resulted in a good approximation of ET<sub>a,Obs</sub>, in particular in 2015 and 2016 growing seasons, characterised by long dry spells. In those growing seasons, the differences between cumulative ET<sub>a</sub> from simulations and observations were below 10 mm. In the other two growing seasons, more frequent rainfalls and the absence of long dry spell caused cumulative ET<sub>a</sub> underestimation ( $-25$  mm) in 2014 and overestimation (66 mm) in 2017. Differences between shrubland and grassland were enhanced during dry spells, leading to a cumulative ET<sub>a,Sim shrub</sub> more than 100 mm higher than the cumulative ET<sub>a,Sim grass</sub>. In the longest dry spells of the growing seasons, ET<sub>a,Obs</sub> was closer to ET<sub>a,Sim shrub</sub>, confirming the role of deeper roots of shrubs.

The results indicate that the shrub-covered area, expected to increase, plays already a key role in the local hydrological cycle, particularly with changes in water availability.

