

Talaat Abdel-Fattah Abdel-Aty<sup>1</sup>, Eman Yousef Moursi<sup>1</sup>, Eman Tayae Elsayed<sup>2</sup>,  
Marwa Ahmed Salah<sup>1</sup>, Sary Ibrahim Abdelsalam<sup>3</sup>

<sup>1</sup>Faculty of Medicine, Internal Medicine Department, Alexandria University, Alexandria, Egypt

<sup>2</sup>Faculty of Medicine, Clinical and Chemical Pathology, Alexandria University, Alexandria, Egypt

<sup>3</sup>General Organization for Teaching Hospitals and Institutes (GOTHI), Denshal Teaching Hospital, Internal Medicine Department, Dammanhur, Egypt

# Association of Angiopoietin-Like Protein-8 Gene Variant (Rs2278426 (C/T)) in a Cohort of Egyptian Patients with Metabolic Syndrome: A Case-Control Study

## ABSTRACT

**Objective:** This study was conducted to reveal the association of *ANGPTL8* gene variant (Rs2278426 (C/T)) with metabolic syndrome in a cohort of Egyptian patients. **Materials and methods:** This study is a case control study that included 150 patients with metabolic syndrome and 150 healthy control subjects. All subjects were submitted to history taking and thorough physical examination and laboratory analysis. Genomic DNA was extracted and *ANGPTL8* gene SNP (rs2278426) was detected by 5' nuclease assay. The tested genotypes included homozygous genotypes for C allele (CC), homozygous genotypes for T allele (TT) and heterozygous genotypes (CT).

**Results:** A total of 300 subjects were included in the study; group 1 included 150 patients with metabolic syndrome. 21.3% were males and 78.7% were females, and group 2 had 150 healthy subjects. 17.3% were males, and 82.7% were females. In the current study, the metabolic syndrome group showed dysregulation

of lipids and fasting plasma glucose (FPG) with a statistically significant increase in body anthropometric measures. There was no statistically significant difference in the distribution of the heterozygous genotype (C/T) relative to the wild-type genotype (C/C) in each of the two tested groups ( $p = 0.287$  and  $0.245$  in the metabolic syndrome and control groups, respectively). **Conclusions:** There was no statistically significant difference in the genotype distribution of *ANGPTL8* gene variant (Rs2278426) genotypes between the metabolic syndrome and control groups. The homozygous recessive genotype for T allele (TT) was not detected in both tested groups.

**Keywords:** metabolic syndrome, *ANGPTL8*, rs2278426, genotyping, polymerase chain reaction (PCR)

## Introduction

Metabolic syndrome (MetS) is considered a major risk factor for cardiovascular diseases and stroke in both developing and developed countries. Central obesity and type 2 diabetes (T2D) are considered the major endophenotypes of MetS, which are inheritable and genetically based [1].

In Egypt, 22% of men and 48% of women were obese, according to a 2010 World Health Organization (WHO) estimate. The Middle East and North Africa Region is considered to have the highest prevalence of diabetes at 10.9% [2].

## Address for correspondence:

Dr. Sary Ibrahim Abdelsalam  
Denshal Teaching Hospital, General Organization for Teaching Hospitals and Institutes (GOTHI), Egypt  
phone: 00201098997094, 00201117862334  
e-mail: dr\_sary2011@hotmail.com  
Clinical Diabetology  
DOI: 10.5603/DK.a2023.0023  
Received: 8.04.2023 Accepted: 1.06.2023  
Early publication date: 11.08.2023

Angiopoietin-like proteins (ANGPTLs) are proteins that are primarily expressed in the liver and adipose tissue and circulate in human plasma. Expression of ANGPTL8 was found to be reduced by fasting and increased by re-feeding in both mice and humans [3]. ANGPTL8 has been associated with two functionally important processes in the development of T2D, insulin resistance and lipid metabolism, and has also been reported to regulate the replication of  $\beta$ -cells in response to insulin resistance [4, 5].

Overexpression of ANGPTL8 leads to increased plasma levels of triacylglycerol (TAG) and lipoproteins via inhibition of LPL activity [6, 7]. Therefore, ANGPTL8 has been suggested as a potential therapeutic target for dyslipidemia, and inhibition of ANGPTL8 has been highlighted as a novel therapeutic strategy for reducing plasma lipoprotein levels [8].

C > T (rs2278426) is a commonly studied variant in *ANGPTL8* gene located on chromosome 19, resulting from the substitution of C for T at c194, that affects levels of the activated form of ANGPTL3 with a similar effect to complete ANGPTL3 deficiency [9].

Due to this suggested association, the current study was conducted to reveal the association of *ANGPTL8* gene variant (Rs2278426 (C/T)) with metabolic syndrome in a cohort of Egyptian patients.

## Materials and methods

### Study design

This is a case-control study that was conducted at the Diabetes and Metabolism Clinic at the Internal Medicine Department, Alexandria University Hospitals, Alexandria, Egypt, for a duration of 1 year between January 2022 and December 2022.

### Study population

This study included a total of 300 subjects who were divided into two groups: Group 1 (the study group), which included 150 patients with metabolic syndrome, and Group 2 (the control group), which included 150 apparently healthy persons.

### Ethical approval

The study was conducted in accordance with Helsinki Standards as revised in 2013 [10]. The study was conducted after obtaining approval from the local ethics committee, Faculty of Medicine, Alexandria University, and after obtaining a written or oral informed consent from the included cases.

The participants were invited from the outpatient clinics of the diabetes and metabolism unit at the Internal Medicine Department, Alexandria University Hospitals, Alexandria, Egypt, for a duration of 1 year between January 2022 and December 2022.

After explanation of the study purpose an informed written consent was obtained from all the participants. The subjects were informed about their right to withdraw at any time from the study.

### Inclusion criteria

We included the patients from both genders aged more than 18 years who were diagnosed to have metabolic syndrome. According to the IDF definition, metabolic syndrome is diagnosed if anyone has central obesity plus two or more of the following four criteria [11]: Triglyceride concentration > 150 mg/dL, high density lipoprotein cholesterol (HDL-C) < 40mg/dL in men and < 50 mg/dL in women, or receiving lipid lowering drugs, systolic blood pressure > 130 mmHg or diastolic blood pressure > 85 mmHg or receiving treatment for hypertension and fasting plasma glucose > 100 mg/dL or previously diagnosed type 2 diabetes.

### Exclusion criteria

The study excluded cases with the following criteria: age < 18 years, patients with type 1 diabetes, patients with acute infection or illness, malignancy, liver cell failure, kidney failure, heart failure, unstable medical condition during the previous 30 days, patients with associated active inflammatory conditions and gastrointestinal diseases, psychological disease and patients who refused to participate in the study.

The cases were subjected to the following: history taking, a thorough physical examination, and a complete clinical history and clinical examination. Anthropometric measures of all the subjects in the study were obtained, including weight, height, body mass index (BMI), and waist circumference.

Body mass index is defined as the weight in kilograms divided by the square of the height in meters ( $\text{kg}/\text{m}^2$ ). BMI values between 18.5 and 24.99  $\text{kg}/\text{m}^2$  are considered normal; individuals with BMI values from 25 to 29.9  $\text{kg}/\text{m}^2$  are considered overweight; and those with values greater than 30  $\text{kg}/\text{m}^2$  are considered obese [12].

Waist circumference (a measurement taken around the abdomen at the level of the umbilicus) The NIH was the first to use the threshold values for waist circumference ( $\geq 88$  cm in women and  $\geq 102$  cm in men) as suggested by Michael Lean and colleagues in combination with a classification of overall obesity as assessed by BMI [13].

### Laboratory data

Lipid profile [triglyceride (TG) (normal: less than 150 milligrams per deciliter (mg/dL), total cholesterol (TC) (normal: less than 200 mg/dL), high density lipo-

protein cholesterol (HDL-C) (60 mg/dL or above), and low-density lipoprotein cholesterol (LDL-C) (the normal range is 50 to 100 mg/dL) and diabetic profile [fasting plasma glucose (FPG) (a fasting plasma glucose level of 99 mg/dL or lower is normal), glycosylated hemoglobin (HbA1c) (normal HbA1c level is  $\leq 5.6$ , pre-diabetes 5.7–6.4 and diabetes  $\geq 6.5$  or below), and fasting serum insulin (normal fasting insulin levels range between 5 and 15  $\mu\text{U/mL}$ )].

Assessment of the insulin resistance (IR) index by using Homeostatic Model Assessment of insulin resistance (HOMA-IR) equation:  $\text{HOMA-IR} = \text{fasting insulin } (\mu\text{U/mL}) \times \text{fasting plasma glucose (mg/dL)} / 405$  [14].

### Blood sampling

All blood samples were divided into two aliquots: the first part was collected in a vacutainer tube containing Na<sub>2</sub>-EDTA for the assay of HbA1c; second part was collected in a plain vacutainer tube and centrifuged (3000 rpm) for serum preparation. Serum was used to measure total cholesterol, triglycerides, LDL cholesterol, HDL cholesterol and glucose. Serum was used to measure total cholesterol, triglycerides, LDL and HDL cholesterol, glucose. Low-density lipoprotein cholesterol was calculated by using Friedewald formula:

$$\text{TC} - (\text{HDL-C}) - (\text{TG}/5) \text{ [14].}$$

Fasting serum insulin [ $\mu\text{IU/mL}$ ] was measured by ELISA from the serum sample. Assessment of insulin resistance (IR) index was made using Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) equation:

$$\text{HOMA-IR} = \text{fasting insulin } (\mu\text{U/mL}) \times \text{fasting glucose (mg/dL)} / 405.$$

### Genetic analysis

***ANGPTL8* gene SNP (rs2278426) detection by allelic discrimination using a fluorogenic 5' nuclease assay**

#### a) DNA extraction:

Genomic DNA was extracted from an EDTA blood sample using the QIAamp DNA Mini Kit according to the manufacturer's instructions (Qiagen, Germany). The extracted DNA concentration and purity were assessed using a NanoDrop 2000/2000c spectrophotometer (Thermo Scientific, USA). Ratios of 260/280 and 260/230 = 1.7–2 indicated highly pure DNA.

#### b) 5' nuclease assay:

*ANGPTL8* gene SNP (rs2278426) was detected by 5' nuclease assay using TaqMan<sup>TM</sup> Genotyping Master Mix, TaqMan<sup>®</sup> SNP Genotyping Assay 40x (Assay ID: C\_15965472\_10) (Thermo Fisher Scientific, USA), and

Rotor-gene Q real time polymerase chain reaction (PCR) system (Qiagen, Germany).

### Principle

Real-time PCR utilizes the 5' to 3' exonuclease activity of Taq polymerase as the enzyme cleaves the 5' terminal nucleotide of double stranded DNA. A fluorescent oligonucleotide TaqMan probe is designed to anneal between the two amplification primers in the PCR reaction. This probe is labeled with two fluorescent dyes, a reporter dye at the 5' end and a quencher dye at the 3' end. The quencher exerts a quenching effect on the fluorescence of the reporter dye as long as the probe is intact and the two dyes are in close proximity.

During extension, Taq polymerase partially displaces the hybridized probe and then cleaves it at the 5' end. The cleavage of the probe between the reporter and the quencher physically separates the two dyes, resulting in increased reporter fluorescence, which indicates that the probe-specific target has been amplified.

For allelic discrimination, two probes with different reporters were used: one gave a green color called "FAM" and the other gave a yellow color called "VIC". An increase in the fluorescence signals of a particular dye indicates homozygosity for the allele whose probe was labeled by that dye; an increase in both signals indicates heterozygosity.

### Statistical analysis

The data collected were coded, processed, and analyzed with SPSS version 26 for Windows<sup>®</sup> (Statistical Package for Social Sciences) (IBM, SPSS Inc., Chicago, IL, USA). Qualitative data in the form of numbers (frequency) and percent was presented. The Chi-Square test (or Fisher's exact test) made the comparison between groups. The Kolmogorov-Smirnov test tested quantitative data for normality. Parametric data were shown as median  $\pm$  SD, while non-parametric data were expressed as median (interquartile range).

To compare two groups with normally distributed quantitative variables, an independent sample (Student's) t-test was used, and a Mann-Whitney U-test was used if the data were abnormally distributed. The Hardy-Weinberg Equilibrium equation was used to assess the genotype frequency distribution in each of the two tested groups. For all tests, P values < 0.05 are considered significant.

The sample size was estimated based on the Epi Info V 7.0 program with a power of 80 and a level of significance of 95%, considering the percent of exposed to the outcome of 16% and the percent of unexposed to the outcome of 7.4%. The program revealed

**Table 1. Demographic Data, Anthropometric Measures, Comorbidities Blood Pressure and Laboratory Data in Cases and Control**

	Group 1 (Cases group)	Group 2 (Control group)	P
Age [years] (Mean $\pm$ SD)	42.3 $\pm$ 10.6	39.9 $\pm$ 12.8	0.07
Female gender	118 (78.7%)	124 (82.7%)	0.380
BMI [kg/m <sup>2</sup> ] (mean $\pm$ SD)	39.6 $\pm$ 6.3	22.4 $\pm$ 1.6	< 0.001
Waist circumference [cm] (mean $\pm$ SD)	115.3 $\pm$ 9.4	88.7 $\pm$ 7.0	< 0.001
Comorbidities			
Type 2 diabetes	44 (29.3%)		
Diabetics on oral	35 (23.3%)		
Diabetics on Insulin	6 (4%)		
Hypertension	100 (66.7%)		
Hypertension on treatment	63 (42%)		
Family history of diabetes	58 (38.7%)		
Systolic blood pressure [mmHg] (mean $\pm$ SD)	129.4 $\pm$ 17.9	116.5 $\pm$ 9.5	< 0.001*
DBP [mmHg] (mean $\pm$ SD)	86.6 $\pm$ 11.7	75.4 $\pm$ 5.6	< 0.001*
Fasting insulin [ $\mu$ IU/mL] [median (IQR)]	15.0 (12.5 – 18.0)	8.0 (6.8–9.6)	< 0.001*
FPG [mg/dL] (mean $\pm$ SD)	116.5 $\pm$ 39.4	87.4 $\pm$ 9.3	< 0.001*
HbA1c [%] (mean $\pm$ SD)	6.3 $\pm$ 1.3	5.0 $\pm$ 0.3	< 0.001*
HOMA-IR [median (IQR)]	2.0 (1.7–2.5)	1.0 (0.8–1.2)	< 0.001*
TGs [mg/dL] [median (IQR)]	139.0 (104.0–175.0)	101.5 (76.0–122.0)	< 0.001*
TC [mg/dL] (mean $\pm$ SD)	207.1 $\pm$ 34.9	176.3 $\pm$ 19.7	< 0.001*
LDL-C [mg/dL] (mean $\pm$ SD)	134.3 $\pm$ 33.9	102.6 $\pm$ 18.8	< 0.001*
HDL-C [mg/dL] (mean $\pm$ SD)	43.6 $\pm$ 5.4	53.8 $\pm$ 4.70	< 0.001*

\*Statistically significant at  $p \leq 0.05$

BMI — body mass index; DBP — diastolic blood pressure; FPG — fasting plasma glucose; HbA1c — glycated hemoglobin; HDL-C — high-density lipoprotein cholesterol; HOMA-IR — Homeostatic Model Assessment for Insulin Resistance, IQR — interquartile range; LDL-C — low-density lipoprotein cholesterol; SBP — systolic blood pressure; SD — standard deviation; TC — total cholesterol; TG — triglycerides

a minimum sample size of 246 individuals, 123 for each group. However, this study will be conducted on a sample size of 300 subjects for the purpose of improving the validity of the research results.

## Results

Table 1 shows that there was no statistically significant difference between the two groups regarding the age and sex. The BMI and waist circumference were statistically significantly higher in the metabolic syndrome group as compared to the control. In the metabolic syndrome group, the prevalence of diabetes was 29.3% and the prevalence of hypertension was 66.7%.

The mean systolic blood pressure and the mean diastolic blood pressure were statistically significantly higher in the metabolic syndrome group as compared to the control. The mean fasting plasma glucose level, postprandial plasma glucose level, HbA1c, and HOMA-IR were statistically significantly higher in the metabolic syndrome group as compared to the control.

The mean triglyceride level, total cholesterol level, and low density lipoprotein level were statistically significantly higher in the metabolic syndrome group as compared to the control. The mean high density lipoprotein level was statistically significantly lower in the metabolic syndrome group as compared to the control.

Regarding the distribution of the genotype of the tested gene in the two groups; there were 126 cases (84%) with wild type form and 24 (16%) with heterotype form in the metabolic syndrome group, while in the control group, wild type was detected in 82.7% and heterotype was detected in 17.3%, with no statistically significant difference between the two groups ( $p = 0.757$ ).

Also, by using Hardy-Weinberg Equilibrium for genotype analysis, there was no statistically significant difference in the distribution of the heterozygous genotype (C/T) relative to the wild type genotype (C/C) in each of the two tested groups ( $p = 0.287$  and  $0.245$  in the metabolic syndrome and control groups, respectively).

**Table 2. Association between Genotype with Different Parameters in Group 1 (n = 150)**

	Genotype		P
	Wild (n = 126)	Hetero (C/T) (n = 24)	
Gender			
Female	100 (79.4%)	18 (75.0%)	0.63
Age [years] (mean ± SD)	42.9 ± 10.8	39.4 ± 8.9	0.14
BMI [kg/m <sup>2</sup> ] (mean ± SD)	39.8 ± 6.2	38.8 ± 6.6	0.45
Waist circumference [cm] (mean ± SD)	115.5 ± 9.3	114.4 ± 9.5	0.56
SBP [mmHg] (mean ± SD)	130.1 ± 18.9	125.8 ± 11.8	0.29
DBP [mmHg] (mean ± SD)	86.8 ± 12.0	85.6 ± 10.1	0.66
Fasting insulin (mean ± SD)	15.4 ± 4.3	14.7 ± 4.4	0.45
FPG [median (IQR)]	107.0 (71.0–361.0)	100.0 (63.0–172.0)	0.18
HbA1c [%] (mean ± SD)	6.3 ± 1.3	6.2 ± 1.1	0.59
HOMA-IR [median (IQR)]	2.1 (0.6–6.6)	1.8 (1.0–3.5)	0.29
TGs [median (IQR)]	139.0 (53.0–298.0)	145.5 (71.0–301.0)	0.95
TC (mean ± SD)	207.6 ± 35.5	204.4 ± 32.16	0.68
LDL-C (mean ± SD)	135.1 ± 34.3	129.8 ± 32.8	0.48
HDL-C (mean ± SD)	43.2 ± 5.3	46.1 ± 5.4	0.01*

\*Statistically significant at  $p \leq 0.05$

BMI — body mass index; DBP — diastolic blood pressure; FPG — fasting plasma glucose; HbA1c — glycated hemoglobin; HDL-C — high-density lipoprotein cholesterol; HOMA-IR — Homeostatic Model Assessment for Insulin Resistance, IQR — interquartile range; LDL-C — low-density lipoprotein cholesterol; SBP — systolic blood pressure; SD — standard deviation; TC — total cholesterol; TG — triglycerides

Table 2 shows that there was no statistically significant difference between the wild type and heterotype genotypes in all the tested clinical and laboratory parameters except for HDL-C.

The mean level of HDL-C in the hetero (C/T) genotype was  $46.13 \pm 5.45$  mg/dL, which was statistically significantly higher compared to the wild type genotype ( $43.17 \pm 5.33$  mg/dL) ( $p = 0.014$ ).

## Discussion

The current study included 300 subjects in two groups; group 1 (which included 150 patients diagnosed as having metabolic syndrome) and group 2 (which included 150 healthy subjects as a control group).

In the current study, the prevalence of metabolic syndrome was higher in females, who represented 78.7% of the cases with metabolic syndrome. This agreed with Sliem et al., who ascertained in their study that women had a higher prevalence of the MetS than men, especially in Iran, India, Oman, Pakistan, Saudi Arabia, and Egypt; this might be due to cultural barriers to physical activity that have been reported among women [15]. Similarly, in a study in Egypt in 2010, Nasr et al., claimed that the prevalence of obesity was 70.9% among Egyptian patients included in their study and that women had an elevated prevalence of the MetS compared to men [16].

On the other hand, in Finland, the prevalence of MetS was higher in males compared with females in a study conducted on subjects between 45 and 64 years old, as the occurrence of MetS was 52.6% among males and 45.3% among females [17]. The variations could be explained by variations in population characteristics.

Consistent with previous studies, BMI was found to be significantly correlated with waist circumference in individuals with MetS [18]. BMI showed robust performance in estimating visceral fat measured using computed tomography compared with waist circumference [19]. A credible body of evidence supports waist circumference as a better predictor of MetS. In fact, BMI cannot account for body fat distribution [20].

In the current study, the prevalence of hypertension in the metabolic syndrome group was 66.7%, and 42% only received hypertension treatment. The mean SBP and DBP were statistically significantly higher in the cases with metabolic syndrome compared to the controls. This agreed with Abd Elaziz et al. [2] who included 220 middle-aged and senior Egyptians who were voluntarily screened in an Egyptian private hospital. The authors observed that ~ 76.6% of hypertensive patients had MetS.

In the current study, the prevalence of T2D in the metabolic syndrome group was 29.3%. Also, the fasting insulin level, fasting plasma glucose, HbA1c, and HOMA-IR were statistically significantly higher in the

metabolic syndrome group. This agreed with Abd Elaziz et al., who showed that the prevalence of MetS among people with diabetes in their study was 85.6%. This was also similar to a study done on Pakistani patients with T2D, where 85.5% of the people with diabetes had MetS [21].

In the current study, the mean TGs level, TC level, and LDL-C were statistically significantly higher in the cases with metabolic syndrome, while the HDL-C was statistically significantly lower in the cases group compared to the control. This came in accordance with Refaat et al. [22], who included 100 Egyptian patients selected by random sampling technique (males and females), diagnosed as having metabolic syndrome, and fifty apparently healthy volunteers recruited from paramedical personnel, served as controls. They showed higher levels of TGs, LDL, and cholesterol in the cases with metabolic syndrome and a lower level of HDL compared to the controls.

The circulating level of *ANGPTL8* was significantly higher in subjects with MetS (metabolic syndrome) as well as subjects with an increasing number of MetS components such as insulin resistance and central obesity. *ANGPTL8* also showed a significant association with hsCRP, BMI, TG, LDL, HOMA-IR, and FPG and was associated with a 2.4-fold increase in having MetS [23].

However, there is a shortage of studies concerning the association between gene polymorphisms and metabolic syndrome.

In the current study, regarding the distribution of the genotype of *ANGPTL8* gene variant (Rs2278426) in the two groups; there were 126 cases (84%) with wild type form and 24 (16%) with heterotype form in the metabolic syndrome group, while in the control group, wild type was detected in 82.7% and heterotype was detected in 17.3%, with no statistically significant difference between the two groups ( $p = 0.757$ ). The current results were similar to those of Alenad et al., who included a total of 905 unrelated Saudi adults (580 healthy controls and 325 MetS). The genotype and allele frequency distribution of rs737337 (T/C) and rs2278426 (C/T) polymorphisms in *ANGPTL8* gene were studied. The results showed that the genotype/allele frequency distribution of rs2278426(C/T) was comparable in both study groups ( $p > 0.05$ ) [24]. The current results disagreed with El-Lebedy, who included 272 subjects classified into 68 patients with T2DM, 68 patients with T2D + CVD, and 136 control subjects. The results showed that *ANGPTL8* c194C>T (rs2278426), the frequency of the variant allele (T), was significantly higher in T2DM patients than in control subjects (16% vs. 7.4%,  $p = 0.001$ ) and was associ-

ated with a 2.33-times increased risk of T2DM (95% CI 1.33–4.08) [25].

This also contradicts those of Guo et al., who included a total of 217 T2D patients and 201 healthy control subjects with normal glucose tolerance (NGT) in their study. The authors showed that the frequency of the CT + TT genotype in the T2D group (45.6%) was significantly higher than in the NGT group (34.8%). Relative-risk analysis showed that the CT + TT genotype was a risk factor for T2DM (OR 1.57, 95% CI 1.05–2.32) [26].

Also, the current study disagreed with those of Huo et al., who included 1,460 participants in this case-control study: healthy controls,  $n = 524$ ; pre-DM,  $n = 460$ ; and T2D,  $n = 460$ . The results showed that in T2D patients, the frequency of the CT (48%) and TT (15%) genotypes was higher than that in the control group (CT: 40.1% and TT: 6.8%). After adjusting for age, sex, and BMI, the ORs were 1.384 for the CT genotype (95% CI = 1.013–1.890) and 2.530 for the TT genotype (95% CI = 1.476–4.334) [27].

Other studies reached the same conclusion opposite to ours. A study found that, in the Iranian population, the frequency of the CT gene at this locus was significantly higher in patients with T2D than in the control group, suggesting that this genotype increased the risk of T2D [28]. Additionally, a similar study of the rs2278426 polymorphism in the Japanese population found that a higher proportion of patients with T2D and impaired glucose tolerance carried the CT or TT genotypes compared with the CC genotype [29].

The variations may be mostly due to ethnic differences, different grouping indicators, and variations in MAF (minor allele frequency).

In the current study, there was no statistically significant difference between the wild type and heterotype genotypes in all the tested clinical and laboratory parameters except for HDL-C. The mean level of HDL-C in the hetero (C/T) genotype was  $46.13 \pm 5.45$  mg/dL, which was statistically significantly higher compared to the wild type genotype ( $43.17 \pm 5.33$  mg/dL) ( $p = 0.014$ ).

The results of the association between the genotypes and the diabetic and lipid profiles were contradictory.

Ethnic variation has been associated with variable minor allele frequency (MAF) rates for this variant with respect to LDL-C and HDL-C plasma levels. In the Dallas Heart Study (DHS), a multiethnic population-based study of Dallas County, Hispanics had the highest MAF of 26%, followed by African-Americans (18%), and the least was reported in European-Americans (5%). *ANGPTL8* rs2278426 was associated with lower LDL-C and HDL-C in African-Americans and Hispanics but

not in European-Americans, and was not associated with fasting plasma glucose or homeostatic model assessment-insulin resistance in any ethnic group [9]. However, in a genome-wide association study that predominantly included individuals of European ancestry, this variant was associated with both HDL-C and LDL-C [30]. Meanwhile, in a study on non-diabetic Arabs, individuals who carried this variant had higher FPG, but did not show significant differences in their LDL and HDL levels compared to the wild type carriers [23].

In addition, Hanson et al. revealed that the T allele of the rs2278426 (C/T) variant is associated with decreased levels of total cholesterol and HDL-C in Pima Indians and Mexican Americans [31].

Furthermore, Ghasemi et al. demonstrated that the CT heterozygote genotype of the rs2278426 (C/T) polymorphism had higher levels of FPG, HbA1c, insulin, and insulin resistance compared to the CC and TT genotypes. In other words, CT genotype is associated with the risk factors of T2D. On the other hand, there was no significant relationship between this variant and HDL-C or LDL-C [28].

There is a disparity between reported results about *ANGPTL8* rs2278426 (C/T) polymorphism and biochemical factors, which is mainly due to the differences in the races of the studied populations. In this regard, Guo et al. showed that differences in race can affect the allelic and genotypic frequency of *ANGPTL8* rs2278426 (C/T) and also their relationship with other biochemical parameters [32].

Regarding the significant association between the heterotype genotype and higher HDL-C levels, previous studies have shown that *ANGPTL3* inhibits lipoprotein lipase (LPL) activity [33, 34]. In addition, another study showed that inactivation of *ANGPTL3* causes a significant reduction in HDL-C levels [35]. Therefore, it can be concluded that *ANGPTL3* activity is necessary for HDL-C biogenesis. Quagliarini et al. demonstrated that *ANGPTL3* activity required its interaction with *ANGPTL8*. Therefore, it is possible that any factor interfering with this interaction can decrease HDL-C levels [9].

This study had the limitation that it was conducted in a single center, "at the Diabetes and Metabolism Clinic at the Internal Medicine Department, Alexandria University Hospitals", and therefore, generalization is limited.

## Conclusions

Metabolic syndrome is a common health problem that affects Egyptian society. There was no statistically significant difference in the genotype distribution of *ANGPTL8* gene variant (Rs2278426) genotypes between the metabolic syndrome and control groups. The ho-

mozygous recessive genotype for the T allele (TT) was not detected in both tested groups.

## Funding

This study is self-funded.

## Conflict of interest

None declared.

## REFERENCES

- Lind L, Sundström J, Ärnlöv J, et al. A longitudinal study over 40 years to study the metabolic syndrome as a risk factor for cardiovascular diseases. *Sci Rep.* 2021; 11(1): 2978, doi: [10.1038/s41598-021-82398-8](https://doi.org/10.1038/s41598-021-82398-8), indexed in Pubmed: [33536506](https://pubmed.ncbi.nlm.nih.gov/33536506/).
- Abd Elaziz KM, Gabal MS, Aldafrawy OA, et al. Prevalence of metabolic syndrome and cardiovascular risk factors among voluntary screened middle-aged and elderly Egyptians. *J Public Health (Oxf).* 2015; 37(4): 612–617, doi: [10.1093/pubmed/ufdu097](https://doi.org/10.1093/pubmed/ufdu097), indexed in Pubmed: [25503580](https://pubmed.ncbi.nlm.nih.gov/25503580/).
- Ren G, Kim JY, Smas CM. Identification of RIFL, a novel adipocyte-enriched insulin target gene with a role in lipid metabolism. *Am J Physiol Endocrinol Metab.* 2012; 303(3): E334–E351, doi: [10.1152/ajpendo.00084.2012](https://doi.org/10.1152/ajpendo.00084.2012), indexed in Pubmed: [22569073](https://pubmed.ncbi.nlm.nih.gov/22569073/).
- Kugelberg E. Diabetes: Betatrophin--inducing  $\beta$ -cell expansion to treat diabetes mellitus? *Nat Rev Endocrinol.* 2013; 9(7): 379, doi: [10.1038/nrendo.2013.98](https://doi.org/10.1038/nrendo.2013.98), indexed in Pubmed: [23648871](https://pubmed.ncbi.nlm.nih.gov/23648871/).
- Lickert H. Betatrophin fuels  $\beta$  cell proliferation: first step toward regenerative therapy? *Cell Metab.* 2013; 18(1): 5–6, doi: [10.1016/j.cmet.2013.06.006](https://doi.org/10.1016/j.cmet.2013.06.006), indexed in Pubmed: [23823472](https://pubmed.ncbi.nlm.nih.gov/23823472/).
- Zhang R. Lipasin, a novel nutritionally-regulated liver-enriched factor that regulates serum triglyceride levels. *Biochem Biophys Res Commun.* 2012; 424(4): 786–792, doi: [10.1016/j.bbrc.2012.07.038](https://doi.org/10.1016/j.bbrc.2012.07.038), indexed in Pubmed: [22809513](https://pubmed.ncbi.nlm.nih.gov/22809513/).
- Zhang R. The *ANGPTL3*-4-8 model, a molecular mechanism for triglyceride trafficking. *Open Biol.* 2016; 6(4): 150272, doi: [10.1098/rsob.150272](https://doi.org/10.1098/rsob.150272), indexed in Pubmed: [27053679](https://pubmed.ncbi.nlm.nih.gov/27053679/).
- Fu Z, Abou-Samra AB, Zhang R. A lipasin/Angptl8 monoclonal antibody lowers mouse serum triglycerides involving increased postprandial activity of the cardiac lipoprotein lipase. *Sci Rep.* 2015; 5: 18502, doi: [10.1038/srep18502](https://doi.org/10.1038/srep18502), indexed in Pubmed: [26687026](https://pubmed.ncbi.nlm.nih.gov/26687026/).
- Quagliarini F, Wang Y, Kozlitina J, et al. Atypical angiopoietin-like protein that regulates *ANGPTL3*. *Proc Natl Acad Sci U S A.* 2012; 109(48): 19751–19756, doi: [10.1073/pnas.1217552109](https://doi.org/10.1073/pnas.1217552109), indexed in Pubmed: [23150577](https://pubmed.ncbi.nlm.nih.gov/23150577/).
- World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA.* 2013; 310(20): 2191–2194, doi: [10.1001/jama.2013.281053](https://doi.org/10.1001/jama.2013.281053), indexed in Pubmed: [24141714](https://pubmed.ncbi.nlm.nih.gov/24141714/).
- Assaad-Khalil SH, Mikhail MM, Aati TA, et al. Optimal waist circumference cutoff points for the determination of abdominal obesity and detection of cardiovascular risk factors among adult Egyptian population. *Indian J Endocrinol Metab.* 2015; 19(6): 804–810, doi: [10.4103/2230-8210.167556](https://doi.org/10.4103/2230-8210.167556), indexed in Pubmed: [26693432](https://pubmed.ncbi.nlm.nih.gov/26693432/).
- Hergenroeder AL, Wert DM, Hile ES, et al. Association of body mass index with self-report and performance-based measures of balance and mobility. *Phys Ther.* 2011; 91(8): 1223–1234, doi: [10.2522/ptj.20100214](https://doi.org/10.2522/ptj.20100214), indexed in Pubmed: [21680770](https://pubmed.ncbi.nlm.nih.gov/21680770/).
- Lean ME, Han TS, Morrison CE. Waist circumference as a measure for indicating need for weight management. *BMJ.* 1995; 311(6998): 158–161, doi: [10.1136/bmj.311.6998.158](https://doi.org/10.1136/bmj.311.6998.158), indexed in Pubmed: [7613427](https://pubmed.ncbi.nlm.nih.gov/7613427/).
- Matthews DR, Hosker JP, Rudenski AS, et al. Homeostasis model assessment: insulin resistance and beta-cell function from fasting

- plasma glucose and insulin concentrations in man. *Diabetologia*. 1985; 28(7): 412–419, doi: [10.1007/BF00280883](https://doi.org/10.1007/BF00280883), indexed in Pubmed: [3899825](https://pubmed.ncbi.nlm.nih.gov/3899825/).
15. Sliem HA, Ahmed S, Nemr N, et al. Metabolic syndrome in the Middle East. *Indian J Endocrinol Metab*. 2012; 16(1): 67–71, doi: [10.4103/2230-8210.91193](https://doi.org/10.4103/2230-8210.91193), indexed in Pubmed: [22276254](https://pubmed.ncbi.nlm.nih.gov/22276254/).
  16. Nasr G, Sliem H, Gamal A, et al. Screening for diabetes and cardiovascular risk factors among Egyptian population. *Clin Diabetes*. 2010; 9: 127–35.
  17. Hu G, Lindström J, Jousilahti P, et al. The increasing prevalence of metabolic syndrome among Finnish men and women over a decade. *J Clin Endocrinol Metab*. 2008; 93(3): 832–836, doi: [10.1210/jc.2007-1883](https://doi.org/10.1210/jc.2007-1883), indexed in Pubmed: [18073296](https://pubmed.ncbi.nlm.nih.gov/18073296/).
  18. Gierach M, Gierach J, Ewertowska M, et al. Correlation between Body Mass Index and Waist Circumference in Patients with Metabolic Syndrome. *ISRN Endocrinol*. 2014; 2014: 514589, doi: [10.1155/2014/514589](https://doi.org/10.1155/2014/514589), indexed in Pubmed: [24729884](https://pubmed.ncbi.nlm.nih.gov/24729884/).
  19. Gharipour M, Sadeghi M, Dianatkhah M, et al. Predictors of metabolic syndrome in the Iranian population: waist circumference, body mass index, or waist to hip ratio? *Cholesterol*. 2013; 2013: 198384, doi: [10.1155/2013/198384](https://doi.org/10.1155/2013/198384), indexed in Pubmed: [23634297](https://pubmed.ncbi.nlm.nih.gov/23634297/).
  20. Brenner DR, Tepylo K, Eny KM, et al. Comparison of body mass index and waist circumference as predictors of cardiometabolic health in a population of young Canadian adults. *Diabetol Metab Syndr*. 2010; 2(1): 28, doi: [10.1186/1758-5996-2-28](https://doi.org/10.1186/1758-5996-2-28), indexed in Pubmed: [20459858](https://pubmed.ncbi.nlm.nih.gov/20459858/).
  21. Mohsin A, Zafar J, Nisar YB, et al. Frequency of the metabolic syndrome in adult type 2 diabetics presenting to Pakistan Institute of Medical Sciences. *J Pak Med Assoc*. 2007; 57(5): 235–239, indexed in Pubmed: [17571478](https://pubmed.ncbi.nlm.nih.gov/17571478/).
  22. Refeat MM, Hassan NAM, Ahmad IH, et al. Correlation of circulating miRNA-33a and miRNA-122 with lipid metabolism among Egyptian patients with metabolic syndrome. *J Genet Eng Biotechnol*. 2021; 19(1): 147, doi: [10.1186/s43141-021-00246-8](https://doi.org/10.1186/s43141-021-00246-8), indexed in Pubmed: [34611771](https://pubmed.ncbi.nlm.nih.gov/34611771/).
  23. Abu-Farha M, Abubaker J, Al-Khairi I, et al. Circulating angiopoietin-like protein 8 (betatrophin) association with HsCRP and metabolic syndrome. *Cardiovasc Diabetol*. 2016; 15: 25, doi: [10.1186/s12933-016-0346-0](https://doi.org/10.1186/s12933-016-0346-0), indexed in Pubmed: [26850725](https://pubmed.ncbi.nlm.nih.gov/26850725/).
  24. Alenad A, Alenezi MM, Alokail MS, et al. Association of ANGPTL8 (Betatrophin) Gene Variants with Components of Metabolic Syndrome in Arab Adults. *Sci Rep*. 2020; 10(1): 6764, doi: [10.1038/s41598-020-63850-7](https://doi.org/10.1038/s41598-020-63850-7), indexed in Pubmed: [32317770](https://pubmed.ncbi.nlm.nih.gov/32317770/).
  25. El-Lebedy D. Interaction between endothelial nitric oxide synthase rs1799983, cholesteryl ester-transfer protein rs708272 and angiopoietin-like protein 8 rs2278426 gene variants highly elevates the risk of type 2 diabetes mellitus and cardiovascular disease. *Cardiovasc Diabetol*. 2018; 17(1): 97, doi: [10.1186/s12933-018-0742-8](https://doi.org/10.1186/s12933-018-0742-8), indexed in Pubmed: [29973202](https://pubmed.ncbi.nlm.nih.gov/29973202/).
  26. Guo C, Zhao Li, Ding Yi, et al. Gene Polymorphism rs2278426 Is Related to Carotid Intima-Media Thickness in T2DM. *Diabetes Metab Syndr Obes*. 2020; 13: 4519–4528, doi: [10.2147/DMSO.S274759](https://doi.org/10.2147/DMSO.S274759), indexed in Pubmed: [33244249](https://pubmed.ncbi.nlm.nih.gov/33244249/).
  27. Hou G, Tang Y, Ren L, et al. The 8 rs2278426 (C/T) Polymorphism Is Associated with Prediabetes and Type 2 Diabetes in a Han Chinese Population in Hebei Province. *Int J Endocrinol*. 2020; 2020: 1621239, doi: [10.1155/2020/1621239](https://doi.org/10.1155/2020/1621239), indexed in Pubmed: [33343659](https://pubmed.ncbi.nlm.nih.gov/33343659/).
  28. Ghasemi H, Karimi J, Khodadadi I, et al. Association between rs2278426 (C/T) and rs892066 (C/G) variants of ANGPTL8 (betatrophin) and susceptibility to type2 diabetes mellitus. *J Clin Lab Anal*. 2019; 33(1): e22649, doi: [10.1002/jcla.22649](https://doi.org/10.1002/jcla.22649), indexed in Pubmed: [30191588](https://pubmed.ncbi.nlm.nih.gov/30191588/).
  29. Liu J, Yagi K, Nohara A, et al. High frequency of type 2 diabetes and impaired glucose tolerance in Japanese subjects with the angiopoietin-like protein 8 R59W variant. *J Clin Lipidol*. 2018; 12(2): 331–337, doi: [10.1016/j.jacl.2017.12.011](https://doi.org/10.1016/j.jacl.2017.12.011), indexed in Pubmed: [29397342](https://pubmed.ncbi.nlm.nih.gov/29397342/).
  30. Teslovich TM, Musunuru K, Smith AV, et al. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature*. 2010; 466(7307): 707–713, doi: [10.1038/nature09270](https://doi.org/10.1038/nature09270), indexed in Pubmed: [20686565](https://pubmed.ncbi.nlm.nih.gov/20686565/).
  31. Hanson RL, Leti F, Tsinajinnie D, et al. The Arg59Trp variant in AN-GPTL8 (betatrophin) is associated with total and HDL-cholesterol in American Indians and Mexican Americans and differentially affects cleavage of ANGPTL3. *Mol Genet Metab*. 2016; 118(2): 128–137, doi: [10.1016/j.ymgme.2016.04.007](https://doi.org/10.1016/j.ymgme.2016.04.007), indexed in Pubmed: [27117576](https://pubmed.ncbi.nlm.nih.gov/27117576/).
  32. Guo T, Yin RX, Wu J, et al. Association of the angiopoietin-like protein 8 rs2278426 polymorphism and several environmental factors with serum lipid levels. *Mol Med Rep*. 2015; 12(3): 3285–3296, doi: [10.3892/mmr.2015.3825](https://doi.org/10.3892/mmr.2015.3825), indexed in Pubmed: [26004022](https://pubmed.ncbi.nlm.nih.gov/26004022/).
  33. Lee EC, Desai U, Gololobov G, et al. Identification of a new functional domain in angiopoietin-like 3 (ANGPTL3) and angiopoietin-like 4 (ANGPTL4) involved in binding and inhibition of lipoprotein lipase (LPL). *J Biol Chem*. 2009; 284(20): 13735–13745, doi: [10.1074/jbc.M807899200](https://doi.org/10.1074/jbc.M807899200), indexed in Pubmed: [19318355](https://pubmed.ncbi.nlm.nih.gov/19318355/).
  34. Shimizugawa T, Ono M, Shimamura M, et al. ANGPTL3 decreases very low density lipoprotein triglyceride clearance by inhibition of lipoprotein lipase. *J Biol Chem*. 2002; 277(37): 33742–33748, doi: [10.1074/jbc.M203215200](https://doi.org/10.1074/jbc.M203215200), indexed in Pubmed: [12097324](https://pubmed.ncbi.nlm.nih.gov/12097324/).
  35. Shimamura M, Matsuda M, Yasumo H, et al. Angiopoietin-like protein3 regulates plasma HDL cholesterol through suppression of endothelial lipase. *Arterioscler Thromb Vasc Biol*. 2007; 27(2): 366–372, doi: [10.1161/01.ATV.0000252827.51626.89](https://doi.org/10.1161/01.ATV.0000252827.51626.89), indexed in Pubmed: [17110602](https://pubmed.ncbi.nlm.nih.gov/17110602/).