



Physiological maturity and its influence on seed yield and quality in cucumber (*Cucumis sativus*) under different growing environments

GIRISH KADDI¹, B S TOMAR², BALRAJ SINGH³ and S S PARIHAR⁴

Indian Agricultural Research Institute, New Delhi 110 012

Received: 16 December 2014; Accepted: 11 February 2015

ABSTRACT

To study the physical and physiological changes occurred during maturation in fruit and seeds of cucumber (*Cucumis sativus* L.) (cv. Pant Shankar Khira-1) grown under naturally ventilated polyhouse, insect proof net house and open field conditions at Centre for Protected Cultivation Technology (CPCT) farm, Indian Agricultural Research Institute (IARI), New Delhi during summer and *kharif* seasons 2011. The fruits were harvested at 25, 30, 35, 40 and 45 days from anthesis (DFA) and seeds were extracted following fermentation. Various physical parameters, viz. fruit weight, fruit length, fruit width, number of filled seeds, number of unfilled seeds, 100 seed weight, seed yield/fruit, seed fresh weight, seed dry weight were recorded. The physiological maturity was evaluated by germinating seeds before and after drying. There were only small, non-significant difference in germinability among seed harvested at 35, 40 and 45 DFA, all of which were of higher quality than seeds which had been harvested at 25 and 30 DFA. The analysis of data showed that the seeds attained physiological maturity at 35 DFA under naturally ventilated poly house and insect proof net house, whereas at 40 DFA under open field condition.

Key words: Cucumber, Germination, Insect proof net house, Naturally ventilated polyhouse, Pant Shankar Khira-1, Physiological maturity

The low productivity of the cucumber (*Cucumis sativus* L.) crop in India is due to lesser use of quality seed, high incidence of pests and diseases and low availability of high yielding hybrid/varieties. The production of quality seeds depend upon optimal harvest time and seed extraction technique in fleshy fruit. The physiological maturity of seed and their harvest time are closely related, promoting the preservation of the physiological quality of seed. This step is variable between species and even among the species, and that is not always easy to detect (Anon. 2011). Since cucumber is a fleshy fruit that can serve as a morphological marker of the point of harvest due to changes in colour of fruit over time. However, maturity of fruit could be identified by change in colour but variable responses of cultivar towards the colour of the fruits it has become difficult to estimate the optimal timing of seed maturation. The information on the physiological maturity of seeds of cucumber crop grown under protected growing conditions is not available. Thus, the present investigation was planned to standardize the optimum time of harvest of fruit for quality seed under

different growing environments, viz. naturally ventilated poly house, insect proof net house and open condition.

MATERIALS AND METHODS

The field experiment was conducted during summer 2011 and *kharif* 2011 at Centre for Protected Cultivation Technology (CPCT) and laboratory observations were carried out at Division of Seed Science and Technology, Indian Agricultural Research Institute (IARI), New Delhi. The experiment was laid out under three growing conditions, viz. naturally ventilated poly house, insect proof net house and open field.

The seedlings of the parental lines of cucumber cv. Pant Shankar Khira 1, was raised in plug tray with soil less medium in a modern nursery at CPCT farm. The 20 days old seedlings were transplanted at a spacing of 1.00 m × 0.60 m, with a ratio 3:1 of (female: male) following block method. The cross flowers were tagged, in seed parent block aiming to get fruits at different stages of maturity. Ten cross fruits of each age group 25, 30, 35, 40 and 45 days after anthesis (DFA) were selected from different growing environments. The fruit characters, viz. average weight (g), length (cm) and width (cm) were recorded on ten randomly selected fruits in each age group. The seeds were manually extracted and subjected to fermentation for one day at room temperature in order to eliminate the mucilage surrounding following thorough washing with water. Number of seeds/

¹ Ph D scholar (e mail: girish.iari@gmail.com), ² Principal Scientist (e mail: bst_spu_iari@rediffmail.com), Seed Production Unit, IARI, New Delhi; ³ Director, NRCSS, Ajmer; ⁴ Ex Head and Professor, Division of Seed Science and Technology, IARI, New Delhi.

fruit, number of filled seeds/fruit, number of unfilled seeds/fruit, 100 seed weight (g), seed fresh weight and dry weight and seed yield/fruit (g) were recorded. The high constant temperature oven dry method at 130° C for one hour using 50 seeds of three replications was followed to estimate the seed moisture content (Anon 2011). The seed germination test was conducted as per ISTA procedure by adopting Between Paper (BP) method on freshly harvested 100 seeds in four replications. The first count on 4th day and final count on 8th day were made. To study the desiccation tolerance, seeds were targeted to 10 % moisture content using silica gel following procedure of Sun (2002). Four replications of 50 seeds were dried by silica gel and once they reached the targeted moisture content, they were sealed in Plastic containers with parafilm and seed germination was conducted following ISTA procedure (Anon 2011).

The topographical tetrazolium chloride test (TZ test) was conducted according to ISTA procedure (Anon 2011). Seeds were moistened for 18 hr and seeds were cut longitudinally to remove the seed coat and exposing the

embryo. The excise embryos were soaked in TZ solution for 6 hr at 30°C. The analysis of the viable and non-viable seeds is done. The 100 % colored embryos are viable and 50% and/ or less coloured embryos as non-viable were categorized.

RESULTS AND DISCUSSION

Fruit characters

Analysis of variance showed that there were significant difference in fruit weight, fruit length and fruit width under different growing environments at different stages of harvest in both summer and *kharif* 2011, presented in Table 1. The fruit weight, length and width showed increasing trend from 25 DFA to 45 DFA, whereas non-significant difference was recorded between 35 DFA and 45 DFA. The highest fruit weight, length and width (735 g, 21.83 cm and 8.23 cm respectively) at 40 DFA during *kharif* 2011, whereas highest fruit weight, length and width (641 g, 21.66 cm and 8.23 cm respectively) at 45 DFA during summer were recorded under insect proof net house. It

Table 1 Effect of time of harvest and growing conditions on fruit attributes of cucumber (Pant Shankar Khira 1)

Days from anthesis	Summer 2011				<i>Kharif</i> 2011			
	Open	Net house	Polyhouse	Mean	Open	Net house	Polyhouse	Mean
<i>Fruit weight (g)</i>								
25 DFA	290.00	415.00	368.33	357.78 _c	312.66 _g	503.66 _e	495.00 _e	437.11 _c
30 DFA	380.00	595.00	471.66	482.22 _b	446.66 _f	720.00 _{ab}	645.00 _d	603.88 _b
35 DFA	483.33	616.66	518.33	539.44 _a	514.33 _e	732.33 _a	688.33 _c	645.00 _a
40 DFA	576.66	633.33	526.66	578.89 _a	515.33 _e	735.00 _a	687.00 _c	645.44 _a
45 DFA	533.33	641.66	535.00	570.00 _a	511.66 _e	729.00 _a	696.66 _{bc}	645.77 _a
Mean	452.67 _b	580.33 _a	484.00 _b		459.93 _c	684.00 _a	642.40 _b	
	Days	Condition	Interaction		Days	Condition	Interaction	
CD (P=0.05)	51.08	39.57	NS		14.15	10.96	24.49	
<i>Fruit length (cm)</i>								
25 DFA	16.00	16.86	16.66	16.51 _d	15.66 _f	20.33 _d	20.33 _d	18.77 _c
30 DFA	18.33	19.33	19.66	19.11 _c	19.33 _e	21.00 _a	20.66 _{cd}	20.33 _b
35 DFA	20.16	20.83	19.83	20.27 _b	20.66 _{cd}	21.83 _a	21.33 _{abc}	21.27 _a
40 DFA	21.33	21.50	20.33	21.05 _a	21.00 _{bcd}	21.83 _a	22.00 _a	21.61 _a
45 DFA	21.50	21.66	20.33	21.16 _a	21.00 _{bcd}	21.66 _a	22.00 _a	21.55 _a
Mean	19.46 _b	20.04 _a	19.36 _b		19.53 _b	21.53 _a	21.26 _a	
	Days	Condition	Interaction		Days	Condition	Interaction	
CD (P=0.05)	0.62	0.48	NS		0.39	0.30	0.69	
<i>Fruit width (cm)</i>								
25 DFA	5.26 _g	6.93 _e	7.03 _e	6.41 _d	5.30 _f	6.13 _e	6.26 _e	5.90 _c
30 DFA	5.80 _f	7.63 _d	7.90 _{cd}	7.11 _c	5.40 _f	8.06 _a	7.16 _c	6.87 _b
35 DFA	7.06 _e	8.33 _{ab}	8.30 _{abc}	7.90 _b	6.30 _e	8.13 _a	7.63 _b	7.38 _a
40 DFA	7.90 _{cd}	8.60 _a	8.30 _{abc}	8.26 _a	6.63 _d	8.23 _a	7.73 _b	7.53 _a
45 DFA	8.13 _{bc}	8.70 _a	8.33 _{ab}	8.38 _a	6.66 _d	8.13 _a	7.76 _b	7.51 _a
Mean	6.83 _b	8.04 _a	7.97 _a		6.06 _c	7.74 _a	7.31 _b	
	Days	Condition	Interaction		Days	Condition	Interaction	
CD (P=0.05)	0.24	0.18	0.41		0.15	0.12	0.26	

Figures sub scripted in interaction with days and conditions or subscripted in main effects by same alphabet are at par at 0.05 level of significance

was evident from the results that harvesting of fruits at 40 DFA resulted in higher fruit development attributes compared to other harvesting stages owing to accumulation of food reserves with the increased period of harvest. The results were in agreement with Barbedo *et al.* (1993) in cucumber cv. Nazare. Similarly, decreased fruit weight in early harvested fruits was also reported by Sureshbabu *et al.* (2003) in brinjal and Hamsaveni *et al.* (2003) in tomato.

Seed yield characters

The seed yield parameters, viz. number of seeds/fruit, number of filled or unfilled seeds, seed yield/fruit and 100 seed weight were significantly higher under naturally ventilated polyhouse and insect proof net house compared to open condition. The similar results were obtained during hybrid seed production by Flemine (2010) in pumpkin, Gograj (2011) in bitter gourd and Kunj Bihari (2012) in summer squash. The number of seeds/fruit could not recorded significant increase between 35 DFA (173.11-summer and 184.44-*kharif*) to 45 DFA (174.11-summer and 190.33.44- *kharif*), but during same period the number of

filled seeds increased significantly (Table 2). The 100 seed weight at 25 DFA (1.61g -summer and 1.93g -*kharif*) was significantly increased and reached maximum at 45 DFA (3.02g -summer and 3.07g -*kharif*). The 100 seed weight, seed yield/fruit were highest at 40 DFA (3.02 g, 5.18 g summer and 3.07 g, 5.67 g, -*kharif*, respectively). Harvesting fruits at 40 DFA established complete development of fruits and seeds on account of continuous supply and accumulation of metabolites (food reserves) from mother plant into fruits and seeds. These results are in conformity with the reports of Biradar (1994) in brinjal, Chaudhari *et al.* (1992) in tomato, Nandeesh *et al.* (1995) in cucumber cv. Japanese Green Long.

Seed quality characters

The seed fresh weight, seed dry weight and moisture content showed that the physiological status of seeds at each maturation stage shown in Fig 1. The seed fresh weight per 50 seeds was highest at 25 DFA (3.79g -summer and 4.34g -*kharif*) decreased significantly at each level and become constant between 40-45 DFA (2.22g -summer and

Table 2 Effect of time of harvest and growing conditions on seed parameters of cucumber (Pant Shankar Khira 1)

Days from anthesis	Summer 2011				<i>Kharif</i> 2011			
	Open	Net house	Poly house	Mean	Open	Net house	Poly house	Mean
<i>Number of seeds/fruit</i>								
25 DFA	103.66	165.00	130.33	133.00 _c	113.33 _g	171.00 _c	151.33 _e	145.22 _c
30 DFA	132.33	184.33	167.66	161.44 _b	134.66 _f	206.00 _a	185.00 _b	175.22 _b
35 DFA	152.33	190.33	176.66	173.11 _a	153.66 _{de}	202.66 _a	197.00 _a	184.44 _a
40 DFA	154.33	184.66	185.00	174.66 _a	168.33 _c	201.33 _a	202.66 _a	190.77 _a
45 DFA	150.66	186.00	185.66	174.11 _a	165.33 _c	201.66 _a	204.00 _a	190.33 _a
Mean	138.66 _c	182.06 _a	169.06 _b		147.06 _c	196.53 _a	188.00 _b	
	Days	Condition	Interaction		Days	Condition	Interaction	
CD (P=0.05)	9.08	7.03	NS		6.92	5.36	11.97	
<i>Number of filled seeds/fruit</i>								
25 DFA	45.00 _h	86.33 _g	71.66	67.66 _d	45.33 _e	85.66 _d	81.00 _d	70.66 _d
30 DFA	71.66 _g	149.33 _{cd}	133.00 _e	118.00 _c	80.00 _d	162.66 _b	157.00 _b	133.22 _c
35 DFA	113.00 _f	179.33 _a	163.00 _{bc}	151.77 _b	131.00 _c	199.33 _a	195.00 _a	175.11 _b
40 DFA	147.00 _{de}	178.00 _{ab}	180.00 _a	168.33 _a	164.33 _b	199.33 _a	200.66 _a	188.11 _a
45 DFA	147.33 _{de}	179.33 _a	182.00 _a	169.55 _a	163.33 _b	199.66 _a	202.33 _a	188.44 _a
Mean	104.80 _c	154.46 _a	145.93 _b		116.80 _b	169.33 _a	167.20 _a	
	Days	Condition	Interaction		Days	Condition	Interaction	
CD (P=0.05)	8.93	6.9	15.46		6.93	5.37	11.99	
<i>100 seed weight (g)</i>								
25 DAA	1.38 _f	1.76 _e	1.68 _e	1.61 _d	1.39 _g	2.23 _f	2.18 _f	1.93 _c
30 DAA	1.74 _e	2.82 _c	2.70 _d	2.42 _c	2.69 _e	2.92 _{bc}	2.82 _d	2.81 _b
35 DAA	2.67 _d	3.10 _b	3.09 _b	2.95 _b	2.91 _c	3.11 _a	3.08 _a	3.03 _a
40 DAA	2.69 _d	3.12 _{ab}	3.19 _{ab}	3.00 _{ab}	2.97 _{bc}	3.11 _a	3.11 _a	3.06 _a
45 DAA	2.69 _d	3.18 _{ab}	3.20 _a	3.02 _a	2.98 _b	3.11 _a	3.11 _a	3.07 _a
Mean	2.24 _b	2.80 _a	2.77 _a		2.58 _b	2.89 _a	2.86 _a	
	Days	Condition	Interaction		Days	Condition	Interaction	
CD (P=0.05)	0.05	0.04	0.09		0.04	0.03	0.05	

Figures sub scripted in interaction with days and conditions or subscripted in main effects by same alphabet are at par at 0.05 level of significance




Fig 1 Effect of time of harvest and growing conditions seed development of cucumber (Pant Shankar Khira 1) during 2011

2.24g -kharif). The seed dry weight of 50 seeds increased significantly from 25 DFA (0.79g -summer and 0.98g -kharif) to 45 DFA (1.52g -summer and 1.56g -kharif). Seed water content (fresh weight basis) declined from 77% to 30% between 25 DFA and 40 DFA (when maximum dry weight was attained). The results are in corroboration with Marcos (2005) in cucurbits and Pamela *et al.* (2011) in cucumber.

Germination and viability

The significant increase in germination (%) was recorded at each maturation stage. Onset of germination occurred after 25 DFA and reached to 70 % at 35 DFA under naturally ventilated poly house and insect proof net house however, to attain 70% germination in open condition it took 40 DFA (Fig 2). Sanhewe and Ellis (1996) stated that immature bean seeds had acquired the ability to germinate




Fig 2 Effect of time of harvest and growing conditions on onset of germination and viability in cucumber (Pant Shankar Khira 1) during 2011

well before mass maturity. The view that water loss (i.e. desiccation) alone or in combination with time from ovule abscission enables developing seeds to switch from a developmental mode to a mode in which germination is possible (Misra and Bewley 1985, Bewley *et al.* 1989 and Jiang and Kennode 1994). The results of tetrazolium test revealed that the viability at 25 DFA was 32 % in summer and 40 % in kharif, but it increased rapidly between 25-35 DFA and reached to 86 % in both the seasons. The results are in conformity with Nerson H (2002) in water melon, Maria *et al.* (2010) in gherkin and Pervej *et al.* (2010) in tomato.

The physical and physiological analysis inferred that the superior physiological quality of seeds of hybrid cucumber Pant Shankar Khira 1 could be obtained at 35 DFA under naturally ventilated polyhouse and insect proof net house and at 40 DFA under open field condition during both summer and kharif season.

REFERENCES

Anonymous. 2011. International Rules for Seed Testing. Seed Science and Technology, Zurich, Switzerland.

- Barbedo C J, Coelho A S, Zanin A C W and Nakagawa J. 1993. Influence of age of cucumber fruits on seed quality. *Hort. Brasileira* **11**: 18–21.
- Barbedo, Claudio J B, Adelianna S C, Nakagawa and Sato O. 1999. Effect of fruit age and post-harvest period of cucumber on stored seeds. *Pesquisa Agropecuária*: 66–74.
- Bewley J D, Kermode A R and Mbra S. 1989. Desiccation and minimal drying treatments of seeds of castor bean and *Phaseolus vulgaris* which terminate development and promote germination cause changes in protein and messenger RNA synthesis. *Annals of Botany* **63**: 3–17.
- Biradar G S. 1994. Studies on physiological maturity and harvesting .of green fruits on seed yield and quality of brinjal (*Solanum melongena* L.). M Sc (Agri.) thesis, UAS, Dharwad, Karnataka.
- Chaudhari R V, Meshram L D and Kukade B K. 1992. Relationship between maturity and seed quality in tomato. *Agricultural Science Digest* **12**: 38–40.
- Flemine X. 2010. Studies on hybrid seed production in pumpkin under insect proof net house and open field conditions. M Sc thesis, IARI, New Delhi.
- Hamsaveni M R, Kurdikeri M B, Shekargouda M, Shashidhara S D and Dharmatti P R. 2003. Influence of harvesting stages and post harvest ripening periods on seed quality in tomato (cv. Megha). *Karnataka Journal of Agricultural Science* **16(4)**: 597–9.
- Jat Gograj Singh. 2011. Studies on hybrid seed production in pumpkin under insect proof net house and open field conditions. M Sc thesis, IARI, New Delhi.
- Jiang L and Kennode A R. 1994. Role of desiccation in the termination of expression of genes for storage proteins. *Seed Science Research* **4**: 149–73.
- Kunj Bihari. 2012. Effect of Growing environments on hybrid seed production in summer squash cv. Pusa Alankar. M Sc thesis, IARI, New Delhi.
- Marcos Filho. 2005. Physiology of seeds of cultivated plants. *Piracicaba: Fealq* 495.
- Maria A M, Costa L G, Barros S T and Ana V D. 2010. Physiological maturity in gherkin (*Cucumis anguria* L.) seeds. *Journal of Seed* **32**: 3.
- Misra S and Bewley J D. 1985. Reprogramming of protein synthesis from a developmental to a germinative mode induced by desiccation of the axes of *Phaseolus vulgaris*. *Plant Physiology* **78**: 876–2.
- Nandeesh, Javare Gowda S and Rame Gowd G. 1995. Studies on the stage of harvest and post harvest ripening on seed quality in cucumber (*Cucumis sativus* L.). *Seed Research* **23(2)**: 113–5.
- Nerson H. 2002. Effect of seed maturity, extraction practice and storage duration on germinability in watermelon. *Sciatica Horticultural* **93**: 245–56.
- Pamela G N, John A O M Larissa C M, Luiz A G and Edila V R. 2011. Physiological and biochemical performance of cucumber seeds at different maturity stages. *Journal of Seed* **33**: 1.
- Parvej M R, Khan M A H and Awal M A. 2010. Phenological development and production potentials of tomato under polyhouse climate. *Journal of Agricultural Science* **5**:1: 19–31.
- Sanhewe A J and Ellis R H. 1996. Seed development and maturation in *Phaseolus vulgaris* I. Ability to germinate and to tolerate desiccation. *Journal of Experimental Botany* **47**: 959–65.
- Sun W Q. 2002. Methods for the study of water relations under desiccation stress. (*In*) Desiccation and Survival in Plants Drying without Dying, pp 48– 91. Black M, and Prichard H W (Eds) CABI.
- Sureshababu T, Kurdikeri M B, Shekargouda M, Shashidhara S D and Dharmatti P R. 2003. Influence of fruit maturity stages and post harvest ripening on seed yield and quality in brinjal. *Seed Research* **31(2)**: 204–8.