

## Tap water perceptions and water filter use vary with sociodemographic characteristics and influence water consumption in university students

Melissa J. Slotnick<sup>1</sup>, Julia A. Wolfson<sup>2</sup>, and Cindy W. Leung<sup>3</sup>

<sup>1</sup>Department of Nutritional Sciences, The University of Michigan School of Public Health, Ann Arbor, MI, 48108, USA

<sup>2</sup>Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, 21205, USA Department of Health Policy and Management, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, 21205, USA

<sup>3</sup>Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, 02115, USA

**Corresponding Author:** Melissa J. Slotnick, PhD, MPH, MEd, Research Associate, Department of Nutritional Sciences, The University of Michigan School of Public Health, 1415 Washington Heights, Ann Arbor, MI 48109, USA, [slotnick@umich.edu](mailto:slotnick@umich.edu), ORCID: 0000-0002-1380-2577

**Short Title:** Tap water perceptions and beverage intake



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S1368980023001659

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

## Abstract

**Objective.** The goal of this study is to evaluate university students' perceptions of tap water safety and water filter use, and determine how these perceptions and behaviors affect water and sugar sweetened beverage intake.

**Design.** Cross-sectional; online survey conducted in Fall 2021.

**Setting.** A large, public Midwestern university in the United States

**Participants.** 793 undergraduate students

**Results.** Students who experienced food insecurity, were on a Pell grant, were first generation college students or were racial/ethnic minorities were less likely to trust tap water safety. Tap water filtration behavior also varied by age and race/ethnicity. Students who did not agree with the statement “my local tap water is safe to drink” had lower odds of consuming  $\geq 3$  cups of total water per day (OR=0.45, 95% CI: 0.32, 0.62), lower odds of consuming tap water  $\geq 3$  times/day (OR=0.46, 95% CI: 0.34, 0.64), higher odds of drinking bottled water  $\geq 1$  time per day (OR=1.80, 95% CI: 1.22, 2.66), and higher odds of drinking SSBs  $\geq 1$  time per day (OR=1.47, 95% CI: 1.01, 2.14) than those who agreed. Students who always or sometimes filtered their tap water had lower odds of consuming  $\geq 3$  cups of total water per day (OR=0.59, 95% CI: 0.39, 0.90) than students who never filtered their tap water.

**Conclusions.** Tap water perceptions and behaviors affect tap and bottled water and SSB intake among university students. Tap water perceptions and behaviors in this demographic provide important context for university programming promoting healthy beverage initiatives.

**Key words:** Tap water, SSBs, water insecurity, bottled water, water filter, university students

## Introduction

Approximately 20% of the population in the United States (US) does not drink tap water regularly.<sup>(1)</sup> Lower tap water consumption in the US is associated with lower income and education levels, identifying as Black, Asian or Hispanic races, and being born outside of the US.<sup>(1, 2)</sup> Distrust in water safety in these populations<sup>(3, 4)</sup> likely contributes to these notable differences in water consumption behavior.<sup>(1)</sup>

Disparities in sugar sweetened beverage (SSB) intake, a major source of sugar intake for young adults,<sup>(5)</sup> mimic disparities in tap water consumption. SSB intake is higher in young adults from minority racial/ethnic groups and low-income populations.<sup>(5)</sup> Food insecurity among university students has also been associated with higher SSB consumption<sup>(6)</sup> and added sugar consumption from SSBs.<sup>(7)</sup> Higher water intake has been associated with lower SSB intake; however, these benefits are not observed across all socioeconomic subgroups.<sup>(8)</sup>

Water filtration could be a strategy to increase tap water consumption. Water filter use at home has been associated with higher odds of drinking tap water and lower odds of SSB consumption in US adults.<sup>(9, 10)</sup> However, information on water filter use in the US is limited,<sup>(10)</sup> and no known studies have investigated how attitudes towards water safety and water filtration use affect beverage intake among young adults or in university settings.

The university environment is a unique setting for studying dietary behaviors and perceptions among young adults. Students are often living away from home for the first time and independently making food and beverage choices. Such choices have potential to influence their future dietary patterns and purchasing decisions.<sup>(11-13)</sup>

The current study has two objectives: (1) to determine how perceptions of tap water safety and water filter use vary with sociodemographic variables in university students, and (2) to evaluate how perceptions affect water and SSB intake. It is hypothesized that distrust of tap water safety will be associated with lower water and higher SSB intake, and water filter use will be associated with higher water and lower SSB intake. Understanding how perceptions, behaviors and disparities affect beverage consumption in this population can help to shape healthy beverage initiatives in secondary institutions and lend understanding to health disparities.

## Methods

### Study Participants

In Fall 2021, an online (Qualtrics) survey was fielded to students recruited from the university registrar's office, oversampling first-generation and racial/ethnic minority students. The study purpose was "[to] improve our understanding of students' demographics and health behaviors." 3782 students were contacted and 885 students completed the survey, yielding a response rate of 23%. Students received a \$10 Amazon gift card for survey completion. Those with incomplete responses were excluded from analysis, leaving an analytic sample of 793 participants.

### Measures

**Tap Water Perceptions and Filter Use.** Tap water perceptions were assessed based on questions modified from the 2010 ConsumerStyles and Youth-Styles mail surveys.<sup>(4)</sup> Students were asked the question "my local tap water is safe to drink." For analysis, response options were grouped into two categories: agree/strongly agree and "do not agree" (strongly disagree/disagree/neutral). Water filter use was assessed with the following question: "I filter my tap water before drinking it." For analysis, response choices of "always" and "sometimes" were grouped together, and "never" was coded as "no."

**Sociodemographic covariates.** Students self-reported age, receipt of Pell grant in the current academic year (yes/no), gender (man, woman, other/non-binary), and race/ethnicity (Asian/PI, Black, Hispanic, Other/Multiracial, White/MENA). MENA was grouped with White according to US Census guidelines.<sup>(14)</sup> Food security was assessed using the 10-item U.S. Adult Food Security Survey Module, and low and very low food security were collectively termed food insecurity.<sup>(15)</sup> Federal Pell grants are usually awarded to students with exceptional financial need and are therefore an indicator of income level.

**Beverage Intake.** Beverage consumption was assessed for 15 types of beverages using the validated BEVQ-15,<sup>(16)</sup> modified as previously described.<sup>(7)</sup> Beverages assessed included SSBs (regular soda, energy/sports drinks, fruit-flavored drinks, sweetened tea, sweetened coffee, flavored milk), and other beverages (diet soda/pop/fruit drinks/energy drinks/sport drinks, 100% pure fruit juice, plain/unflavored milk, artificially sweetened or unsweetened tea and coffee, plain bottled water, plain tap water, and plain/flavored unsweetened sparkling water). Students

self-reported intake frequencies ranging from “never” to “6 or more times per day,” and the usual volume consumed each time ranging from “less than 6 fl. oz. (0.75 cup)” to “more than 20 fl. oz. (2.5 cups)” for beverages and total water (plain tap water, bottled water, and sparkling water). Ounces consumed per day were calculated for each beverage and total water. SSB intake (oz./day) was calculated by summing intake of regular soda, energy/sports drinks, fruit-flavored drinks, sweetened tea, sweetened coffee, and flavored milk. Ounces were converted to L/day for presentation of results. For the BEVQ-15, volume consumed is not recorded for tap water and bottled water, so frequency of intake (times/day) was used for analysis of tap and bottled water. For logistic regression analysis, to compare with previous research<sup>(10)</sup> and to allow for separation of regular consumers and non-consumers due to variable distribution in the current study, beverage consumption was classified into the following categories: total water intake  $\geq 3$  cups (710 ml)/day, SSB intake  $\geq 1$  serving (12 oz. or 355 ml)/day, bottled water intake  $\geq 1$  time/day, tap water intake  $\geq 3$  times/day.

### Statistical analysis

Chi-square tests were used to evaluate differences in water perceptions and behaviors by sociodemographic characteristics. Differences in mean beverage intake for each water perception/behavior were calculated using two-sided t-tests. Associations between water perceptions and behaviors and beverage intake were assessed using multivariate logistic regression. Models were adjusted for age, gender, race/ethnicity, first generation status, food insecurity status and Pell grant status. Interactions between sociodemographic variables and total water and SSB intake were evaluated. Missing values (n=17) were assigned an indicator value. All statistical tests were two-sided, performed using SAS, version 9.4,<sup>(17)</sup> and significance was considered at  $P < 0.05$ .

### Results

Younger students were less likely to agree that their tap water was safe to drink than older students ( $p < 0.001$ ), and Asian/PI, Black, and Hispanic students were less likely to agree that their tap water was safe to drink than White/MENA students ( $p < 0.001$ ) (**Table 1**). A higher percentage of students who did not agree their tap water was safe to drink were first generation (57%) versus not first generation (41%) ( $p < 0.001$ ), and on a Pell grant (56%) versus not on a Pell

grant (44%) ( $p=0.001$ ). Food insecure students were less likely to agree that their tap water is safe to drink than food secure students ( $p<0.001$ ).

Tap water filtration behavior also varied by age and race/ethnicity. Students  $<21$  years old were more likely to filter their tap water than those  $\geq 21$  ( $p=0.03$ ). A higher percentage of Asian students filtered their water (43%) than did not (23%), but a higher percentage of Black students did not filter their water (13.5%) than filtered their water (5.3%) ( $p<0.001$ ).

Mean total water intake was higher for students who agreed their tap water was safe (mean =  $54.5 \pm 40.1$  oz./day or  $1.61 \pm 1.19$  L/day), versus those who did not (mean =  $42.5 \pm 48.9$  oz./day or  $1.26 \pm 1.44$  L/day) ( $p<0.001$ ). Frequency of tap water intake demonstrated a similar pattern. Mean SSB intake was lower for students who agreed that their tap water was safe to drink (mean =  $7.7 \pm 13.2$  oz./day or  $0.23 \pm 0.39$  L/day) than for those who did not (mean =  $12.2 \pm 23.9$  oz./day or  $0.36 \pm 0.71$  L/day) ( $p=0.003$ ). Mean frequency of bottled water intake was also lower for students who agreed that their water was safe to drink (mean = 0.5 times/day, SD = 1.2) compared with those who did not (mean = 1.0 times/day, SD = 1.7) ( $p<0.001$ ).

Students who filtered their tap water had higher mean total water intake (mean =  $57.4 \pm 41.6$  oz./day or  $1.70 \pm 1.23$  L/day) than students who did not (mean =  $48.1 \pm 44.3$  oz./day or  $0.26 \pm 0.39$  L/day) ( $p=0.01$ ). Similarly, students who filtered their tap water had a higher mean frequency of tap water intake (mean = 4.4 times/day, SD = 2.1) than students who did not (mean = 4.0 times/day, SD = 2.5) ( $p=0.03$ ). Mean SSB intake and mean frequency of bottled water intake did not differ by tap water filtration.

Students who did not agree with the statement “my local tap water is safe to drink” had lower odds of consuming  $\geq 3$  cups of total water per day (OR=0.45, 95% CI: 0.32, 0.62), lower odds of consuming tap water  $\geq 3$  times/day (OR=0.46, 95% CI: 0.34, 0.64), higher odds of drinking  $\geq 1$  SSB serving per day (OR=1.47, 95% CI: 1.01, 2.14), and higher odds of drinking bottled water  $\geq 1$  time per day (OR=1.80, 95% CI: 1.22, 2.66) than those who agreed (**Table 2**). Students who filtered their tap water had lower odds of consuming  $\geq 3$  cups of total water per day (OR=0.59, 95% CI: 0.39, 0.90) than students who did not. Tests for interactions between sociodemographic variables and total water intake were not significant ( $P_s > 0.05$ ).

## Discussion

Students who were food insecure, on a Pell grant, first generation, or racial/ethnic minorities were less likely to trust tap water safety. In the US, higher tap water mistrust has been observed among people who are Hispanic, non-Hispanic Black, lower-income, and immigrants.<sup>(3, 18, 19)</sup> Furthermore, tap water avoidance has been associated with food insecurity in the US.<sup>(20)</sup> Our results confirm that similar attitudes persist in this university student/young adult population; however, it should be noted that due to sampling methods, this sample had a higher percentage of Asian (38.9 vs 19%), Black (7.0 vs 4%), Hispanic (10.3 vs 8%), Pell grant recipients (47.0 vs 19%), and first-generation students (43.9 vs 16%) than the general student body.

In a previous study, SSB warning labels on the current university campus led to a decrease in SSB intake, but no change in water intake.<sup>(7)</sup> We observe that tap water distrust is associated with lower total water and tap water intake, and thus plays a role in student beverage choices. Students distrusting tap water had higher mean SSB intake, and tap water distrust was associated with higher SSB intake in the multivariate adjusted model.

We hypothesized that tap water filtration could be a strategy to increase tap water intake, decrease bottled water intake, and/or decrease SSB intake. Previously, US adults using a water filter had a higher odds of drinking tap water and lower odds of consuming SSBs,<sup>(10)</sup> and filter use was associated with increased water and decreased SSB intake in a recent intervention,<sup>(21)</sup> yet our results do not corroborate the findings. It is possible that filtering tap water does not negate safety concerns in this population. Given a large percentage of the study population drinks filtered water (78%), other factors – such as access, palatability, or beverage availability – may be influencing results. Further research is necessary to understand beverage choice behaviors in this population.

Colleges present an opportune setting to improve beverage consumption habits in young adults, and some university initiatives to improve water access, reduce SSB intake, and promote sustainability have proven effective.<sup>(22-24)</sup> To improve campus sustainability, bottled water reduction strategies have been implemented at several universities<sup>(25-29)</sup> yet do not always result in increased tap water intake.<sup>(26, 27)</sup> The current study emphasizes the need to consider how socioeconomic disparities in tap water safety perceptions affect consumption, and to account for

such disparities in healthy beverage initiatives. For example, future studies focusing on water quality messaging, access, and perceptions driving beverage choices have potential to improve effectiveness of current initiatives.

Strengths of this study include large sample size and refined beverage intake measurements. This study has several limitations. Amounts consumed per serving were not available for tap and bottled water, resulting in different intake measures for different water sources. Tap water quality data were also unavailable. The survey did not distinguish between types of water filters or assess awareness of campus water filters. This study is limited to university students, and may not be representative of all young adults. Lastly, as students in Michigan, attitudes in this population may be disproportionately affected by the Flint Water Crisis.<sup>(30)</sup>

This study lends important insight into how water quality perceptions and behaviors affect tap water, bottled water and SSB intake in young adults and university students. The findings provide a launching point for university programming promoting healthy beverage initiatives and drinking water access in diverse student populations.

**Financial Support:** This study was supported by a grant from the University of Michigan Office of Research (MS and CL), and by the National Institutes of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health (JW Award #K01DK119166).

**Conflicts of Interest:** None

**Authorship:** The authors responsibilities were as follows – MS: data analysis, interpretation of results, writing of the manuscript, critical revisions of the manuscript; JW: review of results, critical revisions of the manuscript; CL: data collection, interpretation of results, critical revisions of the manuscript

**Ethical Standards Disclosure:** This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the University of Michigan Institutional Review Board. Digital informed consent was obtained from all subjects.



**References**

1. Rosinger AY, Patel AI Weaks F (2022) Examining recent trends in the racial disparity gap in tap water consumption: NHANES 2011–2018. *Public health nutrition* 25, 207-213.
2. Patel AI, Hecht CE, Cradock A *et al.* (2020) Drinking water in the United States: implications of water safety, access, and consumption. *Annual Review of Nutrition* 40, 345-373.
3. Park S, Onufrak S, Patel A *et al.* (2019) Perceptions of drinking water safety and their associations with plain water intake among US Hispanic adults. *Journal of Water and Health* 17, 587-596.
4. Onufrak SJ, Park S, Sharkey JR *et al.* (2014) The relationship of perceptions of tap water safety with intake of sugar- sweetened beverages and plain water among US adults. *Public Health Nutrition* 17, 179-185.
5. Han E & Powell LM (2013) Consumption patterns of sugar-sweetened beverages in the United States. *Journal of the Academy of Nutrition and Dietetics* 113, 43-53.
6. Ryan RA, Murphy B, Deierlein AL *et al.* (2022) Food Insecurity, Associated Health Behaviors, and Academic Performance Among Urban University Undergraduate Students. *Journal of Nutrition Education and Behavior* 54, 269-275.
7. Leung CW, Wolfson JA, Hsu R *et al.* (2021) Warning labels reduce sugar-sweetened beverage intake among college students. *The Journal of Nutrition* 151, 179-185.
8. Vieux F, Maillot M, Rehm CD *et al.* (2020) Opposing consumption trends for sugar-sweetened beverages and plain drinking water: analyses of NHANES 2011–16 data. *Frontiers in nutrition* 7, 233.
9. Rosinger AY, Herrick KA, Wutich AY *et al.* (2018) Disparities in plain, tap and bottled water consumption among US adults: National Health and Nutrition Examination Survey (NHANES) 2007–2014. *Public health nutrition* 21, 1455-1464.
10. Park S, Onufrak SJ, Cradock AL *et al.* Factors Related to Water Filter Use for Drinking Tap Water at Home and Its Association With Consuming Plain Water and Sugar-Sweetened Beverages Among U.S. Adults. *American Journal of Health Promotion* 0, 08901171211073304.

11. Nelson MC, Story M, Larson NI *et al.* (2008) Emerging adulthood and college-aged youth: an overlooked age for weight-related behavior change. *Obesity* 16, 2205.
12. Sawyer SM, Afifi RA, Bearinger LH *et al.* (2012) Adolescence: a foundation for future health. *The lancet* 379, 1630-1640.
13. Dunn JE, Liu K, Greenland P *et al.* (2000) Seven-year tracking of dietary factors in young adults: the CARDIA study. *American Journal of Preventive Medicine* 18, 38-45.
14. Bureau USC (2022) About the topic of race. <https://www.census.gov/topics/population/race/about.html> (accessed 06/26/2022)
15. USDA (2022) Food Security in the US Survey Tools. <https://www.ers.usda.gov/topics/food-nutrition-assistance/food-security-in-the-u-s/survey-tools/> (accessed April 14 2022)
16. Hedrick VE, Savla J, Comber DL *et al.* (2012) Development of a Brief Questionnaire to Assess Habitual Beverage Intake (BEVQ-15): Sugar-Sweetened Beverages and Total Beverage Energy Intake. *Journal of the Academy of Nutrition and Dietetics* 112, 840-849.
17. SAS (2013) Version 9.4. SAS Institute, Inc. Cary, NC.
18. Pierce G & Gonzalez S (2017) Mistrust at the tap? Factors contributing to public drinking water (mis) perception across US households. *Water Policy* 19, 1-12.
19. Onufrak SJ, Park S, Sharkey JR *et al.* (2014) Perceptions of tap water and school water fountains and association with intake of plain water and sugar-sweetened beverages. *Journal of school health* 84, 195-204.
20. Rosinger AY, Bethancourt HJ Young SL (2022) Tap Water Avoidance Is Associated with Lower Food Security in the United States: Evidence from NHANES 2005-2018. *Journal of the Academy of Nutrition and Dietetics*.
21. Reese AC, Burgos-Gil R, Cleary SD *et al.* (2023) Use of a Water Filter at Home Reduces Sugary Drink Consumption among Parents and Infants/Toddlers in a Predominantly Hispanic Community: Results from the Water Up!@ Home Intervention Trial. *Journal of the Academy of Nutrition and Dietetics* 123, 41-51.
22. Patel AI & Schmidt LA (2021) Healthy beverage initiatives in higher education: an untapped strategy for health promotion. *Public health nutrition* 24, 136-138.

23. Malan H, Amsler Challamel G, Silverstein D *et al.* (2020) Impact of a scalable, multi-campus “Foodprint” seminar on college students’ dietary intake and dietary carbon footprint. *Nutrients* 12, 2890.
24. Kalair N, Mousli LM, Jacobs LM *et al.* (2022) Work site access to fluoridated tap water and retail beverages: An assessment of the University of California, San Francisco campuses. *The Journal of the American Dental Association* 153, 201-207.
25. Cunningham J, Honeyman C, Prygoski A *et al.* (2010) Options for Reducing Bottled Water Use in the Michigan Union and on Campus: University of Michigan, Michigan, MI, USA.
26. Mikhailovich K & Fitzgerald R (2014) Community responses to the removal of bottled water on a university campus. *International Journal of Sustainability in Higher Education* 15, 330-342.
27. Berman ER & Johnson RK (2015) The unintended consequences of changes in beverage options and the removal of bottled water on a university campus. *American journal of public health* 105, 1404-1408.
28. Choate B, Davis BY Verrecchia J (2018) Campus bottled water bans, not always the solution. *International Journal of Sustainability in Higher Education*.
29. Meisterling K, Vo J, Garvey KA *et al.* (2022) Healthy beverage initiatives: A case study of scenarios for optimizing their environmental benefits on a university campus. *Cleaner and Responsible Consumption* 4, 100049.
30. Hanna-Attisha M, LaChance J, Sadler RC *et al.* (2016) Elevated Blood Lead Levels in Children Associated With the Flint Drinking Water Crisis: A Spatial Analysis of Risk and Public Health Response. *American Journal of Public Health* 106, 283-290.

**Table 1.** Sociodemographic characteristics of undergraduate students at a large, public Midwestern university and mean beverage intake, stratified by drinking water safety perception and water filter use\*

	Tap Water is Safe to Drink				Filter Tap Water		
	Total n (%)	Agree n (%)	Do Not Agree† n (%)	p-value‡	Never n (%)	Always/Sometimes n (%)	p-value
<b>Total (%)</b>	793	502	291		171	622	
<b>Age</b>							
≤18 years	98 (12.3)	50 (10.0)	48 (16.5)	<b>&lt;0.001</b>	19 (11.1)	79 (12.7)	<b>0.025</b>
19-20 years	171 (21.6)	83 (16.5)	88 (30.2)		25 (14.6)	146 (23.5)	
≥21 years	524 (66.1)	369 (73.5)	155 (53.3)		127 (74.2)	397 (63.8)	
<b>Gender</b>							
Man	287 (36.2)	195 (38.8)	92 (31.6)	0.102	65 (38.0)	222 (35.7)	0.746
Other/Non-Binary	24 (3.0)	16 (3.2)	8 (2.7)		4 (2.3)	20 (3.2)	
Woman	482 (60.8)	291 (58.0)	191 (65.6)		102 (59.7)	380 (61.2)	
<b>Race/ethnicity</b>							
Asian/PI	309 (38.9)	179 (35.7)	130 (44.7)	<b>&lt;0.001</b>	39 (22.8)	270 (43.4)	<b>&lt;0.001</b>
Black	56 (7.0)	32 (6.4)	24 (8.2)		23 (13.5)	33 (5.3)	
Hispanic	82 (10.3)	32 (6.4)	50 (17.2)		16 (9.4)	66 (10.6)	
Other/Multi-racial§	59 (7.4)	41 (8.2)	18 (6.2)		11 (6.4)	48 (7.7)	
White or MENA <sup>†</sup>	287 (36.2)	218 (43.4)	69 (23.7)		82 (48.0)	205 (33.0)	
<b>First generation<sup>†</sup></b>							
No	435 (56.1)	315 (62.7)	120 (41.2)	<b>&lt;0.001</b>	101 (60.5)	334 (54.8)	0.194
Yes	341 (43.9)	176 (35.1)	165 (56.7)		66 (39.5)	275 (45.2)	
<b>Pell grant<sup>**</sup></b>							
No	420 (53.0)	292 (58.2)	128 (44.0)	<b>&lt;0.001</b>	96 (56.1)	324 (52.1)	0.347
Yes	373 (47.0)	210 (41.8)	163 (56.0)		75 (43.9)	298 (47.9)	
<b>Food insecure</b>							
No	631 (79.6)	428 (85.3)	203 (69.8)	<b>&lt;0.001</b>	134 (78.4)	497 (79.9)	0.658
Yes	162 (20.4)	74 (14.7)	88 (30.2)		37 (21.6)	125 (20.1)	
	<b>Total Mean (SD)</b>	<b>Mean (SD)</b>	<b>Mean (SD)</b>	<b>p-value</b>	<b>Mean (SD)</b>	<b>Mean (SD)</b>	<b>p-value</b>
Total water intake (oz/day)	50.0 (43.8)	54.5 (40.1)	42.5 (48.9)	<b>&lt;0.001<sup>††</sup></b>	48.1 (44.3)	57.4 (41.6)	<b>0.013<sup>‡‡</sup></b>

	(L/day)	1.48 (1.30)	1.61 (1.19)	1.26 (1.44)		1.42 (1.31)	1.70 (1.23)	
SSBs intake	(oz/day)	9.2 (17.9)	7.7 (13.2)	12.2 (23.9)	<b>0.003<sup>††</sup></b>	9.5 (19.1)	8.7 (13.2)	0.531 <sup>††</sup>
	(L/day)	0.27 (0.53)	0.23 (0.39)	0.36 (0.71)		0.28 (0.56)	0.26 (0.39)	
		<b>Total Mean (SD)</b>	<b>Mean (SD)</b>	<b>Mean (SD)</b>	<b>p-value</b>	<b>Mean (SD)</b>	<b>Mean (SD)</b>	<b>p-value</b>
Bottled water intake (times/day)		0.7 (1.5)	0.5 (1.2)	1.0 (1.7)	<b>&lt;0.001<sup>††</sup></b>	0.7 (1.4)	0.8 (1.5)	0.501 <sup>‡‡</sup>
Tap water intake (times/day)		3.3 (2.3)	3.7 (2.2)	2.5 (2.4)	<b>&lt;0.001<sup>‡‡</sup></b>	3.6 (2.3)	3.2 (2.3)	<b>0.031<sup>‡‡</sup></b>

\* Column percentages represent percent of students falling into drinking water perception category or filter use category, total percentages represent percent of total sample population. Missing data were removed from analyses

<sup>†</sup>Includes those who neither agree nor disagree with the statement “My local tap water is safe to drink”.

<sup>‡</sup>P-value indicates statistical significance of comparisons between groups using  $X^2$  tests.

<sup>§</sup>Includes students reporting race/ethnicity as Other (self-defined) or Multiracial/Multiethnic

<sup>¶</sup>Middle Eastern/North African

<sup>¶</sup>Students were classified as first-generation college students if the highest level of education of both parents was high school/GED. A missing data indicator was used for n=17 participants missing data on first generation status.

<sup>\*\*</sup>Students receiving a Federal Pell grant; Federal Pell grants are usually awarded to students with exceptional financial need and are therefore an indicator of lower income level.

<sup>††</sup>Two-sample t-test for equality of means, unequal variances

<sup>‡‡</sup>Two-sample t-test for equality of means, equal variances

**Table 2.** Adjusted odds ratios (OR) with 95% confidence intervals (CI) and mean intake data for drinking water behaviors and beverage consumption among undergraduate students at a large, public Midwestern university (n=793)<sup>\*</sup>

Beverage Consumption	Tap Water is Safe to Drink				Filter Tap Water			
	Agree (n=502)		Do Not Agree (n=291)		No (n=171)		Always/Sometimes (n=622)	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Total water intake $\geq 3$ cups/day	Ref.	-	<b>0.45</b>	<b>(0.32, 0.62)</b>	Ref.	-	<b>0.59</b>	<b>(0.39, 0.90)</b>
SSB intake $\geq 1$ serving/day <sup>†</sup>	Ref.	-	<b>1.47</b>	<b>(1.01, 2.14)</b>	Ref.	-	1.11	(0.71, 1.73)
Bottled water intake $\geq 1$ time/day	Ref.	-	<b>1.80</b>	<b>(1.22, 2.66)</b>	Ref.	-	0.70	(0.44, 1.10)
Tap water intake $\geq 3$ times/day	Ref.	-	<b>0.46</b>	<b>(0.34, 0.64)</b>	Ref.	-	0.79	(0.55, 1.13)

<sup>\*</sup>All models were adjusted for age, gender, race/ethnicity, first generation status, Pell grant status, and food insecurity status. N=17 respondents were missing data on first generation status and were retained in the model using a missing value indicator.

<sup>†</sup>Serving defined as 12 fluid ounces (355 ml)