

REVIEW

Comparative reproduction of the female horse, elephant and rhinoceros: implications for advancing assisted reproductive technologies

Janine Meuffels-Barkas^{1,2}, Sandra Wilsher³, W R Twink Allen³, Cyrillus Ververs⁴ and Imke Lueders^{1,5}

¹Cryovault, Rhino Force SA NPC, Tokai, South Africa

²Department of Production Animal Studies, Faculty of Veterinary Science, University of Pretoria, Onderstepoort, South Africa

³The Paul Mellon Laboratory, Brunswick, Newmarket, Suffolk, UK

⁴VetRepSol, Rhenen, Utrechtsestraatweg, TW, The Netherlands

⁵Mammal Research Institute, Faculty of Natural and Agricultural Sciences, University of Pretoria, Hatfield, South Africa

Correspondence should be addressed to J Meuffels-Barkas; Email: Meuffels.janine@gmail.com

Abstract

Recent loss of rhinoceros subspecies has renewed interest in using more advanced assisted reproductive technologies (ART) in rhinoceroses and elephants. Currently, only semen collection, semen preservation and artificial insemination (AI) have been used repeatedly with success in these species. Although ovum pick-up (OPU) and intra-cytoplasmic sperm injection (ICSI) have been reported recently in rhinoceroses, the techniques are not yet optimised. In contrast, multiple ART applications are routinely used in the horse. Since elephants and rhinoceroses share some reproductive features with equids, we postulate that procedures such as OPU, ICSI, *in vitro* fertilisation (IVF) and embryo transfer (ET), which are well established in the horse, may represent a basis to develop protocols for endangered pachyderms. In this review, we summarise current knowledge on reproductive physiology relevant to ART. We discuss the current state of ART in all three families and the requirements for the successful implementation of OPU, ICSI, IVF and ET in these species.

Lay summary

Wild rhinoceros and elephant populations are facing ongoing threats; therefore, additional measures are required to protect these species for future generations. Assisted reproductive technologies (ART) include the collection of semen to directly inseminate females or to fertilise oocytes (eggs) in a laboratory to produce embryos, which can be transferred into a recipient female at a later date. While these techniques are routinely used in humans and domestic animals such as the horse, more research is needed to incorporate such technologies into the breeding of elephants and rhinoceroses. As the horse is the closest related domestic species to the rhinoceros, it may serve as the best possible role model. We discuss the current state of ART in the horse, elephant and rhinoceros and the possibilities for future use of these techniques in breeding such endangered animals.

Keywords: ▶ artificial reproduction ▶ ovum pick-up ▶ embryo transfer ▶ elephantidae ▶ rhinocerotidae

Reproduction and Fertility (2023) 4 e230020

Introduction

Captive breeding populations of endangered species need a certain number of fertile animals with genetic diversity to remain stable and/or become self-sustaining. However, natural breeding in captive animals often poses difficulties. For example, genetically diverse mating partners may be located at different facilities in other countries or continents, and fertility may be sub-optimal when the social needs of captive animals are not met. The use of assisted reproductive technologies (ART) would appear to be one solution to help overcome these problems (Comizzoli 2015) and preserve fertility beyond the reproductive lifespan of an individual. However, currently the use of ART in pachyderms is far from being a routine procedure that can be used to increase an individual's breeding prospects and/or increase numbers within a population.

The application of any ART to increase numbers and diversity in captive breeding programmes should never be viewed as the single answer. It is important to remember that many other factors that influence reproduction in these animals have not been studied in enough detail, including the physiology and hormonal control of reproduction, husbandry needs *per se* and how the structure of their societies and social relations affects breeding and fertility (Lueders & Allen 2020).

Hence, the current rather low success rate of ART in the rhinoceros and elephant is due, in part, to a lack of fundamental reproductive knowledge. Coupled with this are the intractability of these species compared to domestic animals, low numbers of individuals available to undertake research on and the substantial cost involved with developing these techniques in new species (Lueders & Allen 2020). To complicate the matter still further, there are several species, and even subspecies, of elephants and rhinoceroses, each with their own individual reproductive characteristics.

Nevertheless, for both the elephant and rhinoceros, ART has a role to play in captive wildlife breeding programmes (Hermes *et al.* 2007, 2013, Stoops *et al.* 2016). Most of the advancements in the modern reproductive technologies in pachyderms have been made in *ex situ* breeding of Asian and African elephants, primarily by establishing protocols for semen freezing and artificial insemination (AI).

Over a decade ago, Hermes *et al.* (2007) reviewed ART being undertaken in elephants and rhinoceroses. Since then, improvements in AI techniques in both families

and developments in other more advanced reproductive technologies such as ovum pick-up (OPU) and *in vitro* embryo production in rhinoceroses have been made (Hermes *et al.* 2009b, Hildebrandt *et al.* 2012, 2018, Stoops *et al.* 2016, Pennington *et al.* 2020). However, the use of ART in pachyderms is far from routine. Since ART are more advanced in the horse (Hinrichs 2018), can we use this species as a model for the elephant and rhinoceros, despite variations in their reproductive anatomy and physiology?

Elephants, rhinoceroses and horses are all large terrestrial mammals. However, taxonomic classification of these three species varies. Elephants are classified in the order Proboscidea and family Elephantidae and consist of three existing species: African savannah (*Loxodonta africana*), African forest (*Loxodonta cyclotis*) and Asian elephant (*Elephas maximus*) (Roca *et al.* 2015) and are far removed from the rhinoceros and horse on the evolutionary tree. Two African species, the black (*Diceros bicornis*) and white rhinoceros (*Ceratotherium simum*), and three Asian species, the Indian (*Rhinoceros unicornis*), Sumatran (*Dicerorhinus sumatrensis*) and Javan rhinoceros (*Rhinoceros sondaicus*) (Amin *et al.* 2006), are classified in the same order as the horse, Perissodactyla, and although in different families (Rhinocerotidae and Equidae), they share more reproductive characteristics than the horse and elephant. Nevertheless, certain reproductive similarities between the elephant and horse are present (Allen 2010).

Hence, the aim of this review is to highlight similarities and differences in the reproductive characteristics of the domestic horse mare, elephant cow and white rhinoceros cow and discuss how these relate to our ability to practically apply ART in pachyderms, and to what degree the horse is a suitable model.

Reproductive anatomy

Understanding the reproductive anatomy of a species is essential to undertake reproductive procedures, interpret clinical findings and allow the development of species-specific equipment.

Schematic drawings of the female reproductive tract of the horse, elephant and rhinoceros *in situ* and *ex situ* are compared in Fig. 1. Similarities, like a bicornuate uterus and ovarian functional structures, differences and special features are summarised in Table 1 and briefly described below.




Figure 1 Schematic drawing of the female reproductive organs *in situ* (A, B, C) and *ex situ* (D, E, F) in the mare (A and D), elephant (B and E) and rhinoceros (C and F). B, bursa; B lig, broad ligament; Ce, cervix; Cl, clitoris; IC Lig, intercornuate ligament; K, kidney; M, mammary gland; O, ovary; Ov, oviduct; OvF, ovulation fossa; R, rectum; UB, urinary bladder; UC, uterine corpus; UH, uterine horn; UOs, urethral os; Va, vagina; Ve, vestibule; VOs, vaginal os; Vu, Vulva. Note that the full vestibule of the elephant *ex situ* (D) is not displayed due to the length. Adapted from [Allen et al. 2003](#), [Schaffer et al. 2001](#).

Mare

The internal reproductive tract of the mare is positioned horizontally, extending into the pelvic cavity and abdomen ([Fig. 1A](#)). The vulva forms the gateway to the vestibule which continues cranially as the vagina; a vestibule-vaginal ring is present at the junction of the two, and a hymen may be present in unmated females ([Kainer 2011](#)). The relatively straight cervix consists of a thick muscle layer and multiple longitudinal mucosal folds, but no transverse cervical rings, making it easy to pass catheters through for AI or embryo transfer (ET) ([Stout 2020](#)). The external os of the cervix protrudes into the cranial vagina and is surrounded by the vaginal fornix, an internal os opens into the uterus

([Fig. 1D](#)). The Y-shaped uterus consists of a prominent uterine body and two horns suspended within the pelvic cavity and abdomen by the broad ligament ([Kainer 2011](#)). The relatively long oviducts of 20–30 cm have a fimbriated infundibulum positioned over the ovulation fossa to guide the oocyte into the ampulla of the oviduct and are attached to the lateral surface of the ovaries ([Kainer 2011](#)). The ovaries can be readily palpated per rectum, are comparatively large and consist of an outer medulla and an inner cortex containing the functional structures, such as follicles and corpora lutea. Follicles can reach up to 50 mm prior to ovulation, which can only occur at the ovulation fossa as the rest of the ovarian surface is covered by the tough tunica albuginea ([Kainer 2011](#)).

Table 1 Overview of special reproductive anatomy characteristics of mare, African elephant and white rhinoceros cow with average measurements.

	Horse	Elephant	Rhinoceros
Vestibule and vagina	Horizontally positioned, separated by vestibule-vaginal ring, vestibule 10-12 cm, vagina 15-20 cm long	Extremely long (100-150 cm) vestibule, opens between the hind legs; vagina (20-40 cm) has many longitudinal folds and opens into the vestibule through small vaginal os (0.5-2 cm) within the intact hymenal membrane in nulliparous females, additional vaginal opening(s) and /or blind pouches on either side of the vaginal os possible; vaginal os in parous females wide at 5-19 cm	Similar conformation to the mare, transversal, thin vaginal folds in the cranial vagina, vestibule 14-20 cm, vagina 20-30 cm long
Cervix	Short (5-7 cm), diameter: 3.5-4 cm; consists of a thick muscle layer and longitudinal folds	Short (6-10 cm), longitudinal folds, cranio-ventral direction of pelvis	Long (13-20 cm), tortuous, with firm, right angle folds of connective tissue, narrow lumen, prominent cervical os and fornix similar to mare
Uterus	Bicornuate; Y-shaped, prominent uterine body (18-20 cm), horns (20-25 cm)	Bicornuate; U-shaped, short body (5-15 cm); long horns (30-80 cm), connected for 50-70 cm up to the uterine bifurcation	Bicornuate; short body (3.5-7.5 cm); long, straight horns (40-50 cm) of equal diameter (3-4 cm) over the entire length
Oviduct	Relatively long (20-30 cm), fimbriated infundibulum positioned over the ovulation fossa	Short (10 cm); enveloped by the infundibulum of the oviduct which is incorporated in the ovarian bursa, forms serosal pouch around ovary	Short, fimbriated at the cranial opening nearest the ovaries
Ovaries	<ul style="list-style-type: none"> • Comparably large (6-8 × 3-4 cm) cranially of the tip of the uterine horns • Outer medulla and inner cortex • Ovulation fossa where cortical tissue extends to the ventral surface of the ovary 	<ul style="list-style-type: none"> • Relatively small (7 × 5 × 2.5 cm), cranially of uterine horn • Inner medulla and outer cortex • Surface has irregular ridges • Entire ovarian surface is enveloped by ovarian bursa 	<ul style="list-style-type: none"> • Comparably large (8 × 4 × 2cm), when active, cranially of the tip of the uterine horns, caudo-laterally to the kidneys • Inner medulla and outer cortex
Functional structures	<ul style="list-style-type: none"> • Antral follicles of various sizes always present • Pre-ovulatory follicle size: 3.5-5 cm • One CL from ovulation • Accessory CLs during pregnancy 	<ul style="list-style-type: none"> • Tiny antral follicles only during follicular phase visible (0.1-1.5 cm) • Pre-ovulatory follicle size: 2 cm • One CL from ovulation (4-5 cm) • Accessory CLs formed from luteinised unovulated follicles prior to each ovulation (2-3.5 cm) 	<ul style="list-style-type: none"> • Antral follicles of various sizes always present • Pre-ovulatory follicle size: 3-4 cm • Only one CL during pregnancy, no accessory CLs

Elephant cow

The reproductive tract of the female elephant, from vestibule to ovary, is the longest of the terrestrial animals, reaching a length of approximately 3.5 m in the African elephant (Balke *et al.* 1988). Some unique anatomical features (Figs. 1B and E) complicate the development and use of ART in these species (Balke *et al.* 1988, Hildebrandt *et al.* 1999). The extremely long vestibule curves from the caudal pelvis to just under the anus before arching downwards to open between the hind legs (Fig. 1B). The pelvic part of the vestibule is separated from the vagina through a hymen-like structure which breaks only during the first parturition, and not during copulations as in the mare and the rhinoceros. Unique to elephants is a small

opening in the hymen and two blind pouches on either side of the vaginal opening (Hildebrandt *et al.* 2006), although variations are seen between individuals (i.e., two vaginal openings, no blind pouch; Balke *et al.* 1988 and I. Lueders, personal observation; Fig. 2D-F). A short cervix with longitudinal folds opens into a short uterine body. Similar to the horse, the two horns are initially positioned straight before diverging laterally from each other towards the ovaries (Fig. 1E). However, in contrast to the mare, the long horns are connected by a short intercornuate ligament for 0.5-0.7 m and aligned in an anterior direction (Allen *et al.* 2003, Hermes *et al.* 2006). The ovaries are relatively small, lying cranial to the uterine horn at the termination of short oviducts, and the entire ovarian surface is encapsulated by an ovarian bursa (Fig. 2). The




Figure 2 Photographs of special anatomical features of the rhinoceros (A) and elephant (B–F). (A) White rhinoceros cervix opened, exposing the folded structure. utb, uterine body; uh, uterine horn; (B) African elephant ovary within the ovarian bursa (ovb) and uterine horn tip (uh); (C) same ovary, ovarian bursa (ovb) removed and ovary turned over, note the corpora lutea (CLs) and a flaccid cystic structure (Cy); (D–F) endoscopic view (*in situ*, from vestibulum) onto the hymen-like structure separating vestibule and vagina in the Asian elephants: (D) with AI catheter placed through the vaginal os; (E) two small ostia opening into the vagina, (F) variation of the larger vaginal os with tissue bridge and vaginal mucosa shining through.

elephant ovary is composed of an inner medulla and outer cortex, meaning ovulation can occur at any location on the ovaries' surface (Hildebrandt *et al.* 2000). In contrast to the mare and rhinoceros cow, which have smooth plum-like ovaries, the surface of an elephants' ovaries show irregular ridges (Allen *et al.* 2003). Antral follicles are visible only during the follicular phase and at no other stage. Dominant follicles are considerably smaller than those of the mare and rhinoceros cow, reaching only about 2.0 cm prior to ovulation (Hildebrandt *et al.* 2011).

Rhinoceros cow

The uterus, vagina and cervical os of the rhinoceros resemble those of the mare (Figs. 1C and F). In maiden females, a hymenal membrane is present (Hermes *et al.*

2006). The cervix is particularly long and has firm folds of connective tissue (Figs. 1F and 2) leaving a narrow, tortuous lumen (Godfrey *et al.* 1991, Hermes *et al.* 2007) which is challenging to catheterise. The uterine horns are straight with their tips pointing cranially and not diverging laterally as in the mare and elephant cow (Fig. 1F). Like those of the horse, the oviducts consist of a small tubular structure with fimbriae at the cranial openings nearest the ovaries. The ovaries are positioned within an ovarian bursa (Godfrey *et al.* 1991, Schaffer *et al.* 2001). Inactive ovaries are small and flat, while active ovaries are large and round with follicles, and/or a corpus luteum or corpus haemorrhagicum (Schaffer *et al.* 2001, Hermes *et al.* 2006). Unlike the mare, rhinoceros ovaries show an inner medulla and outer cortex. Despite this difference, antral follicles, present at any cycle stage develop similar to the horse with a single dominant follicle prior to ovulation (Radcliffe *et al.* 1997).

Reproductive physiology

A precondition for undertaking ART is a thorough understanding of the hormonal changes of the oestrous cycle and early gestation. The mare and rhinoceros cow share several hormonal oestrous cycle features, whereas elephants present some unique aspects (Fig. 3). While extensive literature is available for the mare (as reviewed by Satué & Gardon 2020), only relatively recently did we understand the reproductive physiology of the elephant (as reviewed by Hildebrandt *et al.* 2011) and rhinoceros (as reviewed by Roth 2006, Stoops *et al.* 2016) in more detail. The most relevant aspects are summarised in Table 2 and briefly described here.

Oestrous cycle

Mare

The mare is a long day, seasonally polyoestrous breeder with regular oestrous cycles of 20–23 days occurring between spring and autumn. Each cycle consists of 5–6 days of oestrus or sexual receptivity induced by oestrogens secreted by enlarging follicles in the ovaries and which is followed by 14–15 days of dioestrus under the dominance of progesterone (Fig. 3A) the source of which is a, usually single, corpus luteum (CL). Luteinising hormone (LH) levels rise during late oestrus and peak after, not before, ovulation (Ginther 1992, Aurich 2008). Regression of




Figure 3 Model of changes in known reproductive hormones during the oestrous cycle of the mare (A), elephant (B) and (Indian) rhinoceros cow (C). Day 0, day of ovulation; P4, progesterone (5 α -reduced metabolites in elephants); LH, luteinising hormone; FSH, follicle-stimulating hormone; E2, oestrogen; PGF2 α , prostaglandin F2alpha. Adapted from [Stoops *et al.* 2004](#), [Ginther 2007](#), [Aurich 2011](#), [Hildebrandt *et al.* 2011](#), [Lueders *et al.* 2011](#).

other follicles in a cohort occurs when one dominant follicle begins to secrete inhibin as it enlarges causing suppression of follicle-stimulating hormone (FSH) from the pituitary, helping to ensure only a single follicle matures and ovulates ([Fig. 3A](#)). Progesterone dominates until around day 14 of dioestrus when spike-like bursts of prostaglandin F2alpha (PGF2 α), released from the luminal epithelium of the endometrium, travel to the ovaries via the maternal circulation causing luteolysis of the corpus luteum ([Ginther 1992](#)).

Elephant cow

The female elephant exhibits a long oestrous cycle of 13–17 weeks, composed of a follicular phase of 6–8 weeks, followed by a luteal phase of 6–10 weeks ([Lueders *et al.* 2010b](#)). Instead of a single ovulation-inducing LH peak, the elephant expresses two LH peaks ([Fig. 3B](#)), of which only the second results in ovulation ([Brown *et al.* 1999](#)). During a first follicular wave, which commences when progesterone of the previous luteal phase reaches baseline levels, a cohort of follicles develop over the course of approximately 3 weeks ([Lueders *et al.* 2010b](#)). This first wave ends with a first, anovulatory LH peak in the absence of overt oestrous signs, although interest in the bull and even mating have been observed at this time ([Lueders *et al.* 2010b](#), [Hildebrandt *et al.* 2011](#)). Typically, 1–10 follicles of >10 mm luteinise after the first LH peak but only become ultrasonographically visible as luteinised unovulated follicles (LUFs) about 10 days later. These LUFs form slowly growing accessory CLs in each oestrous cycle. A second wave of follicular growth then commences and a second LH peak occurs approximately 3 weeks (19–21 days) later (as reviewed by [Hildebrandt *et al.* 2011](#)). Only 12–24 h after the second LH peak does one single follicle of the second cohort mature and ovulate when the follicle reaches approximately 2.0 cm ([Hildebrandt *et al.* 2000](#), [Lueders *et al.* 2010b](#)). The ovulated follicle develops into the largest CL, which along with the accessory CLs, produces metabolites of 5- α reduced progestagen ([Lueders *et al.* 2012](#)). The serum progestagen rise occurs within 24 h after the second LH surge ([Fig. 3B](#)). The ovulatory CL is significantly larger than the anovulatory CLs derived from LUFs and remains discernible into the next follicular phase ([Lueders *et al.* 2010b](#)). All CLs persist throughout the dioestrous period after which they regress in the absence of pregnancy during the new follicular phase ([Brown *et al.* 1999](#), [Hildebrandt *et al.* 2006, 2011](#), [Lueders *et al.* 2010b](#)).

The LUFs do not produce measurable progestagens in the peripheral circulation prior to the second LH peak but are a source of inhibin ([Kaewmanee *et al.* 2011](#), [Yamamoto *et al.* 2012](#)), which rises measurably only after the first LH peak ([Fig. 3B](#)) and in parallel with LUF growth. Therefore, LUF formation may be a precondition for the selection of a single, dominant follicle at the second LH peak ([Lueders *et al.* 2011](#)).

Rhinoceros cow

The five living rhinoceros' species show slight variation in their oestrous cycle length, dominant follicle size and

Table 2 Overview of key features of the reproduction physiology of the mare, elephant cow and the rhinoceros species.

	Horse	Elephant	Rhinoceros				
			White	Black	Indian	Sumatran	Javan
Female maturity (years)	1	6-10	5-6	4-7	6-7	4-6	5-6
Oestrous cycle length (days)	20-23	95-120	30-35 or 65-70	27	40-50	21-24	?
Oestrous duration (days)	5-6	2-4	1-3				
Repeated reproductive ultrasound	Yes	Yes	Yes	Yes	Yes	Yes	No
Ovulation type	Spontaneous	Spontaneous	Partially induced	Spontaneous	Spontaneous	Induced	?
Ovulatory follicle size (cm)	3.5-5.0	2.0	3.0-4.0	4.7-5.1	12.0-14.0	2.0-2.5	?
Pregnancy detection							
TU at	14 days	55 days	15 days				
Serum	eCG	Prolactin at 120 days	Progesterone				
Embryonic vesicle	Spherical, mobile	Spherical, immobile	Spherical, mobile (?)*				
Gestation length (days)	330	600-680	450-490	450-490	450-490	450-490	490-580
Intercalving interval (years)	1	4-8	2.5-3.5	2.5-3.5	3	3	?
Placenta type	Diffuse epitheliochorial	Endotheliochorial zony	Diffuse epitheliochorial with villus-free areas*				

TU, transrectal ultrasound.
*this is for all sub-species of rhinoceros

hormone secretion, but overall they are akin to the mare with a 3- to 5-day period of oestrous behaviour (Radcliffe *et al.* 1997) that accompanies the growth and ovulation of one single dominant follicle (as reviewed by Roth *et al.* 2018). Ovulation of the dominant follicle (30-40 mm in white rhinoceroses) at the end of the follicular phase takes place within the cortex of the ovary and is induced by a single pre-ovulatory LH peak (Hermes *et al.* 2007, Radcliffe *et al.* 2001, Fig. 3C).

Oestrous cycle length differs, and shorter or longer cycles have been reported in all captive species (Roth *et al.* 2018, Table 2). Unique to the white rhinoceros, two distinctive cycle lengths of 30-35 and 65-70 days have been recognised in captive animals (Patton *et al.* 1999, Hermes *et al.* 2012, Pennington *et al.* 2020). Although ovulations followed the follicular phase in cycles of both lengths, the longer cycle was originally thought to be associated with in- or subfertility (Schwarzenberger *et al.* 1998, Patton *et al.* 1999, Brown *et al.* 2001). However, more recently, ovulation induction and AI in rhinoceroses presenting with long luteal phases prior to the new cycle resulted in pregnancy (Pennington *et al.* 2020). In the white rhinoceros and its African relative, the black rhinoceros, longer cycles are associated with longer luteal phases which have not been identified in the Asian species. Instead, longer cycles in Asian rhinoceroses appear to be related to a delayed development of a dominant follicle in a follicular wave or ovarian cysts (Stoops *et al.* 2004).

As reported in the horse, follicles may become haemorrhagic (Pennington *et al.* 2019), frequently referred to as haemorrhagic anovulatory follicles (HAFs) in the mare (Ginther 2007). These follicles typically grow larger than the normal pre-ovulatory size and develop echogenic patterns and fibrinous, spider-web-like structures within the fluid-filled lumen. Like HAFs in mares, they may luteinise and secrete some progesterone or slowly organise and regress without any progestagen production, as noted in white rhinoceroses (Roth *et al.* 2018).

Oestradiol, progesterone and LH (Fig. 3C) all play a part in orchestrating the reproductive cycle in the white rhinoceros, but other reproductive hormone (especially proteohormones such as FSH, inhibin, prolactin, PGF2 α) profiles have not been elucidated.

Placentation and the hormonal control of early gestation

A sound understanding of early embryonic development, maternal recognition of pregnancy (MRP) and the

hormonal changes involved during early gestation is required in order to undertake procedures such as ET successfully. While the mare has been studied in detail (as reviewed by [Allen 2001](#)), and over the last decade much has been learnt about early pregnancy in the elephant (as reviewed by [Allen & Stansfield 2021](#)), there remains a paucity of information regarding the rhinoceros.

Mare

Pregnancy in the mare includes a number of fascinating and apparently equine-specific events. Only fertilised oocytes enter the uterus, with the early embryo secreting prostaglandin E2 (PGE2) from day 5 to day 6 ([Weber *et al.* 1991](#)) to cause relaxation and contraction of the muscles of the oviduct and, hence, passage through the utero-tubal junction into the uterus. Unfertilised oocytes remain within the oviduct and either degenerate over time or pass into the uterus in the presence of an embryo. The equine embryo has a long oviductal transport time of 144–156 h post-ovulation ([Battut *et al.* 1998](#)) and arrives in the uterus as a late morula or early blastocyst. Practically, this means embryo recovery for transfer, bisection or cryopreservation in the mare can only be undertaken from day 6.5, with optimum recovery rates from day 7.

The spherical equine embryo sheds the zona pellucida around day 7, underneath which is a tough acellular glycoprotein capsule, at least partly secreted by the trophoblast cells, that will encapsulate the embryo until around day 25 ([Betteridge 2000](#)). The equine embryo also shows mobility during early pregnancy, continually moving through the uterine lumen until around day 16 when it lodges at the base of one of the uterine horns ([Ginther 1983](#)). This mobility is believed to be driven by prostaglandins secreted by the embryo ([Stout & Allen 2001](#)) and is considered essential to liberate the MRP signal over the uterine epithelial surface to prevent the cyclic release of PGF2 α from the endometrium and a return to oestrus by the mare. In addition, copious amounts of oestradiol are produced by the early equine conceptus from at least day 12 of pregnancy, and although their precise role has not been elucidated, they have been proposed to play a role in the establishment of pregnancy ([Heap *et al.* 1982](#)).

Throughout the mobility stage, and even after the conceptus becomes stationary at the base of a uterine horn, it relies on protein-rich secretions from the endometrial glands, the so-called histotrophe, which contain numerous proteins (e.g. uterocalin and uteroferrin) ([Ellenberger *et al.* 2008](#)), to supply its nutritional needs. Interdigitation of the allantochorionic

placenta does not commence until around day 45 and this is initially rudimentary ([Samuel *et al.* 1974](#)), making histotrophic nutrition very important in the horse. Indeed, secretions from the endometrial glands play an important role throughout gestation in the horse ([Ellenberger *et al.* 2008](#)).

The equine conceptus also produces a chorionic gonadotrophin (eCG) during early pregnancy with the only other mammals known to do so being humans and the higher primates. In the horse, a specialised band of tissue called the chorionic girdle develops around the conceptus. This invades the endometrium around days 36–38 of pregnancy to form the endometrial cups which secrete eCG into the maternal bloodstream between 40 and 120 days of gestation ([Allen & Moor 1972](#)).

Hence, following the development of the endometrial cups, progesterone levels in the mare's bloodstream, which had been falling steadily from around day 20, rise sharply and remain elevated thereafter until days 120–140 as a result of the continuing 'collaborative' actions of the pituitary FSH-stimulated follicular waves and luteinising properties of the eCG secreted by the endometrial cups causing accessory CL to develop. Interestingly, eCG also causes an increase in peripheral oestrogens which are secreted by luteal tissue within the ovaries ([Daels *et al.* 1991](#)).

Continuing interdigitation of the allantochorion from day 45 results in the formation of complex microcotyledons over the entire surface of the allantochorion which interdigitate with the endometrium in corresponding crypts with no breakdown of the maternal tissue ([Samuel *et al.* 1974](#)). Hence, placentation in the horse is described as diffuse, non-invasive and epitheliochorial. In addition to providing the means for haemotrophic nutrition during pregnancy, the fetoplacental unit also synthesises steroid hormones during the second half of pregnancy. To do so, it utilises fetal C-19 precursors produced by the enlarged fetal gonads to produce oestrogens from around day 80, peaking at 220 days before falling as parturition approaches. In addition, maternal C-21 precursors are synthesised by the placenta to 5 α -reduced progestagens ([Allen 2001](#)).

In summary, principal endocrinological changes during pregnancy in equids are unusual with many being specific to this genus. Luteal progesterone production declines during the first 40 days but thereafter shows a secondary rise during the next 100 days due to the development of secondary or accessory CLs in the maternal ovaries stimulated by the combined actions of continuing 10- to 12-day releases of pituitary FSH and the

LH-like properties of eCG secreted by the endometrial cups between 40 and 120 days of gestation. Thereafter maternal serum progesterone concentrations remain low and constant until a further rise occurs during the last 6–8 weeks of gestation as a result of secretion by the fetal adrenal glands. Oestrogens also play a role in equine pregnancy with initial production by the early embryo, then the luteal tissue and in later pregnancy synthesis by the placenta from precursors from the fetal gonads (Allen 2001).

Elephant cow

A number of aspects of the 590–680 days gestation in the elephant are of particular interest due to their similarity to events in the pregnant mare, while other features vary greatly (Allen & Stansfield 2021).

As mentioned previously, the LUFs from the first wave of follicular growth persist in the ovaries well beyond the second follicular wave and the associated ovulation that can result in pregnancy, if accompanied by a fertile mating. If conception has taken place, all CLs enlarge significantly around the time of implantation (40–50 days). These CLs persist in the ovaries for the remainder of gestation and secrete the metabolites of 5α -reduced progestagens needed to maintain pregnancy, since the placenta of the elephant does not synthesise progestagens (Lueders *et al.* 2012).

Implantation of the embryo occurs at the base of the uterine horn, always ipsilateral to the side of ovulation (Lueders *et al.* 2012) with no observable free embryonic migration as described for the mare. Consistent with the long oestrous cycle and gestation length, embryonic development is rather slow. Compared to the horse and rhinoceros, the embryonic vesicle and the embryo proper only become apparent ultrasonically at 40–50 days (Fig. 4C) (Lueders *et al.* 2010a) and 65 days (Fig. 4D), respectively, resembling the stages of day 14 (Fig. 4A) and day 21 (Fig. 4B) pregnancy in the mare. At this stage, a band of trophoblast in the equatorial region of the spherical conceptus extrudes ‘fingers’ of trophoblast cells, which undermine and dislodge the luminal epithelium of the endometrium to replace it with a layer of trophoblast, which now lies on the luminal surface of the endometrial stroma. The endotheliochorial zonary placenta will subsequently develop from increasingly elongated upgrowths of the trophoblast-lined stroma. The trophoblast begins to secrete placental lactogen (Yamamoto *et al.* 2012), which with its prolactin-like action appears to stimulate the enlargement of the all persisting CLs at around 40–50 days of gestation. Unlike the horse, no chorionic gonadotrophin (CG) has

been detected during pregnancy in the elephant. The zonary-girdle placenta is unable to produce progestagens to support the pregnancy but probably contributes to its maintenance through the indirect effect of the prolactin/placental lactogens it produces (Yamamoto *et al.* 2017, Lueders *et al.* 2012).

The progesterone metabolite-secreting ability of the fetal gonads, which in parallel with the situation in equids, enlarge greatly during gestation, may add to the pool of progestagens and, hence, assist the long-lived maternal CLs to maintain the pregnancy state until term (Stansfield & Allen 2012). Thus, as in the horse, the elephant fetus appears to play a hormonal role in orchestrating the endocrinology of pregnancy (Allen *et al.* 2003).

Rhinoceros cow

There is a paucity of information on pregnancy in the rhinoceros compared to the horse and elephant. The gestation length averages 450–500 days for the four captive rhinoceros species (Hermes *et al.* 2007, Schwarzenberger & Hermes 2023).

Progesterone rises only 5–7 days after ovulation and is at luteal phase levels for the first 2–5 months with an increase to pregnancy levels thereafter (Berkeley *et al.* 1997, Patton *et al.* 1999, Brown *et al.* 2001, Pennington *et al.* 2020, Schwarzenberger & Hermes 2023). Only a single CL has been described in pregnant rhinoceroses. The rhinoceroses share some features of early pregnancy with the horse, as the early embryonic vesicle is spherical (Fig. 4) and appears to be mobile during the early stages (Radcliffe *et al.* 1997, Hermes & Hildebrandt 2011). Furthermore, the vesicle can be visualised by ultrasound from 15 days post-conception (Fig. 4E), the embryo itself around 23 days (Fig. 4F) and an embryonic heartbeat at 26 days (Radcliffe *et al.* 1997, Ververs *et al.* 2015), mirroring the timing of these stages in the equine conceptus (Fig. 4). Another similarity with the horse is the diffuse, epitheliochorial placenta, although the villus surface of the rhinoceros placenta has villus-free lines, sometimes referred to as ‘streets’ (Benirschke & Lowenstine 1995).

No CG nor other fetoplacental hormones have been described in the rhinoceros that could play a role in maintaining the single CL, which appears to remain throughout pregnancy with no accessory CL formation (Sherman *et al.* 1997). However, between 60 and 120 days of gestation, there appears to be a transition from luteal to fetoplacental progesterone secretion since faecal pregnane metabolite levels rise significantly above luteal phase concentrations. These faecal pregnanes are




Figure 4 Ultrasonographic appearance of the early pregnancy of the mare (A, B), Asian elephant (C, D) and white rhinoceros (E, F) presented at similar stages of pronounced, spherical embryonic vesicle (EV) and appearance of the embryo proper (→). The EV appears very similar in all three families, although development in the elephant is considerably longer than that of the same stage embryos as in mare and rhinoceros are reached; DAY, day from ovulation; (?), day from observed last breeding, exact day of ovulation was not determined in this rhinoceros; sizes normalised to bar (= 1 cm).

similar to pregnancies in the plasma of pregnant mares, where the source is known to be the foeto-placental unit (Schwarzenberger & Hermes 2023). An endocrinological difference with the horse is the low oestrogen level throughout gestation in at least three of the five rhinoceros species in which this hormone has been measured (Schwarzenberger *et al.* 2000, Stoops *et al.* 2004, Roth 2006).

ART - current state

The development, advancement and implementation of ART in many wildlife species, including elephants and rhinoceroses, is limited by several challenges, including restricted access to individuals, individual differences and intractability as well as low commercial drive (Lueders & Allen 2020). Additionally, some of the unique anatomical features of the rhinoceros and elephant, as described above, complicate routine implementation of ART into captive breeding programmes.

Reports on the successful application of ART in pachyderms are mostly limited to a few individuals, thus bringing the sustainability and repeatability of these techniques into question (as reviewed by Hermes *et al.* 2007, Ververs *et al.* 2015, Pennington & Durrant 2018). However, equine reproductive physiology and many of the ARTs in the mare have been studied for decades resulting in a wealth of knowledge and the ability to apply them

routinely in clinical practice (Hinrichs 2018). To-date applied ARTs in the mare, elephant and rhinoceros species are indicated in Table 3.

Ultrasonography of the female reproductive tract

Ultrasonographic visualisation of the reproductive tract is a requirement to directly undertake ART and to manage or monitor associated reproductive cycles. Due to a thick rectal wall in both, elephant and rhinoceros, the ovaries are not palpable. Like the more tractable mare, both captive elephants and rhinoceroses can be trained to enter a restraining chute to facilitate transrectal manipulation and/or ultrasound of the ovaries and uterus (Fig. 5). Heavy-duty restraining chutes offer the safest option for repeated examinations in pachyderms. However, since the use of chutes requires financial investment and time-consuming training and not all zoos are equipped with these, a standing sedation is an alternative option (Hermes *et al.* 2005, 2007).

Useful descriptions of the female reproductive sono-anatomy are available for horses (Kähn & Volkmann 2004), elephants (Hildebrandt *et al.* 2000) and rhinoceroses (black: Schaffer *et al.* 2001, Sumatran rhinoceros: Roth 2001, white: Hermes *et al.* 2005, Indian: Roth *et al.* 2018).

In contrast to equine transrectal ultrasound, where a linear probe is commonplace, in elephants and rhinoceroses a convex, 3–7 MHz ultrasound probe is more

Table 3 Summary of to-date reported and successfully performed assisted reproductive technologies (ART) in the mare, elephant and rhinoceros species.

	Horse	Elephant	Rhinoceros				
			White	Black	Indian	Sumatran	Javan
Semen collection	AV	TM, EE	EE, UC	EE, UC	EE	EE	None
Semen cryopreservation	Yes	Yes	Yes	Yes	Yes	Yes	No
Hormonal oestrous induction	Yes	No	Yes	No	Yes	No	No
Artificial insemination							
Fresh semen	Yes	Yes	Yes	No	Yes	No	No
Frozen-thawed semen	Yes	Yes	Yes	No	Yes	No	No
Sperm sex sorting	Yes	Yes	Yes	Yes	Yes	No	No
Successful AI with sex-sorted semen	Yes	No	No	No	No	No	No
Ovum pick-up	Yes (TV)	No	Yes (TR)	Yes (TR)	No	Yes (TR)	No
<i>In vitro</i> maturation	Yes	Yes (EM), no (LA)	Yes	Yes	No	Yes	No
IVF	Yes	No	No	2- and 4-cell	No	No	No
ICSI	Yes	No	Blastocyst	No	No	No	No
<i>In vitro</i> embryo production	Yes	No	Yes	Yes	No	No	No
Embryo cryopreservation	Yes	No	Yes	No	No	No	No
Successful embryo transfer	Yes	No	No	No	No	No	No

AV, artificial vagina; EE, electro-ejaculation; EM, *Elephas maximus* (Asian elephant); ICSI, intra-cytoplasmic sperm injection; TM, transrectal massage; TV, transvaginal; TR, transrectal; UC, urethral catheterisation.

appropriate due to the size and depth of the reproductive organs in these substantially larger patients.

Artificial insemination (AI)

Collecting and maintaining semen are pre-requisites for successful AI. Improvements in semen collection in chemically restrained elephants and rhinoceroses using electro-ejaculation (Saragusty *et al.* 2016b, Hermes *et al.* 2018) and urethral catheterisation (rhinoceroses only: Meuffels 2022) or manual stimulation by transrectal massage (elephants only: Hildebrandt *et al.* 1998) as well as advances in semen cryopreservation protocols (African

elephant: Hermes *et al.* 2013, Asian elephant: Arnold 2014, white rhinoceros: Hermes *et al.* 2018, and black rhinoceros: Meuffels 2022) have been made in the last decade. AI with chilled or fresh semen is the easiest approach to breeding mares and is also a practical option in captive elephants and rhinoceroses, if distances allow for the movement of semen within a restricted time frame (Hildebrandt *et al.* 1999, 2007).

The first report of an African elephant calf born following AI with fresh semen originates from the late 1990s (Hildebrandt *et al.* 1999), but it took the same group more than another decade to repeat this success using frozen-thawed semen (Hildebrandt *et al.* 2012).




Figure 5 Ultrasonographic appearance of the dominant, pre-ovulatory follicle and concurrent uterine horn in cross-section of mare (A, B), Asian elephant (C, D), white rhinoceros (E, F); this kind of ultrasonic appearance (large oval follicle, uterine horn with oedema, slight intraluminal fluid accumulation maybe present) would indicate oestrus and the correct timing for AI with fresh semen. CL, corpus luteum; EM, endometrium; Fo, pre-ovulatory follicle; { = myometrium; → = fluid within uterine lumen; sizes normalised to bar (= 1 cm).

In Asian elephants, at least one successful conception has occurred following frozen–thawed semen AI (Thongtip *et al.* 2009), although this did not result in the birth of a live calf. However, results using fresh semen AI are more encouraging and >40 Asian and African elephant calves have been born worldwide (Saragusty *et al.* 2009) using a non-surgical endoscopic AI method whereby semen is placed intravaginally (Brown *et al.* 2004).

The first living white rhinoceros' calves after successful AI with fresh semen were reported in 2007 (Hildebrandt *et al.* 2007), and frozen–thawed semen has since successfully been used in white (Hermes *et al.* 2009b) and Indian rhinoceroses (Stoops *et al.* 2016). However, the exact number of white and Indian rhinoceros calves born following AI is not known, and we estimate only a handful worldwide. No reports of successful AI in black, Sumatran or Javan rhinoceros were found by the authors.

Timing of AI

The detection of oestrus and prediction of ovulation is crucial for successful AI particularly with frozen semen. Knowledge on the longevity of spermatozoa in the female tract is also helpful although this is lacking in the elephant and rhinoceros.

Behavioural observations, endocrine measurements and ovarian ultrasonography (Fig. 5) all play a role in oestrous detection in horses, elephants and rhinoceroses. Similar to the mare, female rhinoceroses may urinate small volumes, lift the tail and accept the sexual behaviour of males by showing a standing reflex which allows mounting (Jenikejew *et al.* 2021). However, behavioural signs may be misleading in captive animals, and even in the domesticated horse, the absence of social cues can make detection of oestrus difficult.

Non-invasive monitoring of faecal or urinary progesterone (P4) levels may retrospectively define when oestrus has occurred. For elephants and rhinoceroses, ovulation takes place around 3 to 1 days (Brown *et al.* 2001, Lueders *et al.* 2010b) and at least 1 week prior to P4 rising from the baseline (Stoops *et al.* 2004).

Another endocrine monitoring option is the determination of the LH peak which, in rhinoceroses and elephants, occurs approximately 24 h prior to the ovulation as opposed to the mare, where ovulation frequently takes place up to 24 h before LH peaks (Fig. 3). However, daily collection of serum is required to measure LH in elephants (Brown *et al.* 1999, 2001) and rhinoceroses; in the latter species, urine LH determination alone is also an option (Stoops *et al.* 2004). Since elephants express two

quite exactly spaced LH peaks, the identification of the first, anovulatory LH surge allows prediction of timing of the second, ovulatory LH surge 19–21 days later and therefore of ovulation (Brown *et al.* 1999, Thitaram *et al.* 2008, Hildebrandt *et al.* 2012). The ability to measure LH in elephants is, in part, the reason for the relatively good success rate following AI. If only LH determination is feasible in the training routine but not ultrasound, the best results are obtained using fresh semen AI on the day of the LH peak, since fresh semen is better placed into the vagina/uterus well in advance to ovulation (I. Lueders, personal observation). In both rhinoceroses and elephants, a commercial in-house serum LH snap test for cats and dogs (Witness[®]LH, Zoetis) may be used with qualitative results within 20 min. In a report of successful frozen–thawed semen AI in an African elephant, this in-house LH test was performed every 12 h to optimise the timing (Hildebrandt *et al.* 2012). Daily inseminations for 2–4 days around the day of ovulation with fresh or frozen semen have resulted in pregnancies in elephants (Brown *et al.* 2004a, Thongtip *et al.* 2009, Hildebrandt *et al.* 2012). One may speculate that similar to the mare, frozen–thawed semen AI in elephant and rhinoceros cows may yield the highest chances of success when the semen is placed no earlier or later than 6 h prior to or after ovulation.

In the mare, the standard procedure to judge the timing for insemination is monitoring follicular development during oestrus via transrectal ultrasound and, since most mares are easily restrained and tractable, this may be repeated daily, or more frequently when frozen semen is used, to measure the size of the pre-ovulatory follicle. Fresh semen AI in mares is usually performed when a follicle of >3.5 cm is present on an ovary in combination with uterine oedema (Fig. 5A and B). Ovulation inducing drugs (e.g. GnRH agonists or human chorionic gonadotrophin (hCG)) are usually administered to ensure ovulation occurs some 36–40 h later. The timing of fresh semen AI within this time window is often operator preference and is, in part, dictated by the availability and/or expected longevity of the semen (Aurich 2012).

As the longevity of frozen equine semen is decreased, AI is usually undertaken as close as possible to ovulation. The increasing use of low doses of frozen semen has led to most mares being inseminated post-ovulation, with semen deposition at the tip of the horn ipsilateral to ovulation.

For precise determination of follicular development and ovulation for AI in rhinoceroses and elephants, daily ultrasound is the only feasible option in trained pachyderms in captivity. Like in the mare, the pre-ovulatory follicle changes shape from round to more

flaccid (Fig. 5A, C and E), and obvious uterine tone (convoluted uterine appearance) and endometrial oedema (Fig. 5B, D and F) sometimes with intraluminal fluid accumulation occur close to ovulation. Fresh semen AI has been performed when a dominant follicle was >2.0 cm in Asian and African elephants (Brown *et al.* 2004a, Thongtip *et al.* 2009) and >3.5 cm in white rhinoceroses (Hermes *et al.* 2009b).

Control of the oestrous cycle via exogenous hormones has not yet been described for elephants and appears challenging given the two LH peaks necessary to prepare the ovary with sufficient luteal structures to maintain pregnancy and with the first LH peak needed to provide inhibin secretion from the LUFs. An attempt to induce the second LH peak by administering a GnRH agonist intravenously during the late, anovulatory follicular phase, resulted in a low non-ovulatory LH peak and a natural, second LH peak and/or ovulation 15–22 days later (Thitaram *et al.* 2008).

In rhinoceroses, particularly white rhinoceroses, oestrous induction or synchronisation may be carried out with a combination of oral progestagen supplementation for at least 30 days followed by GnRH injections or implants (Hermes *et al.* 2006, Pennington & Durrant 2018, Roth *et al.* 2018).

Practicalities of AI

During natural mating in elephants, the semen is deposited intra-vaginally, whereas in the mare most of the ejaculate ends up in the uterus, even though the tip of the stallion penis does not penetrate the cervix. In the rhinoceros, due to the shape of the penis and the structure of the cervix, only a fraction of the ejaculate may reach the cervical canal.

For AI, the mare's short cervix is easily passed during oestrus making intra-uterine insemination easy and straightforward. In contrast, the white rhinoceros' cervix is quite long and consists of three to five tight interdigitated folds of fibrous connective tissue leaving a very thin lumen (Fig. 1F and 2A), which can only be passed during oestrus using a species-specific semi-rigid AI-catheter to allow intra-uterine semen deposition under standing sedation or general anaesthesia. Reported sperm doses were 5800 and $10,560 \times 10^6$ total sperm for fresh (Hildebrandt *et al.* 2007) and approx. 500×10^6 motile sperm for frozen-thawed semen (Hermes *et al.* 2009b).

In elephants, the unique anatomical features complicate AI. Although the cervical canal is short with folds like those in the mare, the long vestibule and the

hymen-like structure, which narrows the vestibule-vaginal opening to <1 cm in nulliparous females (Figs. 2C, D, E and F), have to be passed to deposit semen as close as possible to the cervix (Hermes *et al.* 2007). Initially, a flexible tube was guided through the curvature of the urogenital canal (Balke *et al.* 1988), but endoscopic guidance to visualise the vaginal os (Fig. 2C, D, E and F) with confirmation of AI catheter placement into the vagina by trans-rectal ultrasound is now commonplace (Brown *et al.* 2004, Thongtip *et al.* 2009, Hildebrandt *et al.* 2012). In African elephants, AIs with frozen-thawed semen were successful when total doses of 720×10^6 motile sperm/AI were used (Hildebrandt *et al.* 2012).

Surgical elephant AI has also been described, whereby an incision is made ventral of the anus into the proximal part of the vestibule through which a speculum with a light source is inserted and an equine insemination pipette guided through the vaginal os (Schmitt 2006). Bypassing the long vestibulum vaginae by directly entering the pelvic part gives straight access to the vagina and cervix at arms-length. The necessary postsurgical care of the incision wound leaves the use of this technique for routine, fresh semen AI impractical and questionable. For deep intrauterine AI with small doses of frozen-thawed semen, embryo recovery or ET, it may provide an interesting option.

Advanced art – future perspectives

Embryo recovery and transfer

In the mare, embryos are generally flushed from the uterus on day 7 or 8 post-ovulation (day 0=ovulation). Embryos do not enter the uterus until late day 6 and rapid expansion of the blastocoele means that by late day 9 or 10, embryos are more fragile and easily damaged using commercial flushing equipment. Embryo recovery is undertaken using a sterile flexible two-way 24Fr embryo flushing catheter which is inserted through the dioestrous cervix. The uterus is infused with 1–2 L of a physiological flushing medium (e.g. lactated Ringer solution) and following gentle trans-rectal massage of the uterus, the flushing medium is recovered by gravity through an in-line embryo filter. Searching for the embryo is done using a binocular dissecting microscope. Equine embryos are spherical (200–800 μm diameter at days 7 and 8) at this stage and easily visualised. Recovered embryos can be transferred trans-cervically using a non-surgical technique into a synchronised surrogate mare with a >80% success rate (Oguri & Tsutsumi 1972, Wilsher & Allen 2004).

The mare also boasts an unusually wide window for synchrony between donor and recipient with +1 to –5 days with respect to the donor achieving acceptable pregnancy rates (Stout 2006).

The timing of entry of the embryo into the uterus in the elephant and rhinoceros can currently only be speculated upon, so if embryo recovery, either for transfer or cryopreservation, is to progress in pachyderms, the timing of entry of the embryo into the uterus will need to be determined for these two species. As mentioned previously, in the white rhinoceros female, a clear embryonic vesicle (EV) is detectable at a similar time (15 days) to the mare (Radcliffe *et al.* 1997), although it is possible to visualise the EV ultrasonographically in some mares as early as day 9 post-ovulation. In elephants, an ultrasonographically detectable, tiny EV appeared no earlier than 42 days post-ovulation when elephants were scanned every other day (Lueders *et al.* 2010a, 2012).

From maturation of rhinoceros oocytes *in vitro*, we know that early embryonic development is slower than in the horse (Galli *et al.* 2007), but this may not translate to a longer oviductal transport time in this species as the timing of first cleavage, subsequent cell divisions and oviductal transport varies between species in general. Trials for embryo flushing would probably need to take place on days 7 and 8 post-ovulation in the rhinoceros and not before day 14 post-ovulation or potentially even later in elephants.

In pachyderms, close and repeated monitoring of the reproductive cycle and/or hormonal manipulation to synchronise recipients will only be practical in captive animals. In the rhinoceros family, different cycle lengths, the occurrence of long and short oestrous cycles in white rhinoceros individuals (Pennington *et al.* 2019) as well as endocrine differences, complicate the development of reliable and universal synchronisation protocols. In the elephant, the few cycles that occur throughout a year and two LH peaks with an exact timing make synchronisation with exogenous hormones challenging to say the least. However, the double LH peak in elephants does on the other hand allow for accurate timing of natural heat and ovulation, meaning that cryopreservation of recovered embryos would appear to be a more promising option, with subsequent transfer to a recipient elephant that has ovulated following a natural oestrous cycle.

Embryo transfer equipment for rhinoceros cows could involve using Wilsher forceps developed for the non-surgical mare ET (Card 2018) to fixate and straighten the cervix and a specially designed semi-rigid catheter to overcome the long folded cervix through which a smaller

diameter ET catheter may be passed to dispose fresh or frozen–thawed embryos. In the elephant, a minimally invasive access through the cranial part of the vestibulum as described previously for AI (Schmitt 2006) may be an option to ensure access to the cervix and placement of the embryo into the ipsilateral horn of ovulation.

Oocyte recovery and ovum pick-up (OPU)

In the mare, oocytes can be harvested from ovaries recovered *post-mortem* (genetic salvage). This method has been used to good effect to produce live offspring in the horse and protocols for recovery and transport of ovaries are well established. For example, Hatzel *et al.* (2021) reported a 19% blastocyst rate from 620 oocytes that reached metaphase II from 1524 oocytes harvested from 168 sets of ovaries. Results are usually lower than when oocytes are recovered via follicle puncture and aspiration in live animals, because compromised mares (e.g., those with colic that are euthanised) are invariably the majority in which genetic salvage is undertaken. Likewise, *post-mortem* oocyte recovery has been reported in black and Sumatran rhinoceroses (Stoops *et al.* 2011a,b), and in the Asian (Hermes *et al.* 2007) but not in the African elephant, although the African elephant ovarian cortex and folliculogenesis have been described in detail (Stansfield *et al.* 2011) and ovarian tissue preservation has been tested successfully (Gunasena *et al.* 1998).

Oocyte recovery from live animals requires specific equipment and skills. Transvaginal OPU in the horse was first reported in the early 1990s (Brück *et al.* 1992) and has gained interest in large commercial breeding programs over the last decade (Stout 2020). Due to the lack of commercially available gonadotropins to stimulate the development of multiple preovulatory follicles, ultrasound-guided transvaginal aspiration of immature oocytes is performed in the standing sedated mare. Immature equine cumulus–oocyte complexes (COCs) in the mare are anchored to the follicle wall (Hawley *et al.* 1995), so successful recovery of oocytes requires vigorous scraping of the follicular wall and repeated flushing of the follicle with a double-lumen needle (12 gauge). Even with this approach, the oocyte recovery rates for punctured follicles are considered good if $\geq 50\%$ (Galli *et al.* 2007).

OPU by laparoscopy or transvaginal ultrasound guidance is impractical in white rhinoceroses and elephants due to their body size and anatomical features, including the distance from the cranial vagina to the cranial located ovaries (1.2–1.4 m from the anus in rhinoceroses – Hermes *et al.* 2007, Hildebrandt *et al.* 2018).

To overcome the anatomical problems of the transvaginal approach, transrectal ultrasound-guided OPU has been used with success in black and white rhinoceroses leading to the recovery of multiple oocytes (Hermes *et al.* 2007, 2009a). Different devices have been developed to facilitate the procedure (Hildebrandt *et al.* 2007, 2018, Hermes *et al.* 2009a). One device was described as 1.5 m long with a double-lumen needle system at an angle to the instrument axis, which could be guided by ultrasound through the cleaned and disinfected rectal wall into the ovarian tissue. Follicles identified ultrasonographically with a diameter of at least 1 cm (Hildebrandt *et al.* 2018) are aspirated and flushed by repeated refilling of the follicle with a commercial bovine or equine flushing media through the outer needle channel, and aspiration of fluid through the inner channel (Hildebrandt *et al.* 2018). The aspirated fluid is collected into sterile tubes and kept at 37°C with the excess passed through a nylon mesh filter before the remainder is searched for recovered oocytes with a stereomicroscope. Recovered immature oocytes are washed and placed in holding media prior to undergoing *in vitro* maturation (IVM). Recovery rates of <30% per aspirated follicle have been reported (Hildebrandt *et al.* 2018), which is lower than in the horse but may improve with further modification of the technique.

Successful OPU has not been described for elephants to date and comes with additional challenges. Similar to the rhinoceros cow, the reproductive tract of the female elephant is too large to undertake OPU as in the mare, and a vaginal approach is unfeasible due to the anatomy of the vestibulum. Therefore, transrectal, ultrasound-guided follicle aspiration would be the method of choice using a similar set-up to that described for rhinoceroses (Hildebrandt *et al.* 2018). OPU in elephants may be complicated by the thick rectal wall, the ovarian bursa covering the ovary (Fig. 2B) and the small size and number of antral follicles (1–15 mm), which are only present during the follicular phase prior to and between the two LH surges (Lueders *et al.* 2010b).

Superovulation protocols as used in cattle, but ineffective in the mare, may be worth investigating to potentially produce more/larger follicles for aspiration. In one report, white rhinoceroses were treated with a GnRH analogue to boost follicle growth. Intramuscular injections of 3.0 mL Histrelin (0.5 mg/mL BioRelease®, Bet Pharm LLC) were given every other day three to four times and OPU performed 24 or 48 h after the last injection (Hildebrandt *et al.* 2018). Superovulation protocols for elephants have not been considered and it is not known

which exogenous hormone would stimulate elephant follicular development.

In vitro maturation (IVM)

In humans, where adequate superstimulation and superovulation allow for the collection of mature oocytes, IVM is not routinely used. However, in the mare, IVM of oocytes is the first step required prior to undertaking IVF or intra-cytoplasmic sperm injection (ICSI) and the production of an embryo. With the significant increase in the popularity of ICSI in horses and, hence, the need for IVM (Claes & Stout 2022), commercial IVM media are now available and may prove suitable for the rhinoceros or elephant. In the horse, shipping of immature oocytes collected via OPU at 20–22°C to specialised ICSI centres can be easily accomplished. One review gives the average maturation rate of equine oocytes shipped to an ICSI lab as 59% (Claes & Stout 2022), which usually involves culture for 24–30 h for viable oocytes to reach metaphase II (attain nuclear maturation).

Developing effective protocols and maturation media for IVM in new species, like the rhinoceros, has been described as difficult, especially when there is limited availability of oocytes (Pennington & Durrant 2018). Nevertheless, cell culture media (e.g. Medium 199; Dulbecco's Modified Eagle's Medium -F12) supplemented with porcine or ovine luteinising hormone FSH, oestradiol, and insulin-like growth factor 1 (Stoops *et al.* 2011b), southern white rhinoceros follicular fluid (Pennington & Durrant 2018) and white rhinoceros oestrus serum (Hermes *et al.* 2009a, Hildebrandt *et al.* 2018, Ruggeri *et al.* 2022), have been used as IVM media. In a recent study, a media containing FSH, LH, oestradiol, somatotropin and horse follicular fluid achieved better maturation results compared to white rhinoceros oestrous serum (Ruggeri *et al.* 2022).

Rhinoceros oocytes have been typically incubated at 37.5–38.5°C in a humidified atmosphere containing 5% CO₂ for 32–44 h prior to being denuded and checked for a polar body to confirm if they had reached metaphase II (Hermes *et al.* 2009a, Stoops *et al.* 2011b, Hildebrandt *et al.* 2018, Pennington & Durrant 2018). In the few existing reports on IVM of rhinoceros oocytes, maturation rates were low (<30%) for *post-mortem* collected Sumatran and black rhinoceros oocytes after culture times of 24–32 h (Stoops *et al.* 2011a,b) and <40% for oocytes recovered by OPU from White rhinoceroses after culture times of 36–44 h (Hildebrandt *et al.* 2018).

In the Asian elephant, IVM of *post-mortem* collected oocytes with a success rate of 60% was reported (Hermes *et al.* 2007).

IVF and ICSI

IVF and ICSI of oocytes for embryo production have been described with success in many species including the horse (Hinrichs 2018).

In the horse, ICSI as opposed to IVF is the method of choice for *in vitro* production of embryos, as only very recently has conventional IVF produced efficient and repeatable results following prolonged sperm incubation to induce its capacitation (Felix *et al.* 2022). IVF has been attempted in black and white rhinoceroses for more than a decade, but to date no blastocysts have been produced (Pennington & Durrant 2018). In black rhinoceroses, IVF of *post-mortem* collected oocytes with frozen–thawed epididymal semen collected *post-mortem*, resulted in a two-cell embryo (Stoops *et al.* 2011b), while a four-cell embryo was obtained from an oocyte collected via transrectal OPU fertilised with fresh, chilled semen (Hermes *et al.* 2009a).

After maturation, COCs were rinsed and incubated in a humidified atmosphere of 5% CO₂ in air for 1–4 h at 39°C or for 6.5–48 h at 38.6°C before co-incubation with sperm in IVF media (synthetic oviduct fluid supplemented with heparin (20 mg/mL) and 5% (white) rhinoceros oestrous serum (Hermes *et al.* 2009a)) or IVF–TALP (Stoops *et al.* 2011b), respectively. Homologous and heterologous (Indian rhinoceros) frozen–thawed sperm were capable of binding, penetrating and fertilising black rhinoceros oocytes recovered from *post-mortem* ovaries (Stoops *et al.* 2011b).

Results from ICSI in horses do vary between laboratories, but laboratories with skilled technicians can produce good results. For example, from a total of 515 OPU session in 2021, in which recovered oocytes were shipped overnight to a specialist laboratory, there was a 78% chance of obtaining one or more blastocysts per session following ICSI, with a mean of 2.1 embryos per session. Approximately 60% of transferred embryos result in a live foal (Claes & Stout 2022).

In rhinoceroses, ICSI was performed by injecting a single-frozen–thawed motile sperm of northern and southern white rhinoceroses, with normal morphology and selected through the swim-up procedure, into the cytoplasm of southern white rhinoceros' oocytes using a piezo-driven micromanipulator (Hildebrandt *et al.* 2018). COCs were treated with hyaluronidase and the cumulus

cells were mechanically removed after 36 h of culture prior to ICSI. Injected oocytes were cultured in a modified SOF medium supplemented with BSA and MEM amino acids until blastocyst development after 9–12 days. Of 32 injected oocytes, 12 underwent cleavage and 7 developed to blastocysts (Hildebrandt *et al.* 2018).

Oocyte and embryo cryopreservation

Although live offspring have been produced from cryopreserved oocytes in the mare (De Coster *et al.* 2019), blastocyst rates remain poor and the technique is not routinely undertaken in clinical practice.

Oocyte cryopreservation in rhinoceroses and elephants is likely to be equally challenging, and current protocols are insufficient for routine application (Pereira & Marques 2008). African elephant ovarian tissue has been successfully cryopreserved (Gunasena *et al.* 1998), but to date oocytes have only been obtained at *post-mortem*. Although it was recently reported that *intra vitam*-aspirated oocytes of rhinoceroses were frozen using a conventional freezing method (Pennington & Durrant 2018), information on their post-thaw quality and viability has not been published.

Embryos, although sensitive to chilling, freezing and thawing injury (Pereira & Marques 2008), offer more prospect as a means to cryopreserve genetics than oocytes do. Historically, embryo cryopreservation in the horse was problematic. The reasons typically cited for this were the late entry of the embryo into the uterus when it has already begun to blastulate, the quick development of a large blastocoele cavity and inhibition of the passage of cryoprotectants by the equine capsule. Hence, only morulae or early blastocysts $\leq 300 \mu\text{m}$ were originally a prospect for slow freezing or vitrification with any degree of success (Eldridge-Panuska *et al.* 2005). More recently, considerable success has been achieved in vitrifying larger embryos by micro-manipulator-assisted or manual puncture of the blastocoele cavity and its collapse (Wilsher *et al.* 2019, 2021) with pregnancy rates upon warming and transfer only marginally below or equivalent to those reported for transfer of fresh embryos. For *in vivo*-produced equine embryos, vitrification would appear to be the method of choice, whereas *in vitro* (ICSI)-produced ones, with their smaller blastocoele cavity and poorly developed capsule, have excellent survival rates when slow freezing or vitrification methods are used (Claes & Stout 2022, Spanner *et al.* 2022).

To date, no early elephant embryos have been recovered/produced and thus, their cryopreservation

properties remain unknown. However, in recent years, equine reproduction specialist Dr Cesare Galli succeeded in producing white rhinoceros blastocysts from *in vivo* derived oocytes and frozen-thawed rhinoceros semen which were subsequently cryopreserved using equine-based protocols (Hildebrandt *et al.* 2018). Blastocysts with a clear detectable inner cell mass (11–13 days after successful ICSI), were cryopreserved in 0.25 mL straws after equilibration in media containing glycerol (5% for 5 min followed by 10% for 10 min). A slow-freezing vitrification protocol developed for cattle and horse embryos was followed using a freezing machine that allowed a controlled freezing curve to be achieved, prior to their long-term storage in liquid nitrogen (Hildebrandt *et al.* 2018).

The recovery of *in vivo*-derived embryos from both rhinoceroses and elephants will require the development of embryo recovery techniques in these species. The challenge in doing this lies with catheterising the cervix and, as previously mentioned, determining the day of entry of the embryo into the uterus. If these hurdles can be overcome, the technique should, in theory, be achievable. Furthermore, given the spherical nature of the rhinoceros and elephant embryonic vesicle, it is interesting to speculate that an embryonic capsule may also be present in these species. If this is indeed the case, the work already undertaken in the horse to successfully cryopreserve capsulated embryos could potentially be feasible for rhinoceros and elephant embryos, too.

Tissue and stem cell preservation and cloning

Cryopreservation of ovarian tissue has been suggested as an alternative for oocyte and embryo preservation, since primordial follicles respond better to the challenges of cryopreservation and therefore more generic protocols appear to be successful in freezing ovarian tissue (Gunasena *et al.* 1998, Santos *et al.* 2010). In African elephants, pieces of ovarian tissue (1–2 cm³) were preserved under field conditions and xenografted into mice where antral follicles developed in the elephant ovarian tissue graft (Gunasena *et al.* 1998).

In vitro culture of fibroblasts from tissue can be used to establish and cryopreserve somatic cell lines and establish induced pluripotent stem cells (iPSCs), which has been achieved for the northern white rhinoceros (Ben-Nun *et al.* 2011, Saragusty *et al.* 2016a, Hildebrandt *et al.* 2018, Korody *et al.* 2017). Like embryonic stem cells, iPSCs have the potential to differentiate into gametes, which could then be used for IVF or ICSI (Saragusty *et al.* 2016a). In white rhinoceroses, stem cell derivation from

embryos (Hildebrandt *et al.* 2018) and the differentiation of spermatogonial stem cells, isolated from testicular tissue, into mature spermatozoa, have been reported (Gomez *et al.* 2018). Cloning by interspecies somatic cell nuclear transfer (iSCNT), is especially interesting for (functionally) extinct species, like the northern white rhinoceros or the mammoth (Saragusty *et al.* 2016b). iSCNT was successfully used to produce blastocysts from Asian elephant's fibroblast cell lines, established from ear tissues and porcine oocytes. However despite containing the elephant genome, when transferred into sows these embryos appeared non-viable (Nguyen *et al.* 2022) and it appears questionable if these blastocysts were truly elephants.

Conclusion

Although the complex problems wild elephant and rhinoceros populations are facing cannot be solved with ART alone, these techniques certainly have a role to play in contributing to captive breeding programmes. Main advantages of implementing these techniques are the ability (i) to extend the individual reproductive lifespan by gamete preservation, (ii) to enhance genetic diversity by introducing individuals that are not reproducing naturally and (iii) to exchange genetics without moving entire animals which may come with welfare issues and disease risks.

In horses, ART has advanced tremendously in recent years and rhinoceros and elephant ART are likely to profit from the establishment of new protocols and the successes in equids. Basic applications, such as semen collection and preservation, and AI are currently applicable to elephants and rhinoceroses, whereas more advanced methods need further research. The recent achievements in rhinoceros OPU, oocyte maturation, fertilisation and embryo cryopreservation show that even in large pachyderms, these techniques can be successfully implemented. Although we are far from achieving consistent results, in white rhinoceroses the mare has served as a model, specifically for laboratory protocols. In elephant advanced ART, we may need to overcome additional obstacles.

Problems with availability of study animals, the handling of large, and thus potentially dangerous mammals, and sufficient funding remain challenging. Clever adaptations to routine equine protocols and equipment are necessary and can be feasible, as discussed and shown in this review. We need to attract a larger community and colleagues from the domestic animal

reproductive disciplines to assist in this elephant-sized task.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the manuscript.

Funding

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Author contribution statement

All authors reviewed the literature and contributed to the preparation of the manuscript. JMB and IL prepared the figures.

Acknowledgements

This review is dedicated to the late Professor WR (Twink) Allen for his contributions to understanding reproduction in the horse and elephant, who sadly died before the manuscript was submitted for publication.

References

- Allen WR** 2001 Fetomaternal interactions and influences during equine pregnancy. *Reproduction* **121** 513–527. (<https://doi.org/10.1530/rep.0.1210513>)
- Allen WR** 2010 Sex, science and satisfaction: a heady brew. *Animal Reproduction Science* **121** S262–S278. (<https://doi.org/10.1016/j.anireprosci.2010.04.178>)
- Allen WR & Moor RM** 1972 The origin of the equine endometrial cups. I. Production of PMSG by fetal trophoblast cells. *Journal of Reproduction and Fertility* **29** 313–316. (<https://doi.org/10.1530/jrf.0.0290313>)
- Allen WRT & Stansfield FJ** 2021 Placentation in the African elephant (*Loxodonta africana*). *Advances in Anatomy, Embryology, and Cell Biology* **234** 181–204. (https://doi.org/10.1007/978-3-030-77360-1_9)
- Allen WR, Mathias S, Wooding FBP & van Aarde RJ** 2003 Placentation in the African elephant (*Loxodonta africana*): II morphological changes in the uterus and placenta throughout gestation. *Placenta* **24** 598–617. ([https://doi.org/10.1016/s0143-4004\(03\)00102-4](https://doi.org/10.1016/s0143-4004(03)00102-4))
- Amin R, Thomas K, Emslie RH, Foote TJ & Strien NV** 2006 An overview of the conservation status of and threats to rhinoceros species in the wild. *International Zoo Yearbook* **40** 96–117. (<https://doi.org/10.1111/j.1748-1090.2006.00096.x>)
- Arnold D** 2014 *Successful Cryopreservation of Asian Elephant (Elephas maximus) Semen Using Simple Techniques*. Masters thesis. University of Guelph, Ontario, Canada
- Aurich C** 2008 Recent advances in cooled-semen technology. *Animal Reproduction Science* **107** 268–275. (<https://doi.org/10.1016/j.anireprosci.2008.04.015>)
- Aurich C** 2011 Reproductive cycles of horses. *Animal Reproduction Science* **124** 220–228. (<https://doi.org/10.1016/j.anireprosci.2011.02.005>)
- Aurich JE** 2012 Artificial insemination in horses—more than a century of practice and research. *Journal of Equine Veterinary Science* **32** 458–463. (<https://doi.org/10.1016/j.jevs.2012.06.011>)
- Balke JME, Boever WJ, Ellersieck MR, Seal US & Smith DA** 1988 Anatomy of the reproductive tract of the female African elephant (*Loxodonta africana*) with reference to development of techniques for artificial breeding. *Journal of Reproduction and Fertility* **84** 485–492. (<https://doi.org/10.1530/jrf.0.0840485>)
- Battut I, Colchen S, Fieni F, Tainturier D & Bruyas JF** 1998 Success rates when attempting to nonsurgically collect equine embryos at 144, 156 or 168 hours after ovulation. *Equine Veterinary Journal. Supplement* **30** 60–62. (<https://doi.org/10.1111/j.2042-3306.1997.tb05102.x>)
- Benirschke K & Lowenstine LJ** 1995 The placenta of the Rhinocerotidae. In *Internationales Symposium über die Erkrankungen der Zoo- und Wildtiere*. Dresden, Deutschland: Institut für Zoo- und Wildtierforschung. Verhandlungsbericht Erkrankungen der Zootiere. 15–23.
- Ben-Nun IF, Montague SC, Houck ML, Tran HT, Garitaonandia I, Leonardo TR, Wang YC, Charter SJ, Laurent LC, Ryder OA, *et al.*** 2011 Induced pluripotent stem cells from highly endangered species. *Nature Methods* **8** 829–831. (<https://doi.org/10.1038/nmeth.1706>)
- Berkeley EV, Kirkpatrick JF, Schaffer NE, Bryant WM & Threlfall WR** 1997 Serum and fecal steroid analysis of ovulation, pregnancy and parturition in the black rhinoceros (*Diceros bicornis*). *Zoo Biology* **16** 121–132. ([https://doi.org/10.1002/\(SICI\)1098-2361\(1997\)16:2<121::AID-ZOO3>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1098-2361(1997)16:2<121::AID-ZOO3>3.0.CO;2-C))
- Betteridge KJ** 2000 Comparative aspects of equine embryonic development. *Animal Reproduction Science* **60–61** 691–702. ([https://doi.org/10.1016/s0378-4320\(00\)00075-0](https://doi.org/10.1016/s0378-4320(00)00075-0))
- Brown JL, Schmitt DL, Bellem A, Graham LH & Lehnhardt J** 1999 Hormone secretion in the Asian elephant (*Elephas maximus*): characterization of ovulatory and anovulatory luteinizing hormone surges. *Biology of Reproduction* **61** 1294–1299. (<https://doi.org/10.1095/biolreprod61.5.1294>)
- Brown JL, Bellem AC, Fouraker M, Wildt DE & Roth TL** 2001 Comparative analysis of gonadal and adrenal activity in the black and white rhinoceros in north America by noninvasive endocrine monitoring. *Zoo Biology* **20** 463–486. (<https://doi.org/10.1002/zoo.10028>)
- Brown JL, Goritz F, Pratt-Hawkes N, Hermes R, Galloway M, Graham LH, Gray C, Walker SL, Gomez A, Moreland R, *et al.*** 2004 Successful artificial insemination of an Asian elephant at the national zoological park. *Zoo Biology* **23** 45–63. (<https://doi.org/10.1002/zoo.10116>)
- Brück I, Raun K, Synnestvedt B & Greve T** 1992 Follicle aspiration in the mare using a transvaginal ultrasound-guided technique. *Equine Veterinary Journal* **24** 58–59. (<https://doi.org/10.1111/j.2042-3306.1992.tb02780.x>)
- Card C** 2018 Non-surgical embryo transfer technique and recipient mare pregnancy rate. *Veterinary Record* **183** 320–322. (<https://doi.org/10.1136/vr.k3700>)
- Claes A & Stout TAE** 2022 Success rate in a clinical equine in vitro embryo production program. *Theriogenology* **187** 215–218. (<https://doi.org/10.1016/j.theriogenology.2022.04.019>)
- Comizzoli P** 2015 BioBanking efforts and new advances in male fertility preservation for rare and endangered species. *Asian Journal of Andrology* **17** 640–645. (<https://doi.org/10.4103/1008-682X.153849>)
- Daels PF, DeMoraes JJ, Stabenfeldt GH, Hughes JP & Lasley BL** 1991 The corpus luteum: source of oestrogen during early pregnancy in the mare. *Journal of Reproduction and Fertility. Supplement* **44** 501–508.
- De Coster T, Velez DA, Van Soom A, Woelders H & Smits K** 2019 Cryopreservation of equine oocytes: looking into the crystal ball.

- Reproduction, Fertility, and Development* **32** 453–467. (<https://doi.org/10.1071/RD19229>)
- Eldridge-Panuska WD, di Brienza VC, Seidel GE, Squires EL & Carnevale EM** 2005 Establishment of pregnancies after serial dilution or direct transfer by vitrified equine embryos. *Theriogenology* **63** 1308–1319. (<https://doi.org/10.1016/j.theriogenology.2004.06.015>)
- Ellenberger C, Wilsher S, Allen WR, Hoffman C, Kölling M, Bazer FW, Klug J, Schoon D & Schoon HA** 2008 Immunolocalisation of the uterine secretory proteins uterocalin, uteroferrin and uteroglobulin in the mare's uterus and placenta throughout gestation. *Theriogenology* **70** 746–757. (<https://doi.org/10.1016/j.theriogenology.2008.04.050>)
- Felix MR, Turner RM, Dobbie T & Hinrichs K** 2022 Successful in vitro fertilization in the horse: production of blastocysts and birth of foals after prolonged sperm incubation for capacitation. *Biology of Reproduction* **107** 1551–1564. (<https://doi.org/10.1093/biolre/iaoc172>)
- Galli C, Colleoni S, Duchi R, Lagutina I & Lazzari G** 2007 Developmental competence of equine oocytes and embryos obtained by in vitro procedures ranging from in vitro maturation and ICSI to embryo culture, cryopreservation and somatic cell nuclear transfer. *Animal Reproduction Science* **98** 39–55. (<https://doi.org/10.1016/j.anireprosci.2006.10.011>)
- Ginther OJ** 1983 Mobility of the early equine conceptus. *Theriogenology* **19** 603–611. ([https://doi.org/10.1016/0093-691x\(83\)90180-2](https://doi.org/10.1016/0093-691x(83)90180-2))
- Ginther OJ** 1992 *Reproductive Biology of the Mare, Basic and Applied Aspects*, 2nd ed. Equiservices: Cross Plains.
- Ginther OJ, Gastal EL, Gastal MO & Beg MA** 2007 Incidence, endocrinology, vascularity, and morphology of hemorrhagic anovulatory follicles in mares. *Journal of Equine Veterinary Science* **27** 130–139. (<https://doi.org/10.1016/j.jevs.2007.01.009>)
- Godfrey RW, Pope CE, Dresser BL & Olsen JH** 1991 Gross anatomy of the reproductive tract of female black *Diceros bicornis michaeli* and white rhinoceros *Ceratotherium simum simum*. *Zoo Biology* **10** 165–175. (<https://doi.org/10.1002/zoo.1430100208>)
- Gomez MC, Cates Y, Stansfield DB, Young C, Klee R & Durrant B** 2018 187 Morphological appearance and expression of spermatogonial stem cell markers in white rhinoceros testicular tissue. *Reproduction, Fertility and Development* **30** 233. (<https://doi.org/10.1071/RDv30n1Ab187>)
- Gunasena KT, Lakey JRT, Villines PM, Bush M, Raath C, Critser ES, McGann LE & Critser JK** 1998 Antral follicles develop in xenografted cryopreserved african elephant (*Loxodonta africana*) ovarian tissue. *Animal Reproduction Science* **53** 265–275. ([https://doi.org/10.1016/S0378-4320\(98\)00132-8](https://doi.org/10.1016/S0378-4320(98)00132-8))
- Hatzel JN, Lederman J & Stokes J** 2021 Equine post-mortem oocyte recovery: a retrospective analysis. *Proceeding of American Association Equine Practitioners* **67** 266–267.
- Hawley LR, Enders AC & Hinrichs K** 1995 Comparison of equine and bovine oocyte-cumulus morphology within the ovarian follicle. *Biology of Reproduction* **52** 243–252. (https://doi.org/10.1093/biolreprod/52.monograph_series1.243)
- Heap R, Hamon M & Allen W** 1982 Studies on the oestrogen synthesis by the preimplantation equine conceptus. *Journal of Reproduction and Fertility. Supplement* **32** 343–352.
- Hermes R & Hildebrandt TB** 2011 Rhinoceros theriogenology, Fowler's Zoo and Wild Animal Medicine: current Therapy. Elsevier Inc. (<https://doi.org/10.1016/b978-1-4377-1986-4.00071-8>)
- Hermes R, Hildebrandt TB, Blottner S, Walzer C, Silinski S, Patton ML, Wibbelt G, Schwarzenberger F & Göritz F** 2005 Reproductive soundness of captive southern and northern white rhinoceroses (*Ceratotherium simum simum*, C.s. cottoni): Evaluation of male genital tract morphology and semen quality before and after cryopreservation. *Theriogenology* **63** 219–238. (<https://doi.org/10.1016/j.theriogenology.2004.04.007>)
- Hermes R, Hildebrandt TB, Walzer C, Göritz F, Patton ML, Silinski S, Anderson MJ, Reid CE, Wibbelt G, Tomasova K, et al.** 2006 The effect of long non-reproductive periods on the genital health in captive female white rhinoceroses (*Ceratotherium simum simum*, C.s. cottoni). *Theriogenology* **65** 1492–1515. (<https://doi.org/10.1016/j.theriogenology.2005.09.002>)
- Hermes R, Göritz F, Streich WJ & Hildebrandt TB** 2007 Assisted reproduction in female rhinoceros and elephants - Current status and future perspective. *Reproduction in Domestic Animals* **42**(Supplement 2) 33–44. (<https://doi.org/10.1111/j.1439-0531.2007.00924.x>)
- Hermes R, Göritz F, Portas TJ, Bryant BR, Kelly JM, Maclellan LJ, Keeley T, Schwarzenberger F, Walzer C, Schnorrenberg A, et al.** 2009a Ovarian superstimulation, transrectal ultrasound-guided oocyte recovery, and IVF in rhinoceros. *Theriogenology* **72** 959–968. (<https://doi.org/10.1016/j.theriogenology.2009.06.014>)
- Hermes R, Göritz F, Saragusty J, Sós E, Molnar V, Reid CE, Schwarzenberger F & Hildebrandt TB** 2009b First successful artificial insemination with frozen-thawed semen in rhinoceros. *Theriogenology* **71** 393–399. (<https://doi.org/10.1016/j.theriogenology.2008.10.008>)
- Hermes R, Hildebrandt TB, Walzer C, Göritz F, Gray C, Niemuller C & Schwarzenberger F** 2012 Estrus induction in white rhinoceros (*Ceratotherium simum*). *Theriogenology* **78** 1217–1223. (<https://doi.org/10.1016/j.theriogenology.2012.05.015>)
- Hermes R, Saragusty J, Göritz F, Bartels P, Potier R, Baker B, Streich WJ & Hildebrandt TB** 2013 Freezing African elephant semen as a new population management tool. *PLoS One* **8** e57616. (<https://doi.org/10.1371/journal.pone.0057616>)
- Hermes R, Hildebrandt TB & Göritz F** 2018 Cryopreservation in rhinoceros—setting a new benchmark for sperm cryosurvival. *PLoS One* **13** e0200154. (<https://doi.org/10.1371/journal.pone.0200154>)
- Hildebrandt TB, Göritz F, Pratt NC, Schmitt DL, Quandt S, Raath J & Hofmann RR** 1998 Reproductive assessment of male elephants (*Loxodonta africana* and *Elephas maximus*) by ultrasonography. *Journal of Zoo and Wildlife Medicine* **29** 114–128.
- Hildebrandt TB, Göritz F, Hermes R, Schmitt DL, Brown JL, Schwammer H, Loskutoff N, Pratt NC, Lehnhardt JL, Montali RJ, et al.** 1999 Artificial insemination of African (*Loxodonta africana*) and Asian (*Elephas maximus*) elephants. Proceedings of American Association Zoo Veterinarians, Columbus, OH, USA, pp. 83–86.
- Hildebrandt TB, Hermes R, Pratt NC, Fritsch G, Blottner S, Schmitt DL, Ratanahkorn P, Brown JL, Rietschel W & Göritz F** 2000 Ultrasonography of the urogenital tract in elephants (*Loxodonta africana* and *Elephas maximus*): an important tool for assessing male reproductive function. *Zoo Biology* **19** 333–345. ([https://doi.org/10.1002/1098-2361\(2000\)19:5<333::AID-ZOO5>3.0.CO;2-A](https://doi.org/10.1002/1098-2361(2000)19:5<333::AID-ZOO5>3.0.CO;2-A))
- Hildebrandt TB, Ritz FG, Hermes R, Reid C, Dehnhard M & Brown JL** 2006 Aspects of the reproductive biology and breeding management of Asian and African elephants, *Elephas maximus*, *Loxodonta africana*. *International Zoo Yearbook* **40** 20–40. (<https://doi.org/10.1111/j.1748-1090.2006.00020.x>)
- Hildebrandt TB, Hermes R, Walzer C, Sós E, Molnar V, Mezösi L, Schnorrenberg A, Silinski S, Streich J, Schwarzenberger F, et al.** 2007 Artificial insemination in the anaestrous and the postpartum white rhinoceros using GnRH analogue to induce ovulation. *Theriogenology* **67** 1473–1484. (<https://doi.org/10.1016/j.theriogenology.2007.03.005>)
- Hildebrandt TB, Lueders I, Hermes R, Goeritz F & Saragusty J** 2011 Reproductive cycle of the elephant. *Animal Reproduction Science* **124** 176–183. (<https://doi.org/10.1016/j.anireprosci.2010.08.027>)
- Hildebrandt TB, Hermes R, Saragusty J, Potier R, Schwammer HM, Balfanz F, Vielgrader H, Baker B, Bartels P & Göritz F** 2012 Enriching the captive elephant population genetic pool through artificial insemination with frozen-thawed semen collected in the wild. *Theriogenology* **78** 1398–1404. (<https://doi.org/10.1016/j.theriogenology.2012.06.014>)

- Hildebrandt TB, Hermes R, Colleoni S, Diecke S, Holtze S, Renfree MB, Stejskal J, Hayashi K, Drukker M, Loi P, *et al.*** 2018 Embryos and embryonic stem cells from the white rhinoceros. *Nature Communications* **9** 2589. (<https://doi.org/10.1038/s41467-018-04959-2>)
- Hinrichs K** 2018 Assisted reproductive techniques in mares. *Reproduction in Domestic Animals* **53**(Supplement 2) 4–13. (<https://doi.org/10.1111/rda.13259>)
- Jenikejew J, Wauters J, Dehnhard M & Scheumann M** 2021 Linking socio-sexual and vocal behaviour with faecal progesterone and oestrogenmetabolite levels in Southern white rhinoceros females. *Conservation Physiology* **9** coab098. (<https://doi.org/10.1093/conphys/coab098>)
- Kaewmanee S, Watanabe G, Kishimoto M, Jin WZ, Yamamoto Y, Yamamoto T, Nagaoka K, Narushima E, Komiya T & Taya K** 2011 Secretion of inhibin during the estrous cycle in the female Asian elephant (*Elephas maximus*). *Journal of Veterinary Medical Science* **73** 77–82. (<https://doi.org/10.1292/jvms.10-0267>)
- Kähn W & Volkmann D** 2004 **Ultrasonography in the mare. In Veterinary Reproductive Ultrasonography.** W Kähn & D Volkmann Eds. Hannover: Schlütersche Verlagsgesellschaft mbH, pp. 11–80.
- Kainer RA** 2011 **Internal reproductive anatomy. In Equine Reproduction.** AO McKinnon, EL Squires, WE Vaala & DD Varner Eds. Philadelphia, London: Wiley-Blackwell, pp. 1582–1597.
- Korody ML, Pivaroff C, Nguyen TD, Peterson SE, Ryder OA & Loring J** 2017 Four new induced pluripotent stem cell lines produced from Northern White Rhinoceros with non-integrating reprogramming factors. *Biorxiv*. (<https://doi.org/10.1101/202499>)
- Lueders I & Allen WRT** 2020 Managed wildlife breeding-an undervalued conservation tool? *Theriogenology* **150** 48–54. (<https://doi.org/10.1016/j.theriogenology.2020.01.058>)
- Lueders I, Drews B, Niemuller C, Gray C, Rich P, Fickel J, Wibbelt G, Göritz F & Hildebrandt TB** 2010a Ultrasonographically documented early pregnancy loss in an Asian elephant (*Elephas maximus*). *Reproduction, Fertility, and Development* **22** 1159–1165. (<https://doi.org/10.1071/RD09305>)
- Lueders I, Niemuller C, Gray C, Rich P & Hildebrandt TB** 2010b Luteogenesis during the estrous cycle in Asian elephants (*Elephas maximus*). *Reproduction* **140** 777–786. (<https://doi.org/10.1530/REP-10-0022>)
- Lueders I, Taya K, Watanabe G, Yamamoto Y, Yamamoto T, Kaewmanee S, Niemuller C, Gray C, Streich WJ & Hildebrandt TB** 2011 Role of the double luteinizing hormone peak, luteinizing follicles, and the secretion of inhibin for dominant follicle selection in Asian elephants (*Elephas maximus*). *Biology of Reproduction* **85** 714–720. (<https://doi.org/10.1095/biolreprod.110.090167>)
- Lueders I, Niemuller C, Rich P, Gray C, Hermes R, Goeritz F & Hildebrandt TB** 2012 Gestating for 22 months: luteal development and pregnancy maintenance in elephants. *Proceedings. Biological Sciences* **279** 3687–3696. (<https://doi.org/10.1098/rspb.2012.1038>)
- Meuffels J** 2022 *Improved Methods for Collection and Cryopreservation of Semen from African Rhinoceroses (Ceratotherium simum simum; Dicerus bicornis) and African Elephants (Loxodonta africana)*. Pretoria: University of Pretoria.
- Nguyen TD, Li H, Zhuang Y, Chen B, Kinoshita K, Jamal MA, Xu K, Guo J, Jiao D, Tanabe K, *et al.*** 2022 In vitro and in vivo development of interspecies Asian elephant embryos reconstructed with pig enucleated oocytes. *Animal Biotechnology* 1–10. (<https://doi.org/10.1080/10495398.2022.2058005>)
- Oguri N & Tsutsumi Y** 1972 Non-surgical recovery of equine eggs, and an attempt at non-surgical egg transfer in horses. *Journal of Reproduction and Fertility* **31** 187–195. (<https://doi.org/10.1530/jrf.0.0310187>)
- Patton ML, Swaisgood RR, Czekala NM, White AM, Fetter GA, Montagne JP, Rieches RG & Lance VA** 1999 Reproductive cycle length and pregnancy in the Southern White Rhinoceros (*Ceratotherium simum simum*) as determined by fecal pregnane analysis and observations of mating behavior. *Zoo Biology* **18** 111–127. ([https://doi.org/10.1002/\(SICI\)1098-2361\(1999\)18:2<111::AID-ZOO3>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1098-2361(1999)18:2<111::AID-ZOO3>3.0.CO;2-0))
- Pennington PM & Durrant BS** 2018 Assisted reproductive technologies in captive rhinoceros. *Mammal Review* **49** 1–15.
- Pennington PM, Marshall KL, Capiro JM, Felton RG & Durrant BS** 2019 Ovulation induction in anovulatory southern white rhinoceros (*Ceratotherium simum simum*) without altrenogest. *Conservation Physiology* **7** coz033. (<https://doi.org/10.1093/conphys/coz033>)
- Pennington PM, Marshall KL, Capiro JM, Howard L & Durrant BS** 2020 Pregnancies following long luteal phases in southern white rhinoceros (*Ceratotherium simum simum*). *Zoo Biology* **39** 141–144. (<https://doi.org/10.1002/zoo.21529>)
- Pereira RM & Marques CC** 2008 Animal oocyte and embryo cryopreservation. *Cell Tissue Bank* **9** 267–277. (<https://doi.org/10.1007/s10561-008-9075-2>)
- Radcliffe RW, Czekala NM & Osofsky SA** 1997 Combined serial ultrasonography and fecal progesterin analysis for reproductive evaluation of the female white rhinoceros (*Ceratotherium simum simum*): preliminary results. *Zoo Biology* **16** 445–456. ([https://doi.org/10.1002/\(SICI\)1098-2361\(1997\)16:5<445::AID-ZOO7>3.0.CO;2-A](https://doi.org/10.1002/(SICI)1098-2361(1997)16:5<445::AID-ZOO7>3.0.CO;2-A))
- Radcliffe RW, Eyres AI, Patton ML, Czekala NM & Emslie RH** 2001 Ultrasonographic characterization of ovarian events and fetal gestational parameters in two southern black rhinoceros (*Dicerus bicornis minor*) and correlation to fecal progesterone. *Theriogenology* **55** 1033–1049. ([https://doi.org/10.1016/s0093-691x\(01\)00464-2](https://doi.org/10.1016/s0093-691x(01)00464-2))
- Roca AL, Ishida Y, Brandt AL, Benjamin NR, Zhao K & Georgiadis NJ** 2015 Elephant natural history: a genomic perspective. *Annual Review of Animal Biosciences* **3** 139–167. (<https://doi.org/10.1146/annurev-animal-022114-110838>)
- Roth T** 2001 Ultrasound and endocrine evaluation of the ovarian cycle and early pregnancy in the Sumatran rhinoceros, *Dicerorhinus sumatrensis*. *Reproduction* **121** 139–149. (<https://doi.org/10.1530/reprod/121.1.139>)
- Roth TL** 2006 A review of the reproductive physiology of rhinoceros species in captivity. *International Zoo Yearbook* **40** 130–143. (<https://doi.org/10.1111/j.1748-1090.2006.00130.x>)
- Roth TL, Schook MW & Stoops MA** 2018 Monitoring and controlling ovarian function in the rhinoceros. *Theriogenology* **109** 48–57. (<https://doi.org/10.1016/j.theriogenology.2017.12.007>)
- Ruggeri E, Young C, Ravidia N, Sirard MA, Krisher R, de la Rey M, Herbst C & Durrant B** 2022 Glucose consumption and gene expression in granulosa cells collected before and after in vitro oocyte maturation in the southern white rhinoceros (*Ceratotherium simum simum*). *Reproduction, Fertility, and Development* **34** 875–888. (<https://doi.org/10.1071/RD22071>)
- Samuel CA, Allen WR & Steven DH** 1974 Studies on the equine placenta. I. Development of the microcotyledons. *Journal of Reproduction and Fertility* **41** 441–445. (<https://doi.org/10.1530/jrf.0.0410441>)
- Santos RR, Amorim C, Cecconi S, Fassbender M, Imhof M, Lornage J, Paris M, Schoenfeldt V & Martinez-Madrid B** 2010 Cryopreservation of ovarian tissue: an emerging technology for female germline preservation of endangered species and breeds. *Animal Reproduction Science* **122** 151–163. (<https://doi.org/10.1016/j.anireprosci.2010.08.010>)
- Saragusty J, Hildebrandt TB, Behr B, Knieriem A, Kruse J & Hermes R** 2009 Successful cryopreservation of Asian elephant (*Elephas maximus*) spermatozoa. *Animal Reproduction Science* **115** 255–266. (<https://doi.org/10.1016/j.anireprosci.2008.11.010>)
- Saragusty J, Diecke S, Drukker M, Durrant B, Friedrich Ben-Nun I, Galli C, Göritz F, Hayashi K, Hermes R, Holtze S, *et al.*** 2016a Rewinding the process of mammalian extinction. *Zoo Biology* **35** 280–292. (<https://doi.org/10.1002/zoo.21284>)

- Saragusty J, Prieto MT, Courtiol A, Potier R, Göritz F, Hildebrandt TB & Hermes R** 2016b Sperm rescue in wild African elephants. *Reproduction, Fertility, and Development* **28** 1433–1442. (<https://doi.org/10.1071/RD14378>)
- Satué K & Gardon JC** 2020 **Physiological and clinical aspects of the endocrinology of the estrous cycle and pregnancy in mares.** In *Animal Reproduction in Veterinary Medicine*. **F Aral, R Payan-Carreira & M Quaresma** Eds. London: Intech Open Book Series, pp. 5–20.
- Schaffer NE, Foley GL, Gill S & Pope CE** 2001 Clinical implications of rhinoceros reproductive tract anatomy and histology. *Journal of Zoo and Wildlife Medicine* **32** 31–46. ([https://doi.org/10.1638/1042-7260\(2001\)032\[0031:CIORRT\]2.0.CO;2](https://doi.org/10.1638/1042-7260(2001)032[0031:CIORRT]2.0.CO;2))
- Schmitt D** 2006 **Reproductive system.** In *Biology, Medicine, and Surgery of Elephants*. **ME Fowler & S Mikota** Eds. IA: Aimes, pp. 347–356.
- Schwarzenberger F & Hermes R** 2023 Comparative analysis of gestation in three rhinoceros species (*Diceros bicornis*; *Ceratotherium simum*; *Rhinoceros unicornis*). *General and Comparative Endocrinology* **334** 114214. (<https://doi.org/10.1016/j.ygcen.2023.114214>)
- Schwarzenberger F, Walzer C, Tomasova K, Vahala J, Meister J, Goodrowe KL, Zima J, Strauß G & Lynch M** 1998 Faecal progesterone metabolite analysis for non-invasive monitoring of reproductive function in the white rhinoceros (*Ceratotherium simum*). *Animal Reproduction Science* **53** 173–190. ([https://doi.org/10.1016/S0378-4320\(98\)00112-2](https://doi.org/10.1016/S0378-4320(98)00112-2))
- Schwarzenberger F, Rietschel W, Vahala J, Holeckova D, Thomas P, Maltzan J, Baumgartner K & Schaftenaar W** 2000 Faecal progesterone, estrogen, and androgen metabolites for noninvasive monitoring of reproductive function in the female Indian rhinoceros, *Rhinoceros unicornis*. *General and Comparative Endocrinology* **119** 300–307. (<https://doi.org/10.1006/gcen.2000.7523>)
- Sherman GB, Lund LA, Bunick D & Winn RJ** 1997 Characterization and phylogenetic significance of rhinoceros luteinizing hormone beta (LHbeta) subunit messenger RNA structure, complementary DNA sequence and gene copy number. *Gene* **195** 131–139. ([https://doi.org/10.1016/S0378-1119\(97\)00080-2](https://doi.org/10.1016/S0378-1119(97)00080-2))
- Spanner J, Pedersen J & Lorenzen E** 2022 Transport of vitrified-warmed ICSI blastocysts in straws for up to 5 hours does not affect the pregnancy rate. *Journal of Equine Veterinary Science* **113**. (<https://doi.org/10.1016/j.jevs.2022.103986>)
- Stansfield FJ & Allen WR** 2012 Luteal maintenance of pregnancy in the African elephant (*Loxodonta africana*). *Reproduction* **143** 845–854. (<https://doi.org/10.1530/REP-12-0032>)
- Stansfield FJ, Picton HM & Nöthling JO** 2011 Early primary—rather than primordial follicles constitute the main follicular reserve in the African elephant (*Loxodonta africana*). *Animal Reproduction Science* **123** 112–118. (<https://doi.org/10.1016/j.anireprosci.2010.11.003>)
- Stoops MA, Pairan RD & Roth TL** 2004 Follicular, endocrine and behavioural dynamics of the Indian rhinoceros (*Rhinoceros unicornis*) oestrous cycle. *Reproduction* **128** 843–856. (<https://doi.org/10.1530/rep.1.00328>)
- Stoops MA, Bateman H, Campbell MK & Roth TL** 2011a Attempted in vitro maturation and fertilization of postmortem Sumatran rhinoceros (*Dicerorhinus sumatrensis*) oocytes. *American Association of Zoo Veterinarians* **42** 723–726. (<https://doi.org/10.1638/2010-0186.1>)
- Stoops MA, O'Brien JK & Roth TL** 2011b Gamete rescue in the African black rhinoceros (*Diceros bicornis*). *Theriogenology* **76** 1258–1265. (<https://doi.org/10.1016/j.theriogenology.2011.05.032>)
- Stoops MA, Campbell MK, DeChant CJ, Hauser J, Kottwitz J, Pairan RD, Shaffstall W, Volle K & Roth TL** 2016 Enhancing captive Indian rhinoceros genetics via artificial insemination of cryopreserved sperm. *Animal Reproduction Science* **172** 60–75. (<https://doi.org/10.1016/j.anireprosci.2016.07.003>)
- Stout TA** 2006 Equine embryo transfer: review of developing potential. *Equine Veterinary Journal* **38** 467–78. (<https://doi.org/10.2746/042516406778400529>)
- Stout TA & Allen WR** 2001 Role of prostaglandins in intrauterine migration of the equine conceptus. *Reproduction* **121** 771–775. (<https://doi.org/10.1530/rep.0.1210771>)
- Stout TAE** 2020 Clinical application of in vitro embryo production in the horse. *Journal of Equine Veterinary Science* **89** 103011. (<https://doi.org/10.1016/j.jevs.2020.103011>)
- Thitaram C, Brown JL, Pongsupawijit P, Chansitthiwet S, Wongkalasin W, Daram P, Roongsri R, Kalmapijit A, Mahasawangkul S, Rojansthiem S *et al.*** 2008 Seasonal effects on the endocrine pattern of semi-captive female Asian elephants (*Elephas maximus*): Timing of the anovulatory luteinizing hormone surge determines the length of the estrous cycle. *Theriogenology* **69** 237–244. (<https://doi.org/10.1016/j.theriogenology.2007.09.018>)
- Thongtip N, Mahasawangkul S, Thitaram C, Pongsopavijitr P, Kornkaewrat K, Pinyopummin A, Angkawanish T, Jansittiwate S, Rungsri R, Boonprasert K, *et al.*** 2009 Successful artificial insemination in the Asian elephant (*Elephas maximus*) using chilled and frozen-thawed semen. *Reproductive Biology and Endocrinology* **7** 75. (<https://doi.org/10.1186/1477-7827-7-75>)
- Ververs C, Van Zijl Langhout M, Govaere J & Van Soom A** 2015 Features of reproduction and assisted reproduction in the white (*Ceratotherium simum*) and black (*Diceros bicornis*) rhinoceros. *Vlaams Diergeneeskundig Tijdschrift* **84** 175–187. (<https://doi.org/10.21825/vdt.v84i4.16593>)
- Weber JA, Freeman DA, Vanderwall DK & Woods GL** 1991 Prostaglandin E2 secretion by oviductal transport-stage equine embryos. *Biology of Reproduction* **45** 540–543. (<https://doi.org/10.1095/biolreprod45.4.540>)
- Wilsher S & Allen WR** 2004 An improved method for nonsurgical embryo transfer in the mare. *Equine Veterinary Education* **16** 39–44. (<https://doi.org/10.1111/j.2042-3292.2004.tb00265.x>)
- Wilsher S, Rigali F, Couto G, Camargo S & Allen WR** 2019 Vitrification of equine expanded blastocysts following puncture with or without aspiration of the blastocoele fluid. *Equine Veterinary Journal* **51** 500–505. (<https://doi.org/10.1111/evj.13039>)
- Wilsher S, Rigali F, Kovacs S & Allen WT** 2021 Successful vitrification of manually punctured equine embryos. *Equine Veterinary Journal* **53** 1227–1233. (<https://doi.org/10.1111/evj.13400>)
- Yamamoto Y, Nagaoka K, Kamite Y, Watanabe G, Allen T, Stansfield F & Taya K** 2017 Different origins of two corpora lutea recovered from a pregnant African elephant (*Loxodonta africana*). *Reproduction in Domestic Animals* **52** 1138–1141. (<https://doi.org/10.1111/rda.13010>)
- Yamamoto Y, Yuto N, Yamamoto T, Kaewmanee S, Shiina O, Mouri Y, Narushima E, Katayanagi M, Sugimura K, Nagaoka K, *et al.*** 2012 Secretory pattern of inhibin during estrous cycle and pregnancy in African (*Loxodonta africana*) and Asian (*Elephas maximus*) elephants. *Zoo Biology* **31** 511–522. (<https://doi.org/10.1002/zoo.20415>)

Received 4 April 2023

Accepted 13 April 2023

Available online 13 April 2023

Version of Record published 7 August 2023