

Est.  
1841

YORK  
ST JOHN  
UNIVERSITY

Sinani, Charikleia ORCID:

<https://orcid.org/0000-0001-8942-8780>, Wilmut, Kate, Mierzwinski, Mark ORCID: <https://orcid.org/0000-0002-9751-5865>, Williams, Jacqueline, Licari, Melissa, Purcell, Catherine, Gentle, Judith and Wood, Greg (2023) The Impact of Developmental Coordination Disorder in the UK. In: DCD-UK 2023, 7-8 July 2023, Manchester Metropolitan University, Manchester.

Downloaded from: <http://ray.yorks.ac.uk/id/eprint/8333/>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

# RaY

Research at the University of York St John

For more information please contact RaY at [ray@yorks.ac.uk](mailto:ray@yorks.ac.uk)

# The Impact of Developmental Coordination Disorder in the UK



Dr. Charikleia Sinani, Dr. Mark Mierzwinski, Dr. Jacqueline Williams, Dr. Mellissa Licari, Dr. Catherine Purcell, Dr. Judith Gentle, Dr. Greg Wood & Professor Kate Wilmut.

Est.  
1841

YORK  
ST JOHN  
UNIVERSITY

National Survey Evaluating the Impact of  
Developmental Coordination Disorder in Australia

impact  
FOR DCD

# Background

- In 2020, an Australian research group carried out one of the largest surveys conducted in the world to identify the challenges experienced by children with DCD and their families -  
<https://www.telethonkids.org.au/projects/impact-for-dcd/>
- Key findings:
  - delays in diagnosis,
  - the use of inconsistent terminology,
  - limited funding for support,
  - lack of awareness amongst teachers and health professionals
  - high rates of mental health issues

# The UK study - aims

Following on the Australian study there was a clear need for this study to be replicated in the UK to:

- i) understand the scale and breadth of issues experienced in DCD *and*
- ii) to identify and prioritise areas in which children with DCD and their families require support.



# The UK study

- Questionnaire adapted based on the Australian study
- Included questions around the following areas:
  - family and diagnostics,
  - movement & difficulties,
  - activity,
  - schooling,
  - therapy,
  - social and emotional impact on the child (some questions based on the Strengths and Difficulties Questionnaire (SDQ) (Goodman, 1987) <https://www.sdqinfo.org/>)
  - Demographics, socioeconomic status,
  - parents to identify priorities and further comments.




# The UK study

- Questionnaire adapted based on the Australian study
- Included questions around the following areas:
  - family and diagnostics,
  - movement & difficulties,
  - activity,
  - schooling,
  - therapy,
  - **social and emotional impact on the child** (some questions based on the Strengths and Difficulties Questionnaire (SDQ) (Goodman, 1987) <https://www.sdqinfo.org/>)
  - Demographics, socioeconomic status,
  - parents to identify priorities and further comments.



# Who took part in the study



**Figure 1.** Distribution of age


- 240 parental responses included:
  - 5-18 years (mean age of 11.10 years).
  - 71.1% (N=173) male and 27.9% (N= 67) female.
  - Ethnicity: 93.3% (white), 3.33% (mixed heritage), 3.33% (Asian).
- 79.17% had a diagnosis for their movement difficulties
  - 55% had been diagnosed with co-occurring conditions
  - 45% had been diagnosed with more than one

# Social and Emotional Impact on the child

**Parental concerns** (89.6% concerned, 4.2% not sure)

1. Poor mental health including anxiety and depression (N=44)
2. Friendships (N=41)
3. Low confidence (N=39)
4. Poor self-esteem (N=37)
5. Difficulty with keeping up (N=31)
6. Acceptance (N=17)
7. Bullying (N=15)
8. Isolation and withdrawal (N=12)
9. Exclusion by peers (N=7)
10. Suicidal thoughts / self-harm (N=3)

*"Masking difficulties, his anxiety has started to come out by self harming, we only discovered this a few weeks ago."*






# Social and Emotional In

**Parental concerns** (89.6% concerned, 4.2% not sure)

1. Poor mental health including anxiety and depression (N=44)
2. Friendships (N=41)
3. Low confidence (N=39)
4. Poor self-esteem (N=37)
5. Difficulty with keeping up (N=31)
6. Acceptance (N=17)
7. Bullying (N=15)
8. Isolation and withdrawal (N=12)
9. Exclusion by peers (N=7)
10. Suicidal thoughts / self-harm (N=3)

*“Since his diagnosis he has repeatedly struggled with being different, and has on occasion said he doesn’t belong here and doesn’t deserve to be alive. His internal dialogue is set to negative thoughts, and he often berates himself and tells me he is terrible and stupid.”*




# Social and Emotional Impact of

## Parental concerns (89.6% concerned, 4.2% not sure)

1. Poor mental health including anxiety and depression (N=44)
2. Friendships (N=41)
3. Low confidence (N=39)
4. Poor self-esteem (N=37)
5. Difficulty with keeping up (N=31)
6. Acceptance (N=17)
7. Bullying (N=15)
8. Isolation and withdrawal (N=12)
9. Exclusion by peers (N=7)
10. Suicidal thoughts / self-harm (N=3)

*“Low confidence, people sometimes judge him (been voted most clumsy in school year book) . Not socially confident at all outside his friendship group”*



# Social and Emotional Impact on the child





Figure 2. Frequency of socioemotional problems as reported by parents.

# Socioemotional Symptoms – measured using the SDQ



**Figure 3.** Illustration of the outcome of the Strengths and Difficulties Questionnaire using the categories supplied with the questionnaire.

# Socioemotional symptoms by age – measured using the SDQ




Figure 4. Breakdown of the emotional problems experienced by age group




Figure 5. Breakdown of the peer problems experienced by age group

# Socioemotional symptoms by age – measured using the SDQ (continued)




Figure 6. Breakdown of prosocial problems experienced by age group

# Socioemotional symptoms by occurrence – measured using the SDQ




Figure 7. Breakdown of the **emotional problems** experienced by co-occurrence




Figure 8. Breakdown of the **peer problems** experienced by co-occurrence




Figure 9. Breakdown of the **pro-social behaviour** experienced by co-occurrence

# Conclusions




- 89.6% of parents were concerned about the impact of their child's movement difficulties on their social and emotional health
- Two thirds of children scored in clinically concerning ranges in relation to emotional difficulties and peer problems.
- These problems seem to get worse the older the child gets and when a motor diagnosis is accompanied by more than one other diagnosis.
- Peer and prosocial problems either peaked or increased within the age ranges of 9-10 and 15 -16 years old.
- Over half of families reported that their child is 'always' or 'very often' **anxious learning or performing movement related activities** and **withdraws from such activities**.






# Conclusions



- **Mental health was one of the major parental concerns and one of the main priority areas for their child.**
- Assessments of co-occurring conditions and mental health problems should be incorporated.
- Intervention should address mental health problems as well as motor development.
- There is a clear need to educate not only families but also professionals working with children with DCD about the signs of poor mental health and the services available to support families and child but also increase the resources available.


*“There needs to be far better recognition and acceptance of the mental health needs of young children and how mental health is deeply intertwined with a child’s physical and overall well being. We have been repeatedly ignored and let down by CAMHS, pushed off onto the medical side who in turn offer no supportive therapies, leaving it to the school to pick up and provide what little they can manage. The entire system is broken and letting children down, not just those with DCD, but all children with extra needs.”*



**One of the major parental  
one of the main priority areas for**

- Interventions of co-occurring conditions and mental health problems should be incorporated.
- Intervention should address mental health problems as well as motor development.
- There is a clear need to educate not only families but also professionals working with children with DCD about the signs of poor mental health and the services available to support families and child but also increase the resources available.


*“Mental health support is very hard to access and is often focused on general anxiety with very little support focusing on the causes”*



- **Mental health was one of the major parental concerns and one of the main priority areas for their child.**
- Assessments of co-occurring conditions and mental health problems should be incorporated.
- Intervention should address mental health problems as well as motor development.
- There is a clear need to educate not only families but also professionals working with children with DCD about the signs of poor mental health and the services available to support families and child but also increase the resources available.

Cor

*"The government cite Mental Health awareness, but provide no support and the NHS are under-funded for this. A neurodiverse condition, such as DCD, can lead to mental health issues due to not understanding why you can't do the same things others can do and the frustration and anger that arises from this. People can feel very isolated and unworthy as a result."*



**impact**  
FOR DCD

... was one of the major parental concerns and one of the main priority areas for child.

- Assessments of co-occurring conditions and mental health problems should be incorporated.
- Intervention should address mental health problems as well as motor development.
- There is a clear need to educate not only families but also professionals working with children with DCD about the signs of poor mental health and the services available to support families and child but also increase the resources available.

# Acknowledgements

- Children and families who participated in this survey
- Dyspraxia Foundation
- Funders: York St John University and Oxford Brookes University
- Dr. Mark Mierzwinski, Dr. Jacqueline Williams, Dr. Mellissa Licari, Dr. Catherine Purcell, Dr. Judith Gentle, Dr. Greg Wood & Professor Kate Wilmut.

You can find us on twitter [@impactdcduk](#) and on Facebook [ImpactDCDUK | Facebook](#)

Email: [C.Sinani@yorksja.ac.uk](mailto:C.Sinani@yorksja.ac.uk)



# References

- Goodman R. (1997) The Strengths and Difficulties Questionnaire: a research note. *Journal of Child Psychology and Psychiatry*, 38, 581-586

