## Check for updates

## **OPEN ACCESS**

EDITED AND REVIEWED BY Carl-Olav Stiller, Karolinska Institutet (KI), Sweden

\*CORRESPONDENCE Slavica Kvolik, Skvolik@mefos.hr Nenad Neskovic, nneskov@gmail.com

RECEIVED 30 June 2023 ACCEPTED 31 July 2023 PUBLISHED 22 August 2023

#### CITATION

Neskovic N, Mandic D, Marczi S, Skiljic S, Kristek G, Vinkovic H, Mraovic B, Debeljak Z and Kvolik S (2023), Corrigendum: Different pharmacokinetics of tramadol, *O*demethyltramadol and *N*demethyltramadol in postoperative surgical patients from those observed in medical patients. *Front. Pharmacol.* 14:1250698. doi: 10.3389/fphar.2023.1250698

#### COPYRIGHT

© 2023 Neskovic, Mandic, Marczi, Skiljic, Kristek, Vinkovic, Mraovic, Debeljak and Kvolik. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this

journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms. Corrigendum: Different pharmacokinetics of tramadol, O-demethyltramadol and N-demethyltramadol in postoperative surgical patients from those observed in medical patients

Nenad Neskovic<sup>1,2</sup>\*, Dario Mandic<sup>2,3</sup>, Saska Marczi<sup>2,4</sup>, Sonja Skiljic<sup>1,2</sup>, Gordana Kristek<sup>1,2</sup>, Hrvoje Vinkovic<sup>1,2</sup>, Boris Mraovic<sup>5</sup>, Zeljko Debeljak<sup>2,3</sup> and Slavica Kvolik<sup>1,2</sup>\*

<sup>1</sup>Department of Anesthesiology, Resuscitation and ICU, Osijek University Hospital, Osijek, Croatia, <sup>2</sup>Faculty of Medicine, University Josip Juraj Strossmayer, Osijek, Croatia, <sup>3</sup>Department of Clinical and Laboratory Diagnostics, Osijek University Hospital, Osijek, Croatia, <sup>4</sup>Laboratory for Molecular and HLA Diagnostic, Department of Transfusion Medicine, Osijek University Hospital, Osijek, Croatia, <sup>5</sup>University of Missouri, Department of Anesthesiology and Perioperative Medicine, School of Medicine, Columbia, MO, United States

### KEYWORDS

postoperative analgesia, postoperative pain, inflammation, cholinesterase, CYP2D6, O-demethyltramadol, N-demethyltramadol, tramadol

## A Corrigendum on

Different pharmacokinetics of tramadol, *O*-demethyltramadol and *N*-demethyltramadol in postoperative surgical patients from those observed in medical patients

by Neskovic N, Mandic D, Marczi S, Skiljic S, Kristek G, Vinkovic H, Mraovic B, Debeljak Z and Kvolik S (2021). Front. Pharmacol. 12:656748. doi: 10.3389/fphar.2021.656748

In the published article, there was an error in Figure 1 as published. The Figures 1B, C were incorrect and appeared inverted in the figure caption.

The corrected Figure 1 and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

# Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



## FIGURE 1

Concentrations of tramadol (A), N-demethyltramadol (B), and O-demethyltramadol (C) in the first 24 postoperative hours. Concentrations were measured 1, 2, 4 h after the first dose of 100 mg tramadol iv, and just before the second (time point 6 h), third (time point 12 h), and fifth (time point 24 h) doses of tramadol. PM, poor metabolizer; IM, intermediate metabolizer; EM, extensive metabolizer; UM, ultrafast metabolizers; Dot, tramadol 100 mg IV injections; \* statistically significant differences (Mann-Whitney U test) between PM and EM/IM (B), and EM and IM/PM (C).