

Number of Jacobi quartic curves over finite fields *

Rongquan Feng¹, Hongfeng Wu²

¹ LMAM, School of Mathematical Sciences, Peking University,
Beijing 100871, P.R. China

² Academy of Mathematics and Systems Science, Chinese Academy of Sciences,
Beijing 100190, P.R. China

fengrq@math.pku.edu.cn, whfmath@gmail.com

Abstract

In this paper the number of $\overline{\mathbb{F}}_q$ -isomorphism classes of Jacobi quartic curves, i.e., the number of Jacobi quartic curves with distinct j -invariants, over finite field \mathbb{F}_q is enumerated.

Keywords: Elliptic curves, Jacobi quartic, isomorphism classes, cryptography

1 Introduction

Elliptic curve cryptosystems were proposed by Miller (1986) and by Koblitz (1987) which relies on the difficulty of the elliptic curve discrete logarithmic problem. No sub-exponential algorithms have been found for solving the discrete logarithm problem based on elliptic curves is one of main advantages of this system. One basic operation required to implement the system is the point multiplication, that is, the computation of kP for an integer k and a point P on the curve. To obtain faster operations, more efforts have been

*Supported by NSF of China (No. 10990011)

done in representing the elliptic curves in special forms which provide faster addition, doubling and tripling in the last decades.

The Jacobi quartic curves is one of the most important curves in cryptography. Recent works have shown that arithmetics on the Jacobi quartic elliptic curves can be performed more efficiently. The reader is referred to [1] for the comparison analysis of computational costs for all kinds of curves. A Jacobi quartic elliptic curve over a field K is defined by $y^2 = x^4 + 2ax^2 + 1$, where $a \in K$ with $a^2 \neq 1$. Such curves were first proposed by Chudnovsky and Chudnovsky [3] in 1986. After that, Billet and Joye [2], Duquesne [4], and Hisil, etc. [5] gave more improvements for the arithmetics on Jacobi quartic curves.

In order to study the elliptic curves cryptosystem, we need first to answer how many curves there are up to isomorphism, because two isomorphic elliptic curves are the same in the point of cryptographic view. In this paper the number of $\overline{\mathbb{F}}_q$ -isomorphism classes of Jacobi quartic curves, i.e., the number of Jacobi quartic curves with distinct j -invariants, over a finite field is enumerated.

Throughout the paper, \mathbb{F}_q denotes the finite field with q elements and $\overline{\mathbb{F}}_q$ is the algebraic closure of \mathbb{F}_q .

2 Background

A curve means a projective variety of dimension 1. There are several ways to define elliptic curves. In this paper, an irreducible curve is said to be an elliptic curve if it is birationally equivalent to a non-singular plane cubic curve.

It is well-known that every elliptic curve E over a field K can be written as a Weierstrass equation

$$E : Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6$$

with coefficients $a_1, a_2, a_3, a_4, a_6 \in K$. The discriminant $\Delta(E)$ and the j -invariant $j(E)$ of E are defined as

$$\Delta(E) = -b_2^2b_8 - 8b_4^3 - 27b_6^2 + 9b_2b_4b_6$$

and

$$j(E) = (b_2^2 - 24b_4)^3 / \Delta(E),$$

where

$$\begin{aligned} b_2 &= a_1^2 + 4a_2, \\ b_4 &= 2a_4 + a_1a_3, \\ b_6 &= a_3^2 + 4a_6, \\ b_8 &= a_1^2a_6 - a_1a_3a_4 + 4a_2a_6 + a_2a_3^2 - a_4^2. \end{aligned}$$

Two projective varieties V_1 and V_2 are isomorphic if there exist morphisms $\phi : V_1 \rightarrow V_2$ and $\varphi : V_2 \rightarrow V_1$, such that $\varphi \circ \phi$ and $\phi \circ \varphi$ are the identity maps on V_1 and V_2 , respectively. Two elliptic curves are said to be isomorphic if they are isomorphic as projective varieties. Let $E_1 : Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6$ and $E_2 : Y^2 + a'_1XY + a'_3Y = X^3 + a'_2X^2 + a'_4X + a'_6$ be two elliptic curves defined over K . It is known [6] that E_1 and E_2 are isomorphic over \overline{K} if and only if $j(E_1) = j(E_2)$, where \overline{K} is the algebraic closure of K . However, E_1 and E_2 are isomorphic over \overline{K} if and only if there exists $u, r, s, t \in \overline{K}$ and $u \neq 0$ such that the change of variables

$$(X, Y) \rightarrow (u^2X + r, u^3Y + u^2sX + t)$$

maps the equation of E_1 to the equation of E_2 (see [6]). Therefore, E_1 and E_2 are isomorphic over \overline{K} if and only if there exists $u, r, s, t \in \overline{K}$ and $u \neq 0$ such that

$$\begin{cases} ua'_1 = a_1 + 2s, \\ u^2a'_2 = a_2 - sa_1 + 3r - s^2, \\ u^3a'_3 = a_3 + ra_1 + 2t, \\ u^4a'_4 = a_4 - sa_3 + 2ra_2 - (t + rs)a_1 + 3r^2 - 2st, \\ u^6a'_6 = a_6 + ra_4 + r^2a_2 + r^3 - ta_3 - t^2 - rta_1. \end{cases}$$

For the simplified Weierstrass equations where $a_1 = a_3 = a'_1 = a'_3 = 0$, then E_1 and E_2 are isomorphic over \overline{K} if and only if there exists $u, r \in \overline{K}$ and $u \neq 0$ such that

$$\begin{cases} u^2a'_2 = a_2 + 3r, \\ u^4a'_4 = a_4 + 2ra_2 + 3r^2, \\ u^6a'_6 = a_6 + ra_4 + r^2a_2 + r^3. \end{cases} \quad (1)$$

For more results on the isomorphism of elliptic curves, see [6] for more details.

In order to enumerate the number of elliptic curves with distinct j -invariants, one need to study the value distribution of the j -invariant as a function of curve parameters. However it is effective for only low degree j -invariant function, but very difficult for counting the value set of the j -invariant function of Jacobi quartic curves where a polynomial of degree 6 is

involved, especially for the general Jacobi quartic curves where polynomials with 2 variables are associated. In this paper, this number is enumerated by studying the $\overline{\mathbb{F}}_q$ -isomorphism classes of those curves.

3 Enumeration for Jacobi quartics curves

Let $E_a : y^2 = x^4 + ax^2 + 1$ ($a^2 \neq 4$) be a Jacobi quartic curve defined over a field K of characteristic not 2. It is clear that the j -invariant of E_a is $\frac{16(a^2+12)^3}{(a^2-4)^2}$.

Lemma 3.1. *Let K be a field of characteristic > 3 , let $a \in K$ with $a^2 \neq 4$. Then the curve*

$$E_a : y^2 = x^4 + ax^2 + 1$$

is birationally equivalent to the elliptic curve

$$W_a : v^2 = u(u-1) \left(u - \frac{2-a}{4} \right)$$

via the change of variables $\varphi(x, y) = (u, v)$, where

$$u = \frac{x^2 - y + 1}{2}, \quad v = \frac{x(2x^2 - 2y + a)}{4}.$$

The inverse change is $\psi(u, v) = (x, y)$, where

$$x = \frac{4v}{4u + a - 2}, \quad y = \left(\frac{4v}{4u + a - 2} \right)^2 - 2u + 1.$$

Proof. In order to prove

$$v^2 = u(u-1) \left(u - \frac{2-a}{4} \right),$$

it is sufficient to prove $64v^2 = 4u(4u-4)(4u-(2-a))$. Since $4u-(2-a) = 2x^2 - 2y + a$ and $64v^2 = 4x^2(2x^2 - 2y + a)^2$, it is sufficient to show that $4x^2(2x^2 - 2y + a) = 4u(4u - 4)$. The result then follows immediately from

$$4x^2(2x^2 - 2y + a) = 8x^4 - 8x^2y + 4ax^2,$$

and

$$\begin{aligned}
4u(4u - 4) &= 4(x^2 - y + 1)(x^2 - y - 1) \\
&= 4(x^4 + y^2 - 2x^2y - 1) \\
&= 4(x^4 + x^4 + ax^2 + 1 - 2x^2y - 1) \\
&= 8x^4 - 8x^2y + 4ax^2.
\end{aligned}$$

On the other hand, from $v^2 = u(u^2 - 2au + a^2 - 4b)$, $x = \frac{4v}{4u+a-2}$, and $y = \left(\frac{4v}{4u+a-2}\right)^2 - 2u + 1$, we have $y^2 = x^4 + ax^2 + 1$ by a direct computation. Obviously, the maps φ and ψ are mutually inverse to each other. \square

Lemma 3.2. *Let $E_a : y^2 = x^4 + ax^2 + 1$ ($a^2 \neq 4$) and $E_b : y^2 = x^4 + bx^2 + 1$ ($b^2 \neq 4$) be two Jacobi quartics curves defined over a field K of characteristic not 2. Then $j(E_a) = j(E_b)$ if and only if $\frac{2-b}{4} \in \left\{\frac{2-a}{4}, \frac{4}{2-a}, \frac{2+a}{4}, \frac{4}{2+a}, \frac{2-a}{2+a}, \frac{a+2}{a-2}\right\}$.*

Proof. The curve $E_a : y^2 = x^4 + ax^2 + 1$ is birational equivalent to the curve $W_a : y^2 = x(x-1)(x - \frac{2-a}{4})$ by Lemma 3.1. Therefore $j(E_a) = j(W_a)$. Furthermore, it is well known that for two Legendre curves $L_\lambda : y^2 = x(x-1)(x-\lambda)$ and $L_\mu : y^2 = x(x-1)(x-\mu)$, they have the same j -invariant if and only if $\mu \in \left\{\lambda, \frac{1}{\lambda}, 1-\lambda, \frac{1}{1-\lambda}, \frac{\lambda}{1-\lambda}, \frac{\lambda-1}{\lambda}\right\}$. Thus the lemma follows. \square

Theorem 3.3. *Let N_a be the number of $\overline{\mathbb{F}}_q$ -isomorphism classes of Jacobi quartic curves defined over the finite field \mathbb{F}_q . Then we have*

$$N_a = \begin{cases} \frac{q+5}{6}, & \text{if } q \equiv 1, 7 \pmod{12}, \\ \frac{q+1}{6}, & \text{if } q \equiv 5, 11 \pmod{12}. \end{cases}$$

Proof. From Lemma 3.2, we know that for the elliptic curve $L_\lambda : y^2 = x(x-1)(x-\lambda)$ ($\lambda \neq 0, 1$), the map $\lambda \mapsto j(L_\lambda)$ is exactly six-to-one unless when $\lambda \in \{-1, 2, \frac{1}{2}\}$, the map is three-to-one, or when $\lambda^2 - \lambda + 1 = 0$, the map is two-to-one. Note that $\lambda^2 - \lambda + 1 = 0$ has a root in \mathbb{F}_q if and only if \mathbb{F}_q^* has an element of order 3, which is equivalent to $q \equiv 1$ or $7 \pmod{12}$. Therefore, we have

$$N_a = \begin{cases} \frac{q-2-3-2}{6} + 1 + 1 = \frac{q+5}{6}, & \text{if } q \equiv 1, 7 \pmod{12}, \\ \frac{q-2-3}{6} + 1 = \frac{q+1}{6}, & \text{if } q \equiv 5, 11 \pmod{12}. \end{cases}$$

\square

4 Enumeration for general Jacobi quartics curves

In this section, we consider the general Jacobi quartics curve $E_{a,b} : y^2 = x^4 + ax^2 + b$ with $(a^2 - 4b)b \neq 0$ defined over \mathbb{F}_q of characteristic > 3 . A Jacobi quartics curve is a special one of $W_{a,b}$ with $b = 1$. The j -invariant of $E_{a,b}$ is $j(E_{a,b}) = \frac{16(a^2+12b)^3}{b(a^2-4b)^2}$. Note that $y^2 = bx^4 + ax^2 + 1$ can be changed to $y^2 = x^4 + ax^2 + b$ by $x \mapsto 1/x$. So we consider only the form $y^2 = x^4 + ax^2 + b$ for convenience.

The following lemma can be proved by a direct computation similar as that in Lemma 3.1.

Lemma 4.1. *Let K be a field of characteristic > 3 , let $a, b \in K$ with $(a^2 - 4b)b \neq 0$. Then the curve*

$$E_{a,b} : y^2 = x^4 + ax^2 + b$$

is birationally equivalent to the elliptic curve

$$W_{a,b} : v^2 = u(u^2 - 2au + a^2 - 4b)$$

via the change of variables

$$u = 2x^2 - 2y + a, \quad v = 2x(2x^2 - 2y + a).$$

The inverse change is

$$x = \frac{v}{2u}, \quad y = \left(\frac{v}{2u}\right)^2 - \frac{u-a}{2}.$$

For the elliptic curve $E_{a,b}$, we know that $j(E_{a,b}) = \frac{16(a^2+12b)^3}{b(a^2-4b)^2}$. Therefore $j(E_{a,b}) = 0$ if and only if $a^2 + 12b = 0$. Moreover, we have the following proposition.

Proposition 4.2. *Let $E_{a,b} : y^2 = x^4 + ax^2 + b$ be a general Jacobi quartics curve defined over the finite field \mathbb{F}_q of characteristic > 3 , where $b(a^2 - 4b) \neq 0$. Then $j(E_{a,b}) = 1728$ if and only if $a(a^2 - 36b) = 0$, that is $a = 0$ or $a^2 = 36b$.*

Proof. By Lemma 4.1, $E_{a,b}$ is birational equivalent to the curve $W_{a,b} : y^2 = x^3 - 2ax^2 + (a^2 - 4b)x$, and $W_{a,b}$ is isomorphic to $S_{a,b} : y^2 = x^3 + (-4a - \frac{a^2}{3})x + (\frac{2a^3}{27} - \frac{8ab}{3})$. It is clear that the j -invariant of $S_{a,b}$ is equal to 1728 if and only if $\frac{2a^3}{27} - \frac{8ab}{3} = 0$, that is $a(a^2 - 36b) = 0$. Thus $j(E_{a,b}) = 1728$ if and only if $a(a^2 - 36b) = 0$. \square

Corollary 4.3. *Let $(a^2 - 4b)b \neq 0$ and let N be the number of curves of the form $E_{a,b}$ with $j(E_{a,b}) \neq 0, 1728$. Then*

$$N = \begin{cases} \frac{(q-1)(q-7)}{2}, & \text{if } q \equiv 1, 7 \pmod{12}, \\ \frac{(q-1)(q-5)}{2}, & \text{if } q \equiv 5, 11 \pmod{12}, \end{cases}$$

when b is a square and

$$N = \begin{cases} \frac{(q-1)^2}{2}, & \text{if } q \equiv 1, 7 \pmod{12}, \\ \frac{(q-1)(q-3)}{2}, & \text{if } q \equiv 5, 11 \pmod{12}. \end{cases}$$

when b is not a square.

Proof. Assume first that b is a square in \mathbb{F}_q . Then $a^2 - 4b = 0$ has two roots. Hence the number of curves of the form $E_{a,b}$ over \mathbb{F}_q is $(q-2) \cdot (q-1)/2 = (q-1)(q-2)/2$. If $j(E_{a,b}) = 0$, then $a^2 + 12b = 0$ which has two roots in \mathbb{F}_q if $q \equiv 1, 7 \pmod{12}$, but has no root if $q \equiv 5, 11 \pmod{12}$. Therefore the number of curves of the form $E_{a,b}$ over \mathbb{F}_q with $j(E_{a,b}) = 0$ is $2 \cdot \frac{q-1}{2} = q-1$ if $q \equiv 1, 7 \pmod{12}$, and is 0 if $q \equiv 5, 11 \pmod{12}$. If $j(E_{a,b}) = 1728$, then $a = 0$ or $a^2 = 36b$. Thus the number of curves of the form $E_{a,b}$ with $j(E_{a,b}) = 1728$ is $\frac{q-1}{2} + 2 \cdot \frac{q-1}{2} = \frac{3(q-1)}{2}$. By subtraction, we get that

$$N = \begin{cases} \frac{(q-1)(q-2)}{2} - (q-1) - \frac{3(q-1)}{2} = \frac{(q-1)(q-7)}{2}, & \text{if } q \equiv 1, 7 \pmod{12}, \\ \frac{(q-1)(q-2)}{2} - 0 - \frac{3(q-1)}{2} = \frac{(q-1)(q-5)}{2}, & \text{if } q \equiv 5, 11 \pmod{12}. \end{cases}$$

The number N can be computed similarly when b is not a square. In this case the number of curves $E_{a,b}$ is $q \cdot (q-1)/2 = q(q-1)/2$ and $a^2 + 12b = 0$ has two roots in \mathbb{F}_q if $q \equiv 5, 11 \pmod{12}$, has no root if $q \equiv 1, 7 \pmod{12}$. \square

Now consider curves $E_{a,b}$ over \mathbb{F}_q with $j(E_{a,b}) \neq 0, 1728$. Suppose that two elliptic curves $E_{a,b}$ and $E_{m,n}$ are isomorphic over $\overline{\mathbb{F}_q}$. Then $j(E_{a,b}) = j(E_{m,n})$, and then $j(W_{a,b}) = j(W_{m,n})$ by Lemma 4.1, which is equivalent to $W_{a,b}$ and $W_{m,n}$ are isomorphic over $\overline{\mathbb{F}_q}$. Moreover, by (1), the last statement holds if and only if there exist $u, r \in \overline{\mathbb{F}_q}$ with $u \neq 0$ such that

$$\begin{cases} 2mu^2 = 2a - 3r, \\ (m^2 - 4n)u^4 = 3r^2 - 4ar + (a^2 - 4b), \\ r(r^2 - 2ar + a^2 - 4b) = 0. \end{cases} \quad (2)$$

Proposition 4.4. *Let $(a^2 - 4b)b \neq 0$ and let b be a square element. Then for every general Jacobi quartic curve $E_{a,b} : y^2 = x^4 + ax^2 + b$, there is a Jacobi quartic curve $E_m : y^2 = x^4 + mx^2 + 1$ which is $\overline{\mathbb{F}}_q$ -isomorphic to it.*

Proof. Assume that $b = d^2$ for some $d \in \mathbb{F}_q^*$. Let $m = ad^{-1}$, $u = d$ and $r = 0$. Then $E_{a,b}$ and E_m are isomorphic over $\overline{\mathbb{F}}_q$ by (2). \square

Proposition 4.5. *Let $E_{a,b} : y^2 = x^4 + ax^2 + b$ be a general Jacobi quartic curve defined over the finite field \mathbb{F}_q of characteristic > 3 , where $b(a^2 - 4b) \neq 0$. Assume that $j(E_{a,b}) \neq 1728$, then $E_{a,b}$ and the curve $E_{m,bm^2/a^2} : y^2 = x^4 + mx^2 + (bm^2/a^2)$ are isomorphic over $\overline{\mathbb{F}}_q$ for any $m \in \mathbb{F}_q^*$.*

Proof. Since $j(E_{a,b}) \neq 1728$, we have $a \neq 0$. From (2), for any $m \in \mathbb{F}_q^*$, let $u = \sqrt{\frac{a}{m}} \in \overline{\mathbb{F}}_q$ and $r = 0$, we know that $E_{a,b}$ and $E_{m,bm^2/a^2}$ are isomorphic over $\overline{\mathbb{F}}_q$. \square

For the elliptic curves $E_{a,b}$ with $j(E_{a,b}) \neq 0, 1728$, assume that the curve $E_{a,n}$ is $\overline{\mathbb{F}}_q$ -isomorphic to $E_{a,b}$, then there exist $u, r \in \overline{\mathbb{F}}_q$ with $u \neq 0$ such that

$$\begin{cases} 2au^2 = 2a - 3r, \\ (a^2 - 4n)u^4 = 3r^2 - 4ar + (a^2 - 4b), \\ r(r^2 - 2ar + a^2 - 4b) = 0. \end{cases} \quad (3)$$

just by replacing m to a in (2). Thus $r = 0$ or $r^2 - 2ar + a^2 - 4b = 0$. When $r = 0$, we have immediately that $n = b$. In the following, assume that $r \neq 0$. So we have $r = a + 2\sqrt{b}$ or $r = a - 2\sqrt{b}$, Therefore

$$n = \frac{a^2}{4} - \frac{3r^2 - 4ar + (a^2 - 4b)}{4u^4} = \frac{a^2(a - 2\sqrt{b})^2}{4(a + 6\sqrt{b})^2}. \quad (4)$$

or

$$n = \frac{a^2(a + 2\sqrt{b})^2}{4(a - 6\sqrt{b})^2}. \quad (5)$$

by substituting $u^2 = \frac{2a-3r}{2a}$ and $r = a + 2\sqrt{b}$ or $r = a - 2\sqrt{b}$ in the second equation of (3).

Assume first that b is not a square in \mathbb{F}_q . We claim that neither $\frac{a^2(a-2\sqrt{b})^2}{4(a+6\sqrt{b})^2}$ nor $\frac{a^2(a+2\sqrt{b})^2}{4(a-6\sqrt{b})^2}$ is an element of \mathbb{F}_q which is contradicted to $n \in \mathbb{F}_q$. In fact, if

$$\frac{a^2(a - 2\sqrt{b})^2}{4(a + 6\sqrt{b})^2} = \frac{a^2}{4(a^2 - 36b)^2} \cdot ((a - 2\sqrt{b})(a - 6\sqrt{b}))^2 \in \mathbb{F}_q,$$

then

$$((a - 2\sqrt{b})(a - 6\sqrt{b}))^2 = (a^2 - 8a\sqrt{b} + 12b)^2 \in \mathbb{F}_q.$$

Therefore, we must have $16a^3 + 192ab = 16a(a^2 + 12b) = 0$. Contradicts to the assumptions that $j(E_{a,b}) \neq 0$, 1728. Similarly, if $\frac{a^2(a+2\sqrt{b})^2}{4(a-6\sqrt{b})^2} \in \mathbb{F}_q$, one can get a contradiction again. This proves that if b is not a square in \mathbb{F}_q , and $E_{a,n}$ is $\overline{\mathbb{F}}_q$ -isomorphic to $E_{a,b}$, then we must have $n = b$. Therefore, by Proposition 4.5, when b is not a square element, the number of elliptic curves of the form $E_{a,b}$ with $j(E_{a,b}) \neq 0$, 1728 in each of its $\overline{\mathbb{F}}_q$ -isomorphism class is $q - 1$. Thus when b is not a square, the number of $\overline{\mathbb{F}}_q$ -isomorphism classes of curves of the form $E_{a,b}$ with $j(E_{a,b}) \neq 0$, 1728 is

$$\begin{cases} \frac{q-1}{2}, & \text{if } q \equiv 1, 7 \pmod{12}, \\ \frac{q-3}{2}, & \text{if } q \equiv 5, 11 \pmod{12}, \end{cases} \quad (6)$$

by Corollary 4.3.

On the other hand, assume that $b = d^2$ is a square in \mathbb{F}_q . We claim that neither $\frac{a^2(a-2\sqrt{b})^2}{4(a+6\sqrt{b})^2}$ nor $\frac{a^2(a+2\sqrt{b})^2}{4(a-6\sqrt{b})^2}$ is equal to b . In fact, if

$$\frac{a^2(a-2\sqrt{b})^2}{4(a+6\sqrt{b})^2} = b,$$

then

$$a^2(a-2d)^2 = 4d^2(a+6d)^2.$$

Thus

$$a(a-2d) = 2d(a+6d) \quad \text{or} \quad a(2d-a) = 2d(a+6d).$$

So

$$(a+2d)(a-6d) = 0 \quad \text{or} \quad a^2 + 12d^2 = 0,$$

that is $a^2 - 4b = 0$ or $a^2 = 36b$ or $a^2 + 12b = 0$, which is contradict to the assumptions that $j(E_{a,b}) \neq 0$, 1728. Similarly, if $\frac{a^2(a+2\sqrt{b})^2}{4(a-6\sqrt{b})^2} = b$, then one can get a contradiction again. Furthermore, we can check easily that $\frac{a^2(a-2\sqrt{b})^2}{4(a+6\sqrt{b})^2} \neq \frac{a^2(a+2\sqrt{b})^2}{4(a-6\sqrt{b})^2}$. This proves n has 3 choices in this case. Therefore, when b is a square element, the number of elliptic curves of the form $E_{a,b}$ with $j(E_{a,b}) \neq 0$, 1728 in each of its $\overline{\mathbb{F}}_q$ -isomorphism class is $3(q-1)$ by

Proposition 4.5, and then the number of $\overline{\mathbb{F}}_q$ -isomorphism classes of curves of the form $E_{a,b}$ with $j(E_{a,b}) \neq 0$, 1728 is

$$\begin{cases} \frac{q-7}{6}, & \text{if } q \equiv 1, 7 \pmod{12}, \\ \frac{q-5}{6}, & \text{if } q \equiv 5, 11 \pmod{12}, \end{cases} \quad (7)$$

when b is a square, by Corollary 4.3.

Adding together the numbers in (6), (7) above and 2 which corresponding the two special classes of curves with $j(E_{a,b}) = 0$ and $j(E_{a,b}) = 1728$, respectively, we have the following theorem.

Theorem 4.6. *Let $N_{a,b}$ be the number of $\overline{\mathbb{F}}_q$ -isomorphism classes of general Jacobi quartic curves $E_{a,b} : y^2 = x^4 + ax^2 + b$ with $(a^2 - 4b)b \neq 0$ defined over the finite field \mathbb{F}_q . Then we have*

$$N_{a,b} = \begin{cases} \frac{4q+2}{6}, & \text{if } q \equiv 1, 7 \pmod{12}, \\ \frac{4q-2}{6}, & \text{if } q \equiv 5, 11 \pmod{12}. \end{cases}$$

Remark 4.7. *We know from Proposition 4.4 that the number N_a of $\overline{\mathbb{F}}_q$ -isomorphism classes of Jacobi quartic curves E_a is equal to the number of $\overline{\mathbb{F}}_q$ -isomorphism classes of general Jacobi quartic curves $E_{a,b}$ with square b . Thus we have*

$$N_a = \frac{q-7}{6} + 2 = \frac{q+5}{6}$$

when $q \equiv 1, 7 \pmod{12}$ but

$$N_a = \frac{q-5}{6} + 1 = \frac{q+1}{6}$$

when $q \equiv 5, 11 \pmod{12}$ from the numbers in (7) since $a^2 + 12b = 0$ has no root, i.e., $j(E_{a,b}) \neq 0$, when b is a square and $q \equiv 5, 11 \pmod{12}$. Therefore we get the result in Theorem 3.3 again.

References

- [1] D.J. Bernstein and T. Lange, Analysis and optimization of elliptic-curve single-scalar multiplication. Contemp. Math., Vol 461, 1-20, Amer. Math. Soc., 2008.

- [2] O. Billet and M. Joye, The Jacobi model of an elliptic curve and side-channel analysis, AAEECC 2003, LNCS 2643, 34-42, Springer-Verlag, 2003.
- [3] D. V. Chudnovsky, and G. V. Chudnovsky, Sequences of numbers generated by addition in formal groups and new primality and factorization tests, *Advances in Applied Mathematics* 7, 385-434, 1986.
- [4] S. Duquesne, Improving the arithmetic of elliptic curves in the Jacobi model, *Information Processing Letters* 104(3), 101-105, 2007.
- [5] H. Hisil, G. Carter and E. Dawson, New formulae for efficient elliptic curve arithmetic, in *INDOCRYPT 2007*, LNCS 4859, 138-151, Springer-Verlag, 2007.
- [6] J.H. Silverman. *The Arithmetic of Elliptic Curves*, GTM 106, Springer-Verlag, Berlin, 1986.