

University of Louisville

## ThinkIR: The University of Louisville's Institutional Repository

---

Electronic Theses and Dissertations

---

5-2023

### The association of long working hours and the use of prescription sedatives among U.S. workers.

Emmanuel U. Ezekekwa  
*University of Louisville*

Follow this and additional works at: <https://ir.library.louisville.edu/etd>



Part of the [Clinical Epidemiology Commons](#), [Community Health and Preventive Medicine Commons](#), [Health Services Research Commons](#), [Occupational Health and Industrial Hygiene Commons](#), [Pharmacoeconomics and Pharmaceutical Economics Commons](#), and the [Pharmacy Administration, Policy and Regulation Commons](#)

---

#### Recommended Citation

Ezekekwa, Emmanuel U., "The association of long working hours and the use of prescription sedatives among U.S. workers." (2023). *Electronic Theses and Dissertations*. Paper 4117.  
Retrieved from <https://ir.library.louisville.edu/etd/4117>

This Doctoral Dissertation is brought to you for free and open access by ThinkIR: The University of Louisville's Institutional Repository. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of ThinkIR: The University of Louisville's Institutional Repository. This title appears here courtesy of the author, who has retained all other copyrights. For more information, please contact [thinkir@louisville.edu](mailto:thinkir@louisville.edu).

THE ASSOCIATION OF LONG WORKING HOURS AND THE USE OF  
PRESCRIPTION SEDATIVES AMONG U.S. WORKERS

By

Emmanuel U. Ezekekwa  
BPharm., University of Nigeria, 2010  
MHA., Western Kentucky University, 2017

A Dissertation  
Submitted to the Faculty of the  
School of Public Health and Information Sciences at the University of Louisville  
in Partial Fulfillment of the Requirements  
for the Degree of

Doctor of Philosophy in Public Health Sciences

Department of Health Management and System Sciences  
University of Louisville  
Louisville, Kentucky

May 2023

Copyright 2023 by Emmanuel U. Ezekekwa  
All rights reserved.



THE ASSOCIATION OF LONG WORKING HOURS AND THE USE OF  
PRESCRIPTION SEDATIVES AMONG U.S. WORKERS

By

Emmanuel U. Ezekekwa  
BPharm., University of Nigeria, 2010  
MHA., Western Kentucky University, 2017

A Dissertation Approved on

April 11, 2023

By the following Dissertation Committee:

---

Christopher E. Johnson, PhD, Dissertation Chair

---

Seyed M. Karimi, PhD, Dissertation Committee Member

---

Demetra Antimisiaris, PharmD, BCGP, FASCP, Dissertation Committee Member

---

Doug Lorenz, PhD, Dissertation Committee Member

## DEDICATION

This dissertation is dedicated to my parents for their unequivocal love and support. To the love of my life, Rahel, for her unwavering love, and my siblings, Dr. Chinedu and Emeka, you all made this possible.

## ACKNOWLEDGEMENTS

I express my gratitude to Dr. Christopher Johnson, my dissertation chair, for his support, mentorship, and encouragement throughout the program. I extend my appreciation to Dr. Seyed Karimi for his invaluable and relentless guidance, feedback, and assistance throughout the dissertation process. I would like to give special thanks to Dr. Demetra Antimisiaris for her amazing and indispensable mentorship, support, and kindness throughout the program and dissertation. Also, I would like to thank Dr. Doug Lorenz for his kindness, meticulous and critical assistance in completing this dissertation.

I am grateful to Drs. Judah Thornewill, Bob Esterhay (of blessed memory), J'Aime Jennings, Lee Bewley, all the Faculty, colleagues, and staff in the School of Public Health and Information Sciences for their support in the program. Lastly, many thanks to Dr. Ekene Ezeani, Dr. Raphael Fumey, Dr. Umar Kabir, Dr. Anthony Okafor, Brandon Mitchell, extended family, and friends for their encouragement through the years.

## ABSTRACT

### THE ASSOCIATION OF LONG WORKING HOURS AND THE USE OF PRESCRIPTION SEDATIVES AMONG U.S. WORKERS

Emmanuel U. Ezekekwa

April 11, 2023

**BACKGROUND:** Meeting the needs of a round-the-clock and globalized society has led to an increase in long working hours. This trend has been accompanied by a corresponding rise in sleep disorders and subsequent use of sedating medications.

Overtime hours have been associated with adverse health outcomes such as cardiovascular diseases, symptoms of psychological distress, and health behaviors, including risky intake of alcohol and smoking.

Hence, the main objectives of this three-paper dissertation were to examine the multi-faceted relationship between working hours, the use of prescription sleep aids, the onset of psychological distress, and the use of health care services.

**METHODS:** The 2010-2019 Medical Expenditure Panel Survey (MEPS) data was utilized.

The first paper investigated the relationship between working hours and the use of prescribed sedating medications. Different regression models were employed, ranging



from multivariable linear regression, Tobit regression, Heckman regression, and multivariable logistic regression. The second paper utilized a fixed-effect linear model in evaluating the relationship between working hours and the onset of psychological distress. The third paper also employed fixed-effect estimators in modeling the association between working hours, health care utilization, and the use of sedating medications.

**FINDINGS:** Long working hours were associated with increased odds of using sleep aids and medications with sedative properties. Females had a higher likelihood of using sleep aids when compared to males. Also, professional services had the highest likelihood of using sleep medications. Over time hours was associated with the onset of psychological distress, with differences in the risk of onset across gender. Respondents working very long hours had the highest odds of using outpatient medical services. This association between very long hours and the use of outpatient services was significantly more pronounced in respondents using medications with sedating side effects.

**CONCLUSIONS:** Long working hours were associated with an elevated risk of using sedating medications, onset of psychological distress, and healthcare utilization. This highlights the probable negative impact of overtime hours on the health status of individuals. Implementing policies that encourage work-life balance and aid interventions that decrease work-related stress may help in mitigating risks associated with long work hours.

## TABLE OF CONTENTS

|  | PAGE |
|--|------|
| DEDICATION.....  | iii  |
| ACKNOWLEDGEMENTS.....  | iv   |
| ABSTRACT.....  | v    |
| LIST OF TABLES.....  | ix   |
| LIST OF FIGURES.....   | x    |
| CHAPTER ONE: EXAMINING THE RELATIONSHIP BETWEEN LONG WORKING HOURS AND THE USE OF PRESCRIPTION SEDATIVES AMONG U.S. WORKERS.....         | 7    |
| 1.0. OVERVIEW.....   | 7    |
| 1.1. INTRODUCTION.....   | 9    |
| 1.2. METHODS.....  | 13   |
| 1.3. RESULTS.....  | 21   |
| 1.4. DISCUSSION.....   | 29   |
| 1.5. CONCLUSION.....   | 34   |
| CHAPTER TWO: LONG WORKING HOURS AND ONSET OF PSYCHOLOGICAL DISTRESS: A LONGITUDINAL ANALYSIS.....  | 46   |
| 2.0. OVERVIEW.....   | 46   |
| 2.1. INTRODUCTION.....   | 48   |
| 2.2. METHODS.....  | 53   |
| 2.3. RESULTS.....  | 62   |
| 2.4. DISCUSSION.....   | 66   |
| 2.5. CONCLUSION.....   | 72   |
| CHAPTER THREE: ASSOCIATIONS BETWEEN WORKING HOURS, HEALTH CARE UTILIZATION, AND PRESCRIPTION MEDICATIONS WITH SEDATIVE SIDE EFFECTS..... | 80   |
| 3.0. OVERVIEW.....   | 80   |
| 3.1. INTRODUCTION.....   | 82   |

|                               |     |
|-------------------------------|-----|
| 3.2. METHODS .....            | 84  |
| 3.3. RESULT .....             | 92  |
| 3.4. DISCUSSION.....          | 97  |
| 3.5. CONCLUSION.....          | 101 |
| CHAPTER FOUR: CONCLUSION..... | 103 |
| 4.1. FUTURE RESEARCH .....    | 107 |
| REFERENCES .....              | 117 |
| APPENDICES .....              | 139 |
| APPENDIX CHAPTER 1: .....     | 140 |
| APPENDIX CHAPTER 2.....       | 187 |
| APPENDIX CHAPTER 3.....       | 197 |
| CURRICULUM VITAE.....         | 212 |

| LIST OF TABLES   | PAGE |
|--|------|
| Table 1. Descriptive statistics of work hours and use of prescription medications .....  | 37   |
| Table 2. Bivariate Analysis .....  | 40   |
| Table 3. Regressions for the Use of Prescription Sleep Aids.....   | 42   |
| Table 4. Regressions for the Use of Prescription Medications with Sedative Properties .  | 44   |
| Table 5. Descriptive statistics of work hours and psychological distress. ....   | 75   |
| Table 6. Association Between long working Hours and the onset of psychological<br>distress: Fixed effects regression analysis. ....          | 77   |
| Table 7. Descriptive statistics of work hours and health care utilization.....   | 110  |
| Table 8. Association Between Working Hours and Health Care Utilization: Fixed Effects<br>Regression Analysis – Overall Sample .....          | 112  |
| Table 9. Association Between working Hours and Outpatient Healthcare Services – ....   | 113  |
| Table 10. Association Between working Hours and Health care Utilization: Fixed effects<br>regression analysis – Among Medication Users ..... | 115  |

## LIST OF FIGURES

|   | PAGE |
|---|------|
| Figure 1. Conceptual framework depicting the influence of working hours on the use of sleep aid drugs (adapted from Aday & Andersen, 1974).....   | 36   |
| Figure 2. Conceptual model of the possible causal relationship between exposure to long working hours and psychological distress (adapted from Pega et al., 2021; Rugulies et al., 2021).....                         | 74   |
| Figure 3. Onset of psychological distress according to working hours from the fixed effects regression model of the overall sample. ....  | 78   |
| Figure 4. Onset of psychological distress according to working hours from the fixed effects regression model of all men sample. ....  | 79   |
| Figure 5. Onset of psychological distress according to working hours from the fixed effects regression model of all women sample. ....  | 79   |
| Figure 6. Conceptual framework depicting the association between working hours, the use of prescription medications with sedative side effects, and health care utilization (adapted from Aday & Andersen, 1974)..... | 109  |
| Figure 7. Use of outpatient services according to working hours from the fixed effects regression model of the overall sample. ....   | 114  |
| Figure 8. Use of outpatient services according to working hours from the fixed effects regression model among users of medications with sedative side effects. ....   | 116  |

## INTRODUCTION

Globally, long working hours persist as a prevalent phenomenon, and the United States is no exception. According to a report by the International Labor Organization, the U.S. ranked as the fourth-highest country in terms of the proportion of workers who work overtime weekly, indicating the widespread practice of extended working hours in the country (Bannai et al., 2015).

Several factors are responsible for this trend of long working hours in the U.S., such as motivations for increased wages, career development, and strong cultural influences (Li et al., 2019). One of the primary reasons is the motivation for increased wages, as many employees earn more financial rewards from working overtime hours. Also, the incentive for career development is another crucial factor driving the trend, as some employees believe that working longer hours could help them climb the corporate ladder more quickly. Additionally, cultural influences such as the "workaholic" mentality also contribute to the practice of working long hours in the U.S.

It is crucial to note that while long working hours may seem like an efficient way to increase productivity, it can have deleterious consequences on employees' physical and mental well-being, leading to sleep disorders, fatigue, burnout, and stress (Cheng et al., 2014; Lunde et al., 2016; Skogstad et al., 2019). Moreover, the quality of work produced during extended working hours may not be up to par with that of regular working hours, which can have adverse effects on organizational performance.

## **Long working hours, Sedating Medications, and Health Outcomes**

The continuous rise in the prevalence of long working hours has been accompanied by an increase in sleep disorders (Virtanen et al., 2009; Nakashima et al., 2011; Cheng et al., 2014). Sleep disorders, while being under-recognized, remain a persistent public health challenge (Hale, Troxel, & Buysse, 2020). The Institute of Medicine estimates that between 50 to 70 million people experiences sleep disorders in the U.S. (IOM, 2006). The American psychiatric association defines sleep disorders as challenges with the timing, quantity, and quality of sleep, which creates problems associated with distress and functioning during the daytime (APA, 2019). Insomnia is the most common sleep disorder, and others include obstructive sleep apnea, narcolepsy, and restless leg syndrome (APA, 2019). Sleep disorders have been associated with several chronic ailments such as cardiovascular diseases, depression, obesity, diabetes, and increased risk of mortality (Roane & Taylor, 2008; Chandola, Ferrie, Perski, Akbaraly, & Marmot, 2010; Cappuccio FP et al., 2010; CDC, 2018). It also extends to increased risk for accidents at the workplace and vehicle crashes and has been estimated to cost about \$ 150 billion annually through indirect costs. These include workplace accidents, presenteeism, and absenteeism (WEF, 2010).

Sleep and substance abuse disorders have been closely linked (Roane & Taylor, 2008; Mahfoud, Stroom, & Budur, 2009). Also, it is common practice to use medications, including prescribed and controlled medicines, for their side effects, such as sedating effects, and for recreational purposes (Lopez-Quintero et al., 2021). Thus, the use of medications with secondary sedating side effects, such as pain medications, anti-histaminic drugs, and anti-depressants, has increased (Votaw et al., 2019). From 2003 to

2012, there was a notable surge of more than 65% in the treatment of individuals with sedative-tranquilizer use disorder (Abuse, 2014). Additionally, there has been a substantial increase in the mortality rate associated with benzodiazepine-related overdoses, with the numbers soaring over 400% from 1999 to 2013 (Bachhuber et al., 2016). Sedating medications have been linked with adverse health outcomes, including increased risk of cardiovascular diseases, morbidity, and mortality (Mailliet, Galloux & Poisson, 2001; Mallon, Broman, & Hetta, 2009; Kivimäki et al., 2015; Kim et al., 2018; Skogstad et al., 2019).

The prevalence of long working hours in the U.S. working population is expected to lead to a continuous increase in the use of sedating medications and other associated substance abuse to aid sleep. However, while many studies have highlighted the relationship between long working hours and high risky consumption of alcohol, there is a paucity of studies examining the influence of working hours on the use of sedating medications in the working population.

Also, the studies examining the existing relationship between working hours and psychological distress have been mixed (Afonso et al., 2017; Li et al., 2019; Virtanen et al., 2011; Wong et al., 2019 Vs. Angrave & Charlwood, 2015; Lee et al., 2017; Rugulies et al., 2019) mainly due to study designs challenges. Despite the high prevalence of long working hours and psychological distress in the U.S., longitudinal studies examining the relationship between working hours and psychological distress among U.S. workers are sparse.

Furthermore, the increase in the incidence and prevalence of chronic diseases associated with long working hours is expected to reflect in the elevated use of health



services by employees. Also, the literature informs that long working hours are linked with reduced utilization of preventive health services among employees. This is primarily attributed to the lack of time available to workers putting in overtime hours (Fell et al., 2007). Preventive health services are important in sustaining the health of individuals, especially those with increased risk of chronic diseases. However, studies investigating the associations between long working hours, healthcare utilization, and the use of medications in the U.S. is sparse.

## **Overview**

This dissertation presents a unique approach in its three-staged strategy in investigating the multi-faced relationships between working hours, the use of prescription sleep aids, the onset of psychological distress, and the use of health care services among U.S. workers. A nationally representative panel of full-time workers in the U.S. to examine these relationships.

The first manuscript applied the Andersen healthcare utilization model to conceptualize the relationship between long working hours and the use of prescription sleep aids. This is the first study to examine the relationship between long working hours and the use of sleep aid medications, along with the use of medications with sedative properties. Methodologically, different regression models were utilized, including multivariable linear regression, Tobit regression, Heckman regression, and multivariable logistic regression, to ensure consistency, robustness, and reliability of observed associations. The independent variable and outcome variables were working hours and the use of sedating medications, respectively. This paper provides an initial insight into

characterizing the important association between working hours on the use of prescribed sedating medications.

The second manuscript utilized the longitudinal structure of the data to investigate the relationship between long working hours and psychological distress. The study utilized a modified WHO logic model (Pega et al., 2021; Rugulies et al., 2021) to conceptualize this relationship. The model informs that long working hours influence the onset of psychological distress through two causal pathways—psycho-physiological and health-behavioral pathways.

This study represents an improvement over previous research that was hampered by problems of reverse causality and cross-sectional data. Methodologically, the study assessed this relationship in individuals without psychological distress at baseline. The assumption here is that individuals working overtime hours will have a higher risk of psychological distress. This provides longitudinal evidence of the existing relationship between working hours and the onset of psychological distress among U.S. workers.

The concluding or third manuscript of this dissertation examined the relationship between working hours, health care utilization, and the use of sedative medications in the U.S. It utilized the longitudinal structure of the data to investigate these relationships. Employing the Andersen healthcare utilization model to conceptualize these various associations. Fixed effect estimators were utilized in modeling the relationship between working hours, the use of sedating medications, and health care services among full-time employees in the U.S. An assumption here is that individuals working long hours will have a higher use of healthcare services, especially healthcare services with less time requirement. Also, individuals working overtime hours burdened with chronic diseases

are expected to have higher use of health care services. It operationalized the use of health care services through five dependent variables: annual outpatient visits, office-based visits, emergency department visits, hospital discharges, and dental visits. Furthermore, further analyses were performed on characterizing annual outpatient visits to further understand the observed relationships.

Together, these manuscripts address crucial gaps in our current knowledge of the multi-dimensional relationship between working hours and health outcomes. The findings from this research offer a solid foundation of evidence and valuable insights into the relationship between working hours, the use of sleep aids, and their negative impact on health outcomes. Therefore, these findings can serve as a guiding tool for employers, health systems, and researchers in creating interventions and making recommendations regarding work-based policies, with the ultimate aim of enhancing the health and well-being of employees.

## CHAPTER ONE: EXAMINING THE RELATIONSHIP BETWEEN LONG WORKING HOURS AND THE USE OF PRESCRIPTION SEDATIVES AMONG U.S. WORKERS

### 1.0. OVERVIEW

**OBJECTIVES:** The demands of a round-the-clock service and globalized society have led to an increase in long working hours. This trend has been accompanied by a corresponding rise in sleep disorders. Additionally, sedative-tranquilizers have been reported as the third most commonly misused drug class in the U.S. Given the prevalence of long working hours among the U.S. working population, it is expected that there will be a continued increase in the use of sleep aid medications among this population, along with an associated increase in substance abuse to aid sleep.

**METHODS:** The 2010-2019 Medical Expenditure Panel Survey (MEPS) data from the University of Minnesota's Integrated Public Use Microdata Series (IPUMS) was utilized. Sleep aids and medications with sedation as a side effect were identified. Furthermore, we employed different regression models ranging from multivariable linear regression, Tobit regression, Heckman regression, and multivariable logistic regression, to ensure consistency, robustness, and reliability of associations between working hours and the use of medications.

**RESULTS:** Overall, a sample of 81,518 observations of full-time workers was analyzed. Working 56 hours or more per week was significantly associated ( $p < 0.05$ ) with an increased odds of using sleep aids and medications with sedative properties by 13% (Adjusted Odds Ratio, aOR = 1.13, 95% Confidence Interval, CI=1.01:1.26) and 9% (aOR=1.09, 95% CI=1.03:1.16), respectively more than that among those who worked fewer hours. Females in our study had a higher likelihood (aOR=1.11, 95% CI=1.05:1.19) of using sleep aids when compared to males. Also, professional services had the highest likelihood (aOR=1.31, 95% CI=1.14:1.50) of using sleep medications.

**CONCLUSION AND POLICY IMPLICATION:** We found that long working hours were significantly associated with an elevated use of sleep aids and medications with sedative properties among U.S. workers. Specifically, female workers and individuals working in professional services had the highest likelihood of using sleep medications. Employees should be enlightened on the implications of long working hours on their health.

## 1.1. INTRODUCTION

Long working hours have increased with the continuous attempts at meeting the needs of 24-hour service and a globalized society. The rise in poor sleep quality has accompanied this increase. Also, sleep and substance abuse disorders have been closely linked (Mahfoud et al., 2009; Roane & Taylor, 2008; Weissman et al., 1997). A previous study found that one in three Americans frequently gets less than the recommended amount of sleep (Liu et al., 2016). Furthermore, long working hours have been linked to adverse health outcomes and behaviors such as cardiovascular diseases (Bannai & Tamakoshi, 2014; Cheng et al., 2014; Grosch et al., 2006; Kim et al., 2016; Lunde et al., 2016; Wong et al., 2019), depression and anxiety (Cheng et al., 2014; Lallukka et al., 2008; Lunde et al., 2016; Skogstad et al., 2019), risky alcohol use, and smoking (Bannai & Tamakoshi, 2014; Virtanen et al., 2015).

Sedative and hypnotic medications, commonly referred to as “sleep aids,” are used to initiate or maintain sleep due to their known ability to suppress the central nervous system (Chong et al., 2013). Sedative – tranquilizers have been reported as the third most misused drug class by adolescents and adults in the United States population (Votaw et al., 2019). For example, the prescription of Benzodiazepines – a predominant class of sedatives, doubled from 2003 to 2015 (Agarwal & Landon, 2019).

Furthermore, it is common practice to use medications, including prescribed and controlled medicines, for their side effects, such as sedating effects, and for recreational purposes (Lopez-Quintero et al., 2021). As such, the use of medications with secondary sedating side effects, such as pain medications, anti-histaminic drugs, and anti-

depressants has increased (Votaw et al., 2019). Between 2003 to 2012, treatment for sedative – tranquilizers use disorder increased by over 65% (Abuse, 2014). Also, there has been a significant rise in benzodiazepine-related overdose mortality by over 400% increase between 1999 and 2013 (Bachhuber et al., 2016). Recently, a study (Lopez-Quintero et al., 2021) found that 40% of their study population that used prescription medications extra-medically used pain relievers as sleep aids. Other medical literature regarding the off-label use of medications not indicated for insomnia endorses the concept that people utilize a variety of products with sedative properties for sleep. The often referenced “Up to Date” peer reviewed clinical practice reference includes a section on medications that can be used for insomnia off label, such as several antidepressants (Neubauer et al., 2021). A national study of prescription medications for insomnia reported that 55% of persons using medications for insomnia also concurrently used sedating medications, with 10% taking three or more other sedating medications concurrently (Bertisch et al., 2014).

The prevalence of long working hours in the U.S. working population is expected to lead to a continuous increase in the use of sleep aid medications in this population with other associated substance abuse to aid sleep. However, despite the availability of studies highlighting the relationship between long working hours and the consumption of alcohol, the relationship between long working hours and the use of sleep aid medications has rarely been examined. Notably, there is a higher prevalence of the extra-medical use of prescription medications for sleep in the U.S. compared to several European countries (Lehne et al., 2018; Schepis et al., 2018). Also, studies have linked the use of hypnotics, over-the-counter sleep medications to adverse health outcomes such as cardiovascular

diseases (Jehle et al., 2013; Kim et al., 2018; Mailliet et al., 2001; Mallon et al., 2009) in the general population. With benzodiazepines associated with an increased incidence and risk of mortality from cardiovascular diseases (Kim et al., 2018).

To our knowledge, this is the first study examining the relationship between long working hours and the use of sleep aid medications, along with the use of medications with sedative properties. Hence, the key objective of this study was to characterize the relationship between working hours on the use of sleep aids and medications with sedative properties. We applied the Andersen healthcare utilization model to conceptualize our research questions (Andersen, 1995). This model is relevant to our research as it posits that a person's health behavior and healthcare service use are significantly influenced by enabling, predisposing, and needs factors.

Our conceptual model (*figure 1*) explains that working hours influence the mutable population characteristics (specifically the need factors such as health status) and recurrently influence individuals' health behaviors. Working hours represent the environment, influenced by the occupation, and have been reported to affect health behaviors such as the consumption of alcohol (Virtanen et al., 2015; Wong et al., 2019). At the same time, the health behavior constructs in our model represent the use of sleep aids or medications with sedative properties.

The population characteristics include predisposing, enabling, need, and motivating factors. The predisposing construct represents individuals' factors or conditions that influence their use of health services (Andersen & Newman, 1973; Andersen, 1995). This construct is classified mainly as demographic and social factors. Demographic factors include age, gender, and race. Health disparities exist in the U.S.



based on predisposing constructs such as race, socioeconomic status, age, and gender (Lee, Black, & Held, 2019). At the same time, the social structure includes various factors that define an individual status in the community, such as marital status, education, and occupation.

Additionally, the enabling factors refer to the means individuals have available to use these services. This includes resources like income, health insurance, and the individual's region (Andersen and Newman, 1973). An individual's insurance status will likely determine if the individual will use health services and obtain prescription medications for sleep. However, health insurance is an outcome variable as individuals working 30 or more hours per week are eligible for employee health insurance in the U.S. (Carroll & Miller, 2019). Furthermore, the level of income increases the ability to access health care. The need factor is represented by health status, which was self-reported and thus, a perceived need.

Also, motivating factors influences the amount of time an individual spends working. Professionals and persons with higher socioeconomic status are motivated to work long hours due to its potential financial and career rewards. However, individuals might be co-opted to work long hours due to the financial strains they face, even if they earn meager wages. As measures of motivation, we used the family size and the receipt of food stamps. The presence of large family sizes may bring about heightened financial demands for families, while receipt of food stamp suggests the presence of financial difficulties associated with food insecurity. Family size and receipt of food stamps have been utilized as indicators of financial strains in prior studies (Bhattarai et al., 2005; Kabir, 2021).

The findings from this research could ground evidence and insights into the relationship between working hours, use of sedative-hypnotic, sleep aid substances, and adverse health outcomes. Hence, the findings can guide employers, health systems, and researchers in designing interventions and recommendations for developing work-based policies to improve the health status of employees.

## **1.2. METHODS**

### **1.2.1. Data Sources**

We utilized the 2010-2019 Medical Expenditure Panel Survey (MEPS) data. This is a longitudinal national probability survey conducted by the Agency for Healthcare Research and Quality (AHRQ) and the National Center for Health Statistics (NCHS) (Quality, 2020). It collects data on access, utilization, and expenditures on health services, among other health information, for civilian non-institutionalized Americans. MEPS draws its sample from the National Health Interview Survey (NHIS) respondents from the preceding year. It uses an overlapping panel design to collect data from the survey respondents. The panel is designed to include five rounds of interviews over two full calendar years. Typically, for each participant in a calendar year, data from interview rounds one and two are collected in the first year, while data for rounds three, four, and five are collected in their second year.

The MEPS-Household Component (MEPS-HC), ongoing since 1996, provides information on the participants' demographic, health and employment status, socioeconomic characteristics, health care utilization, health care expenditures by payer, insurance coverage, and healthcare satisfaction. To increase the precision of estimates, MEPS and NHIS oversample subgroups of minorities such as African Americans and

Hispanics. After considering nonresponse rates from the NHIS, the overall MEPS response rates range between 46% and 71%. The MEPS HC enables the production of national estimates. It can also be analyzed at the person or event level.

MEPS elicit information directly from providers in its Medical Provider Component (MEPS-MPC). The main aim of the MEPS-MPC is to supplement or replace information collected during the MEPS-HC survey. The MEPS-MPC includes a pharmacy component as a subcomponent, where it collects medication detail information, the national drug code (NDC), and the medication's generic name. However, unlike the MEPS-HC, MEPS-MPC is not designed to produce national estimates. Thus, we weighted only the MEPS-HC component of the study in the descriptive statistics for the results in the main results. However, following (Solon et al., 2015), we also conducted our analysis using sample weights and accounting for the complex survey design of the MEPS. The weighted results are reported in *Appendices 6 to 10*.

### **1.2.2. IRB Approval Statement**

The study was approved as exempt by the University of Louisville's Institutional Review Board (IRB # 21.0849). The publicly available de-identified dataset was used for this study, and the research did not meet the "Common Rule" definition of human subjects' research. Thus, it was classified as Non-Human Subjects Research (NHSR).

### **1.2.3. Data Organization**

We employed a pooled sample design, and listwise deletion was used to eliminate missing cases. The dataset from MEPS comes in twenty varying files. Therefore, to ease the data management, we employed the MEPS surveys refined and organized at the University of Minnesota's Integrated Public Use Microdata Series (IPUMS). The

National Institute of Child Health and Human Development (NICHD) funds the IPUMS MEPS. This is designed to aid the researcher's use of the MEPS. It provides integrated data files for each year of the MEPS-HC survey with the ease of preselecting the variable before downloading the dataset (Blewett, 2020).

In the next step, we merged the IPUMS MEPS-HC with the prescription medicine event files in the MEPS Pharmacy component. The prescription medicine event files report prescribed medicines obtained or purchased by the participating households. Each record represents a singular prescribed medicine reported for the person within the survey round. A round is approximately 4 months which is the space between each round of data collection. After downloading prescription data files for 2010-2019, the files were appended and merged to the MEPS-HC component. This linking of the two datasets was done using the "MEPSID," which represents the unique households' identifier in both datasets.

#### **1.2.4. Dependent Variable**

The dependent variables in this study were the use of sleep aids and medications with sleep aid properties, extracted from the prescription medicines event files of the MEPS pharmacy component. We defined a prescription as any prescription of a specific medication with its accompanying supply frequency (Chen et al., 2020). We used the number of prescriptions recorded as a continuous dependent variable rather than the quantity of medications recorded. Prescription for a medication represents a reliable need for the medication and can be compared across drug types. On the other hand, the

quantity of medications may not be directly comparable across drug types as they do not have the same dose-effect relationship.

Using the Multum Lexicon Therapeutic class Codes in MEPS, sleep aids were identified. The Multum variables classification in MEPS prescribed medicine files represented the most recent classification of the medications available when the data was recorded. Examples of these sleep aids medication classes include anxiolytics, sedatives, and hypnotics.

To identify medications with sedation as a side effect, we used Wolters Kluwer's Facts and Comparisons®, also known as "Drug Facts and Comparisons", a referential drug resource to identify medications with adverse effects (Comparisons eAnswers., 2021). Medications with drowsiness, dizziness, feelings of sluggishness, or heaviness listed as common or serious adverse effects were defined as having a potential for sedation as an adverse effect (*Appendix 1*). Facts and Comparisons is a product of Wolters Kluwer Health, which is a compendium offering comprehensive drug, disease, and clinical reports. It gives users access to several databases, modules, and medication reference tools for pharmacists and drug information experts. It has been reported as one of the most preferred drug resource used by drug information experts (Grizzle et al., 2019; Roblek et al., 2015). It was recorded to have a completeness score of 95.8%, computed with index value measures such as adverse drug reactions, drug interactions, contraindications, indications, and unlabeled uses (Clauson et al., 2007; Roblek et al., 2015). Facts and Comparisons® was ranked top alongside Micromedex, Clinical Pharmacology, and Lexi-Comp (Clauson et al., 2007; Hanrahan & Cole, 2014). It was grouped among the most accurate and user-friendly medication identifiers with a score of

96.5% (Jackevicius et al., 2019). Furthermore, it was found to have a sensitivity of 93% and a positive predictive value (PPV) of 95.3% in finding black box warnings (Cheng et al., 2010).

### **1.2.5. Primary Independent Variable**

The primary independent variable is work hours per week. The MEPS-HC recorded this variable if respondents in the survey were sixteen years and older and had a current main job. It measured the number of hours worked per week, excluding the unpaid travel time transiting to and from the job. We used 30 hours/week as a baseline as they are considered full-time workers and eligible for health insurance coverage in the U.S. (Carroll & Miller, 2019). Previous research informs that individuals working less than standard working hours, such as part-time workers, could belong to a different health-selected category, such as pre-existing health conditions, or might work less due to family responsibilities (Kivimäki et al., 2015; Li et al., 2020; Virtanen et al., 2012). As such, these groups of individuals working fewer hours may not serve as appropriate comparators.

To better understand medications and use trends associated with work hours, we generated four alternatively-used dummy variables based on weekly work hours. Specifically, our first dummy variable was assigned 1 if an individual worked 36 hours or more per week (h/week), and 0 otherwise. Our second dummy variable took the value of 1 if an individual worked 46 h/week, and 0 otherwise; the third dummy variable was assigned 1 if an individual worked 56 h/week, and 0 otherwise; the fourth dummy variable took the value of 1 if an individual worked 66 h/week, and 0 otherwise. These stratifications were carried out with the aim of identifying the best threshold to

dichotomize long working hours in relation to the use of sleep aids and medications with sedative side effects.

### **1.2.6. Covariates**

The selection of covariates was guided by the conceptual framework of the study using the Gelberg-Andersen behavioral health model. We used age, race, marital status, family size, education, occupation, and panel year as predisposing factors. We categorized age, race, marital status, and family size into three: respectively, 18-26, 27-64, and  $\geq 65$  years; White, Black, and other; married, divorced (including widowed and separated), and unmarried; and  $\leq 3$ , 4-6, and  $> 6$  members. We used age categories aligned with changes in health insurance coverage. The U.S. Patient Protection and Affordable Care Act stipulates that individuals are eligible to be covered by their parent's health insurance until age 26 (Croteau et al., 2021; Patrick & Yang, 2021; Weir et al., 2018).

Additionally, education, as a covariate, was recoded into two categories: having less than a college education and having some college or more education. Occupation is the specific work category of the respondents and was recoded into six categories: natural resources (including mining and construction), leisure (including hospitality services), trade (including wholesale and retail trade), professional services, manufacturing, or other sectors. Panel year, the first year a respondent was surveyed, accounts for variations in healthcare during the study period that could influence medication use and relative health outcomes.

The enabling factors include region, income, receipt of food stamps, and the first-round health status. Health insurance status is an outcome variable of the primary

independent variable – working hours. Thus, it was excluded as a covariate to avoid multicollinearity, as work hours predicts health insurance status. U.S. workers working 30 or more hours per week are eligible for employee health insurance (Carroll & Miller, 2019). MEPS region variable is the census region recorded as northeast, mid-west (including northcentral), south, and west. We recoded income into two: poor to low income (less than 100% to 199% of the federal poverty line) and middle to high income (200% and above the federal poverty line). The receipt of food stamps is a yes/no variable. We recoded the self-reported health status into four: excellent (combination of excellent and very good), good, fair, and poor.

### **1.2.7. Study Population**

**The inclusion criteria include** participants that are 18 years and above, participated in survey years 2010 to 2019 (panel 14 to 22), and considered full-time workers (worked at least 30 h/week). While the exclusion criteria were part-time workers (worked less than 30 h/week) and participants with any missing outcome or predictor variables.

### **1.2.8. Data Analysis**

The data were analyzed using Stata Statistical Software: Release 15 (StataCorp, 2021). We weighted the MEPS-HC component to estimate the descriptive statistics. Considering the complex survey design of the MEPS, we accounted for the weight, clusters, and strata. We used exploratory, descriptive statistics to examine the distribution of independent variables. We examined the frequency distributions of our study's participant categorical variables. The continuous variables' means, standard deviation,



median, and interquartile ranges were also assessed. This was followed by examining the distribution of the participant characteristics across both dependent variables (use of sleep aids and use of medications with sedative properties) using the Chi-square test.

Furthermore, we employed different regression models ranging from multivariable linear regression, Tobit regression, Heckman regression, and multivariable logistic regression, to ensure consistency, robustness, and reliability of associations between the independent and dependent variables in our study.

We first utilized a multivariate linear probability model to estimate the association between the number of prescriptions of sleep aids or medications with sleep aid properties with working hours. Since the data on the outcome variable (number of prescriptions) were censored at zero due to the short period of observation, a linear regression might not be the best approach for analyzing the data. Data on the number of prescriptions were collected in five rounds over two calendar years. As a result, prescriptions and use of medications that occurred outside of these rounds or survey period would be reported as zero. Additionally, no information is collected for over-the-counter sleep aids or medications with sedative properties. Thus, we employed Tobit regression to identify the factors related to the left-censored nature of the data. The Tobit regression marginal effects were then compared with the multivariate linear probability model.

An important limitation of the Tobit model is assuming a normal distribution for the error terms (Cameron & Trivedi, 2010). A potential remedy for the problem is to estimate a two-part model: the selection into using medications versus the degree of medication utilization among users. The use of a two-part model is justified if the error

terms of the two parts are independent. Therefore, we proceeded to test for independence using the Heckman regression model (Cameron & Trivedi, 2010; Koné et al., 2019).

Subsequently, we conducted an equivalent multivariate logistic regression to model the association between working hours and the use versus nonuse of medications.

Additionally, multivariable linear regression was performed to model the association between working hours and the use of medications among identified users. In this multivariable linear regression, the dependent variables were the number of prescriptions of sleep aids and medications with sedative properties among the identified users. In contrast, the independent variable remained the working hours of the identified users.

### **1.3. RESULTS**

#### **1.3.1. Descriptive Statistics**

*Table 1* describes the characteristics of the study sample stratified by different work hour cutoffs. We stratified work hours into four categories, with a cutoff of ten hours intervals using four dummy variables: 36 h/week, 46 h/week, 56 h/week, and 66 h/week. Overall, the number of records was 81,518, representing 98,995,693 individuals in the United States. Most of the participants worked 36 hours or more per week (87,383,592), while few worked 66 hours or more per week (2,185,941). More respondents utilized medications with sedative properties (34.5%), while the prevalence of prescription sleep aids use was substantially smaller (6.2%). The average numbers of prescriptions of sleep aids use and medications with sedative properties in our sample were 2.83 (standard deviation, SD=2.29) and 2.82 (SD=3.46), respectively, within the

survey round. A round is approximately four months, the space between each round of data collection.

The use of sleep aids was most common among participants that worked 56 hours or more per week, with a prevalence rate of 7.0% and a mean prescription sleep aid number of 3.16 (SD=2.58) among users. This group was followed by those who worked 66 hours or more per week, with a prevalence of 6.6% and a mean prescription sleep aid number of 3.06 (SD=2.79) among users. Similarly, the use of medications with sedative properties was the highest among participants that worked 56 hours or more per week, with a mean prescription sleep aid number of 3.18 (SD=3.54) among users and a prevalence rate of 36.1%. This group were followed by individuals that worked 66 hours or more per week with a mean of 3.13 (SD=3.36) among users and a prevalence rate of 36.4%.

Under predisposing factors, most respondents were between the ages of 27 to 64 years (85.3%), 83.6% of the sample identified as White, followed by 10.1% as Black, and 6.4% as Other. More than half of the sample were female (55.5%), and married respondents accounted for over half of the sample (57.4%). Most of the respondents (68.3%) had at least a college education. By occupation, professional services (39.5%) accounted for the largest proportion of the sample, and the largest portion of the respondents was from the survey year 2014 (11.0%).

By enabling factors, the largest proportion of the participants resided in the southern region of the U.S. (39.3 %), followed by those living in the North Central-Midwest (24.2%). In comparison, the Northeast (15.2%) had the least proportion. Most of the respondents were classified as middle to high-income earners (84.4%) – who earned

at least 200% and above the federal poverty threshold. In comparison, 15.6% were classified as poor to low income or earned below 200% of the federal poverty threshold. Additionally, by need factors, most of the respondents were either in excellent (14.9%) or good health (74.3%). Also, the majority of the respondents had a small family size of less than or equal to 3 members (69.3%), and 94.2 % of the sample did not receive food stamps.

### **1.3.2. Bivariate Analysis**

The Chi-square test of independence analysis in *Table 2* yielded a statistically significant association (with p-values,  $p$ , smaller than 0.05) between working 56 hours or more per week and the use of prescription sleep aids. The corresponding association was statistically significant at the 10% level for 36 or more and 46 or more work hours per week but statistically insignificant for 66 or more work hours per week. In addition, there was a statistically significant association between working 56 hours or more per week and the use of prescription medications with sleep aid properties. There was a significant association between the use of prescription sleep aids and all predisposing, enabling, and need factors. Participants aged between 27-64 years (90.2%,  $p<0.000$ ), White (75.1%,  $p<0.000$ ), female (57.5%,  $p<0.000$ ), married (53.9%,  $p<0.000$ ), with less than or equal to three family members (77.5%,  $p<0.000$ ), have some college or more (67.6%,  $p<0.000$ ), working in professional services (42.9%,  $p<0.000$ ), the southern region of the U.S. (36.5%,  $p<0.000$ ), middle to high income (81.1%,  $p<0.000$ ), did not receive food stamps (92.2%,  $p=0.001$ ), and reported having good health status (73.9%,  $p<0.000$ ) made up the highest proportions of those who used prescription sleep aid.

Furthermore, the use of medication with sedative side effects had similar significant proportions and associations with the covariates compared to the use of prescription sleep aids. However, the covariates, education ( $p=0.072$ ), and income ( $p=0.924$ ) did not show significant associations at 5% level with the use of medications with sedative side effects.

### **1.3.3. Correlates of the Use of Prescription Sleep Aids**

The bivariate analysis showed the strongest association between the 56 hours or more per week category, the use of sleep aids, and medications with sleep aid properties. Subsequently, we provided results only for the 56 hours or more per week category in the result section of our analysis. The results for other work hour categories are provided in *Appendices 2–5* of the paper. Overall, we found a similar pattern of positive associations between work hours and the use of medications across the various work hour categories (36 hours, 46 hours, and 66 hours) when compared to the 56 hours per week category. However, they were largely insignificant ( $p > 0.05$ ).

In the linear regression analysis in *Table 3*, the number of prescriptions was used as the outcome variable. We found that working 56 hours or more per week was associated with a 0.05 unit increase in the sleep aid prescription. The increase translates to an average of additional 14 prescription sleep aid pills within the survey round.

Due to the censored dependent variable data, we employed Tobit regression which resulted in a positive association between working 56 hours or more per week and using prescription sleep aids. We found the significant coefficients ( $p < 0.05$ ) from the

Tobit marginal effect similar and comparable with the linear regression coefficients. The Tobit marginal effects showed that individuals working 56 hours or more per week were associated with a 0.07 unit increase in the prescription of sleep aids, translating to an additional 20 prescription sleep aid pills within the survey round on average.

Nonetheless, the conditional moment test for normality in a Tobit model indicated strong non-normality of the error terms ( $p < 0.001$ ), challenging the validity of the Tobit results (Cameron & Trivedi, 2010; Drukker, 2002).

To test if the analysis can be conducted in two parts (*i.e.*, the selection into using sleep aids versus the extent of sleep aid use among users), we ran a Heckman regression (see *Appendix 2a*). The regression resulted in estimating a statistically insignificant selection hazard ratio, indicating the independence of the two parts (inverse Mills' ratio=8.99,  $p=0.701$ ). Therefore, we proceeded with analyzing the two parts independently. Results from the adjusted logistic regression (aOR) showed that the odds of using prescription sleep aids among respondents who worked 56 hours or more per week was 13% more than that among those who worked fewer hours (aOR=1.13, 95% Confidence Interval, CI=1.01:1.26). The logit marginal effect also yielded a significant ( $p<0.05$ ) positive association between working 56 hours or more per week and the likelihood of using prescription sleep aids. It informed that the predicted probability of using sleep aids is 0.007 (CI=0.000, 0.014) greater for the individual that worked 56 hours or more per week than for one that worked less than 56 hours per week.

Additionally, the linear regression analysis among users of prescription sleep aids showed that respondents who worked 56 hours or more per week had 0.37 more sedative prescription compared to those who worked less than 56 hours per week while

controlling for other covariates. This coefficient represents an average of an additional 110 prescription sleep aid pills for respondents in this category (individuals that worked 56 hours or more per week and used sleep aids).

Individuals aged 27-64 years were 2.2 times more likely to have used prescription sleep aids than respondents aged 18-26 years (aOR=2.20, 95% CI=1.90:2.56). Compared to males, females were 11% (aOR=1.11, 95% CI=1.05:1.19) more likely to use prescription sleep aids. While those who reported being divorced (aOR=1.13, 95% CI=1.05:1.21), had some college or more education (aOR=1.24, 95% CI=1.16:1.32), and residents of North central – Midwest (aOR=1.15, 95% CI=1.04:1.27) had a higher likelihood of using prescription sleep aids.

In addition, respondents that worked in professional services were 31% (aOR=1.31, 95% CI=1.14:1.50) more likely to have used prescription sleep aids than those working in natural resources. Those who reported poor health status were 2.6 times more likely to have used prescription sleep aids than persons with excellent health status (aOR=2.609, 95% CI=2.102:3.238). African Americans/Blacks (aOR=0.46, 95% CI=0.41:0.50), and those with 4-6 family members in the household (aOR=0.67, 95% CI=0.62:0.72) were less likely to report the use of prescription sleep aids.

#### **1.3.4. Correlates of the Use of Prescription Medications with Sedative Properties**

In the linear regression analysis in *Table 4*, we found that working 56 hours or more per week was associated with a 0.19 unit increase in medications with sedative properties prescription. The increase translates to an average of additional 38 pills with sedative properties within the survey round. Likewise, we found the Tobit marginal effects had significant positive coefficients ( $p < 0.001$ ) and were comparable to the linear

regression coefficients. The Tobit marginal effects informed that individuals working 56 hours or more per week were associated with a 0.11 unit increase in the prescription of medications with sedative properties, translating to an average of additional 22 pills with sedative properties within the survey round. Furthermore, the conditional moment test for normality in a Tobit model indicated strong non-normality of the error terms ( $p < 0.001$ ).

Correspondingly, to test if the analysis can be conducted in two parts (*i.e.*, the selection into using medications with sedative properties versus the extent of medications with sedative properties use among users), we ran a Heckman regression (see *Appendix 2b*). The regression produced a statistically insignificant selection hazard ratio, indicating the independence of the two parts (inverse Mills' ratio=33.08,  $p=0.434$ ).

Analyzing the two parts independently, the results from the adjusted logistic regression (aOR) informed that the odds of using medications with sedative properties among respondents who worked 56 hours or more per week was 9% more than that among those who worked fewer hours (aOR=1.09, 95% CI=1.03:1.16). Likewise, the logit marginal effect showed a significant ( $p<0.01$ ) positive association between working 56 hours or more per week and the likelihood of using medications with sedative properties. It informed that respondents who worked 56 hours or more per week had 2% (CI=0.7%, 3.3%) more likelihood to use medications with sedative properties.

Furthermore, the linear regression analysis among users of medications with sedative properties showed that respondents who worked 56 hours or more per week had 0.35 more prescription with sedative properties than respondents who worked less than 56 hours per week. This coefficient represents an additional 70 prescription pills of



medications with sedative properties for respondents in this category within the survey round.

Respondents who reported being divorced (aOR=1.06, 95% CI=1.02:1.10) and receiving food stamps (aOR=1.15, 95% CI=1.08:1.21) had a higher likelihood of using medications with sedative properties. While respondents with race classified as other (aOR=0.89, 95% CI=0.85:0.95) and those with 4-6 family members in the household (aOR=0.82, 95% CI=0.79:0.85) were less likely to report the use of prescription prescriptions with sedative properties.

Like prescription sleep aids use, respondents who reported poor health status were 2.6 times more likely to have used medications with sedative properties than persons with excellent health status (aOR=2.60, 95% CI=2.29:2.94). Also, respondents that worked in professional services were 14% (aOR=1.14, 95% CI=1.07:1.22) more likely to have used medications with sedative properties than those who worked in natural resources.

Unlike the aOR results from the sleep aids use, Individuals aged 65 years and older had the highest likelihood of using medications with sedative properties compared to those between 18-26 years (aOR=1.50, 95% CI=1.37:1.63). Also, covariates: sex, education, region, and income did not show a significant association ( $p > 0.05$ ) between working 56 hours or more per week and using medications with sedative properties.

### **1.3.5. Supplemental Analysis**

Additionally, following (Solon et al., 2015), we also conducted our analysis using sample weights and accounting for the complex survey design of the MEPS (*Appendices 6 to 10*). The analysis yielded similar results. Though the point estimates changed, the associations were similar to the unweighted results. The differences between the two sets

(unweighted vs. weighted) were consistent with the observed heterogeneity in the treatment effects and the reported oversampling for minority groups in the MEPS (Quality, 2020; Schaller & Zerpa, 2019).

#### **1.4. DISCUSSION**

We found a similar pattern of associations between long working hours, the use of prescription sleep aids, and medications with sedative properties. The overall prevalence of prescription sleep aids use (6.2%) was similar to the overall incidence rate (6.3%) of risky alcohol use found in the landmark paper by Virtanen et al. 2015 (Virtanen et al., 2015) – that explored the relationship between long working hours and risky alcohol use. Furthermore, previous national estimates reported the monthly prevalence of prescription sleep aid use at 4% among adults aged 20 years and older (Chong et al., 2013). While other studies estimate the prevalence of hypnotics among the general population to range between 3.5% and 11.7% (Chong et al., 2013; Ohayon et al., 1998; Vaidya et al., 2014).

In the analysis of the use of medications with sedative properties, we found a prevalence of 34.5% across respondents. Likewise, a recent report (Lopez-Quintero et al., 2021) informed that about 40% of individuals aged 12 and older used prescription pain medications in combination with sedatives and tranquilizers to aid sleep. Several studies link the use of opioids and sleep due to associated pain reduction and increased initiation of sleep–drowsiness (Angarita et al., 2016; Cheatle & Webster, 2015; Serdarevic et al., 2017).

The results from the Tobit models, with significant positive coefficients that were comparable to the coefficients from the linear probability model, informed a positive association between working 56 hours or more per week and the use of prescription sleep

aids. This adds to the robustness of our analysis (Cameron & Trivedi, 2010; Foster & Kalenkoski, 2013; Wang & Griswold, 2017). The Tobit marginal effects showed that individuals who worked 56 hours or more per week were associated with an increased likelihood of using sleep aids and medications with sedative properties by increased prescription units of 0.07 and 0.11, respectively. These increases translate to an average of additional 20 prescription sleep aid pills, and 22 pills with sedative properties, respectively within the survey round.

We found that the odds of using prescription sleep aids and medications with sedative properties among respondents who worked 56 hours or more per week was 13% and 9%, respectively more than that among those who worked fewer hours. Likewise, a previous meta-analysis of 63 studies with over 333,000 participants across 14 countries found that working 55 hours or more per week was associated with a 12% increased odds of new-onset of risky alcohol use (Virtanen et al., 2015). This increased odds of the association between long working hours and the use of sedative medications could result from the impact of long working hours on sleep.

Also, we found that among the users of sleep aids and medications with sedative properties, those who worked 56 hours or more per week were associated with higher uses (frequency) of these medications compared to those that worked lesser hours. Previous studies (Artazcoz et al., 2009; Nakashima et al., 2011; Virtanen et al., 2009) have shown significant positive relationships between working long hours and the risk of sleep disorders. Also, national estimates reported that 6% to 10% of individuals in the U.S. had been diagnosed with a sleep disorder (Laposky et al., 2016; Liu et al., 2013). Evidence informs that the use of prescription sleep aids are higher among individuals

with sleep disorders (Chong et al., 2013). Long working hours increase the likelihood of stress and are accompanied by an increased prevalence of sleep disorders (Kivimäki et al., 2012; Virtanen et al., 2009; Virtanen et al., 2015). This increased prevalence of sleep disorders likely results in the subsequent use of medications to aid sleep.

Furthermore, we found that females in our study had a higher likelihood of using sleep aids when compared to males. Similarly, several previous findings report that females have a higher chance of using sleep aids than males (Chong et al., 2013; Lopez-Quintero et al., 2021; Vaidya et al., 2014). A rationale theorized for this is females have a higher risk of developing sleep disorders when exposed to long working hours compared to males (Wong et al., 2019). Additionally, employed females have been noted to have increased exposure to unpaid work such as household chores, which result in higher work hours and reduced sleep duration compared to employed males (Bianchi, 2000; Sayer, 2005). Also, insomnia has been reportedly higher in middle-aged women and was linked to peculiar hormonal and psychological changes in this age group (Cirignotta et al., 1985).

Our study found significant differences in the associations between long working hours and the use of sleep medications across socioeconomic groups. Professionals – individuals involved in education, such as teachers and health care workers, had the highest likelihood of using prescription sleep aids and medications with sedative properties. Earlier literature reports that professionals have an increased likelihood of working long hours as it could result in possible financial and career rewards for them (Romani & Ashkar, 2014; Wisetborisut et al., 2014). On the other hand, long working hours could be a response to dire economic needs and financial strain in a continuous

attempt to make ends meet for individuals of lower socioeconomic status. In our study, respondents who reported receiving food stamps – an indication of financial difficulty, had a 15% higher likelihood of using medications with sedative properties than those who did not receive food stamps (aOR=1.15, 95% CI=1.08:1.21).

However, respondents with 4-6 family members (additional indication of financial motivation) in the household had a lower likelihood of using prescription sleep aids and medications with sedative properties. The association between long working hours, use of sleep medications, and financial need might be additionally influenced by the health status of the individuals.

In our study, individuals who reported poor health status had a significantly increased likelihood of using prescription sleep medications and medications with sedative properties (2.7 and 2.6 times, respectively) compared to those who reported excellent health status. Long working hours have been linked to poor health outcomes. Prior literature found that individuals with poor health status are more likely to use prescription pain medications and sleep medications (Lopez-Quintero et al., 2021).

#### **1.4.1. Policy and Theoretical Implications**

Our study provides evidence of the association between long working hours and the use of medications with sedative properties. The use of sleep medications has several implications. Previous studies have linked sleep aids medications with increased mortality (Kivimäki et al., 2015; Skogstad et al., 2019). However, other studies found no significant associations (Jaussent et al., 2013; Phillips & Mannino, 2005; Rumble & Morgan, 1992). The chronic use of these medications increases the risk of cognitive and psychomotor impairments, car and workplace accidents, and addiction (Kim et al., 2016;

Mizoue et al., 2001). Thus, the residual effects of sleep medications and their resultant effects present safety risks to the workers that use them and the public.

The similar patterns of utilization between prescription sleep aids and medications with sedative properties in our study support earlier findings that many persons use drugs with sedative properties for their off label use to aid sleep (Lopez-Quintero et al., 2021; Neubauer et al., 2021). This further increases the risks associated with the use of these medications, such as poly-drug users who believes that these sleep medications could significantly increase the intoxicating experience of other recreational drugs, such as narcotics (Votaw et al., 2019).

Furthermore, we found that working women had an increased risk of associations between using these medications and working long hours. This adds evidence to previous theories, which inform of additional household chores that female workers undertake and account for increased stress that results from extended working hours (Wong et al., 2019).

Government and private organizations should establish clearly defined standard work hours to protect and maintain the health of their citizens and employees. Also, employees should be enlightened on the implications of long working hours on their physical and mental health. There should be increased education on the associated risks of using sleep medications. Thus, behavioral and cognitive interventions that have been shown to improve and sustain sleep quality and hygiene with non-significant adverse effects (Martínez et al., 2014) should be promoted as a choice and sustainable alternative for workers.

#### **1.4.2. Limitations**

This study had several limitations. The study was largely based on self-reports and could be prone to misclassification bias. For example, self-report estimates of alcohol are known to under-report drinking (Gmel & Rehm, 2004). However, this does not automatically bias estimates of associations. In addition, observations with missing covariates were excluded, which may lead to residual confounding by unmeasured covariates and selection bias. Additionally, with survey data, there is an increased likelihood of recall and response bias. Despite this, we compared changes in the result with adjustments for sociodemographic factors and utilized a Heckman analysis to ensure non-sample bias.

We could not use the panel structure of the data due to the lack of observations and changes across the small sample. Therefore, we mainly inferred associations in the relationship between the variables rather than establishing causal inference. Additionally, MEPS does not collect information on over-the-counter medications (OTC). Thus, many non-prescription sleep aids which could be obtained OTC were likely missed. Notwithstanding, the confirmed prescriptions used in the study were not subjected to recall and social desirability bias. As such, it increased the specificity and reliability of the measurement of sleep aid use.

Lastly, we could not test all the components of our conceptual models due to the limitations of the data. Future studies could explore other patterns in the relationship between the use of sleep aids and other constructs from our conceptual model.

## **1.5. CONCLUSION**

The continuous attempts at meeting the needs of round-the-clock service and globalized society have increased long working hours. An increase in sleep disorders has

accompanied this increase. Also, sleep and substance abuse disorders have been closely linked. Our study found that long working hours were significantly associated with an increased use of sleep aids and medications with sedative properties. Working 56 hours or more per week was significantly associated with an increased odds of using sleep aids and medications with sedative properties by 13% and 9%, respectively more than that among those who worked fewer hours. Working females in our study had a higher likelihood of using sleep aids when compared to males. At the same time, people working in professional services had the highest likelihood of using sleep medications. Also, we found an increased frequency of sleep medication use among users exposed to long working hours.

Our findings can guide employers, health systems, and researchers in designing interventions and recommendations for creating work-based policies focused on improving the health status of employees. Government and private organizations should establish clearly defined standard work hours to protect and maintain the health of workers. Furthermore, there is a need for employee education on the potential health implications of long working hours and using medications for sleep. Along with the continuous expansion or promotion of behavioral and cognitive sleep interventions.



Figure 1. Conceptual framework depicting the influence of working hours on the use of sleep aid drugs (adapted from Aday & Andersen, 1974).




Table 1. Descriptive statistics of work hours and use of prescription medications

| Variables  | Overall    | Cutoffs of Weekly Work Hours |            |            |            |            |            |            |            |
|--|------------|------------------------------|------------|------------|------------|------------|------------|------------|------------|
|  |            | 36                           |            | 46         |            | 56         |            | 66         |            |
|  |            | < 36                         | ≥ 36       | < 46       | ≥ 46       | < 56       | ≥ 56       | < 66       | ≥ 66       |
| <b>Number of observations</b>                        | 81,518     | 10,696                       | 70,822     | 65,920     | 15,598     | 76,193     | 5,325      | 79,943     | 1,575      |
| <b>% Sleep aids use</b>                              | 6.2        | 5.8                          | 6.3        | 6.1        | 6.5        | 6.1        | 7          | 6.2        | 6.6        |
| Mean sleep aids among users (SD)                     | 2.83(2.29) | 2.81(2.12)                   | 2.84(2.30) | 2.84(2.29) | 2.81(2.28) | 2.81(2.26) | 3.16(2.58) | 2.83(2.27) | 3.06(2.79) |
| Median (IQR)   | 2(3)       | 2(3)                         | 2(3)       | 2(3)       | 2(3)       | 2(3)       | 2(4)       | 2(3)       | 2(3)       |
| <b>% Use of medications with sedative properties</b> | 34.5       | 33.5                         | 34.6       | 34.5       | 34.5       | 34.4       | 36.1       | 34.4       | 36.4       |
| Mean sedative properties meds among users (SD)       | 2.82(3.46) | 2.87(3.17)                   | 2.82(3.51) | 2.82(3.52) | 2.83(3.20) | 2.80(3.46) | 3.18(3.54) | 2.82(3.47) | 3.13(3.36) |
| Median (IQR)   | 2(2)       | 2(3)                         | 2(2)       | 2(2)       | 2(2)       | 2(2)       | 2(3)       | 2(2)       | 2(3)       |
| <b>Predisposing Factors</b>                          |            |                              |            |            |            |            |            |            |            |
| <b>Age (years)</b>                                   |            |                              |            |            |            |            |            |            |            |
| 18-26  | 9          | 17.1                         | 8          | 10.1       | 5.3        | 9.4        | 5          | 9.1        | 5.2        |
| 27-64  | 85.3       | 72.7                         | 87         | 84.3       | 89.2       | 85.1       | 88         | 85.3       | 86.6       |
| ≥ 65   | 5.6        | 10.2                         | 5          | 5.7        | 5.5        | 5.5        | 7          | 5.6        | 8.2        |
| <b>Race</b>  |            |                              |            |            |            |            |            |            |            |
| White  | 83.6       | 82.2                         | 83.7       | 82.7       | 86.6       | 83.4       | 85.1       | 83.5       | 83.9       |
| Black  | 10.1       | 11.2                         | 9.9        | 10.7       | 8          | 10.1       | 9.3        | 10.1       | 10.7       |
| Other  | 6.4        | 6.6                          | 6.3        | 6.6        | 5.4        | 6.4        | 5.6        | 6.4        | 5.4        |
| <b>Sex</b>   |            |                              |            |            |            |            |            |            |            |
| Male   | 44.5       | 31.4                         | 46.2       | 40.4       | 59.2       | 42.7       | 66.9       | 43.8       | 74.1       |
| Female   | 55.5       | 68.6                         | 53.8       | 59.6       | 40.8       | 57.3       | 33.1       | 56.2       | 25.9       |
| <b>Marital status</b>                                |            |                              |            |            |            |            |            |            |            |
| Married  | 57.4       | 47.3                         | 58.7       | 55.9       | 62.6       | 57.1       | 61.4       | 57.3       | 60.9       |
| Divorced   | 20         | 21.7                         | 19.8       | 20.2       | 19.3       | 19.9       | 20.7       | 19.9       | 23.1       |
| Unmarried  | 22.6       | 31                           | 21.5       | 23.9       | 18.2       | 23         | 17.8       | 22.8       | 15.9       |
| <b>Family size</b>                                   |            |                              |            |            |            |            |            |            |            |
| ≤ 3 Members  | 69.3       | 67.9                         | 69.5       | 69         | 70.5       | 69.1       | 71         | 69.2       | 72.6       |
| 4-6 Members  | 29.2       | 30.3                         | 29         | 29.4       | 28.4       | 29.3       | 27.8       | 29.3       | 25.7       |

|                              |      |      |      |      |      |      |      |      |      |
|------------------------------|------|------|------|------|------|------|------|------|------|
| > 6 Members                  | 1.6  | 1.8  | 1.5  | 1.7  | 1.1  | 1.6  | 1.2  | 1.6  | 1.7  |
| <b>Education</b>             |      |      |      |      |      |      |      |      |      |
| Less than college            | 31.7 | 38.2 | 30.9 | 32.9 | 27.7 | 31.8 | 30.7 | 31.6 | 40.3 |
| Some college or more         | 68.3 | 61.8 | 69.1 | 67.1 | 72.4 | 68.2 | 69.3 | 68.5 | 59.7 |
| <b>Occupation</b>            |      |      |      |      |      |      |      |      |      |
| Natural resources            | 6.9  | 4.9  | 7.2  | 6    | 10.2 | 6.4  | 13.4 | 6.8  | 14.3 |
| Hospitality services         | 5.9  | 14.7 | 4.7  | 6.2  | 4.5  | 6    | 4.6  | 5.9  | 4.5  |
| Trade                        | 11.3 | 15.9 | 10.7 | 11.4 | 10.7 | 11.3 | 10.9 | 11.3 | 11.7 |
| Professional services        | 39.5 | 42.8 | 39   | 40.2 | 36.7 | 39.8 | 35.4 | 39.6 | 34.3 |
| Manufacturing                | 11   | 3.1  | 12   | 10.5 | 12.8 | 11   | 10   | 11   | 8.1  |
| Other                        | 25.5 | 18.6 | 26.4 | 25.6 | 25.1 | 25.5 | 25.8 | 25.5 | 27   |
| <b>Year</b>                  |      |      |      |      |      |      |      |      |      |
| 2010                         | 9.6  | 9    | 9.7  | 9.6  | 9.5  | 9.6  | 9.6  | 9.6  | 10.3 |
| 2011                         | 10.4 | 10.5 | 10.3 | 10.4 | 10.3 | 10.2 | 11.8 | 10.2 | 16   |
| 2012                         | 10.4 | 10.3 | 10.4 | 10.4 | 10.4 | 10.3 | 10.9 | 10.3 | 13.5 |
| 2013                         | 10   | 9.8  | 10   | 10.1 | 9.4  | 10.1 | 9    | 10   | 8.3  |
| 2014                         | 11   | 11.5 | 11   | 10.9 | 11.3 | 11   | 11.7 | 11   | 11.2 |
| 2015                         | 10.6 | 10.5 | 10.6 | 10.4 | 11.2 | 10.6 | 10.9 | 10.6 | 9.6  |
| 2016                         | 10.2 | 9.2  | 10.4 | 10.2 | 10.2 | 10.2 | 9.8  | 10.3 | 8.4  |
| 2017                         | 9.8  | 10   | 9.8  | 9.8  | 9.8  | 9.9  | 9.1  | 9.9  | 8    |
| 2018                         | 9.6  | 9.7  | 9.6  | 9.5  | 9.8  | 9.6  | 9    | 9.6  | 8.5  |
| 2019                         | 8.5  | 9.5  | 8.4  | 8.6  | 8.1  | 8.5  | 8.2  | 8.6  | 6.2  |
| <b>Enabling Factors</b>      |      |      |      |      |      |      |      |      |      |
| <b>Region</b>                |      |      |      |      |      |      |      |      |      |
| Northeast                    | 15.2 | 20.1 | 14.6 | 15.3 | 14.9 | 15.4 | 13   | 15.3 | 11.2 |
| North Central,<br>Midwest    | 24.2 | 23.6 | 24.3 | 23.7 | 26   | 23.9 | 27.4 | 24.2 | 24.6 |
| South                        | 39.3 | 36.2 | 39.8 | 39.4 | 39.2 | 39.4 | 38.7 | 39.3 | 43   |
| West                         | 21.3 | 20.2 | 21.4 | 21.7 | 19.9 | 21.3 | 20.9 | 21.3 | 21.3 |
| <b>Income</b>                |      |      |      |      |      |      |      |      |      |
| Poor to low income           | 15.6 | 32   | 13.4 | 17.7 | 8    | 16.1 | 9.2  | 15.7 | 10.7 |
| Middle to high income        | 84.4 | 68   | 86.6 | 82.3 | 92   | 83.9 | 90.8 | 84.3 | 89.3 |
| <b>Receipt of food stamp</b> |      |      |      |      |      |      |      |      |      |
| No                           | 94.2 | 87.1 | 95.1 | 93.5 | 96.7 | 94.1 | 95.5 | 94.2 | 94.6 |
| Yes                          | 5.8  | 12.9 | 4.9  | 6.5  | 3.3  | 5.9  | 4.5  | 5.8  | 5.4  |
| <b>Need Factors</b>          |      |      |      |      |      |      |      |      |      |

**Health Status**

|           |      |      |      |      |      |      |      |      |      |
|-----------|------|------|------|------|------|------|------|------|------|
| Excellent | 14.9 | 14.2 | 15   | 14.4 | 16.8 | 14.8 | 16.9 | 14.9 | 16.5 |
| Good      | 74.3 | 73.1 | 74.5 | 74.5 | 73.8 | 74.5 | 72.2 | 74.5 | 68.8 |
| Fair      | 9.6  | 11.1 | 9.4  | 9.9  | 8.3  | 9.6  | 9.6  | 9.5  | 13.4 |
| Poor      | 1.2  | 1.6  | 1.1  | 1.2  | 1.1  | 1.2  | 1.2  | 1.2  | 1.3  |

---

Note: SD=standard deviation; IQR=interquartile range; REF=reference

Table 2. Bivariate Analysis

| Variables                           | Use of Prescription Sleep Aid |      |       |       | Use of Prescription Medications with Sedative Properties |      |       |       |
|-------------------------------------|-------------------------------|------|-------|-------|--|------|-------|-------|
|                                     | Yes                           |      | X     | P     | Yes  |      | X     | P     |
|                                     | Frequency                     | %    |       |       | Frequency  | %    |       |       |
| <b>Part 1: Work Hour Categories</b> |                               |      |       |       |  |      |       |       |
| <b>≥ 36 hours/week</b>              |                               |      |       |       |  |      |       |       |
| 30-35                               | 621                           | 12.3 | 3.3   | 0.071 | 3,611  | 13.0 | 0.9   | 0.337 |
| ≥ 36                                | 4,432                         | 87.7 |       |       | 24,244   | 87.0 |       |       |
| <b>≥ 46 hours/week</b>              |                               |      |       |       |  |      |       |       |
| 31-45                               | 4,041                         | 80.0 | 2.8   | 0.096 | 22,470   | 80.7 | 1.1   | 0.301 |
| ≥ 46                                | 1,012                         | 20.0 |       |       | 5,385  | 19.3 |       |       |
| <b>≥ 56 hours/week</b>              |                               |      |       |       |  |      |       |       |
| 31-55                               | 4,680                         | 92.6 | 6.4   | 0.012 | 25,925   | 93.1 | 10.9  | 0.001 |
| ≥ 56                                | 373                           | 7.4  |       |       | 1,930  | 6.9  |       |       |
| <b>≥ 66 hours/week</b>              |                               |      |       |       |  |      |       |       |
| 31-65                               | 4,949                         | 97.9 | 0.5   | 0.501 | 27,297   | 98.0 | 1.1   | 0.288 |
| ≥ 66                                | 104                           | 2.1  |       |       | 558  | 2.0  |       |       |
| <b>Part 2: Predisposing Factors</b> |                               |      |       |       |  |      |       |       |
| <b>Age (years)</b>                  |                               |      |       |       |  |      |       |       |
| 18-26                               | 201                           | 4.0  | 151.2 | 0.000 | 1,927  | 6.9  | 217.2 | 0.000 |
| 27-64                               | 4,556                         | 90.2 |       |       | 24,185   | 86.8 |       |       |
| ≥ 65                                | 296                           | 5.9  |       |       | 1,743  | 6.3  |       |       |
| <b>Race</b>                         |                               |      |       |       |  |      |       |       |
| White                               | 4,262                         | 75.1 | 259.2 | 0.000 | 21,062   | 75.6 | 24.7  | 0.000 |
| Black                               | 482                           | 17.0 |       |       | 4,771  | 17.1 |       |       |
| Other                               | 309                           | 7.9  |       |       | 2,022  | 7.3  |       |       |
| <b>Sex</b>                          |                               |      |       |       |  |      |       |       |
| Male                                | 1,999                         | 42.5 | 19.4  | 0.000 | 11,667   | 41.9 | 7.2   | 0.007 |
| Female                              | 3,054                         | 57.5 |       |       | 16,188   | 58.1 |       |       |
| <b>Marital Status</b>               |                               |      |       |       |  |      |       |       |
| Married                             | 2,722                         | 53.9 | 67.0  | 0.000 | 15,105   | 54.2 | 143.4 | 0.000 |
| Divorced                            | 1,287                         | 25.5 |       |       | 6,553  | 23.5 |       |       |
| Unmarried                           | 1,044                         | 20.7 |       |       | 6,197  | 22.3 |       |       |
| <b>Family Size</b>                  |                               |      |       |       |  |      |       |       |
| 0 - 3 members                       | 3,914                         | 77.5 | 177.3 | 0.000 | 20,138   | 72.3 | 190.1 | 0.000 |
| 4 - 6 members                       | 1,104                         | 21.9 |       |       | 7,371  | 26.5 |       |       |
| > 6 members                         | 35                            | 0.7  |       |       | 346  | 1.2  |       |       |
| <b>Education Level</b>              |                               |      |       |       |  |      |       |       |
| Less than College                   | 1,636                         | 32.4 | 68.1  | 0.000 | 10,656   | 38.3 | 3.2   | 0.072 |
| Some College or more                | 3,417                         | 67.6 |       |       | 17,199   | 61.7 |       |       |

(continued)

Table 2. (Continued)

| Variables                               | Use of Prescription Sleep Aid |      |       |       | Use of Prescription Medications with Sedative Properties |      |          |       |
|---|-------------------------------|------|-------|-------|--|------|----------|-------|
|   | Yes                           |      | X     | P     | Yes  |      | X        | P     |
|   | Frequency                     | %    |       |       | Frequency  | %    |          |       |
| <b>Occupation</b>                       |                               |      |       |       |  |      |          |       |
| Natural resources                       | 278                           | 5.5  | 55.9  | 0.000 | 1,761  | 6.3  | 21.0     | 0.001 |
| Hospitality services                    | 268                           | 5.3  |       |       | 1,822  | 6.5  |          |       |
| Trade                                   | 520                           | 10.3 |       |       | 3,167  | 11.4 |          |       |
| Professional services                   | 2,166                         | 42.9 |       |       | 11,050   | 39.7 |          |       |
| Manufacturing                           | 564                           | 11.2 |       |       | 3,120  | 11.2 |          |       |
| Other                                   | 1,257                         | 24.9 |       |       | 6,935  | 24.9 |          |       |
| <b>Year</b>                             |                               |      |       |       |  |      |          |       |
| 2010                                    | 474                           | 9.4  | 62.1  | 0.000 | 2,768  | 9.9  | 103.2    | 0.000 |
| 2011                                    | 526                           | 10.4 |       |       | 2,892  | 10.4 |          |       |
| 2012                                    |                               | 10.0 |       |       | 2,918  | 10.5 |          |       |
| 2013                                    | 491                           | 9.7  |       |       | 2,867  | 10.3 |          |       |
| 2014                                    | 531                           | 10.5 |       |       | 2,952  | 10.6 |          |       |
| 2015                                    | 506                           | 10.0 |       |       | 3,046  | 10.9 |          |       |
| 2016                                    | 485                           | 9.6  |       |       | 2,811  | 10.1 |          |       |
| 2017                                    | 501                           | 9.9  |       |       | 2,599  | 9.3  |          |       |
| 2018                                    | 513                           | 10.2 |       |       | 2,596  | 9.3  |          |       |
| 2019                                    | 521                           | 10.3 |       |       | 2,406  | 8.6  |          |       |
| <b>Part 3: Enabling Factors</b>         |                               |      |       |       |  |      |          |       |
| <b>Region</b>                           |                               |      |       |       |  |      |          |       |
| Northeast                               | 673                           | 13.3 | 20.7  | 0.000 | 3,790  | 13.6 | 16.9     | 0.001 |
| North Central/Midwest                   | 1,309                         | 25.9 |       |       | 6,637  | 23.8 |          |       |
| South                                   | 1,846                         | 36.5 |       |       | 10,841   | 38.9 |          |       |
| West                                    | 1,225                         | 24.2 |       |       | 6,587  | 23.7 |          |       |
| <b>Income</b>                           |                               |      |       |       |  |      |          |       |
| Poor to low income                      | 957                           | 18.9 | 37.8  | 0.000 | 6,243  | 22.4 | 0.0      | 0.924 |
| Middle to high income                   | 4,096                         | 81.1 |       |       | 21,612   | 77.6 |          |       |
| <b>Receipt of food stamp (No - REF)</b> |                               |      |       |       |  |      |          |       |
| No                                      | 4,661                         | 92.2 | 10.1  | 0.001 | 25,214   | 90.5 | 12.0041  | 0.001 |
| Yes                                     | 392                           | 7.8  |       |       | 2,641  | 9.5  |          |       |
| <b>Part 4: Need Factors</b>             |                               |      |       |       |  |      |          |       |
| <b>Health Status</b>                    |                               |      |       |       |  |      |          |       |
| Excellent                               | 492                           | 9.7  | 142.2 | 0.000 | 3,106  | 11.2 | 700.7854 | 0.000 |
| Good                                    | 3,733                         | 73.9 |       |       | 20,235   | 72.6 |          |       |
| Fair                                    | 713                           | 14.1 |       |       | 3,949  | 14.2 |          |       |
| Poor                                    | 115                           | 2.3  |       |       | 565  | 2.0  |          |       |

Note: N=Frequency, %=Percentage; X=Chi-Square; P=p-value; REF=reference

Table 3. Regressions for the Use of Prescription Sleep Aids.

| Variables                                   | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Prescription Sleep Aid=YES) |
|---|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                      |                               |                               |                            |                               |   |
| ≥ 56 h/week                                 | 0.046***<br>(0.021, 0.070)    | 0.065*<br>(0.015, 0.114)      | 1.126*<br>(1.008, 1.258)   | 0.007*<br>(0.000, 0.014)      | 0.370**<br>(0.128, 0.611)   |
| <b>Age (18-26years - REF)</b>               |                               |                               |                            |                               |   |
| 27-64 years                                 | 0.104***<br>(0.080, 0.127)    | 0.305***<br>(0.254, 0.355)    | 2.203***<br>(1.895, 2.561) | 0.034***<br>(0.030, 0.039)    | 0.513**<br>(0.180, 0.847)   |
| ≥ 65 years                                  | 0.068***<br>(0.032, 0.103)    | 0.251***<br>(0.179, 0.322)    | 1.988***<br>(1.639, 2.412) | 0.028***<br>(0.020, 0.036)    | 0.256<br>(-0.168, 0.680)  |
| <b>Race (White - REF)</b>                   |                               |                               |                            |                               |   |
| Black                                       | -0.127***<br>(-0.144, -0.109) | -0.313***<br>(-0.349, -0.277) | 0.456***<br>(0.413, 0.504) | -0.037***<br>(-0.040, -0.033) | -0.403***<br>(-0.624, -0.182)                                     |
| Other                                       | -0.086***<br>(-0.109, -0.063) | -0.171***<br>(-0.218, -0.124) | 0.687***<br>(0.610, 0.775) | -0.021***<br>(-0.026, -0.015) | -0.557***<br>(-0.822, -0.292)                                     |
| <b>Sex (Male - REF)</b>                     |                               |                               |                            |                               |   |
| Female                                      | 0.014*<br>(0.001, 0.027)      | 0.046***<br>(0.020, 0.073)    | 1.114***<br>(1.047, 1.186) | 0.006***<br>(0.003, 0.010)    | -0.022<br>(-0.158, 0.113)   |
| <b>Marital status (Married - REF)</b>       |                               |                               |                            |                               |   |
| Divorced                                    | 0.028***<br>(0.012, 0.045)    | 0.053**<br>(0.021, 0.085)     | 1.126**<br>(1.046, 1.211)  | 0.007**<br>(0.003, 0.011)     | 0.064<br>(-0.094, 0.222)  |
| Unmarried                                   | 0.002<br>(-0.015, 0.019)      | 0.009<br>(-0.025, 0.043)      | 1.035<br>(0.955, 1.122)    | 0.002<br>(-0.003, 0.007)      | -0.098<br>(-0.274, 0.079)   |
| <b>Family size (≤ 3 Members REF)</b>        |                               |                               |                            |                               |   |
| 4-6 Members                                 | -0.072***<br>(-0.086, -0.058) | -0.171***<br>(-0.200, -0.141) | 0.670***<br>(0.623, 0.721) | -0.021***<br>(-0.025, -0.018) | -0.233**<br>(-0.393, -0.073)                                      |
| > 6 Members                                 | -0.101***<br>(-0.151, -0.051) | -0.307***<br>(-0.418, -0.197) | 0.437***<br>(0.311, 0.615) | -0.037***<br>(-0.048, -0.027) | 0.071<br>(-0.686, 0.827)  |
| <b>Education (Less than college - REF)</b>  |                               |                               |                            |                               |   |
| Some college or more                        | 0.026***<br>(0.012, 0.039)    | 0.086***<br>(0.059, 0.114)    | 1.235***<br>(1.157, 1.319) | 0.012***<br>(0.008, 0.015)    | -0.078<br>(-0.221, 0.065)   |
| <b>Occupation (Natural resources - REF)</b> |                               |                               |                            |                               |   |
| Hospitality services                        | 0.022<br>(-0.012, 0.056)      | 0.02<br>(-0.049, 0.090)       | 1.037<br>(0.869, 1.237)    | 0.002<br>(-0.007, 0.010)      | 0.234<br>(-0.152, 0.619)  |
| Trade                                       | 0.035*<br>(0.005, 0.065)      | 0.06<br>(-0.001, 0.121)       | 1.12<br>(0.961, 1.305)     | 0.006<br>(-0.002, 0.013)      | 0.293<br>(-0.040, 0.626)  |
| Professional services                       | 0.063***                      | 0.123***                      | 1.308***                   | 0.015***                      | 0.338*  |

|  |                 |                 |                |                  |                  |
|--|-----------------|-----------------|----------------|------------------|------------------|
|  | (0.036, 0.089)  | (0.069, 0.177)  | (1.142, 1.497) | (0.008, 0.021)   | (0.043, 0.632)   |
| Manufacturing                            | 0.049**         | 0.094**         | 1.230**        | 0.011**          | 0.294            |
|  | (0.019, 0.079)  | (0.034, 0.155)  | (1.059, 1.428) | (0.003, 0.019)   | (-0.033, 0.621)  |
| Other                                    | 0.041**         | 0.094***        | 1.238**        | 0.011**          | 0.16             |
|  | (0.014, 0.069)  | (0.039, 0.149)  | (1.079, 1.420) | (0.004, 0.018)   | (-0.140, 0.460)  |
| <b>Year (2010 - REF)</b>                 |                 |                 |                |                  |                  |
| 2011                                     | 0.011           | -0.008          | 0.964          | -0.002           | 0.276            |
|  | (-0.017, 0.038) | (-0.064, 0.047) | (0.847, 1.096) | (-0.010, 0.005)  | (-0.004, 0.556)  |
| 2012                                     | -0.004          | -0.051          | 0.856*         | -0.009*          | 0.363*           |
|  | (-0.031, 0.023) | (-0.106, 0.004) | (0.752, 0.975) | (-0.016, -0.001) | (0.080, 0.645)   |
| 2013                                     | 0.009           | -0.02           | 0.922          | -0.005           | 0.372*           |
|  | (-0.019, 0.037) | (-0.076, 0.036) | (0.809, 1.052) | (-0.012, 0.003)  | (0.087, 0.657)   |
| 2014                                     | -0.006          | -0.02           | 0.953          | -0.003           | 0.004            |
|  | (-0.033, 0.022) | (-0.075, 0.036) | (0.837, 1.084) | (-0.010, 0.005)  | (-0.276, 0.284)  |
| 2015                                     | -0.019          | -0.049          | 0.885          | -0.007           | 0.015            |
|  | (-0.047, 0.008) | (-0.104, 0.006) | (0.777, 1.008) | (-0.014, 0.000)  | (-0.268, 0.298)  |
| 2016                                     | -0.011          | -0.04           | 0.896          | -0.006           | 0.119            |
|  | (-0.039, 0.016) | (-0.096, 0.016) | (0.786, 1.023) | (-0.014, 0.001)  | (-0.167, 0.405)  |
| 2017                                     | 0.022           | 0.011           | 0.986          | -0.001           | 0.401**          |
|  | (-0.006, 0.050) | (-0.045, 0.068) | (0.865, 1.123) | (-0.009, 0.007)  | (0.118, 0.685)   |
| 2018                                     | 0.018           | 0.013           | 0.996          | 0                | 0.297*           |
|  | (-0.011, 0.046) | (-0.044, 0.070) | (0.874, 1.135) | (-0.008, 0.008)  | (0.014, 0.581)   |
| 2019                                     | 0.033*          | 0.083**         | 1.191**        | 0.011**          | 0.053            |
|  | (0.003, 0.062)  | (0.024, 0.141)  | (1.046, 1.357) | (0.003, 0.020)   | (-0.228, 0.335)  |
| <b>Region (Northeast - REF)</b>          |                 |                 |                |                  |                  |
| North central, Midwest                   | 0.028**         | 0.059**         | 1.150**        | 0.008**          | 0.065            |
|  | (0.007, 0.049)  | (0.017, 0.100)  | (1.043, 1.267) | (0.003, 0.013)   | (-0.146, 0.275)  |
| South                                    | 0.015           | 0.034           | 1.09           | 0.005            | 0.032            |
|  | (-0.004, 0.035) | (-0.005, 0.073) | (0.994, 1.196) | (-0.000, 0.010)  | (-0.169, 0.233)  |
| West                                     | -0.01           | 0.004           | 1.036          | 0.002            | -0.223*          |
|  | (-0.031, 0.011) | (-0.037, 0.045) | (0.940, 1.143) | (-0.003, 0.007)  | (-0.437, -0.008) |
| <b>Income (Poor to low income - REF)</b> |                 |                 |                |                  |                  |
| Middle to high income                    | 0.001           | 0.014           | 1.041          | 0.002            | -0.095           |
|  | (-0.016, 0.018) | (-0.021, 0.049) | (0.957, 1.132) | (-0.002, 0.007)  | (-0.282, 0.092)  |
| <b>Receipt of food stamp (No - REF)</b>  |                 |                 |                |                  |                  |
| YES                                      | 0.028*          | 0.046           | 1.092          | 0.005            | 0.231            |
|  | (0.005, 0.052)  | (-0.005, 0.097) | (0.970, 1.228) | (-0.002, 0.012)  | (-0.032, 0.494)  |
| <b>Health status (Excellent - REF)</b>   |                 |                 |                |                  |                  |
| Good                                     | 0.058***        | 0.155***        | 1.469***       | 0.019***         | 0.219*           |
|  | (0.040, 0.076)  | (0.119, 0.192)  | (1.333, 1.618) | (0.015, 0.023)   | (0.005, 0.432)   |
| Fair                                     | 0.157***        | 0.308***        | 1.950***       | 0.037***         | 0.864***         |
|  | (0.132, 0.181)  | (0.257, 0.358)  | (1.729, 2.200) | (0.030, 0.044)   | (0.601, 1.128)   |
| Poor                                     | 0.258***        | 0.469***        | 2.609***       | 0.061***         | 1.082***         |
|  | (0.203, 0.312)  | (0.359, 0.580)  | (2.102, 3.238) | (0.043, 0.080)   | (0.620, 1.545)   |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF=reference



Table 4. Regressions for the Use of Prescription Medications with Sedative Properties

| Variables                                   | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Sedative Properties Medications=Y ES) |
|---|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                      |                               |                               |                            |                               |   |
| ≥ 56 h/week                                 | 0.186***<br>(0.119, 0.253)    | 0.114***<br>(0.065, 0.163)    | 1.094**<br>(1.032, 1.160)  | 0.020**<br>(0.007, 0.033)     | 0.347***<br>(0.187, 0.506)  |
| <b>Age (18-26years - REF)</b>               |                               |                               |                            |                               |   |
| 27-64 years                                 | 0.292***<br>(0.228, 0.356)    | 0.235***<br>(0.191, 0.278)    | 1.291***<br>(1.216, 1.370) | 0.054***<br>(0.042, 0.067)    | 0.515***<br>(0.344, 0.686)  |
| ≥ 65 years                                  | 0.376***<br>(0.279, 0.473)    | 0.334***<br>(0.266, 0.401)    | 1.495***<br>(1.372, 1.629) | 0.088***<br>(0.069, 0.107)    | 0.481***<br>(0.244, 0.718)  |
| <b>Race (White - REF)</b>                   |                               |                               |                            |                               |   |
| Black                                       | -0.201***<br>(-0.248, -0.154) | -0.089***<br>(-0.121, -0.056) | 0.974<br>(0.934, 1.015)    | -0.006<br>(-0.015, 0.003)     | -0.502***<br>(-0.616, -0.389)   |
| Other                                       | -0.214***<br>(-0.277, -0.152) | -0.137***<br>(-0.180, -0.095) | 0.893***<br>(0.845, 0.945) | -0.025***<br>(-0.037, -0.013) | -0.452***<br>(-0.609, -0.295)   |
| <b>Sex (Male - REF)</b>                     |                               |                               |                            |                               |   |
| Female                                      | 0.001<br>(-0.034, 0.037)      | 0.008<br>(-0.017, 0.033)      | 1.016<br>(0.984, 1.049)    | 0.003<br>(-0.004, 0.011)      | -0.019<br>(-0.107, 0.068)   |
| <b>Marital status (Married - REF)</b>       |                               |                               |                            |                               |   |
| Divorced                                    | 0.158***<br>(0.113, 0.202)    | 0.085***<br>(0.054, 0.117)    | 1.060**<br>(1.020, 1.102)  | 0.013**<br>(0.004, 0.022)     | 0.298***<br>(0.193, 0.402)  |
| Unmarried                                   | -0.052*<br>(-0.098, -0.006)   | -0.060***<br>(-0.092, -0.028) | 0.919***<br>(0.882, 0.958) | -0.018***<br>(-0.027, -0.010) | -0.019<br>(-0.132, 0.095)   |
| <b>Family size (≤ 3 Members REF)</b>        |                               |                               |                            |                               |   |
| 4-6 Members                                 | -0.232***<br>(-0.271, -0.193) | -0.184***<br>(-0.211, -0.157) | 0.816***<br>(0.788, 0.845) | -0.045***<br>(-0.052, -0.037) | -0.330***<br>(-0.427, -0.234)   |
| > 6 Members                                 | -0.434***<br>(-0.569, -0.298) | -0.342***<br>(-0.431, -0.253) | 0.682***<br>(0.600, 0.775) | -0.081***<br>(-0.106, -0.056) | -0.704***<br>(-1.071, -0.338)   |
| <b>Education (Less than college - REF)</b>  |                               |                               |                            |                               |   |
| Some college or more                        | -0.001<br>(-0.037, 0.036)     | -0.001<br>(-0.027, 0.025)     | 0.999<br>(0.967, 1.032)    | 0<br>(-0.007, 0.007)          | -0.003<br>(-0.092, 0.085)   |
| <b>Occupation (Natural resources - REF)</b> |                               |                               |                            |                               |   |
| Hospitality services                        | 0.051<br>(-0.041, 0.142)      | 0.046<br>(-0.017, 0.110)      | 1.069<br>(0.984, 1.160)    | 0.014<br>(-0.003, 0.032)      | -0.009<br>(-0.237, 0.219)   |
| Trade                                       | 0.146***<br>(0.065, 0.227)    | 0.093**<br>(0.036, 0.149)     | 1.086*<br>(1.010, 1.169)   | 0.018*<br>(0.002, 0.034)      | 0.251*<br>(0.049, 0.452)  |
| Professional services                       | 0.126***                      | 0.106***                      | 1.138***                   | 0.028***                      | 0.103   |

|  |                  |                  |                |                  |                 |
|--|------------------|------------------|----------------|------------------|-----------------|
|  | (0.053, 0.199)   | (0.056, 0.156)   | (1.067, 1.215) | (0.014, 0.042)   | (-0.078, 0.284) |
| Manufacturing                            | 0.056            | 0.067*           | 1.110**        | 0.023**          | -0.047          |
|  | (-0.025, 0.137)  | (0.011, 0.123)   | (1.032, 1.194) | (0.007, 0.038)   | (-0.248, 0.154) |
| Other                                    | 0.069            | 0.080**          | 1.134***       | 0.028***         | -0.064          |
|  | (-0.005, 0.143)  | (0.029, 0.131)   | (1.062, 1.212) | (0.013, 0.042)   | (-0.247, 0.120) |
| <b>Year (2010 - REF)</b>                 |                  |                  |                |                  |                 |
| 2011                                     | -0.002           | -0.076**         | 0.841***       | -0.039***        | 0.325***        |
|  | (-0.077, 0.073)  | (-0.130, -0.023) | (0.787, 0.898) | (-0.054, -0.024) | (0.147, 0.503)  |
| 2012                                     | -0.111**         | -0.164***        | 0.764***       | -0.060***        | 0.163           |
|  | (-0.185, -0.038) | (-0.216, -0.112) | (0.716, 0.815) | (-0.075, -0.045) | (-0.014, 0.341) |
| 2013                                     | -0.034           | -0.086**         | 0.851***       | -0.037***        | 0.181*          |
|  | (-0.110, 0.041)  | (-0.139, -0.033) | (0.797, 0.909) | (-0.051, -0.022) | (0.003, 0.359)  |
| 2014                                     | -0.059           | -0.107***        | 0.834***       | -0.041***        | 0.133           |
|  | (-0.133, 0.016)  | (-0.159, -0.054) | (0.781, 0.890) | (-0.056, -0.026) | (-0.044, 0.310) |
| 2015                                     | -0.066           | -0.097***        | 0.854***       | -0.036***        | 0.084           |
|  | (-0.140, 0.008)  | (-0.150, -0.045) | (0.800, 0.911) | (-0.050, -0.021) | (-0.091, 0.260) |
| 2016                                     | -0.034           | -0.097***        | 0.831***       | -0.042***        | 0.229*          |
|  | (-0.109, 0.041)  | (-0.150, -0.044) | (0.777, 0.887) | (-0.057, -0.027) | (0.050, 0.408)  |
| 2017                                     | 0.047            | -0.065*          | 0.831***       | -0.042***        | 0.452***        |
|  | (-0.030, 0.124)  | (-0.120, -0.011) | (0.777, 0.889) | (-0.057, -0.027) | (0.269, 0.634)  |
| 2018                                     | 0.068            | -0.05            | 0.844***       | -0.038***        | 0.484***        |
|  | (-0.009, 0.145)  | (-0.105, 0.005)  | (0.789, 0.903) | (-0.054, -0.023) | (0.301, 0.667)  |
| 2019                                     | 0.06             | 0.01             | 0.965          | -0.008           | 0.236*          |
|  | (-0.020, 0.141)  | (-0.048, 0.067)  | (0.900, 1.034) | (-0.024, 0.008)  | (0.049, 0.423)  |
| <b>Region (Northeast - REF)</b>          |                  |                  |                |                  |                 |
| North central, Midwest                   | 0.073*           | 0.049*           | 1.043          | 0.009            | 0.139*          |
|  | (0.016, 0.129)   | (0.010, 0.089)   | (0.993, 1.097) | (-0.002, 0.020)  | (0.002, 0.275)  |
| South                                    | 0.039            | 0.03             | 1.03           | 0.007            | 0.074           |
|  | (-0.013, 0.092)  | (-0.007, 0.067)  | (0.983, 1.079) | (-0.004, 0.017)  | (-0.053, 0.202) |
| West                                     | -0.059*          | -0.029           | 0.984          | -0.004           | -0.126          |
|  | (-0.115, -0.003) | (-0.068, 0.010)  | (0.936, 1.034) | (-0.014, 0.007)  | (-0.263, 0.011) |
| <b>Income (Poor to low income - REF)</b> |                  |                  |                |                  |                 |
| Middle to high income                    | -0.006           | -0.004           | 0.993          | -0.001           | 0               |
|  | (-0.052, 0.040)  | (-0.037, 0.028)  | (0.954, 1.035) | (-0.011, 0.008)  | (-0.112, 0.112) |
| <b>Receipt of food stamp (No - REF)</b>  |                  |                  |                |                  |                 |
| YES                                      | 0.216***         | 0.152***         | 1.147***       | 0.031***         | 0.364***        |
|  | (0.152, 0.280)   | (0.105, 0.199)   | (1.084, 1.213) | (0.018, 0.044)   | (0.211, 0.517)  |
| <b>Health status (Excellent - REF)</b>   |                  |                  |                |                  |                 |
| Good                                     | 0.309***         | 0.259***         | 1.356***       | 0.064***         | 0.508***        |
|  | (0.261, 0.357)   | (0.227, 0.291)   | (1.296, 1.418) | (0.055, 0.073)   | (0.379, 0.638)  |
| Fair                                     | 0.932***         | 0.678***         | 1.957***       | 0.150***         | 1.407***        |
|  | (0.866, 0.999)   | (0.630, 0.727)   | (1.844, 2.076) | (0.137, 0.163)   | (1.245, 1.570)  |
| Poor                                     | 1.841***         | 1.199***         | 2.591***       | 0.219***         | 2.689***        |
|  | (1.693, 1.989)   | (1.070, 1.328)   | (2.287, 2.936) | (0.188, 0.249)   | (2.382, 2.997)  |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF=reference

## CHAPTER TWO: LONG WORKING HOURS AND ONSET OF PSYCHOLOGICAL DISTRESS: A LONGITUDINAL ANALYSIS

### 2.0. OVERVIEW

**OBJECTIVES:** Long working hours have been associated with adverse health outcomes such as cardiovascular diseases and symptoms of psychological distress. Despite the high prevalence of long working hours, and psychological distress in the U.S., longitudinal studies examining the relationship between working hours and psychological distress among U.S. workers are sparse. Thus, the aim of this study was to investigate the relationship between long working hours and psychological distress using a nationally representative panel of full-time workers in the U.S.

**METHODS:** The 2010-2019 Medical Expenditure Panel Survey (MEPS) data from the University of Minnesota's Integrated Public Use Microdata Series (IPUMS) was utilized. The dependent variable was self-reported psychological distress measured and identified using the Kessler-6 scale. Using a subset MEPS, comprised of full-time employees without psychological distress at baseline were identified. Furthermore, fixed-effect linear model were utilized in modeling the relationship between working hours and the onset of psychological distress.

**RESULTS:** Overall, the sample comprised 27,562 full time workers in the U.S with 55,124 observations. The highest significant rate of onset of psychological distress was observed at 7.0% (95% Confidence Interval, CI=0.8%, 13.3%,  $p < 0.05$ ) for individuals working 61 hours or more per week (h/week), with the 30-35 h/week per week category as the reference. Also, we found that men working 51-60 h/week were significantly associated with a 7.2% increase in the onset of psychological distress (CI=0.2%, 14.2%,  $p < 0.05$ ) as compared to men working 30-35 h/week.

**CONCLUSION:** Overall, the results of this study indicate a positive association between working long hours and the onset of psychological distress. The highest risk was found in individuals who worked very long hours, and a risk increase was also observed among those who worked regular, standard hours per week when compared to those working fewer hours per week.

## 2.1. INTRODUCTION

Mental disorders, such as psychological distress (P.D.), are reported to account for about 50% of all long-term disability and morbidity in the working-age population in the Organization for Economic and Development (OECD) countries (Afonso et al., 2017). Psychological distress refers mental illness and may include signs of anxiety and depression (Bessaha, 2017). It is usually measured using the Kessler-6 (K6) scale. (Kabir, 2021; Kessler, Barker, et al., 2003). While some research reports the K6 scale as unidimensional depicting only a single factor of nonspecific psychological distress, other research indicates the K6 depicts two factors of psychological distress, mostly anxiety and symptoms of depression (Lace et al., 2020).

Psychological distress and other mental illnesses continue to be a leading cause of disease burden globally (Collaborators, 2022). A recent study reported that approximately 14% of adults in the United States (U.S.) experienced severe psychological distress in 2020 (McGinty et al., 2020). While this prevalence of psychological distress may have partially been influenced by the COVID-19 pandemic, several studies have shown similar and consistent estimates of the prevalence of psychological distress and related mental disorders, such as depression and anxiety (Bischoff et al., 2017; Collaborators, 2022; Kessler et al., 2009).

Long working hours have been linked with negative health outcomes such as cardiovascular diseases (Bannai & Tamakoshi, 2014; Cheng et al., 2014; Grosch et al., 2006; Lunde et al., 2016; Wong et al., 2019) and symptoms of psychological distress (Cheng et al., 2014; Lallukka et al., 2008; Lunde et al., 2016; Skogstad et al., 2019; Virtanen & Kivimäki, 2012). Additionally, long working hours are more prevalent in the

U.S. and Asian countries (Virtanen et al., 2018; Virtanen & Kivimäki, 2012) when compared to European countries.

A significant proportion of employees in the U.S. work longer than 40 hours per week (Williams, 2001). The International Labor Organization ranked the U.S. as the fourth highest nation in terms of the proportion of workers who worked above 48 hours per week between 2004 and 2005 (Bannai et al., 2015). This prevalence of long working hours among U.S. employees is influenced by several factors, such as incentives for higher earnings, career needs, and strong cultural influences.

Research examining the relationship between working hours and psychological distress has produced mixed results. While some studies have found statistically significant associations between long working hours and psychological distress (Afonso et al., 2017; Li et al., 2019; Virtanen et al., 2011; Wong et al., 2019), others have reported no significant associations (Angrave & Charlwood, 2015; Lee et al., 2017; Rugulies et al., 2019). These differences in findings could be mainly attributed to challenges in study design and analysis, such as cross-sectional designs, reverse causality, and endogeneity.

Cross-sectional designs, which are unable to infer causality, may not account for mental health status at baseline, thereby confounding the relationship between working hours and psychological distress. Reverse causality may arise when part-time workers are included in the analysis, as part-time work could be due to pre-existing health problems such as psychological distress (De Raeye et al., 2009; Virtanen et al., 2011). In addition, while it is crucial to control for relevant mediators and confounders, it is also necessary to be cognizant of the endogeneity that exists in the relationship between working hours and psychological distress. Hence, using analytic methods such as fixed effects, which

considerably account for endogeneity, would be beneficial in addressing this challenge (Cameron & Trivedi, 2010; Mousteri et al., 2020).

Considering the ethical challenges inherent in conducting randomized controlled studies using long working hours as an exposure, various systematic reviews and meta-analyses have been performed in an effort to establish the relationship between working hours and the development of psychological distress (Bannai & Tamakoshi, 2014; Rivera et al., 2020; Rugulies et al., 2021; Theorell et al., 2015; Virtanen et al., 2018; Watanabe et al., 2016). However, many of these studies (Bannai & Tamakoshi, 2014; Rivera et al., 2020; Theorell et al., 2015) have relied heavily on published cross-sectional studies. On the other hand, (Watanabe et al., 2016) reviewed seven cohort studies and found an inconclusive or non-statistically significant relationship between long working hours and the onset of depressive symptoms. While the study had a heterogeneous mix of studies from various countries, none of the studies included in their analysis were from the U.S. due to the scarcity of published longitudinal studies examining the relationship between long working hours and the onset of psychological distress in the U.S. Additionally, (Virtanen et al., 2018) a meta-analysis of 10 published and 18 unpublished prospective cohort studies with individual participant data and found no significant association between long working hours and depressive symptoms in North American studies.

Recently, (Rugulies et al., 2021) a study conducted a systematic and meta-analytic review of 22 cohort studies and found inconclusive evidence for the effect of long working hours on the onset of depression. Another limitation in some existing prospective studies is establishing psychological distress status at baseline. For instance, the study conducted by Dembe & Yao (2016), which evaluated 32 years of work history

(1979 to 2009), did not measure baseline depression. The inability to measure depressive symptoms at baseline was due to limitations in the methodology used in assessing depressive symptoms and chronic illness by the dataset - NLSY79 (National Youth Longitudinal Survey, 1979). This resulted in a missed assessment of baseline depression since information on chronic diseases was collected when the participants of the survey turned 40 years.

Despite the abundance of studies examining long working hours in Asian countries such as Japan and South Korea, comparatively fewer studies have examined long working hours in the U.S. (Li et al., 2019). Hence, in cognizance of the high prevalence of long working hours, psychological distress, and a lack of longitudinal studies investigating the relationship between long working hours and psychological distress among U.S. workers, the objective of this study was to examine the relationship between long working hours and psychological distress using a nationally representative panel of full-time workers in the U.S. The objective was to determine if long working hours predict the future onset of psychological distress in U.S. full-time employees without symptoms of psychological distress at baseline while controlling for endogeneity.

We utilized a modified WHO logic model (Pega et al., 2021; Rugulies et al., 2021) in this study. This model is based on the idea that the relationship between risk factors and outcomes is complex and can be understood through a process-oriented approach (Rehfuess et al., 2018). It is an a priori model, meaning it is constructed before data is collected. Similar models have been previously employed in examining the relationship between risk factors, chronic diseases (Pega et al., 2021; Rugulies et al., 2021), and health behaviors (Pachito et al., 2021).



The conceptual model, shown in *Figure 2* (Pega et al., 2021; Rugulies et al., 2021), informs that impact of long working hours on psychological distress is mediated through two main causal pathways: psycho-physiological changes and health-behavioral changes. The psycho-physiological changes pathway posits that exposure to long working hours causes the release of excessive stress hormones, activating the hypothalamic-pituitary-adrenal (HPA) axis, leading to structural lesions and functional dysregulation of the adrenal systems (Kivimäki & Steptoe, 2018), and sleep disorders (Bannai & Tamakoshi, 2014; Kronfeld-Schor & Einat, 2012). In our study we used participants' chronic disease status as a representative of this mediator. The second pathway occurs through health-related behaviors such as physical inactivity, alcohol consumption, and the use of sleep medications (Pega et al., 2021; Rugulies et al., 2021). In our study we used participants' use of medications with sedative side effects as a representative of this mediator.

We hypothesize that persons working long hours will have a higher likelihood of developing psychological distress.

Furthermore, we controlled for possible confounding factors, such as age, gender, and Social Economic Status (SES) represented by various indicators, including marital status, family size, education, occupation, region, income, and receipt of Supplemental Nutrition Assistance Program benefits. Prior studies have shown that these factors have an effect on the relationship between long working hours and psychological distress. It is known that age, sex, and SES can be linked with psychological distress (Angrave & Charlwood, 2015; Bonde, 2008; Kessler, Berglund, et al., 2003), and it is important to

control for these factors in order to draw valid conclusions about the relationship between long working hours and psychological distress.

## **2.2. METHODS**

### **2.2.1. Data Sources**

We used data from the 2010-2019 Medical Expenditure Panel Survey (MEPS), a longitudinal national probability survey (Blewett, 2020; Quality, 2020). This survey is performed by the Agency for Healthcare Research and Quality (AHRQ) and the National Center for Health Statistics (NCHS). It gathers information on health services access, utilization, expenditures, and other health-related data for non-institutionalized civilians in the U.S. The sample for MEPS is drawn from respondents of the previous year's National Health Interview Survey (NHIS). MEPS uses a panel design with five rounds of interviews over two years, with data from the first two rounds collected in the first year and data from the remaining rounds collected in the second year.

The MEPS Household Component (MEPS-HC) commenced in 1996 and provides data on participant demographics, health and employment status, healthcare utilization, healthcare expenditures by payer, insurance coverage, and satisfaction with healthcare. MEPS and NHIS oversample minority groups such as African Americans and Hispanics to increase the precision of estimates. Response rates for MEPS range from 46-71% after accounting for nonresponse rates from NHIS. MEPS-HC can be used to produce national estimates and analyzed at the individual or event level.

MEPS also includes a Medical Provider Component (MEPS-MPC), which collects additional or replacement information for the MEPS-HC survey. In addition, the MEPS-MPC includes a pharmacy component that gathers medication details, including

the national drug code (NDC) and generic names. However, MEPS-MPC is not designed to produce national estimates, so we only weighted the MEPS-HC component of the study in the descriptive statistics.

### **2.2.2 IRB Approval Statement**

This study received approval as exempt from the University of Louisville's Institutional Review Board (IRB # 21.0849). We used a publicly available, de-identified dataset for the research. It did not meet the definition of human subjects research under the common rule, so it was classified as Non-Human Subjects Research (NHSR).

### **2.2.3 Data Organization**

We used a pooled sample design and applied listwise deletion to eliminate missing cases. The MEPS dataset consists of 20 files, so we used the refined and organized version of the MEPS surveys provided by the University of Minnesota's Integrated Public Use Microdata Series (IPUMS) to facilitate data management (Blewett, 2020). The IPUMS MEPS is funded by the National Institute of Child Health and Human Development (NICHD) and offers integrated data files for each year of the MEPS-HC survey, allowing the researcher to select specific variables before downloading the dataset.

We merged the IPUMS MEPS-HC with the prescription medicine event files in the MEPS Pharmacy component, which contain information on prescribed medications obtained or purchased by participating households. Each record represents a single prescribed medication reported for an individual within a survey round, which is

approximately 4 – 5 months between data collection periods. We downloaded prescription data files from 2010-2019. We merged them with the MEPS-HC component using the identifier variable “mepsid,” which defines distinct households in the H.C. data set and pharmacy component. Thus, the mepsid enables linking the H.C. data and pharmacy component.

#### **2.2.4 Dependent Variable**

The dependent variable is psychological distress. It was recoded as a binary variable (Yes or No). The Kessler-6 (K6) scale was used in this study to identify individuals with psychological distress in the study population. The K6 is a tool for measuring general psychological distress in the community rather than a specific mental health disorder (Kabir, 2021; Kessler, Barker, et al., 2003). It is made up of six questions that ask respondents about their experiences of feelings such as nervousness, hopelessness, restlessness, depression, and perceived effort exertion and worthlessness, over the past 30 days, using a five-point Likert scale. Respective scores from each question are added to create a total K6 score ranging from 0 to 24. Respondents with a score less than five are considered to have no psychological distress; No, while those with a score of 5 or above are considered to have psychological distress; Yes (Kessler, Barker, et al., 2003). The K6 scale is valid and reliable (Kessler et al., 2002) and scores are correlated with anxiety and depression (Kubiak et al., 2012; Lace et al., 2020).

Also, the K6 scale has been validated using the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) and was determined to have good precision (estimated at the 99th percentile range of the population distribution) and consistent psychometric properties in several sociodemographic subgroups. It has also

been more effective than other scales, such as the General Health Questionnaire (GHQ), in identifying DSM-IV mood and anxiety disorders. The K6 scale has been successfully translated and used in various settings and cultures, maintaining a high level of internal consistency with a Cronbach's alpha greater than 0.80 (Easton et al., 2017).

Furthermore, we contacted the MEPS–AHRQ to obtain further information on how psychological distress was measured in the survey. MEPS-AHRQ informed our team that variable K6 is derived from the Self-Administered Questionnaire (SAQ), which is administered once each year in round 2 for the panel in its first year and round 4 for the panel in its second year. Thus, after reviewing the five rounds of the dataset, we reserved only round 2 and round 4 observations, with round 2 (year 1) representing the study's baseline. Individuals with a k6 sum of 5 or above in round 2 were excluded.

### **2.2.5 Primary Independent Variable**

In this study, the primary independent variable is the number of hours worked per week by respondents 18 years or older with a current main job. The data for this variable was collected from the MEPS-HC survey and excluded unpaid travel time to and from work. We used 30 hours per week as a baseline because it is considered full-time employment in the U.S. and enables eligibility for employer-provided health insurance coverage (Carroll & Miller, 2019; Dembe & Yao, 2016).

Previous research has shown that individuals who work fewer hours, such as part-time workers, may have different health characteristics (such as underlying health conditions) or may be working fewer hours due to family responsibilities (Kivimäki et al., 2015; Li et al., 2019; Virtanen et al., 2012). Therefore, these groups of individuals who work fewer hours may not be appropriate comparators.

Number of work hours per week is a count variable in MEPS and was stratified into five distinct categories for this study: 30-35 hours per week (h/week), 36-40 h/week, 41-50 h/week, 51-60 h/week, and 61 h/week or more. This stratification provided an effort to identify the best threshold for long working hours in relation to the onset of psychological distress.

### **2.2.6 Covariates**

The selection of covariates in this study was guided by a conceptual framework based on the World Health Organization's logic model described above. Age, marital status and family size were categorized into two groups: 18-64 years and 65 years or older (65 years representing a benchmark for transitioning to normal retirement age – including start of Medicare eligibility {Bairoliya, 2021 #1355}), married and unmarried (including divorced, widowed, and separated), two members or fewer, and greater than two members, respectively. While race was divided into three categories: White, Black, and other.

Education was divided into two categories: less than a college education and some college education or more. The occupation was divided into two categories: professional services (including management and financial occupations) and non-professional services (including jobs in the natural resources, leisure, trade, and related sectors).

Other socioeconomic factors included region, income, chronic disease status, and receipt of food stamps. The region was divided into four categories: Northeast, Midwest, South, and West. Income was divided into two categories: poor to low income (less than 100%-199% of the federal poverty line) and middle to high income (200% or more of the

federal poverty line). Chronic disease status and receipt of food stamps were recorded as binary variables (yes or no). Chronic diseases include a history of heart conditions and complications, cancer, arthritis, and asthma.

### **2.2.8 Sedative Medications and Medications with Sedative Side Effects:**

Sedative and hypnotic medications, referred to as “sleep aids,” are commonly used to induce or maintain sleep as they can suppress the central nervous system (Chong et al., 2013). It is not uncommon for individuals to use prescribed and controlled medications for their sedating side effects or recreationally. Other medical literature on the off-label use of medications not specifically intended for insomnia suggests that people may use various products with sedative properties to aid sleep. The widely referenced “Up to Date” peer-reviewed clinical practice reference includes a section on medications that can be used off-label for insomnia, such as certain antidepressants (Neubauer et al., 2021).

We utilized data from the prescription medicines event files of the Medical Expenditure Panel Survey (MEPS) pharmacy component to investigate the use of sleep aids and medications with sedative properties. We defined a prescription as any recorded instance of a specific medication and its corresponding supply frequency (Chen et al., 2020). The number of prescriptions was analyzed as a continuous quantitative variable. Utilizing the Multum Lexicon Therapeutic class Codes within MEPS, we identified medications classified as sleep aids. This classification system, integrated within the MEPS prescribed medicine files, represents the most recent categorization of medications available during data collection. Examples of medications classified as sleep aids include anxiolytics, sedatives, and hypnotics.

To identify medications with sedation as a side effect, we used Drug Facts and Comparisons®. This referential drug resource lists common or serious adverse effects of medications, such as drowsiness, dizziness, sluggishness, or heaviness (*Appendix 1*) (Comparisons eAnswers., 2021). This resource, produced by Wolters Kluwer Health, is a comprehensive compendium that provides access to various databases, modules, and medication reference tools for pharmacists and drug information experts (Grizzle et al., 2019; Roblek et al., 2015). It is widely considered a reliable source of drug information, with a completeness score of 95.8% and high rankings alongside other leading medication identifiers (Clauson et al., 2007; Jackevicius et al., 2019; Roblek et al., 2015). Additionally, Drug Facts and Comparisons® has also been found to have a sensitivity of 93% and a positive predictive value of 95.3% in identifying black box warnings (Cheng et al., 2010).

### **2.2.9 Study Population**

We extracted available individuals that participated in the survey between 2010 and 2019 (panel 14 to 22) resulting in 193,750 unweighted individuals. Participants aged 18 years or older were then selected (142,716 individuals over the 10- year period). We then identified full time workers or respondents who worked 30 hours and above per week during the survey period, yielding 77,913 participants. In addition, participants with missing variables and rounds were excluded from this study, this further shrunk the sample to 33,414 respondents. Subsequently, individuals with no recorded psychological distress in the initial round (baseline) were then selected resulting the final sample of 27,562 individuals with 55,124 observations.

### **2.2.10 Data Analysis**



The data were analyzed using Stata Statistical Software: Release 17 (StataCorp, 2021). Weights were applied to the MEPS-HC component to estimate descriptive statistics, taking into account the complex survey design of the MEPS, including the weight, clusters, and strata. Exploratory and descriptive statistics were used to examine the distribution of independent variables. The analysis of our main results was unweighted. However, following (Solon et al., 2015), we also conducted our analysis using sample weights and accounting for the complex survey design of the MEPS. The weighted results are reported in *Appendices 7 to 8*.

### **2.2.11 Statistical Analysis**

We examined the effects of work hours on psychological distress in respondents. The sample sizes and longitudinal transitions across work hours groups (30-35 hours per week, 36-40 hours per week, 41-50 hours per week, 51-60 hours per week, and 61 hours per week or more) are detailed in *Appendices 3-4* of the paper.

Using a longitudinal model, with dependent variable represented by changes in psychological distress status, while the independent variable is working hours. Changes in psychological distress status ( $\Delta PDS$ ) is modeled as a function of changes in exposure to long working hours and changes in social economic variables. We employed a fixed-effect linear regression in modeling the relationship between working hours and the onset of psychological distress. Fixed-effects models take into account the correlations between the independent variables and the error terms specific to each unit in the model. A crucial benefit of using fixed effects is that it enables the ability to control for time-invariant confounding factors that have constant effects (Allison, 2009; Gunasekara et al., 2014).

Psychological distress is then modeled as a function of the individual's work hours and socioeconomic attributes using the following model:

$$PDS_{it} = \alpha + \beta_1 H_{it} + \beta_2 S (X_{1,it} + X_{2,it} + \dots + X_{S,it}) + t_t + \mu_i + \varepsilon_{it}$$

(1)

Where  $PDS_{it}$  is a measure of psychological distress for an individual at time  $t$ ,  $\beta$  is the average effect of interest,  $H_{it}$  is a measure of working hours.  $X_{1,it}, X_{2,it}, \dots, X_{S,it}$  represents other time-varying demographic variables that could influence participants' psychological distress, including marital status, family size, occupation, income, and receipt of food stamps.  $\mu_i$  are participant's specific effects and represents the use of fixed effects estimator,  $t_t$  represents a vector of year indicators, and  $\varepsilon_{it}$  is the random error term.

We examined the prevalent transitions across covariates and only included time-varying covariates with sufficient transitions (5% or more longitudinal changes between categories) during the survey period and shown in (*Appendix 5*). The covariates used in the fixed effect model include age, marital status, income, receipt of food stamps, family size, and occupation.

The fixed effects model enabled the investigation of the relationship between changes in psychological distress, changes in working hours, and other time-varying factors among individuals while controlling for constant individual characteristics. This approach was chosen as previous research and initial analysis suggested that pooled cross-sectional regression models may lead to biased results due to correlation with unobserved time-invariant characteristics (Angrave & Charlwood, 2015; Jindra et al., 2022; Wooden et al., 2009).

Furthermore, the choice of the fixed effect linear model was further influenced by our data which is a rare event data with less than 25 % percent of ones (individuals with psychological distress) as seen in the result section. Research informs that linear model yields more accurate predicted probabilities and results than logistic model (Timoneda, 2021). The analysis was conducted separately for each gender as prior studies suggests differences in vulnerability to psychological distress and work arrangement by gender (Ahn, 2018; Petrongolo, 2004). This decision was also influenced by the higher prevalence of psychological distress among females in the sample.

As a part of the sensitivity analyses, we employed crude fixed effect regressions to investigate the relationship between working hours and psychological distress. Additionally, we examined the within-person variation for the independent variable of working hours and other control variables.

## **2.3. RESULTS**

### **2.3.1 Descriptive Statistics**

Overall, the balanced sample comprised 27,562 full time workers in the United States individuals with 55,124 observations. We classified work hours into five distinct categories (30-35 h/week, 36-40 h/week, 41-50 h/week, 51-60 h/week, and  $\geq 61$  h/week), as presented in *Table 5*. Most of the respondents in our sample reported working between 36-40 h/week, representing 60.8% of the sample. On the other hand, the category of 66 or more hours per week had the smallest representation of the sample, at 2.7%. The majority of our sample consisted of individuals between the ages of 18 and 64 years (96%), with more than half of the participants identifying as male (57.7%).

By socioeconomic factors, the majority of the sample (80.3%) identified as White, with 11.1% identifying as Black and 8.6% identifying as Other. By marital status, 61.3% of the sample reported being married. Over half of the respondents had more than two family members in a household (52%). More than two-thirds of the respondents (67.7%) had at least a college education. Additionally, over half of the respondents worked in non-professional services (54.2%), with the largest representation of respondents being from the survey year 2015 (12.2%). The largest proportion of participants resided in the southern region of the U.S. (37.8%), followed by those living in the North Central-Midwest region (21.8%).

Most respondents in the sample (86.3%) were classified as middle to high-income earners, defined as those earning at least 200% and above the federal poverty threshold. In contrast, 13.7% of respondents were classified as poor to low-income earners, defined as those earning below 200% of the federal poverty threshold. Furthermore, over half of the respondents (53.8%) reported having a chronic disease. Additionally, 15.3% of the respondents reported using medications with sedative side effects, while 2.3% of the respondents reported using sedative prescriptions.

The prevalence of psychological distress among the respondents was 8.5%. In comparison to their representation in the sample, a higher prevalence of psychological distress was observed among females (48.4% vs. 42.3%) In comparison, men were 51.6% of those with psychological distress were men as compared to their 57.7% respective sample representation. Similar trends of higher prevalence of psychological distress versus respective sample representation were also found in the following covariates: non-professional services (75.6% vs. 74.2%), poor to low-income earners (17.9% vs. 13.7%),

food stamp recipients (5.8% vs. 4.1%), individuals with chronic diseases (59.6% vs. 53.2%), those using medications with sedative side effects (23.9% vs. 15.3%), and those using sedatives (5.7% vs. 2.3%).

### **2.3.2 Long Working Hours**

The results of our study indicated a positive association between long working hours and the onset of psychological distress. In *Table 6*, the highest rate of onset was observed at 7% (95% Confidence Interval, CI=0.8%, 13.3%) for individuals working 61 hours or more per week, with the 30-35 h/week per week category as reference and statistical significance at  $p < 0.05$ . Furthermore, these relations are seen in *Figures 3 to 5*.

Respondents classified as middle to high-income earners exhibited a 1.6% (CI=0.5%, 2.7%) increase in the likelihood of developing psychological distress. Conversely, respondents who reported having greater than two members in their household were 2.4% (CI=-4.2%, -0.6%) less likely to develop psychological distress.

The results of our analysis by gender revealed a similar trend in the associations between working hours and the onset of psychological distress. Among male participants, we found that men working 51-60 hours per week were significantly associated with a 7.2% increase in the onset of psychological distress (CI=0.2%, 14.2%,  $p < 0.05$ ). Additionally, positive associations ( $p < 0.1$ ) were found between the 36-40 and 61 or more hours per week work hours categories, and the onset of psychological distress among men, associated with 5% (CI=-0.5%, 10.5%) and 8% (CI=-0.2%, 16.2%) increases respectively.

Among women participants, we found mixed but largely insignificant results. Specifically, working 61 or more hours per week was associated with a 9.4% (CI: -1.3%,

20.1%,  $p < 0.1$ ) increase in the onset of psychological distress in women. However, we found negative associations between the 41-50 hours per week and the development of psychological distress, but they were largely statistically insignificant ( $p > 0.1$ ).

### **2.3.3 Supplemental Analysis**

Crude fixed-effects regression was conducted to examine the relationship between working hours and the onset of psychological distress without accounting for individual time-varying covariates. The results, presented in *Appendix 6*, indicate a consistent trend in the association between working hours and the development of psychological distress. Specifically, when using the 30-35 hours per week category as the reference, individuals working 66 or more hours per week were found to have a 7.3% (CI=1.1%, 13.5%) increase in the development of psychological distress, which was statistically significant at  $p < 0.05$ . Additionally, those who worked 36-40 hours per week were found to have a 3.7% (0.1%, 7.3%) increase in the onset of psychological distress, which was also significant at  $p < 0.05$ .

Additionally, following (Solon et al., 2015), we also conducted our analysis using sample weights and accounting for the complex survey design of the MEPS (*Appendices 7 to 8*). The analysis yielded similar results. Though the point estimates changed, the associations were similar to the unweighted results. The differences between the two sets (unweighted vs. weighted) were consistent with the observed heterogeneity in the treatment effects and the reported oversampling for minority groups in the MEPS (Quality, 2020; Schaller & Zerpa, 2019)

## 2.4. DISCUSSION

We examined long working hours as a predictor of psychological distress in a nationally representative sample of U.S. workers. The prevalence of psychological distress among the respondents was 8.5%. This is similar to the estimated global burden of mental illness at 10% (Collaborators, 2022).

We found that working 61 or more hours per week predicted the subsequent development of psychological distress while controlling for various confounding factors in the fixed effects model. This finding is consistent with prior research that reported a higher rate of psychological distress among individuals who work very high hours per week. For example, a study found that male school teachers in Japan working over 60 hours per week had 4.7 times increased odds of having psychological distress compared to those working 40 hours or less per week (Bannai et al., 2015).

Also, a study examining over 60,000 full-time employees in Australia, found that employees that worked 60 hours per week were associated with the highest odds for severe psychological distress (Hilton et al., 2009). Additionally, another study informed that Japanese white-collar employees working 80 hours or more weekly had a higher prevalence of psychological distress than those working 40 hours or less per week (Hino et al., 2015). Furthermore, the greatest gap in working time mismatch (preferred vs. actual work hours) has been reported in individuals working over 60 hours per week (Gerson & Jacobs, 2004).

Finding the ideal threshold of long working hours, the theoretical minimum level of exposure that would result in the lowest possible risk, has been previously debated (Murray et al., 2003; Rugulies et al., 2021; Virtanen et al., 2009). Interestingly, we found

that individuals who worked 36-40 hours per week had an increased likelihood of developing psychological distress compared to those who worked 30-35 h/week. This finding suggests the minimum level of exposure to long working hours could begin just above the standard weekly work hours for some individuals.

Additionally, underemployment may contribute to the risk of psychological distress, as individuals may experience decreased job satisfaction even if they are working fewer hours. Underemployment occurs when employees are not fully utilizing their qualifications and skills, either in terms of working hours or financial rewards. Prior research (Angrave & Charlwood, 2015) suggests that underemployed men and women working 35-40 h/week and less than 35 hours weekly, respectively, were associated with an increased risk of psychological distress.

Sex-stratified analyses revealed that working 61 h/week predicted a higher incidence of psychological distress in women (9.4%) compared to men (8.0%). The marginal statistical significance of the coefficient in the sex-stratified analyses suggests interpreting this result with caution. However, this weaker statistical significance could mainly have resulted from the relatively small number of samples in the sex-stratified analyses (*Appendices 3-4*). Compared to their representation in the sample, a higher prevalence of psychological distress was observed among females in our sample (48.4% vs. 42.3%). Prior studies have also reported a higher risk of the onset of symptoms of psychological distress among women working long hours (Dembe & Yao, 2016; Shields, 1999; Virtanen et al., 2011; Watanabe et al., 2016).

Women are more likely to work longer unpaid hours, including household chores, when compared to men, as reported in previous studies (Dembe & Yao, 2016; Watanabe



et al., 2016). Furthermore, women tend to experience more negative psychosocial work conditions that cumulatively contribute to poor health (Dembe & Yao, 2016; Denton et al., 2004; Watanabe et al., 2016).

We found that respondents with a greater number of individuals living in a household had a decreased likelihood of developing psychological distress. Previous studies (Grinde & Tambs, 2016; You & Henneberg, 2022) have suggested that larger family sizes can present social benefits such as a sense of belonging and support, which could have a protective and positive influence on mental health. On the other hand, studies (AMBREEN et al., 2016; Sobotka & Beaujouan, 2014) have also reported that larger families have a higher association with psychological distress. However, the relationship between psychological distress is multifaceted and influenced by factors such as cultural identity and financial constraints.

Respondents categorized as middle to high-income earners had an increased risk of developing psychological distress. Middle to high-income earners are more likely to work longer hours and also in professional occupations, which may have further influenced the development of psychological distress in this group (Søvold et al., 2021).

Furthermore, we found that individuals in certain demographic groups had a higher prevalence of psychological distress compared to their representation in the sample. These groups included food stamp recipients, individuals with chronic diseases, and those using sedatives and medications with sedative side effects. The relationship between SES variables and psychological distress among employees has been widely studied and documented in the literature (Angrave & Charlwood, 2015; Bonde, 2008; Mousteri et al., 2020). Individuals with low SES are more likely to face

underemployment (Mousteri et al., 2020), financial difficulties (Koltai et al., 2018), and less psychosocial benefits from work (Angrave & Charlwood, 2015; Paul & Batinic, 2010), which can cumulatively contribute to an increased risk of psychological distress.

Respondents with chronic conditions (other than psychological distress), those using sedative medications, and medications with sedative side effects had a higher prevalence of psychological distress compared to their representative population. Prior research suggests that long working hours may increase the risk of chronic diseases, which in turn can contribute to psychological distress. Studies have linked long working hours to an increased risk of chronic diseases such as cardiovascular diseases, cancer, depression, diabetes, and arthritis (Dembe & Yao, 2016; Rivera et al., 2020). Additionally, long working hours can lead to fatigue and sleep deprivation, which can negatively impact mental health, potentially leading to the use of sedatives and medications with sedative properties.

Additionally, individuals with chronic conditions (excluding psychological distress) using sedative medications and medications with sedative side effects had a higher prevalence of psychological distress compared to their respective representative population. Prior studies inform that long working hours may increase the risk of chronic diseases, which can subsequently contribute to psychological distress (Dembe & Yao, 2016; Rivera et al., 2020). A study (Dembe & Yao, 2016) examined a longitudinal sample of 7,492 individuals over a 32-year period and found long working hours associated with an increased risk of developing chronic diseases such as cardiovascular diseases, cancer, depression, diabetes, and arthritis. Additionally, working long hours can lead to fatigue and sleep deprivation, which can negatively impact mental health (Bannai

et al., 2015; Bergs et al., 2018). This could influence an increase in the use of sedatives and medications with sedative side effects to aid sleep.

#### **2.4.1. Policy and Theoretical Implications**

This study provides evidence of the association between long working hours and the development of psychological distress in the U.S. labor force. If these associations are causal, the present findings suggest that more attention is needed in examining the influence of long working hours in the U.S. labor force. Despite large numbers of Americans working long hours, current regulations on working time arrangements in the U.S. are largely unregulated when compared to those in Europe (Gärtner et al., 2019; Yang et al., 2006). Few existing regulations covering working times mainly focus on the transportation sector and are primarily prescriptive rather than risk or performance-based (Gärtner et al., 2019).

Implementing interventions through national legislations and programs, such as the international labor standards on working time arrangements and working time limits (Pega et al., 2021) could help improve the overall well-being of U.S. workers. These interventions should aim to ensure that long working hours and overtime do not have negative effects on the health of workers (Landsbergis, 2018; Messenger, 2018).

Our findings suggest that the onset of psychological distress is more pronounced in individuals working very long hours. However, it is worth noting that the relationship between long working hours and psychological distress is not always straightforward. Some persons may be able to handle the requirements of long working hours without experiencing psychological distress, while others may be more vulnerable to the deleterious effects of long hours. Attributes such as an individual's coping resources, job

satisfaction, and support system may influence their ability to manage the demands of long working hours.

Additionally, there is a plausibility of increased risk for women working long hours. Therefore, it may be beneficial to further study and implement gender-specific and family-friendly interventions in mitigating the effects of long working hours. Some employers have already implemented targeted interventions for female employees working long hours (Campbell et al., 2002).

Employers can benefit from implementing measures to improve the health status of employees and provide better work hours. This can be achieved through interventions such as health and wellness promotion programs, work site screening for risk factors, and management of chronic conditions. Such measures can lead to improved performance and productivity, reduce workplace accidents caused by worker fatigue (Butler et al., 2009; Grzywacz et al., 2008), and decrease medical expenses and absenteeism due to illness (Baicker et al., 2010; Merrill et al., 2011). Furthermore, reducing long working hours can have a positive impact on employee health and well-being, which can ultimately lead to benefits for both the employer and the employee.

#### **2.4.2 Limitations and Strengths**

This study had several limitations. A limitation of this study is that it relied primarily on self-reported data, which may be subject to recall bias. However, the short intervals between rounds of data collection (4-5 months) may have minimized this bias, which is a common issue in population survey data (Kim, 2013).

Also, the dataset did not include information on several potential confounding variables such as the family history of psychological distress, external stressful life

events, and workplace psychosocial conditions limited the scope of our analysis to investigate the relationship between long working hours and psychological distress.

In addition, observations with missing covariates were excluded, which may lead to residual confounding by unmeasured covariates and selection bias. However, the short duration of measurement reduces the likelihood of missing psychological distress that could occur between the extended periods of the survey period. Also, the short duration of the survey period reduces the occurrence of healthy worker effect bias, which results from healthier workers remaining in the survey while less healthy workers drop out, creating a selection effect where the remaining employees are healthier than expected (Angrave & Charlwood, 2015; Dembe & Yao, 2016).

Our findings established a temporal relationship between long working hours and the onset of psychological distress by utilizing longitudinal data and excluding cases of baseline psychological distress among the workers. Additionally, the use of a representative sample of the working population allows for the generalization of findings to the larger population.

## **2.5. CONCLUSION**

Overall, the result of this study suggests a relationship between working long hours and the onset of psychological distress. The highest risk was found in individuals who worked very long hours, and a risk increase was also observed among those who worked regular, standard hours per week when compared to those that worked fewer hours per week. This suggests that even slight increases in weekly work hours may lead to an increased risk of psychological distress for some individuals.

Consequently, the relationship between long working hours and psychological distress may be due to a combination of factors, such as sleep deprivation, underemployment, reduced social support and leisure time, and increased job strain and work-related stress.

Additionally, the study revealed gender differences in the risk of developing psychological distress among workers and found that social economic status may influence the onset of psychological distress among U.S. workers. Hence, it is crucial to consider the potential gender-specific effects of long working hours and explore the implementation of interventions that are tailored to the needs of female workers. Furthermore, factors such as income, chronic disease status, and the use of sleep aids were identified as relevant contributing factors.

It is important for individuals and organizations to be aware of the risks of long working hours on the development and prevalence of psychological distress and other negative health outcomes. Thereby taking informed steps to mitigate these risks through the implementation of policies that promote work-life balance and utilizing interventions that can help reduce work-related stress. Also, government and employers should implement interventions through laws and programs that promote a balance between work and personal life, decrease job-related stress, and improve and maintain the overall health and well-being of the population and employees.

Figure 2. Conceptual model of the possible causal relationship between exposure to long working hours and psychological distress (adapted from Pega et al., 2021; Rugulies et al., 2021)




Table 5. Descriptive statistics of work hours and psychological distress.

| Variables                   | Overall Frequency | Overall Weighted % | Psychological Distress |         |
|-----------------------------|-------------------|--------------------|------------------------|---------|
|                             |                   |                    | No (%)                 | Yes (%) |
| <b>Weekly Working Hours</b> |                   |                    |                        |         |
| 30-35                       | 2,903             | 8.4                | 0.9                    | 9.4     |
| 36-40                       | 16,695            | 52.6               | 4.8                    | 57.4    |
| 41-50                       | 5,221             | 20.0               | 1.9                    | 21.9    |
| 51-60                       | 1,990             | 7.7                | 0.7                    | 8.3     |
| ≥ 61                        | 753               | 2.7                | 0.2                    | 3.0     |
| <b>Age (years)</b>          |                   |                    |                        |         |
| 18-64                       | 26,556            | 96.0               | 87.7                   | 8.3     |
| ≥ 65                        | 1,006             | 4.0                | 3.8                    | 0.2     |
| <b>Sex</b>                  |                   |                    |                        |         |
| Male                        | 15,664            | 57.7               | 58.3                   | 51.6    |
| Female                      | 11,898            | 42.3               | 41.7                   | 48.4    |
| <b>Race</b>                 |                   |                    |                        |         |
| White                       | 19,866            | 80.3               | 80.4                   | 80      |
| Black                       | 4,612             | 11.1               | 11.2                   | 9.6     |
| Other                       | 3,084             | 8.6                | 8.4                    | 10.3    |
| <b>Marital Status</b>       |                   |                    |                        |         |
| Married                     | 16,221            | 61.3               | 62                     | 53.7    |
| Unmarried                   | 11,341            | 38.7               | 38                     | 46.3    |
| <b>Family Size</b>          |                   |                    |                        |         |
| ≤ 2 Members                 | 11,677            | 48.0               | 47.9                   | 48.6    |
| > 2 Members                 | 15,885            | 52.0               | 52.1                   | 51.4    |
| <b>Education</b>            |                   |                    |                        |         |
| Less than college           | 10,607            | 32.3               | 32.1                   | 33.5    |
| Some college or more        | 16,629            | 67.7               | 67.9                   | 66.5    |
| <b>Occupation</b>           |                   |                    |                        |         |
| Professional services       | 10,796            | 45.8               | 46.0                   | 43.0    |
| Non-professional services   | 16,766            | 54.2               | 54.0                   | 57.0    |

(continued)



Table 5. (Continued).

| Variables   | Overall Frequency | Overall Weighted % | Psychological Distress |         |
|---|-------------------|--------------------|------------------------|---------|
|   |                   |                    | No (%)                 | Yes (%) |
| <b>Year</b>   |                   |                    |                        |         |
| 2010  | 2,773             | 9.8                | 9.7                    | 11.3    |
| 2011  | 3,529             | 11.7               | 11.6                   | 13      |
| 2012  | 3,497             | 10.9               | 10.7                   | 12.3    |
| 2013  | 3,009             | 10.0               | 9.9                    | 10.3    |
| 2014  | 2,992             | 10.6               | 10.6                   | 11      |
| 2015  | 3,206             | 12.2               | 12.3                   | 11.4    |
| 2016  | 3,008             | 11.4               | 11.5                   | 10      |
| 2017  | 2,895             | 12.0               | 12.2                   | 9.3     |
| 2018  | 2,653             | 11.4               | 11.4                   | 11.5    |
| <b>Region</b>   |                   |                    |                        |         |
| Northeast   | 4,169             | 17.5               | 17.7                   | 15      |
| North Central,<br>Midwest                               | 5,498             | 21.8               | 21.7                   | 23.1    |
| South   | 10,536            | 37.8               | 37.8                   | 38.1    |
| West  | 7,359             | 22.9               | 22.8                   | 23.8    |
| <b>Income</b>   |                   |                    |                        |         |
| Poor to low<br>income                                   | 5,876             | 13.7               | 13.3                   | 17.9    |
| Middle to high<br>income                                | 21,686            | 86.3               | 86.7                   | 82.1    |
| <b>Receipt of food stamp</b>                            |                   |                    |                        |         |
| No  | 25,682            | 95.9               | 96.1                   | 94.2    |
| Yes   | 1,880             | 4.1                | 3.9                    | 5.8     |
| <b>Limitation</b>                                       |                   |                    |                        |         |
| No  | 25,007            | 90.2               | 90.4                   | 84.8    |
| Yes   | 2,555             | 9.8                | 9.6                    | 15.2    |
| <b>Chronic Diseases</b>                                 |                   |                    |                        |         |
| No  | 13,318            | 46.2               | 46.8                   | 40.4    |
| Yes   | 14,244            | 53.8               | 53.2                   | 59.6    |
| <b>Medication with Sedative Effects Use<sup>†</sup></b> |                   |                    |                        |         |
| No  | 23,344            | 84.7               | 85.5                   | 76.1    |
| Yes   | 4,218             | 15.3               | 14.5                   | 23.9    |
| <b>Sedatives Use<sup>†</sup></b>                        |                   |                    |                        |         |
| No  | 26,930            | 97.7               | 98.0                   | 94.3    |
| Yes   | 632               | 2.3                | 2.0                    | 5.7     |
| Total   | 27,562            | 100                | 91.5                   | 8.5     |

<sup>†</sup>= unweighted frequency

Table 6. Association Between long working Hours and the onset of psychological distress: Fixed effects regression analysis.

| <b>Variables</b>                             | <b>Overall</b>                | <b>All Men</b>             | <b>All Women</b>              |
|--|-------------------------------|----------------------------|-------------------------------|
| <b>Work hours/week (30-35hrs/week - REF)</b> |                               |                            |                               |
| 36-40 h/week                                 | 0.036**<br>(0.000, 0.072)     | 0.050*<br>(-0.005, 0.105)  | 0.023<br>(-0.025, 0.070)      |
| 41-50 h/week                                 | 0.029<br>(-0.014, 0.072)      | 0.048<br>(-0.013, 0.109)   | 0.007<br>(-0.059, 0.073)      |
| 51-60 h/week                                 | 0.034<br>(-0.017, 0.085)      | 0.072**<br>(0.002, 0.142)  | -0.046<br>(-0.131, 0.040)     |
| ≥ 61 h/week                                  | 0.070**<br>(0.008, 0.133)     | 0.080*<br>(-0.002, 0.162)  | 0.094*<br>(-0.013, 0.201)     |
| <b>Age (18-64years - REF)</b>                |                               |                            |                               |
| ≥ 65 years                                   | 0.061***<br>(0.030, 0.092)    | 0.068***<br>(0.025, 0.111) | 0.052**<br>(0.008, 0.097)     |
| <b>Marital Status (Married - REF)</b>        |                               |                            |                               |
| Not married                                  | -0.009<br>(-0.032, 0.014)     | -0.015<br>(-0.046, 0.015)  | 0.002<br>(-0.033, 0.037)      |
| <b>Family Size (≤ 2 Members REF)</b>         |                               |                            |                               |
| > 2 Members                                  | -0.024***<br>(-0.042, -0.006) | -0.006<br>(-0.028, 0.017)  | -0.046***<br>(-0.075, -0.017) |
| <b>Occupation (Natural resources - REF)</b>  |                               |                            |                               |
| Professional services                        | -0.029*<br>(-0.060, 0.003)    | -0.021<br>(-0.057, 0.016)  | -0.043<br>(-0.098, 0.011)     |
| <b>Income (Poor to low income - REF)</b>     |                               |                            |                               |
| Middle to high income                        | 0.016***<br>(0.005, 0.027)    | 0.014**<br>(0.001, 0.027)  | 0.019*<br>(-0.000, 0.037)     |
| <b>Receipt of food stamp (No - REF)</b>      |                               |                            |                               |
| Yes  | 0.010<br>(-0.009, 0.030)      | 0.022*<br>(-0.000, 0.044)  | -0.004<br>(-0.036, 0.029)     |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Figure 3. Onset of psychological distress according to working hours from the fixed effects regression model of the overall sample.





Figure 4. Onset of psychological distress according to working hours from the fixed effects regression model of all men sample.



Figure 5. Onset of psychological distress according to working hours from the fixed effects regression model of all women sample.



CHAPTER THREE: ASSOCIATIONS BETWEEN WORKING HOURS, HEALTH CARE UTILIZATION, AND PRESCRIPTION MEDICATIONS WITH SEDATIVE SIDE EFFECTS.

**3.0. OVERVIEW**

**OBJECTIVES:**

Long working hours are associated with several chronic diseases and the use of sedative medications. This continuous prevalence and increase in long working hours are expected to influence the consistent rise in chronic diseases and subsequent reflection in healthcare use in the United States. However, there is a lack of literature examining the relationship between working hours, health care utilization, and the use of sedative medications in the U.S. Hence, we employed a nationally representative panel of full-time workers in the U.S. to examine these relationships.

**METHODS:**

This study utilized the 2010-2019 Medical Expenditure Panel Survey (MEPS) data. Prescription medications with sedation as a side effect were identified. Respondents' annual healthcare utilization information was identified in the longitudinal survey and utilized. Fixed-effect estimators were employed in modeling the relationship between working hours, health care utilization, and the use of prescription medications with sedative side effects.

**RESULTS:**

In total, the balanced sample was comprised of 43,993 full-time employees in the U.S. Individuals working 61 or more hours per week had the highest odds of using outpatient medical providers (adjusted Odds Ratio (aOR)=2.01, 95% confidence interval (CI) 1.04, 3.86,  $p < 0.05$ ). Also, respondents working 36-40 hours per week were associated with increased use of outpatient medical providers (aOR=1.34, 95% CI=1.01, 1.78). Also, we found a similar pattern of associations between working hours and the use of outpatient services among individuals using medications with sedative properties.

**CONCLUSION:**

Our findings suggest that working hours are significantly associated with the use of outpatient services, with the highest odds in workers that worked 61 or more hours per week. Long working hours negatively impact the health status of individuals, which can reflect in the increased use of health care services. Implementing policies that encourage work-life balance and aid interventions that decrease work-related stress may help in mitigating risks associated with long work hours.

### **3.1. INTRODUCTION**

Long working hours remain prevalent globally and, in the U.S. The International Labor Organization previously ranked the U.S. as the fourth highest nation in the proportion of workers who work overtime weekly (Bannai et al., 2015). This trend of long working hours in the U.S. is driven by several factors, such as motivations for increased wages, career development, and strong cultural influences (Li et al., 2019).

Overtime hours have been associated with an increase in negative health outcomes, including cardiovascular diseases (Cheng et al., 2014; Lunde et al., 2016; Skogstad et al., 2019) and mental ailments (Kim et al., 2016; Wong et al., 2019). Also, long working hours have been linked with a rise in risky health behaviors such as increased use of sleep medications (Ezekekwa et al., 2023), high alcohol use, and smoking (Bannai et al., 2015; Virtanen et al., 2015).

The increase in the incidence and prevalence of chronic diseases associated with long working hours is expected to reflect in the increased utilization of health resources among employees. Previous studies have reported long working hours associated with a decreased use of preventive health services among employees, mainly due to a decrease in the availability of time among overtime workers in countries like Canada and Korea (Fell et al., 2007). Preventive health services such as routine health checks and recommended vaccinations are crucial in maintaining the health of individuals, especially those with increased risk of chronic morbidities.

Furthermore, studies examining the associations between long working hours, health care utilization, and the use of medications in the U.S. is sparse. Thus, we utilized a nationally representative panel of full-time workers in the U.S. to examine the

relationship between working hours, health care utilization, and the use of medications with sedative side effects.

We utilized the Andersen health care utilization model in this study (Andersen, 1995). This model is relevant to our research as it posits that a person's health behavior and use of health care resources are influenced by enabling, predisposing, and needs factors.

Our conceptual model (Figure 6) explains that working hours influence the mutable population characteristics (specifically the need factors such as health status) and recurrently influence individuals' health behaviors. Work hours represent the work environment, which in turn is influenced by the occupation, and have been reported to influence the use of health services and changes in health behaviors, such as the use of sleep medication and alcohol consumption (Virtanen et al., 2015; Wong et al., 2019).

The population characteristics are composed of predisposing, enabling, need, and motivating factors. The predisposing construct represents individuals' conditions that influence their use of health services (Andersen & Newman, 1973; Andersen, 1995). This construct is primarily categorized into demographic and social factors, with demographic factors including age, gender, and race. In addition, an individual's social status within the community, including factors such as marital status, education, and occupation, is also an important component of the social structure that influences health service utilization.

Furthermore, the enabling factors represent the resources that are available to individuals, such as income, health insurance, and geographic location, which aids the utilization of health care services. An individual's insurance status plays a crucial role in



determining their use of health care services and obtaining prescription medications for sleep. However, it is important to note that health insurance is an outcome variable, as employee health insurance in the United States is typically only available to those working 30 or more hours per week (Carroll & Miller, 2019). The level of income is positively associated with access to health care. In addition, the need factor is measured by an individual's health limitations, representing a perceived need for health care services.

Motivating factors also play a role in the amount of time an individual spends working. Professionals and individuals with higher socioeconomic status may be motivated to work overtime due to the potential financial and career benefits. However, others may be compelled to work long hours due to financial difficulties, even if they earn meager wages. To assess motivation, family size and receipt of food stamps were employed as indicators. Large family sizes may result in increased financial demands, while receipt of food stamps suggests financial difficulties associated with food insecurity. Prior studies have utilized family size and receipt of food stamps as measures of financial strain (Bhattarai et al., 2005; Kabir, 2021).

## **3.2. METHODS**

### **3.2.1 Data Sources**

The data for this study was obtained from the Medical Expenditure Panel Survey (MEPS) conducted between 2010-2019. MEPS is a national probability survey carried out by the Agency for Healthcare Research and Quality (AHRQ) and the National Center for Health Statistics (NCHS) (Quality, 2020) . The survey collects health-related

information on non-institutionalized civilians in the U.S. by evaluating their access to health services, utilization, expenditures, and other relevant data. In creating a sample, MEPS identifies and records respondents from the previous year's National Health Interview Survey (NHIS). It utilizes a panel design with five interview rounds conducted over a two-year period, with the initial two rounds occurring in the first year and the remaining rounds in the second year.

The MEPS Household Component (MEPS-HC) began in 1996 and yields crucial data on participant demographics, health, and employment status, health care utilization and expenditures, insurance coverage, and satisfaction with health care. To improve the accuracy of estimates, MEPS and NHIS oversample minority groups such as African Americans and Hispanics. The response rates for MEPS range from 46-71% after adjusting for nonresponse rates from NHIS. MEPS-HC data can be utilized to generate national estimates and analyzed at both individual and event levels.

MEPS utilizes a Medical Provider Component (MEPS-MPC) that gathers supplementary or alternative data to the MEPS-HC survey. Furthermore, the MEPS-MPC includes a pharmacy component that collects medication information, including the national drug code (NDC) and generic names. However, the MEPS-MPC is not intended to generate national estimates; therefore, we only weighted the MEPS-HC component in the descriptive statistics of the study. However, following Solon et al. (2015), we also conducted our analysis using sample weights and accounting for the complex survey design of the MEPS. The weighted results are reported in *Appendices 6 to 8*.

### **3.2.2 IRB Approval Statement**

The research conducted in this study received exemption approval from the Institutional Review Board (IRB # 21.0849) at the University of Louisville. We utilized a de-identified dataset that was publicly available. The study was classified as Non-Human Subjects Research (NHSR), since the dataset did not satisfy the definition of human subject research.

### **3.2.3 Data Organization**

We utilized a pooled sample design and listwise deletion to remove missing observations. The MEPS dataset comprised 20 files; thus, to aid the data management, we used the University of Minnesota's Integrated Public Use Microdata Series (IPUMS). This series provides organized and refined data from the MEPS surveys (Blewett, 2020). The National Institute of Child Health and Human Development (NICHD) funds IPUMS MEPS, which enables researchers to select individual variables prior to downloading the dataset.

We combined the MEPS pharmacy component's prescription medicine event files with the IPUMS MEPS-HC via merging. The event files possess information on prescribed medications received or purchased by participating households. Every prescription record denotes a single prescribed medication registered for respondents within a survey round, estimated at 4-5 months between data collection periods. We downloaded the prescription data files from 2010-2019 and merged them with the MEPS-HC component using the identifier variable "mepsid," which distinguishes distinct households in the H.C. dataset and pharmacy component.

### **3.2.4 Dependent Variables**

Information on the respondent's annual health care utilization was generated at the end of each year of the longitudinal survey. The total health care services use information of the respondents for the survey's first and second years was utilized. The dependent variables in the data were entered as count variables and were recoded into binary variables, which include annual outpatient visits, office-based visits, emergency department visits, hospital discharges, and dental visits.

The annual outpatient visits measure the total number of visits to outpatient medical providers during the year. It entails consultations with either physicians or non-physician providers in outpatient settings and clinics. Annual office-based visits measure the total number of visits to the medical of medical providers in office-based settings or clinics over the course of a year.

Furthermore, the annual emergency department visits to the emergency room reflect the total number of visits to the emergency room during the course of the year. Likewise, it includes encounters with both physicians and non-physician providers. At the same time, annual hospital discharges represent the total number of hospital discharges experienced by a respondent in a year, including both same-day discharges and hospitalizations lasting one or more nights before discharge. In addition, annual dental visits denote the respondent's total number of dental care consultations involving general dentists and orthodontists.

### **3.2.5 Primary Independent Variable**

The number of hours worked per week by the respondents is the primary independent variable. Respondents aged 18 years and above with a current main job were selected. This information was gathered from the MEPS-HC survey, and it does not

include unpaid travel time to and from work. Individuals working 30 hours or more weekly were classified as full-time workers, similar to the U.S. regulation that enables eligibility for employer-provided health insurance coverage (Carroll & Miller, 2019; Dembe & Yao, 2016).

Previous studies inform that part-time workers and individuals working fewer hours may have different health characteristics, such as underlying health conditions, or may have work fewer hours due to family responsibilities (Kivimäki et al., 2015; Li et al., 2019; Virtanen et al., 2012). Hence, part-time workers or people working fewer times may not be appropriate comparators to standard and overtime workers. To identify the best threshold for long working hours, working hours were divided into five categories: 30-35 hours per week (h/week), 36-40 h/week, 41-50 h/week, 51-60 h/week, and 61 h/week or more.

### **3.2.6 Covariates**

The covariate selection was guided by the Gelberg-Andersen behavioral health model, with predisposing factors such as age, race, marital status, family size, education, occupation, and panel year. Predisposing factors were categorized into distinct age groups of 18-26, 27-64, and  $\geq 65$  years; racial categories of White, Black, and other; marital status categories of married, divorced (including widowed and separated), and unmarried; and family size categories of  $\leq 3$ , 4-6, and  $> 6$  members. Education was recoded into two categories, less than college and some college or more, while the occupation was recoded into professional (including financial and administrative services) and non-professional services (including jobs in the natural resources, leisure, trade, and related sectors).

Enabling factors, such as region and income, were also considered. Health insurance status, as an outcome variable of working hours, was excluded to avoid multicollinearity. U.S. employees working 30 or more hours per week are eligible for employee health insurance. Thus, our study respondents are eligible for employee health insurance. The MEPS region variable was recoded into northeast, mid-west (including northcentral), south, and west. Income was recoded into poor to low income less than 100% - 199% of the federal poverty line) and middle to high-income 200% or more of the federal poverty line).

Motivating factors included receipt of food stamps and limitations. The receipt of food stamps was a binary variable. Limitation is a summary measure of the respondents, instrumental activities of daily living, functional and activity limitations. Also, it was recoded into a yes/no variable.

### **3.2.7. Sedative Medications and Medications with Sedative Side Effects**

Sedative and hypnotic medications usually referred to as "sleep aids," are widely used to initiate or maintain sleep owing to their ability to suppress the central nervous system (Chong et al., 2013). The off-label use of medicines not specifically intended for treating insomnia is also prevalent, with certain antidepressants being among the drugs that can be used for this purpose (Neubauer et al., 2021). The medications used off-label are usually used for the sedating side effects and are sometimes used recreationally (Citrome, 2017; Eugene, 2020; Hägg et al., 2020).

To examine the use of medications with sedative properties, we analyzed data from the prescription medicines event files of the Medical Expenditure Panel Survey

(MEPS) pharmacy component. A prescription was defined as a recorded event of an individual medication (Chen et al., 2020).

Furthermore, we employed Facts and Comparisons®, a comprehensive compendium produced by Wolters Kluwer Health, to identify medications with sedative properties (Comparisons eAnswers., 2021). Facts and Comparisons® lists common or serious adverse effects of medications, such as dizziness, drowsiness, sluggishness, or heaviness. It has been widely considered a reliable source of drug information, with a completeness score of 95.8% and high rankings alongside other leading medication identifiers (Clauson et al., 2007; Jackevicius et al., 2019; Roblek et al., 2015). Also, this drug resource has been reported to have a sensitivity of 93% and a positive predictive value of 95.3% in identifying black box warnings (Cheng et al., 2010).

### **3.2.8. Study Population**

Inclusion criteria for the study participants included being 18 years or older, participated in the survey between 2010 and 2019 (panels 14 to 22), and worked at least 30 hours per week. In contrast, the exclusion criteria included individuals who worked less than 30 hours per week (considered part-time workers) and respondents with missing variables.

### **3.2.9. Data Analysis**

The Stata Statistical Software: Release 17 (StataCorp, 2021) was used to analyze that data. The MEPS-HC component was weighted to estimate descriptive statistics, accounting for the complex survey design of the MEPS, including weight, clusters, and strata, to enable the

generalizability of the study and correct estimation of effect size. Exploratory and descriptive statistics were employed to evaluate the distribution of independent variables.

### 3.2.10. Statistical Analysis

We investigated the effects of work hours on the use of health care resources in participants who transitioned between different work hour categories, including 30-35 hours per week, 36-40 hours per week, 41-50 hours per week, 51-60 hours per week, and 61 hours per week or more. The sample sizes and prevalence of transitions in these workhour categories are described in the Appendices of the paper.

Changes in health care utilization are modeled as a function of changes in exposure to long working hours and changes in social and economic variables. Fixed-effect estimator in modeling the relationship between working hours and health care utilization.

This estimator accounts for the interdependence among independent variables and error terms, which are unique to each unit within the model. An important benefit of employing fixed effects lies in its capability to adjust for time-invariant confounding factors with constant effects. (Allison, 2009; Gunasekara et al., 2014).

Health care utilization is then modeled as a function of the individual's work hours status and socioeconomic attributes using the following model:

$$HU_{it} = \alpha + \beta_1 H_{it} + \beta_2 S (X_{1,it} + X_{2,it} + \dots + X_{S,it}) + t_t + \mu_i + \varepsilon_{it}$$

(1)

Where  $HU_{it}$  is a measure of health care use for an individual at time  $t$ ,  $\beta$  is the average effect of interest,  $H_{it}$  is a measure of changes in working hours.  $X_{1,it}, X_{2,it}, \dots, X_{S,it}$  represents other time-varying demographic variables that could influence participants



psychological distress including age, marital status, family size, occupation, income, and receipt of food stamps.  $\mu_i$  are participant's specific effects and represents the use of fixed effects estimator,  $t_i$  represents a vector of year indicators, and  $\varepsilon_{it}$  is the random error term.

While the fixed effects logit model is defined by the logistic probability of  $HU_{it}$ :

$$Pr(HU_{it}=1) = \frac{\exp(\alpha + \beta_1 H_{it} + \beta_{2s}(X_{1,it} + X_{2,it} + \dots + X_{s,it}))}{1 + \exp(\alpha + \beta_1 H_{it} + \beta_{2s}(X_{1,it} + X_{2,it} + \dots + X_{s,it}))}$$

(2)

Where  $HU_{it}=1$ , represents health care use by an individual at time  $t$ .

The use of fixed effects modeling facilitated the examination of the association between changes in health care utilization, changes in working hours, and other time-varying factors within individuals while adjusting for consistent individual attributes. The fixed effects estimator was preferred based on prior literature and preliminary Analysis, which indicated that employing pooled cross-sectional regression models might produce distorted outcomes due to the association with unobserved time-invariant traits (Angrave & Charlwood, 2015; Jindra et al., 2022; Wooden et al., 2009). Fixed effects estimators were also used to investigate the association between working hours and health care utilization among participants using medications with sedative side effects.

### 3.3. RESULT

#### 3.3.1. Descriptive Statistics

This balanced panel data for the analysis included 43,993 individuals with 87,986 observations. Weekly work hours was classified into five distinct categories (30-35 h/week, 36-40 h/week, 41-50 h/week, 51-60 h/week, and  $\geq 61$  h/week) as presented in *Table 7*.

Most of the participants in our analysis fell within the age range of 27 to 64 years (86.2%), with more than half of the participants male (56.1%). Majority of the sample identified as White (79.7%), while 11.3% identified as Black, and 9% as Other. Also, most of the respondents were married (59.5%), and the majority had some college or higher education (67.2%), while about half of the respondents had over two members living in a household. Non-professional services (55.3%) was the largest occupational category, and the survey years 2015 and 2016 had the highest weighted percentage (11.7%).

The largest proportion of the participants resided in the southern region of the U.S. (37.5%), followed by those living in the North Central-Midwest (21.9%), and the Northeast had the smallest proportion (17.5%). The majority of respondents were classified as middle to high-income earners (85.1%), earning 200% or more of the federal poverty threshold. On the other hand, 14.9% were classified as poor to low-income earners or earned below 200% of the federal poverty threshold. Less than 5% of the respondents reported receiving food stamps (4.7%), while most did not have any reported limitation. Additionally, 27.4% of the respondents reported using medications with sedative side effects.

By health care utilization, few of the workers in our analysis had one or more outpatient visits (14.9%), while most had at least one office-based visit (70.9%). The majority of the respondents (89.5%) had no visits to the emergency department during the survey period, and most had no recorded hospital discharge (95.3%). Also, 43.5% of the workers had one or more dental visits during the survey period.

### **3.3.2. Working Hours**

The results of the fixed effects models for the association between working hours and health care utilization is shown in *Table 8*. Fixed regressions were conducted separately with the dependent variables: annual outpatient visits, office-based visits, emergency department visits, hospital discharges, and dental visits. We found significant results only with the annual outpatient visits. Individuals working 61 or more hours per week had the highest odds of using outpatient medical providers (adjusted Odds Ratio (aOR)=2.01, 95% confidence interval (CI)= 1.04, 3.86), with the 30-35 h/week per week category as a reference and statistical significance at  $p < 0.05$ . Also, respondents working 36-40 hours per week were associated with increased use of outpatient medical providers (aOR=1.34, 95% CI=1.01, 1.78) as compared to those working 30-35 hours per week. Also, we found dose-like response associations between working hours and outpatient use, as seen in *Figure 7*.

In addition, respondents aged 65 years and older had significantly increased odds (aOR=1.74, 95% CI=1.0, 3.0) of using outpatient services as compared to those 18-26 years of age. Also, respondents with limitation had increased odds (aOR=1.43, 95% CI=1.26, 1.63) of using outpatient services as compared to those without recorded limitation. On the other hand, employees with more than two family members living in a household had fewer odds of using outpatient medical providers as compared to those with less than two family members living in a household.

Furthermore, from the fixed logit marginal analysis (*Table 9 and Figure 8*), we found a similar dose-like response association between working hours and outpatient use. The predicted probability of using outpatient medical providers was greater (Predicted probability (PB)=0.07, 95% CI=0.00, 0.14,  $p < 0.05$ ) for an individual that worked 61 or

more hours weekly as compared to one that worked less than 30-35 hours per week. Also, the predicted probability of using an outpatient medical provider was greater for an individual that worked 36-40 hours per week as compared to one that worked 30-35 hours per week, with  $p < 0.05$ . Furthermore, the logit marginal effects were comparable to the fixed linear effect regression of the Analysis. The fixed effect linear model yielded a similar dose-response association between working hours and outpatient use. It showed that individuals working 61 or more hours per week had the highest likelihood at 4.6% (95% CI=0.0%, 9.0%,  $p < 0.05$ ), of using outpatient medical providers as compared the those working 30-35 hours per week. Similarly, individuals working 36-40 hours per week had a 2% (95% CI=0.0%, 4%,  $p < 0.05$ ) increased likelihood of using outpatient medical provider services.

### **3.3.3 Health care Utilization Among Individuals Using Medications with Sedative Effects**

The results of the fixed effects models for the association between working hours and health care utilization among respondents using medications with sedative side effects (*Table 10*) were similar to the fixed effects model results from the overall sample. Likewise, significant results only with the annual outpatient visits. Also, a dose response was found between working hours and outpatient use. However, only 36-40 h/week category were significant at  $p < 0.01$ , while the 41-50h/week (aOR=1.52, 95% CI=0.95, 2.43) and 51-55 h/week (aOR=1.89, 95% CI= 0.91, 3.89) categories were significant at  $p < 0.1$ . Also, the logit marginal results produced an increased predicted probability

(PB=0.13, 95% CI=0.04, 0.22) for an individual working 36-40 hours when compared to a respondent working 30-35 hours per week.

### **3.4 Supplemental Analysis**

We carried out further analyses to delineate the pattern of visits for the outpatients' medical providers. Annual outpatient visits in the data are composed of mainly annual outpatient physician visits and annual outpatient non-physician visits. However, data for the annual outpatient non-physician visits were not available for the years 2017-2019. Thus, we were only able to further analyze annual outpatient physician visits that have full data records.

Fixed effects models were similarly utilized in examining the association between working hours and outpatient physician visits (*Appendices 4 and 7*). We found significant associations between working hours and outpatient physicians visits. This association was most pronounced between individuals working 61 hours or more per week and visits to outpatient physicians. The fixed effects logit model showed that participants working 61 or more hours per week had increased odds (aOR=5.04, 95% CI=0.95, 26.6,  $p < 0.1$ ) of using outpatient services as compared to the participants working 30-35 hours per week. Also, the logit marginal effects and fixed linear models yielded similar associations significant at  $p < 0.05$ , for respondents working 61 or more hours per week as compared to those working 30-35 hours per week. Furthermore, fixed effects regression models and marginal analysis are shown in the *Appendices: 2 to 5*, for other dependent variables: office-based visits, emergency department visits, hospital discharges, and dental visits.

Additionally, following (Solon et al., 2015), we also conducted our analysis using sample weights and accounting for the complex survey design of the MEPS (*Appendices 6 to 8*). The analysis yielded similar results. Though the point estimates changed, the associations were similar to the unweighted results. The differences between the two sets (unweighted vs. weighted) were consistent with the observed heterogeneity in the treatment effects and the reported oversampling for minority groups in the MEPS (Quality, 2020; Schaller & Zerpa, 2019)

### **3.4. DISCUSSION**

We found a high prevalence of the use of medications with sedative side effects among the respondents. Significant associations were also found between working hours and the use of outpatient medical services. Individuals working 61 or more hours per week had the highest odds of using outpatient medical providers. Previous research posits that the deleterious effects of long working hours are most pronounced in those working high overtime hours (Bannai et al., 2015).

Outpatient medical services include alcohol and drug abuse clinics; well-baby clinics; obesity clinics; eye, ear, nose, and throat clinics; family planning clinics; cardiology clinics; internal medicine departments; physical therapy clinics; and radiation therapy clinics. This high use of outpatient medical services in respondents working very long hours may be due the poor health status in individuals working long hours. Previous studies report that long working hours have been linked with an increase in chronic morbidities and negative health outcomes (Cheng et al., 2014; Lunde et al., 2016; Skogstad et al., 2019).

However, increased associations were also found among employees working 36-40 hours per week when compared to those working 30-35 hours per week. This indicates that for some people, the minimum threshold for extended work hours could begin slightly above the usual weekly work hours. Prior research has explored determining the optimal threshold for long working hours that would result in the lowest risk level (Murray et al., 2003; Rugulies et al., 2021; Virtanen et al., 2009).

We did not find significant associations between working hours and office-based visits, emergency department visits, hospital discharges, and dental visits. However, we found significant associations between working times and outpatient medical providers, which suggests that working time influences the use of outpatient medical services. Several literature reports that long working times negatively influence the use of primary and preventive health care services, including office-based services (Fell et al., 2007; Fukuoka et al., 2010; Sato et al., 2011; Seok et al., 2016).

A crucial rationale posited for this decreased use of primary health care services is the lack of time available to full-time workers who work overtime. A study (Seok et al., 2016) examining the relationship between long working hours and unmet health care needs among 8,369 Korean workers found "lack of time" as the most reported reason for unmet health care needs. Also, another study (Sato et al., 2011) reported that employees working very long hours have an increased likelihood of using health care resources with decreased time requirements.

A consistent pattern of associations was found between working hours and the use of outpatient services among respondents that used medications with sedative side effects. Previous research (Bush et al., 2009) examined the long-term effect of

employment on health care use among persons with severe mental illness and co-occurring disorders. They found that stable employment significantly decreases inpatient service use while increasing outpatient service use among the study participants. The association between working hours, use of outpatient services and medications with sedative effects might be part of a dynamic described as “psychiatrization”. Psychiatrization might boost medical interventions for individuals who have minor disturbances in well-being because they are subjected to overdiagnosis and overtreatment (Beeker et al., 2021; CDC, 2016).

In addition, we found that employed older participants in our analysis aged 65 years and older, and workers with limitation had significantly increased odds of using outpatients' medical services when compared to the workers aged 18-24, and those without limitation respectively. Older individuals have a higher likelihood of having chronic diseases with related increased need and use of health care services. Also, previous research examining utilization and expenditure of health care services using MEPS data, reported a similar trend in the highest prevalence of the use of health care services in individuals aged 65 years and above (Karimi et al., 2021).

#### **3.4.1. Policy and Theoretical Implications**

This study finds evidence of the association between long working hours and the use of outpatient services. Long working hours could influence deleterious health outcomes and is expected to reflect in the increased utilization of health resources among employees. Lack of time to access health care services could significantly increase the use of the more expensive and unsustainable services with decreased time requirement such as emergency departments and related services. The results of this study highlight



the importance of further investigating the impact of long working hours on the U.S. labor force.

Despite a significant number of workers in the U.S. reporting long work hours, regulations concerning working time mainly concentrate on the transportation industry and are primarily based on rules rather than being based on performance or risk (Gärtner et al., 2019; Yang et al., 2006).

Also, the high prevalence of the use of medications with sedative properties have several implications. Previous research has reported a potential link between sleep aid medications and increased mortality (Kivimäki et al., 2015; Kripke et al., 2012; Skogstad et al., 2019). Moreover, chronic use of these medications may elevate the risk of cognitive and psychomotor impairments, workplace and car accidents, and addiction (Kim et al., 2016; Mizoue et al., 2001). Therefore, the residual effects of sleep aids and the associated risks present potential hazards to both the workers who use them and the general public.

Interventions aimed at improving the well-being of U.S. workers can be implemented through national legislations and programs, such as international labor standards on working time arrangements and limits (Pega et al., 2021). These interventions should prioritize preventing negative health effects caused by long working hours (Landsbergis, 2018; Messenger, 2018).

Employers can also benefit from initiatives aimed at improving employee health and providing flexible work hours. These measures may include health and wellness programs, screening for risk factors at work sites, and management of chronic illnesses. Such interventions have the potential to enhance work performance and productivity,

reduce workplace accidents caused by fatigue (Butler et al., 2009; Grzywacz et al., 2008), and lower medical expenses and absenteeism due to illness (Baicker et al., 2010; Merrill et al., 2011). Furthermore, reducing long working hours may improve employee health and well-being, leading to benefits for both the employer and the employee.

### **3.4.2. Limitations and Strengths**

Several limitations were identified in this study. Relying primarily on self-reported data on work hours may have introduced recall bias, although the short intervals between data collection rounds (4-5 months) may have helped to minimize this issue, which is common in population survey data (Kim, 2013). Also, excluding observations with missing covariates may have resulted in residual confounding and selection bias.

Despite these limitations, this study established a longitudinal association between long working hours and the use of outpatient services. The use of a representative sample of the working population allows for the generalization of our findings to the larger population. Also, the short duration of the survey may have helped minimize the likelihood of healthy worker effect bias, where healthier workers remain in the survey, creating a selection effect that biases the results (Angrave & Charlwood, 2015; Dembe & Yao, 2016; Kim, 2013).

### **3.5. CONCLUSION**

This study examined the relationship between working hours, health care utilization, and the use of medications with sedative side effects. Our findings suggest that working hours are significantly associated with the use of outpatient services, with the highest odds in workers that worked 61 or more hours per week.

Also, we found a similar pattern of associations between working hours and the use of outpatient services in employees using medications with sedative properties. However, we did not find any significant association between working hours and the use of other health care resources, including office-based visits, emergency department visits, hospital discharges, and dental visits.

Awareness of the risks associated with long working hours is crucial for both individuals and organizations, as it can result in negative health outcomes. To mitigate these risks, it is essential to implement policies that promote work-life balance and utilize interventions that help reduce work-related stress.

## CHAPTER FOUR: CONCLUSION

Long working hours have increased with the continuous attempts at meeting the needs of 24-hour service and globalized society. The increase in overtime hours has been associated with deleterious health outcomes such as cardiovascular diseases, symptoms of psychological distress, and health behaviors, including risky intake of alcohol and smoking.

Prior literature has established the relationship between long working hours and risky alcohol intake. Where workers working overtime consume copious amounts of alcohol in a bid to address the stress that results from the production of excessive stress hormones. Thus, the continuous prevalence and increase in long working hours among the U.S. working population are expected to influence the use of sedating medications, the incidence of chronic diseases, and subsequent reflection in healthcare use in the United States.

However, studies examining the influence of working hours on the use of sedating medications are lacking. This dissertation addressed crucial gaps in our current knowledge of the multi-dimensional relationship between working hours and health outcomes. The three-study approach advanced current knowledge of the interplay between long working hours, the use of sedating medications, and health service utilization.

The first study employed the Andersen healthcare utilization model to conceptualize the relationship between long working hours and the use of sleep aid medications, along with the use of medications with sedative properties. The study found that long working hours were significantly associated with increased use of sleep aids and medications with sedative properties. Also, among users of these medications, those working long hours had higher use when compared to those using and working fewer hours. Females had a higher likelihood of using sleep aids when compared to males. Some probable reasons provided for this association in females ranged from the higher predisposition to developing sleep disorder, extended work hour times arising from home chores, and peculiar physiological changes.

The positive association between overtime hours and the use of sedative medication raises implications in the known harmful effects of long-term use of sedating medication, including increased mortality and morbidity. Significant differences in the associations between long working hours and the use of sleep medications across socioeconomic groups. Individuals employed in professional services had the greatest probability of utilizing sleep medications. At the same time, respondents with large family sizes had a lower likelihood of using prescription sleep aids and medications with sedative properties, suggesting a possible protective factor from the family size.

The second study investigated the relationship between long working hours and psychological distress using a longitudinal approach. The study employed the WHO logic model (Pega et al., 2021; Rugulies et al., 2021) to operationalize this relationship. The model posits that overtime hours influence the onset of psychological distress through two causal pathways—psycho-physiological and health—behavioral pathways. This study

represents an improvement over previous research that relied on cross-sectional data and was plagued by problems of reverse causality. It examined whether full-time employees without psychological distress at the outset have a higher likelihood of experiencing psychological distress in the future if they work long hours.

The findings from this study suggest a relationship between working long hours and the onset of psychological distress. The highest risk was found in individuals who worked very long hours, and a risk increase was also observed among those who worked regular, standard hours per week when compared to those that worked fewer hours per week. Factors such as income, chronic disease status, and the use of sleep aids were identified as relevant contributing factors. Also, the study unveiled variations between genders regarding the likelihood of developing psychological distress in the workforce.

The third study of the dissertation evaluated longitudinal associations between working hours, health care utilization, and the use of sedative medications in the U.S. among a representative panel of full-time workers. This study employed the Andersen healthcare utilization model to conceptualize these various associations. Fixed effect estimators were utilized in modeling the relationship between working hours, the use of sedating medications, and health care services.

Contrary to one of the study's hypotheses, no significant association was found between working hours and the use of emergency department visits. However, the study findings suggest that working hours are significantly associated with the use of outpatient services, with the highest odds in workers that worked 61 or more hours per week. Further analyses revealed a more pronounced and similar pattern of associations between working hours and the use of outpatient services in respondents using sedating

medication. Outpatient medical services include alcohol and drug abuse clinics; well-baby clinics; obesity clinics; eye, ear, nose, and throat clinics; cardiology clinics; internal medicine departments; physical therapy clinics; and radiation therapy clinics.

Prior research posits that the most harmful effects of long working hours are most pronounced in those working high long working hours (Bannai et al., 2015). Thus, this high use of outpatient medical services in respondents suggests a poorer health status of individuals working very long hours. Outpatient medical services include alcohol and drug abuse clinics; well-baby clinics; obesity clinics; eye, ear, nose, and throat clinics; family planning clinics; cardiology clinics; internal medicine departments; physical therapy clinics; and radiation therapy clinics. This high use of outpatient medical services in respondents working very long hours may be due to the poor health status of individuals working long hours. Furthermore, older employees aged 65 years and over had the highest odds of using outpatient medical services. The high burden of chronic diseases in older individuals is a possible risk factor for high health care utilization.

Together, the findings from this dissertation offer a solid foundation of evidence and great insights into the associations between working hours, the use of sleep aids, and their negative impact on health outcomes. It provides a foundation of evidence that the deleterious effect of long working hours goes beyond risky alcohol consumption and smoking. It delineates the relationship between overtime hours, the use of sedating medications, the development of psychological distress, and the use of healthcare services.

Therefore, these findings can serve as a guiding tool for employers, health systems, and researchers in creating interventions and making recommendations

regarding work-based policies, with the ultimate aim of enhancing the health and well-being of employees. Despite a significant portion of Americans working extended hours, there is a lack of comprehensive regulations governing working time arrangements in the U.S. Knowledge of the risks associated with long working hours is crucial for both individuals and organizations, as it can result in negative health outcomes. Implementing policies that encourage work-life balance and aid interventions that decrease work-related stress may help in mitigating risks associated with long work hours.

#### **4.1. FUTURE RESEARCH**

This dissertation sets the stage for future follow-up and exploratory studies. The medications identified and used in these studies were confirmed prescribed medications and did not include over-the-counter (OTC) medications. Thus, future research will need to examine the relationship between working hours and the use of over-the-counter sedating medications. OTC medications are more accessible to individuals. Therefore, a higher prevalence of their use is expected.

This dissertation highlights the probable relations between long working hours, sleep disorders, and the use of sedating medications. Future research will build on this in further investigating the relationship between working hours, negative health outcomes of sedative use, and substance abuse disorders.

Also, in cognizance of external factors such as external stressful life events and workplace stressors having notable probabilities to influence the development of psychological distress individuals. Future studies will evaluate the relationship between working hours, external stressors, and the development of psychological distress.



Furthermore, future studies will need to examine the type of healthcare outpatient services that are most utilized on a granular level. Uncovering the specific types of outpatient services with the highest risk of use in relationship with long working hours will help proffer targeted interventions in addressing the healthcare needs of employees, especially those working overtime hours.

Figure 6. Conceptual framework depicting the association between working hours, the use of prescription medications with sedative side effects, and health care utilization (adapted from Aday & Andersen, 1974)




Table 7. Descriptive statistics of work hours and health care utilization.

| Variables                 | Overall Frequency | Overall Weighted % | Weekly Working Hours |       |       |       |      |
|---------------------------|-------------------|--------------------|----------------------|-------|-------|-------|------|
|                           |                   |                    | 30-35                | 36-40 | 41-50 | 51-60 | ≥ 61 |
| <b>Age (years)</b>        |                   |                    |                      |       |       |       |      |
| 18-26                     | 8,211             | 9.3                | 16.8                 | 9.9   | 6     | 5.6   | 5.6  |
| 27-64                     | 76,185            | 86.2               | 75.1                 | 85.9  | 90.6  | 90.3  | 88.8 |
| ≥ 65                      | 3,590             | 4.5                | 8.1                  | 4.2   | 3.4   | 4.1   | 5.5  |
| <b>Sex</b>                |                   |                    |                      |       |       |       |      |
| Male                      | 48,531            | 56.1               | 39.7                 | 53.4  | 62.4  | 71    | 76.7 |
| Female                    | 39,455            | 43.9               | 60.3                 | 46.6  | 37.6  | 29    | 23.3 |
| <b>Race</b>               |                   |                    |                      |       |       |       |      |
| White                     | 62,624            | 79.7               | 79.5                 | 77.3  | 84.5  | 84    | 81.4 |
| Black                     | 15,300            | 11.3               | 12.5                 | 12.8  | 8     | 9     | 8.9  |
| Other                     | 10,062            | 9                  | 8                    | 9.9   | 7.6   | 7     | 9.7  |
| <b>Marital Status</b>     |                   |                    |                      |       |       |       |      |
| Married                   | 50,230            | 59.5               | 50.1                 | 58.7  | 62.9  | 66.3  | 64.3 |
| Unmarried                 | 37,756            | 40.5               | 49.9                 | 41.3  | 37.1  | 33.7  | 35.7 |
| <b>Family Size</b>        |                   |                    |                      |       |       |       |      |
| ≤ 2 Members               | 37,349            | 48.7               | 49.5                 | 47    | 51.8  | 51.3  | 51.7 |
| > 2 Members               | 50,637            | 51.3               | 50.5                 | 53    | 48.2  | 48.7  | 48.3 |
| <b>Education</b>          |                   |                    |                      |       |       |       |      |
| Less than college         | 35,650            | 32.8               | 37.6                 | 35.1  | 26    | 27.6  | 34.3 |
| Some college or more      | 52,336            | 67.2               | 62.4                 | 64.9  | 74    | 72.4  | 65.7 |
| <b>Occupation</b>         |                   |                    |                      |       |       |       |      |
| Professional services     | 33,215            | 44.7               | 34.8                 | 41.6  | 54.2  | 53.3  | 48.7 |
| Non-professional services | 54,771            | 55.3               | 65.2                 | 58.4  | 45.8  | 46.7  | 51.3 |
| <b>Year</b>               |                   |                    |                      |       |       |       |      |
| 2010                      | 4,454             | 5.1                | 5                    | 5.2   | 4.7   | 5.3   | 5.2  |
| 2011                      | 9,972             | 10.7               | 11.2                 | 10.9  | 10    | 10.5  | 12.4 |
| 2012                      | 10,582            | 10.4               | 10.4                 | 10.7  | 9.8   | 9.9   | 10.5 |
| 2013                      | 9,809             | 10.6               | 10.6                 | 11    | 10    | 10.2  | 9    |
| 2014                      | 9,768             | 11                 | 11.9                 | 10.9  | 10.7  | 10.9  | 12.2 |
| 2015                      | 10,382            | 11.3               | 11.4                 | 11    | 11.8  | 12    | 12   |
| 2016                      | 10,245            | 11.7               | 10.6                 | 11.7  | 12.1  | 11.9  | 11.5 |
| 2017                      | 9,437             | 11.7               | 11.6                 | 11.5  | 12.2  | 11.9  | 10.9 |
| 2018                      | 8,944             | 11.5               | 11.6                 | 11.2  | 12.4  | 11.7  | 10.7 |
| 2019                      | 4,393             | 5.9                | 5.6                  | 5.9   | 6.3   | 5.6   | 5.6  |

(continued)

Table 7: (Continued).

| Variables   | Overall Frequency | Overall Weighted % | Weekly Working Hours |       |       |       |      |
|---|-------------------|--------------------|----------------------|-------|-------|-------|------|
|   |                   |                    | 30-35                | 36-40 | 41-50 | 51-60 | ≥ 61 |
| <b>Region</b>   |                   |                    |                      |       |       |       |      |
| Northeast   | 13,549            | 17.5               | 20.0                 | 17.2  | 17.3  | 18.1  | 14.8 |
| North Central, Midwest                                  | 17,677            | 21.9               | 20.5                 | 20.3  | 25.5  | 24.4  | 24.8 |
| South   | 33,182            | 37.5               | 34.5                 | 38.5  | 35.9  | 37.6  | 40.6 |
| West  | 23,578            | 23.1               | 24.9                 | 24.0  | 21.2  | 19.9  | 19.8 |
| <b>Income</b>   |                   |                    |                      |       |       |       |      |
| Poor to low income                                      | 19,887            | 14.9               | 30.0                 | 15.9  | 8.0   | 8.5   | 10.0 |
| Middle to high income                                   | 68,099            | 85.1               | 70.0                 | 84.1  | 92.0  | 91.5  | 90.0 |
| <b>Receipt of food stamp</b>                            |                   |                    |                      |       |       |       |      |
| No  | 81,139            | 95.3               | 90.6                 | 94.8  | 97.6  | 97.6  | 97.2 |
| Yes   | 6,847             | 4.7                | 9.4                  | 5.2   | 2.4   | 2.4   | 2.8  |
| <b>Limitation</b>                                       |                   |                    |                      |       |       |       |      |
| No  | 78,359            | 88.3               | 85.8                 | 88.6  | 88.9  | 88.1  | 87.2 |
| Yes   | 9,627             | 11.7               | 14.2                 | 11.4  | 11.1  | 11.9  | 12.8 |
| <b>Medication with Sedative Effects Use<sup>†</sup></b> |                   |                    |                      |       |       |       |      |
| No  | 63,918            | 72.7               | 72.1                 | 73.3  | 70.2  | 73.4  | 75.3 |
| Yes   | 24,068            | 27.4               | 27.9                 | 26.7  | 29.8  | 26.6  | 24.7 |
| <b>Outpatient Visit</b>                                 |                   |                    |                      |       |       |       |      |
| No  | 76,675            | 85.1               | 84.7                 | 85.6  | 84.1  | 85.1  | 85.1 |
| ≥ 1 visit   | 11,311            | 14.9               | 15.3                 | 14.4  | 15.9  | 14.9  | 14.9 |
| <b>Office Based visit</b>                               |                   |                    |                      |       |       |       |      |
| No  | 30,768            | 29.1               | 30.0                 | 30.4  | 25.1  | 27.3  | 32.4 |
| ≥ 1 visit   | 57,218            | 70.9               | 70.0                 | 69.6  | 74.9  | 72.7  | 67.6 |
| <b>ER visit</b>   |                   |                    |                      |       |       |       |      |
| No  | 78,777            | 89.5               | 87.9                 | 89.4  | 90.2  | 90.6  | 89.2 |
| ≥ 1 visit   | 9,209             | 10.5               | 12.1                 | 10.6  | 9.8   | 9.4   | 10.8 |
| <b>Hospital discharges</b>                              |                   |                    |                      |       |       |       |      |
| No  | 84,200            | 95.3               | 94.8                 | 95.3  | 95.2  | 95.8  | 95.2 |
| ≥ 1 discharge   | 3,786             | 4.7                | 5.2                  | 4.7   | 4.8   | 4.2   | 4.8  |
| <b>Dental visit</b>                                     |                   |                    |                      |       |       |       |      |
| No  | 54,664            | 56.5               | 58.6                 | 58.1  | 51.0  | 55.9  | 59.5 |
| ≥ 1 visit   | 33,322            | 43.5               | 41.4                 | 41.9  | 49.0  | 44.1  | 40.5 |
| <b>Outpatient physician visit</b>                       |                   |                    |                      |       |       |       |      |
| No  | 87,403            | 99.4               | 99.0                 | 99.4  | 99.6  | 99.4  | 99.3 |
| ≥ 1 visit   | 583               | 0.6                | 1.0                  | 0.6   | 0.4   | 0.6   | 0.7  |

<sup>†</sup>= Unweighted frequency

Table 8. Association Between Working Hours and Health Care Utilization: Fixed Effects Regression Analysis – Overall Sample

| <b>Fixed Effect Logit Model – Overall sample</b> |                            |                            |                            |                            |                           |
|--|----------------------------|----------------------------|----------------------------|----------------------------|---------------------------|
| <b>Variables</b>                                 | <b>Outpatient</b>          | <b>Office Based</b>        | <b>Emergency Room</b>      | <b>Hospital Discharges</b> | <b>Dental</b>             |
| <b>Work hours/week (30-35 h/week - REF)</b>      |                            |                            |                            |                            |                           |
| 36-40 h/week                                     | 1.342**<br>(1.013, 1.779)  | 1.000<br>(0.824, 1.214)    | 0.945<br>(0.751, 1.190)    | 1.31<br>(0.904, 1.899)     | 1.032<br>(0.838, 1.272)   |
| 41-50 h/week                                     | 1.333<br>(0.944, 1.884)    | 1.019<br>(0.791, 1.313)    | 0.815<br>(0.601, 1.106)    | 1.091<br>(0.682, 1.746)    | 1.086<br>(0.836, 1.411)   |
| 51-55 h/week                                     | 1.448<br>(0.901, 2.328)    | 0.933<br>(0.673, 1.295)    | 0.876<br>(0.570, 1.346)    | 1.229<br>(0.598, 2.525)    | 0.818<br>(0.575, 1.166)   |
| ≥ 61 h/week                                      | 2.008**<br>(1.044, 3.861)  | 0.992<br>(0.655, 1.503)    | 0.886<br>(0.514, 1.525)    | 1.007<br>(0.405, 2.509)    | 1.011<br>(0.631, 1.618)   |
| <b>Age (18-26years - REF)</b>                    |                            |                            |                            |                            |                           |
| 27-64 years                                      | 1.234<br>(0.854, 1.783)    | 0.807*<br>(0.646, 1.010)   | 0.744*<br>(0.546, 1.014)   | 1.329<br>(0.822, 2.147)    | 0.913<br>(0.723, 1.154)   |
| ≥ 65 years                                       | 1.737**<br>(1.006, 2.998)  | 0.612**<br>(0.375, 0.999)  | 0.656<br>(0.365, 1.178)    | 2.098*<br>(0.879, 5.007)   | 0.763<br>(0.487, 1.193)   |
| <b>Marital Status (Married - REF)</b>            |                            |                            |                            |                            |                           |
| Unmarried  | 0.952<br>(0.689, 1.315)    | 1.184<br>(0.940, 1.490)    | 0.946<br>(0.721, 1.240)    | 0.679*<br>(0.438, 1.052)   | 0.966<br>(0.765, 1.220)   |
| <b>Family Size (≤ 2 Members - REF)</b>           |                            |                            |                            |                            |                           |
| > 2 Members                                      | 0.770***<br>(0.648, 0.915) | 0.758***<br>(0.661, 0.870) | 0.814**<br>(0.693, 0.956)  | 0.867<br>(0.707, 1.064)    | 0.917<br>(0.801, 1.050)   |
| <b>Occupation (Non-professional - REF)</b>       |                            |                            |                            |                            |                           |
| Professional services                            | 0.839<br>(0.631, 1.114)    | 0.893<br>(0.729, 1.095)    | 0.977<br>(0.761, 1.255)    | 0.744<br>(0.497, 1.114)    | 1.105<br>(0.900, 1.357)   |
| <b>Income (Poor to low income - REF)</b>         |                            |                            |                            |                            |                           |
| middle to high income                            | 0.903<br>(0.781, 1.043)    | 1.000<br>(0.907, 1.103)    | 1.016<br>(0.898, 1.148)    | 0.671***<br>(0.553, 0.815) | 1.116**<br>(1.003, 1.241) |
| <b>Receipt of food stamp (No - REF)</b>          |                            |                            |                            |                            |                           |
| Yes  | 1.078<br>(0.877, 1.326)    | 0.93<br>(0.801, 1.080)     | 0.999<br>(0.838, 1.190)    | 1.244*<br>(0.964, 1.607)   | 0.955<br>(0.807, 1.129)   |
| <b>Limitation (No - REF)</b>                     |                            |                            |                            |                            |                           |
| Yes  | 1.431***<br>(1.259, 1.627) | 1.531***<br>(1.329, 1.765) | 1.361***<br>(1.194, 1.551) | 1.755***<br>(1.456, 2.115) | 1.159**<br>(1.027, 1.309) |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Table 9. Association Between working Hours and Outpatient Healthcare Services – Overall Sample

| <b>Fixed Effects Regressions for Outpatient Healthcare Services – Overall Sample</b> |                               |                               |
|--|-------------------------------|-------------------------------|
| <b>Variables</b>   | <b>Marginal Effect</b>        | <b>Linear Model</b>           |
| <b>Work hours/week (30-35 h/week - REF)</b>  |                               |                               |
| 36-40 h/week   | 0.072**<br>(0.004, 0.141)     | 0.020**<br>(0.000, 0.040)     |
| 41-50 h/week   | 0.071*<br>(-0.013, 0.155)     | 0.018<br>(-0.008, 0.045)      |
| 51-55 h/week   | 0.091<br>(-0.024, 0.205)      | 0.024<br>(-0.010, 0.058)      |
| ≥ 61 h/week  | 0.166**<br>(0.020, 0.313)     | 0.046**<br>(0.001, 0.090)     |
| <b>Age (18-26years - REF)</b>  |                               |                               |
| 27-64 years  | 0.052<br>(-0.038, 0.141)      | 0.012<br>(-0.009, 0.033)      |
| ≥ 65 years   | 0.133**<br>(0.008, 0.258)     | 0.051*<br>(-0.000, 0.102)     |
| <b>Marital Status (Married - REF)</b>  |                               |                               |
| Unmarried  | -0.012<br>(-0.091, 0.067)     | -0.005<br>(-0.028, 0.017)     |
| <b>Family Size (≤ 2 Members - REF)</b>   |                               |                               |
| > 2 Members  | -0.064***<br>(-0.106, -0.021) | -0.023***<br>(-0.038, -0.009) |
| <b>Occupation (Non-professional - REF)</b>   |                               |                               |
| Professional services  | -0.043<br>(-0.112, 0.027)     | -0.012<br>(-0.033, 0.008)     |
| <b>Income (Poor to low income - REF)</b>   |                               |                               |
| middle to high income  | -0.025<br>(-0.060, 0.010)     | -0.007<br>(-0.016, 0.003)     |
| <b>Receipt of food stamp (No - REF)</b>  |                               |                               |
| Yes  | 0.018<br>(-0.031, 0.068)      | 0.007<br>(-0.009, 0.022)      |
| <b>Limitation (No - REF)</b>   |                               |                               |
| Yes  | 0.086***<br>(0.055, 0.116)    | 0.044***<br>(0.029, 0.060)    |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Figure 7. Use of outpatient services according to working hours from the fixed effects regression model of the overall sample.





Table 10. Association Between working Hours and Health care Utilization: Fixed effects regression analysis – Among Medication Users

| <b>Fixed Effect Logit Model – Among Medication Users</b> |                            |                              |                              |
|--|----------------------------|------------------------------|------------------------------|
| <b>Variables</b>   | <b>Logit Model</b>         | <b>Marginal Effect</b>       | <b>Linear Model</b>          |
| <b>Work hours/week (30-35 h/week - REF)</b>              |                            |                              |                              |
| 36-40 h/week   | 1.715***<br>(1.159, 2.539) | 0.127***<br>(0.039, 0.216)   | 0.069***<br>(0.020, 0.118)   |
| 41-50 h/week   | 1.518*<br>(0.949, 2.426)   | 0.099*<br>(-0.009, 0.208)    | 0.051<br>(-0.014, 0.115)     |
| 51-55 h/week   | 1.881*<br>(0.909, 3.892)   | 0.148*<br>(-0.012, 0.307)    | 0.076<br>(-0.016, 0.169)     |
| ≥ 61 h/week  | 1.975<br>(0.783, 4.983)    | 0.158<br>(-0.042, 0.358)     | 0.081<br>(-0.042, 0.204)     |
| <b>Age (18-26years - REF)</b>                            |                            |                              |                              |
| 27-64 years  | 1.403<br>(0.796, 2.475)    | 0.08<br>(-0.050, 0.211)      | 0.036<br>(-0.024, 0.096)     |
| ≥ 65 years   | 2.833**<br>(1.198, 6.698)  | 0.225***<br>(0.069, 0.380)   | 0.146**<br>(0.032, 0.260)    |
| <b>Marital Status (Married - REF)</b>                    |                            |                              |                              |
| Unmarried  | 0.715<br>(0.451, 1.135)    | -0.077<br>(-0.186, 0.033)    | -0.041<br>(-0.091, 0.009)    |
| <b>Family Size (≤ 2 Members REF)</b>                     |                            |                              |                              |
| > 2 Members  | 0.784**<br>(0.616, 0.998)  | -0.055*<br>(-0.113, 0.002)   | -0.036**<br>(-0.070, -0.003) |
| <b>Occupation (Non-professional - REF)</b>               |                            |                              |                              |
| Professional services                                    | 1.022<br>(0.682, 1.530)    | 0.005<br>(-0.087, 0.096)     | 0.002<br>(-0.048, 0.053)     |
| <b>Income (Poor to low income - REF)</b>                 |                            |                              |                              |
| middle to high income                                    | 0.796**<br>(0.647, 0.980)  | -0.051**<br>(-0.099, -0.002) | -0.029**<br>(-0.056, -0.003) |
| <b>Receipt of food stamp (No - REF)</b>                  |                            |                              |                              |
| Yes  | 1.238<br>(0.935, 1.641)    | 0.047<br>(-0.014, 0.109)     | 0.03<br>(-0.010, 0.069)      |
| <b>Limitation (No - REF)</b>                             |                            |                              |                              |
| Yes  | 1.607***<br>(1.362, 1.897) | 0.105***<br>(0.065, 0.146)   | 0.082***<br>(0.054, 0.109)   |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference



Figure 8. Use of outpatient services according to working hours from the fixed effects regression model among users of medications with sedative side effects.



## REFERENCES

- Abuse, S. (2014). Treatment episode data set (TEDS): 2002-2012. National admissions to substance abuse treatment services. In *BHSIS Series S-71, HHS Publication No.(SMA) 14-4850*. Substance Abuse and Mental Health Services Administration Rockville, MD.
- Afonso, P., Fonseca, M., & Pires, J. (2017). Impact of working hours on sleep and mental health. *Occupational Medicine*, *67*(5), 377-382.
- Agarwal, S. D., & Landon, B. E. (2019). Patterns in outpatient benzodiazepine prescribing in the United States. *JAMA Network Open*, *2*(1), e187399-e187399.
- Ahn, S. (2018). Working hours and depressive symptoms over 7 years: evidence from a Korean panel study. *International Archives of Occupational and Environmental Health*, *91*(3), 273-283.
- Allison, P. D. (2009). *Fixed effects regression models*. SAGE publications.
- AMBREEN, S., IQBAL, Z., IQBAL, M., & AHMAD, S. (2016). Determinants of antenatal psychological distress in Pakistani women. *Nöro Psikiyatri Arşivi*, *53*(2), 152.
- American Psychiatric Association (2019). What are sleep disorders? Diagnostic and Statistical Manual of Mental Disorders.

- Andersen, R., & Newman, J. F. (1973). Societal and individual determinants of medical care utilization in the United States. *The Milbank Memorial Fund Quarterly. Health and Society*, 95-124.
- Andersen, R. M. (1995). Revisiting the behavioral model and access to medical care: does it matter? *Journal of health and social behavior*, 1-10.
- Angarita, G. A., Emadi, N., Hodges, S., & Morgan, P. T. (2016). Sleep abnormalities associated with alcohol, cannabis, cocaine, and opiate use: a comprehensive review. *Addiction science & clinical practice*, 11(1), 1-17.
- Angrave, D., & Charlwood, A. (2015). What is the relationship between long working hours, over-employment, under-employment and the subjective well-being of workers? Longitudinal evidence from the UK. *Human Relations*, 68(9), 1491-1515.
- Artazcoz, L., Cortès, I., Escribà-Agüir, V., Cascant, L., & Villegas, R. (2009). Understanding the relationship of long working hours with health status and health-related behaviours. *Journal of Epidemiology & Community Health*, 63(7), 521-527.
- Bachhuber, M. A., Hennessy, S., Cunningham, C. O., & Starrels, J. L. (2016). Increasing benzodiazepine prescriptions and overdose mortality in the United States, 1996–2013. *American Journal of Public Health*, 106(4), 686-688.
- Baicker, K., Cutler, D., & Song, Z. (2010). Workplace wellness programs can generate savings. *Health Affairs*, 29(2), 304-311.
- Bairoliya, N., & McKiernan, K. (2021). Revisiting retirement and social security claiming decisions. Available at SSRN 3896031.

- Bannai, A., & Tamakoshi, A. (2014). The association between long working hours and health: a systematic review of epidemiological evidence. *Scandinavian journal of work, environment & health*, 5-18.
- Bannai, A., Ukawa, S., & Tamakoshi, A. (2015). Long working hours and psychological distress among school teachers in Japan. *Journal of occupational health*, 57(1), 20-27.
- Beeker, T., Mills, C., Bhugra, D., te Meerman, S., Thoma, S., Heinze, M., & von Peter, S. (2021). Psychiatrization of society: a conceptual framework and call for transdisciplinary research. *Frontiers in Psychiatry*, 12, 645556.
- Bergs, Y., Hoofs, H., Kant, I., Slangen, J. J., & Jansen, N. W. (2018). Work–family conflict and depressive complaints among Dutch employees: examining reciprocal associations in a longitudinal study. *Scandinavian journal of work, environment & health*, 69-79.
- Bertisch, S. M., Herzig, S. J., Winkelman, J. W., & Buettner, C. (2014). National use of prescription medications for insomnia: NHANES 1999-2010. *Sleep*, 37(2), 343-349.
- Bhattarai, G. R., Duffy, P. A., & Raymond, J. (2005). Use of food pantries and food stamps in low-income households in the United States. *Journal of Consumer Affairs*, 39(2), 276-298.
- Bianchi, S. M. (2000). Maternal employment and time with children: Dramatic change or surprising continuity? *Demography*, 37(4), 401-414.

- Bischoff, R. J., Springer, P. R., & Taylor, N. (2017). Global mental health in action: Reducing disparities one community at a time. *Journal of Marital and Family Therapy, 43*(2), 276-290.
- Blewett, L. A., Drew, J. A. R., Griffin, R., & Williams, K. C. . (2020). IPUMS Health Surveys: Medical Expenditure Panel Survey. *IPUMS*.
- Bonde, J. P. E. (2008). Psychosocial factors at work and risk of depression: a systematic review of the epidemiological evidence. *Occupational and environmental medicine, 65*(7), 438-445.
- Bush, P. W., Drake, R. E., Xie, H., McHugo, G. J., & Haslett, W. R. (2009). The long-term impact of employment on mental health service use and costs for persons with severe mental illness. *Psychiatric Services, 60*(8), 1024-1031.
- Butler, A. B., Grzywacz, J. G., Ettner, S. L., & Liu, B. (2009). Workplace flexibility, self-reported health, and health care utilization. *Work & Stress, 23*(1), 45-59.
- Cappuccio FP, D'Elia L, Strazzullo P, et al. Sleep duration and all-cause mortality: a systematic review and meta-analysis of prospective studies. *Sleep* 2010;33:585–92.
- Cameron, A. C., & Trivedi, P. K. (2010). *Microeconometrics using stata* (Vol. 2). Stata press College Station, TX.
- Campbell, M. K., Tessaro, I., DeVellis, B., Benedict, S., Kelsey, K., Belton, L., & Sanhueza, A. (2002). Effects of a tailored health promotion program for female blue-collar workers: health works for women. *Prev Med, 34*(3), 313-323.
- Carroll, W. A., & Miller, G. E. (2019). STATISTICAL BRIEF# 511 Differences in Health Insurance Coverage between Part-Time and Full-Time Private-Sector

Workers, 2005 and 2015. *Medical Expenditure Panel Survey, Agency for Healthcare Research and Quality, accessed March, 29.*

CDC, U. (2016). Selected prescription drug classes used in the past 30 days, by sex and age: United States, selected years 1988–1994 through 2011–2014. *Health, United States, 2015-Individual Charts and Tables.*

CDC. (2018). Sleep and Sleep Disorders. Retrieved from <https://www.cdc.gov/sleep/index.html>

Chandola, T., Ferrie, J. E., Perski, A., Akbaraly, T., & Marmot, M. G. (2010). The effect of short sleep duration on coronary heart disease risk is greatest among those with sleep disturbance: a prospective study from the Whitehall II cohort. *Sleep, 33*(6), 739-744.

Cheatle, M. D., & Webster, L. R. (2015). Opioid therapy and sleep disorders: risks and mitigation strategies. *Pain Medicine, 16*(suppl\_1), S22-S26.

Chen, E. M., Kombo, N., Teng, C. C., Mruthyunjaya, P., Nwanyanwu, K., & Parikh, R. (2020). Ophthalmic medication expenditures and out-of-pocket spending: an analysis of United States prescriptions from 2007 through 2016. *Ophthalmology, 127*(10), 1292-1302.

Cheng, C. M., Guglielmo, B. J., Maselli, J., & Auerbach, A. D. (2010). Coverage of FDA medication boxed warnings in commonly used drug information resources. *Archives of internal medicine, 170*(9), 831-833.

Cheng, Y., Du, C.-L., Hwang, J.-J., Chen, I.-S., Chen, M.-F., & Su, T.-C. (2014). Working hours, sleep duration and the risk of acute coronary heart disease: a

- case-control study of middle-aged men in Taiwan. *International journal of cardiology*, 171(3), 419-422.
- Chong, Y., Fryar, C. D., & Gu, Q. (2013). *Prescription sleep aid use among adults: United States, 2005-2010*. US Department of Health and Human Services, Centers for Disease Control and ....
- Cirignotta, F., Mondini, S., Zucconi, M., Lenzi, P. L., & Lugaresi, E. (1985). Insomnia: an epidemiological survey. *Clin Neuropharmacol*, 8(Suppl 1), S49-S54.
- Citrome, L. (2017). Activating and sedating adverse effects of second-generation antipsychotics in the treatment of schizophrenia and major depressive disorder: absolute risk increase and number needed to harm. *Journal of clinical psychopharmacology*, 37(2), 138-147.
- Clauson, K. A., Marsh, W. A., Polen, H. H., Seamon, M. J., & Ortiz, B. I. (2007). Clinical decision support tools: analysis of online drug information databases. *Bmc Medical Informatics and Decision Making*, 7(1), 1-7.
- Collaborators, G. M. D. (2022). Global, regional, and national burden of 12 mental disorders in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet Psychiatry*, 9(2), 137-150.
- Comparisons eAnswers., F. (2021). *ADR (Adverse Drug Reaction) identification tool*. <http://online.factsandcomparisons.com>
- Croteau, S. E., Cook, K., Sheikh, L., Chawla, A., Sammon, J., Solari, P., Kim, B., Hinds, D., & Thornburg, C. D. (2021). Health care resource utilization and costs among adult patients with hemophilia A on factor VIII prophylaxis: an administrative claims analysis. *Journal of managed care & specialty pharmacy*, 27(3), 316-326.

- De Raeve, L., Kant, I., Jansen, N. W., Vasse, R. M., & Van den Brandt, P. A. (2009). Changes in mental health as a predictor of changes in working time arrangements and occupational mobility: results from a prospective cohort study. *Journal of psychosomatic research*, 66(2), 137-145.
- Dembe, A. E., & Yao, X. (2016). Chronic disease risks from exposure to long-hour work schedules over a 32-year period. *Journal of Occupational and Environmental Medicine*, 58(9), 861-867.
- Denton, M., Prus, S., & Walters, V. (2004). Gender differences in health: a Canadian study of the psychosocial, structural and behavioural determinants of health. *Social science & medicine*, 58(12), 2585-2600.
- Drukker, D. M. (2002). Bootstrapping a conditional moments test for normality after tobit estimation. *The Stata Journal*, 2(2), 125-139.
- Easton, S. D., Safadi, N. S., Wang, Y., & Hasson, R. G. (2017). The Kessler psychological distress scale: translation and validation of an Arabic version. *Health and quality of life outcomes*, 15(1), 1-7.
- Eugene, A. R. (2020). Association of sleep among 30 antidepressants: A population-wide adverse drug reaction study, 2004–2019. *PeerJ*, 8, e8748.
- Fell, D. B., Kephart, G., Curtis, L. J., Bower, K., Muhajarine, N., Reid, R., & Roos, L. (2007). The relationship between work hours and utilization of general practitioners in four Canadian provinces. *Health Services Research*, 42(4), 1483-1498.
- Foster, G., & Kalenkoski, C. M. (2013). Tobit or OLS? An empirical evaluation under different diary window lengths. *Applied Economics*, 45(20), 2994-3010.



- Fukuoka, Y., Takeshima, M., Ishii, N., Chikako, M., Makaya, M., Groah, L., Kyriakidis, E., & Dracup, K. (2010). An initial analysis: working hours and delay in seeking care during acute coronary events. *The American journal of emergency medicine*, 28(6), 734-740.
- Gärtner, J., Rosa, R. R., Roach, G., Kubo, T., & Takahashi, M. (2019). Working Time Society consensus statements: Regulatory approaches to reduce risks associated with shift work—a global comparison. *Industrial health*, 57(2), 245-263.
- Gerson, K., & Jacobs, J. A. (2004). The time divide: Work, family, and gender inequality.
- Gmel, G., & Rehm, J. (2004). Measuring alcohol consumption. *Contemporary drug problems*, 31(3), 467-540.
- Grinde, B., & Tambs, K. (2016). Effect of household size on mental problems in children: results from the Norwegian Mother and Child Cohort study. *BMC psychology*, 4, 1-11.
- Grizzle, A. J., Horn, J., Collins, C., Schneider, J., Malone, D. C., Stottlemeyer, B., & Boyce, R. D. (2019). Identifying common methods used by drug interaction experts for finding evidence about potential drug-drug interactions: web-based survey. *Journal of Medical Internet Research*, 21(1), e11182.
- Grosch, J. W., Caruso, C. C., Rosa, R. R., & Sauter, S. L. (2006). Long hours of work in the US: associations with demographic and organizational characteristics, psychosocial working conditions, and health. *American journal of industrial medicine*, 49(11), 943-952.

- Grzywacz, J. G., Carlson, D. S., & Shulkin, S. (2008). Schedule flexibility and stress: Linking formal flexible arrangements and perceived flexibility to employee health. *Community, Work and Family, 11*(2), 199-214.
- Gunasekara, F. I., Richardson, K., Carter, K., & Blakely, T. (2014). Fixed effects analysis of repeated measures data. *International journal of epidemiology, 43*(1), 264-269.
- Hägg, S., Jönsson, A. K., & Ahlner, J. (2020). Current evidence on abuse and misuse of gabapentinoids. *Drug safety, 43*, 1235-1254.
- Hale, L., Troxel, W., & Buysse, D. J. (2020). Sleep Health: An Opportunity for Public Health to Address Health Equity. *Annual Review of Public Health, 41*, 81-99.
- Hanrahan, C. T., & Cole, S. W. (2014). Assessment of drug information resource preferences of pharmacy students and faculty. *Journal of the Medical Library Association: JMLA, 102*(2), 117.
- Hino, A., Inoue, A., Kawakami, N., Tsuno, K., Tomioka, K., Nakanishi, M., Mafune, K., & Hiro, H. (2015). Buffering effects of job resources on the association of overtime work hours with psychological distress in Japanese white-collar workers. *International Archives of Occupational and Environmental Health, 88*(5), 631-640.
- Institute of Medicine. (2006). Sleep disorders and sleep deprivation: an unmet public health problem. Washington (DC): National Academies Press.
- Jackevicius, C. A., Lash, D., Singh, D., Hines, K., & Hata, M. (2019). Accuracy and usability of medication identifiers for solid oral medications. *Journal of the American College of Clinical Pharmacy, 2*(2), 98-107.

- Jaussent, I., Ancelin, M.-L., Berr, C., Pérès, K., Scali, J., Besset, A., Ritchie, K., & Dauvilliers, Y. (2013). Hypnotics and mortality in an elderly general population: a 12-year prospective study. *Bmc Medicine*, *11*(1), 1-11.
- Jehle, J., Ficker, E., Wan, X., Deschenes, I., Kisselbach, J., Wiedmann, F., Staudacher, I., Schmidt, C., Schweizer, P., & Becker, R. (2013). Mechanisms of zolpidem-induced long QT syndrome: acute inhibition of recombinant hERG K<sup>+</sup> channels and action potential prolongation in human cardiomyocytes derived from induced pluripotent stem cells. *British Journal of Pharmacology*, *168*(5), 1215-1229.
- Jindra, C., Li, C., Tsang, R. S., Bauermeister, S., & Gallacher, J. (2022). Depression and memory function—evidence from cross-lagged panel models with unit fixed effects in ELSA and HRS. *Psychological medicine*, *52*(8), 1428-1436.
- Kabir, U. Y. (2021). *Usual Source of Care Among Patients with Psychological Distress and Its Effects on Healthcare Expenditures, Utilization, and Patient Satisfaction* [The University of Memphis].
- Karimi, S. M., Eslami, K., & Johnson, C. E. (2021). US Health Care Expenditure Across Income Groups by Age and Type of Care. *Journal of Health Care for the Poor and Underserved*, *32*(2), 958-970.
- Kessler, R. C., Aguilar-Gaxiola, S., Alonso, J., Chatterji, S., Lee, S., Ormel, J., Üstün, T. B., & Wang, P. S. (2009). The global burden of mental disorders: an update from the WHO World Mental Health (WMH) surveys. *Epidemiology and Psychiatric Sciences*, *18*(1), 23-33.
- Kessler, R. C., Barker, P. R., Colpe, L. J., Epstein, J. F., Gfroerer, J. C., Hiripi, E., Howes, M. J., Normand, S.-L. T., Manderscheid, R. W., & Walters, E. E. (2003).

Screening for serious mental illness in the general population. *Archives of general psychiatry*, 60(2), 184-189.

Kessler, R. C., Berglund, P., Demler, O., Jin, R., Koretz, D., Merikangas, K. R., Rush, A. J., Walters, E. E., & Wang, P. S. (2003). The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *Jama*, 289(23), 3095-3105.

Kim, J. (2013). Depression as a psychosocial consequence of occupational injury in the US working population: findings from the medical expenditure panel survey. *BMC Public Health*, 13(1), 1-10.

Kim, W., Park, E.-C., Lee, T.-H., & Kim, T. H. (2016). Effect of working hours and precarious employment on depressive symptoms in South Korean employees: a longitudinal study. *Occupational and environmental medicine*, 73(12), 816-822.

Kim, Y.-H., Kim, H.-B., Kim, D.-H., Kim, J.-Y., & Shin, H.-Y. (2018). Use of hypnotics and the risk of or mortality from heart disease: a meta-analysis of observational studies. *The Korean Journal of Internal Medicine*, 33(4), 727.

Kivimäki, M., Jokela, M., Nyberg, S. T., Singh-Manoux, A., Fransson, E. I., Alfredsson, L., Bjorner, J. B., Borritz, M., Burr, H., & Casini, A. (2015). Long working hours and risk of coronary heart disease and stroke: a systematic review and meta-analysis of published and unpublished data for 603 838 individuals. *The lancet*, 386(10005), 1739-1746.

Kivimäki, M., Nyberg, S. T., Batty, G. D., Fransson, E. I., Heikkilä, K., Alfredsson, L., Bjorner, J. B., Borritz, M., Burr, H., & Casini, A. (2012). Job strain as a risk

- factor for coronary heart disease: a collaborative meta-analysis of individual participant data. *The lancet*, 380(9852), 1491-1497.
- Kivimäki, M., & Steptoe, A. (2018). Effects of stress on the development and progression of cardiovascular disease. *Nature Reviews Cardiology*, 15(4), 215-229.
- Koltai, J., Bierman, A., & Schieman, S. (2018). Financial circumstances, mastery, and mental health: Taking unobserved time-stable influences into account. *Social science & medicine*, 202, 108-116.
- Kripke, D. F., Langer, R. D., & Kline, L. E. (2012). Hypnotics' association with mortality or cancer: a matched cohort study. *Bmj Open*, 2(1), e000850.
- Kronfeld-Schor, N., & Einat, H. (2012). Circadian rhythms and depression: human psychopathology and animal models. *Neuropharmacology*, 62(1), 101-114.
- Lallukka, T., Lahelma, E., Rahkonen, O., Roos, E., Laaksonen, E., Martikainen, P., Head, J., Brunner, E., Mosdol, A., & Marmot, M. (2008). Associations of job strain and working overtime with adverse health behaviors and obesity: evidence from the Whitehall II Study, Helsinki Health Study, and the Japanese Civil Servants Study. *Social science & medicine*, 66(8), 1681-1698.
- Landsbergis, P. A. (2018). Interventions to reduce job stress and improve work organization and worker health. In *Unhealthy work* (pp. 193-209). Routledge.
- Laposky, A. D., Van Cauter, E., & Diez-Roux, A. V. (2016). Reducing health disparities: the role of sleep deficiency and sleep disorders. *Sleep medicine*, 18, 3-6.
- Lee, K., Suh, C., Kim, J.-E., & Park, J. O. (2017). The impact of long working hours on psychosocial stress response among white-collar workers. *Industrial health*, 55(1), 46-53.

- Lehne, G., Zeeb, H., Pischke, C. R., Mikolajczyk, R., Bewick, B. M., McAlaney, J., Dempsey, R. C., Van Hal, G., Stock, C., & Akvardar, Y. (2018). Personal and perceived peer use and attitudes towards use of non-prescribed prescription sedatives and sleeping pills among university students in seven European countries. *Addictive Behaviors, 87*, 17-23.
- Li, J., Pega, F., Ujita, Y., Brisson, C., Clays, E., Descatha, A., Ferrario, M. M., Godderis, L., Iavicoli, S., & Landsbergis, P. A. (2020). The effect of exposure to long working hours on ischaemic heart disease: A systematic review and meta-analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environment international, 142*, 105739.
- Li, Y., Cheng, F., Ye, R., Zhu, Y., Tian, L., Zheng, H., & Jia, C. (2019). Weekly Working Hours and Mental Health Status in Different Occupations: Evidence From the 2015 to 2016 California Health Interview Survey. *Journal of Occupational and Environmental Medicine, 61*(11), e452-e458.
- Liu, Y., Wheaton, A. G., Chapman, D. P., & Croft, J. B. (2013). Sleep duration and chronic diseases among US adults age 45 years and older: evidence from the 2010 Behavioral Risk Factor Surveillance System. *Sleep, 36*(10), 1421-1427.
- Liu, Y., Wheaton, A. G., Chapman, D. P., Cunningham, T. J., Lu, H., & Croft, J. B. (2016). Prevalence of healthy sleep duration among adults—United States, 2014. *Morbidity and Mortality Weekly Report, 65*(6), 137-141.
- Lopez-Quintero, C., Warren, T., Falise, A., Sharma, V., Bares, C., & Oshri, A. (2021). Prevalence and drug use correlates of extra-medical use of prescription medications for sleep among adults in the United States: Results of the 2015–

- 2018 National Survey on Drug Use and Health. *Pharmacology Biochemistry and Behavior*, 204, 173169.
- Lunde, L.-K., Koch, M., Veiersted, K. B., Moen, G.-H., Wærsted, M., & Knardahl, S. (2016). Heavy physical work: Cardiovascular load in male construction workers. *International journal of environmental research and public health*, 13(4), 356.
- Mahfoud, Y., Talih, F., Stroom, D., & Budur, K. (2009). Sleep disorders in substance abusers: how common are they? *Psychiatry (Edgmont)*, 6(9), 38.
- Mailliet, F., Galloux, P., & Poisson, D. (2001). Comparative effects of melatonin, zolpidem and diazepam on sleep, body temperature, blood pressure and heart rate measured by radiotelemetry in Wistar rats. *Psychopharmacology*, 156(4), 417-426.
- Mallon, L., Broman, J.-E., & Hetta, J. (2009). Is usage of hypnotics associated with mortality? *Sleep medicine*, 10(3), 279-286.
- Martínez, M. P., Miró, E., Sánchez, A. I., Diaz-Piedra, C., Cáliz, R., Vlaeyen, J. W., & Buela-Casal, G. (2014). Cognitive-behavioral therapy for insomnia and sleep hygiene in fibromyalgia: a randomized controlled trial. *Journal of behavioral medicine*, 37(4), 683-697.
- McGinty, E. E., Presskreischer, R., Han, H., & Barry, C. L. (2020). Psychological distress and loneliness reported by US adults in 2018 and April 2020. *Jama*, 324(1), 93-94.
- Merrill, R. M., Aldana, S. G., Garrett, J., & Ross, C. (2011). Effectiveness of a workplace wellness program for maintaining health and promoting healthy behaviors. *Journal of Occupational and Environmental Medicine*, 782-787.

- Messenger, J. (2018). Working time and the future of work. *ILO future of work research paper series*, 6(8), 33-37.
- Mizoue, T., Reijula, K., & Andersson, K. (2001). Environmental tobacco smoke exposure and overtime work as risk factors for sick building syndrome in Japan. *American journal of epidemiology*, 154(9), 803-808.
- Mousteri, V., Daly, M., & Delaney, L. (2020). Underemployment and psychological distress: Propensity score and fixed effects estimates from two large UK samples. *Social science & medicine*, 244, 112641.
- Murray, C. J., Ezzati, M., Lopez, A. D., Rodgers, A., & Vander Hoorn, S. (2003). Comparative quantification of health risks: conceptual framework and methodological issues. *Population health metrics*, 1(1), 1-20.
- Nakashima, M., Morikawa, Y., Sakurai, M., Nakamura, K., Miura, K., Ishizaki, M., Kido, T., Naruse, Y., Suwazono, Y., & Nakagawa, H. (2011). Association between long working hours and sleep problems in white-collar workers. *Journal of sleep research*, 20(1pt1), 110-116.
- Neubauer, D. N., Elmore, J. G., & Benca, R. (2021). Pharmacotherapy for insomnia in adults. In: Visit Source.
- Ohayon, M. M., Caulet, M., Priest, R. G., & Guilleminault, C. (1998). Psychotropic medication consumption patterns in the UK general population. *Journal of Clinical Epidemiology*, 51(3), 273-283.
- Pachito, D. V., Pega, F., Bakusic, J., Boonen, E., Clays, E., Descatha, A., Delvaux, E., De Bacquer, D., Koskenvuo, K., & Kröger, H. (2021). The effect of exposure to long working hours on alcohol consumption, risky drinking and alcohol use disorder:



- A systematic review and meta-analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environment international*, 146, 106205.
- Patrick, J., & Yang, P. Q. (2021). Health Insurance Coverage before and after the Affordable Care Act in the USA. *Sci*, 3(2), 25.
- Paul, K. I., & Batinic, B. (2010). The need for work: Jahoda's latent functions of employment in a representative sample of the German population. *Journal of Organizational Behavior*, 31(1), 45-64.
- Pega, F., Náfrádi, B., Momen, N. C., Ujita, Y., Streicher, K. N., Prüss-Üstün, A. M., Group, T. A., Descatha, A., Driscoll, T., & Fischer, F. M. (2021). Global, regional, and national burdens of ischemic heart disease and stroke attributable to exposure to long working hours for 194 countries, 2000–2016: A systematic analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environment international*, 154, 106595.
- Petrongolo, B. (2004). Gender segregation in employment contracts. *Journal of the European Economic Association*, 2(2-3), 331-345.
- Phillips, B., & Mannino, D. M. (2005). Does insomnia kill? *Sleep*, 28(8), 965-971.
- Quality, A. f. H. R. a. (2020). Medical Expenditure Panel Survey, MEPS Survey Questionnaires, Household Questionnaire Sections. *Dept of Health and Human Services*.
- Rehfuess, E. A., Booth, A., Brereton, L., Burns, J., Gerhardus, A., Mozygemba, K., Oortwijn, W., Pfadenhauer, L. M., Tummers, M., & van der Wilt, G. J. (2018). Towards a taxonomy of logic models in systematic reviews and health technology

- assessments: a priori, staged, and iterative approaches. *Research synthesis methods*, 9(1), 13-24.
- Rivera, A. S., Akanbi, M., O'Dwyer, L. C., & McHugh, M. (2020). Shift work and long work hours and their association with chronic health conditions: a systematic review of systematic reviews with meta-analyses. *Plos One*, 15(4), e0231037.
- Roane, B. M., & Taylor, D. J. (2008). Adolescent insomnia as a risk factor for early adult depression and substance abuse. *Sleep*, 31(10), 1351-1356.
- Roblek, T., Vaupotic, T., Mrhar, A., & Lainscak, M. (2015). Drug-drug interaction software in clinical practice: a systematic review. *European journal of clinical pharmacology*, 71(2), 131-142.
- Romani, M., & Ashkar, K. (2014). Burnout among physicians. *Libyan Journal of Medicine*, 9(1).
- Rugulies, R., Ando, E., Ayuso-Mateos, J. L., Bonafede, M., Cabello, M., Di Tecco, C., Dragano, N., Durand-Moreau, Q., Eguchi, H., & Gao, J. (2019). WHO/ILO work-related burden of disease and injury: Protocol for systematic reviews of exposure to long working hours and of the effect of exposure to long working hours on depression. *Environment international*, 125, 515-528.
- Rugulies, R., Sørensen, K., Di Tecco, C., Bonafede, M., Rondinone, B. M., Ahn, S., Ando, E., Ayuso-Mateos, J. L., Cabello, M., & Descatha, A. (2021). The effect of exposure to long working hours on depression: A systematic review and meta-analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. *Environment international*, 155, 106629.

- Rumble, R., & Morgan, K. (1992). Hypnotics, sleep, and mortality in elderly people. *Journal of the American Geriatrics Society*, 40(8), 787-791.
- Sato, K., Yamazaki, S., Hayashino, Y., Takegami, M., Tokuda, Y., Takahashi, O., Shimbo, T., Hinohara, S., Fukui, T., & Fukuhara, S. (2011). Associations between Hours Worked, Symptoms and Health Resource Utilization among Full-time Male Japanese Workers. *Journal of occupational health*, 53(3), 197-204.
- Sayer, L. C. (2005). Gender, time and inequality: Trends in women's and men's paid work, unpaid work and free time. *Social forces*, 84(1), 285-303.
- Schaller, J., & Zerpa, M. (2019). Short-run effects of parental job loss on child health. *American Journal of Health Economics*, 5(1), 8-41.
- Schepis, T. S., Teter, C. J., Simoni-Wastila, L., & McCabe, S. E. (2018). Prescription tranquilizer/sedative misuse prevalence and correlates across age cohorts in the US. *Addictive Behaviors*, 87, 24-32.
- Seok, H., Won, J.-U., Lee, T. I., Kim, Y.-K., Lee, W., Lee, J.-H., Roh, J., & Yoon, J.-H. (2016). A dose-response relationship between long working hours and unmet need for access to hospital facilities. *Scandinavian journal of work, environment & health*, 135-143.
- Serdarevic, M., Osborne, V., Striley, C. W., & Cottler, L. B. (2017). The association between insomnia and prescription opioid use: results from a community sample in Northeast Florida. *Sleep Health*, 3(5), 368-372.
- Shields, M. (1999). Long working hours and health. *Health Rep*, 11(2), 33-48.
- Skogstad, M., Mamen, A., Lunde, L.-K., Ulvestad, B., Matre, D., Aass, H. C. D., Øvstebø, R., Nielsen, P., Samuelsen, K. N., & Skare, Ø. (2019). Shift work

- including night work and long working hours in industrial plants increases the risk of atherosclerosis. *International journal of environmental research and public health*, 16(3), 521.
- Sobotka, T., & Beaujouan, É. (2014). Two Is best? The persistence of a two-child family ideal in Europe. *Population and Development Review*, 40(3), 391-419.
- Solon, G., Haider, S. J., & Wooldridge, J. M. (2015). What are we weighting for? *Journal of Human resources*, 50(2), 301-316.
- Søvold, L. E., Naslund, J. A., Kousoulis, A. A., Saxena, S., Qoronfleh, M. W., Grobler, C., & Münter, L. (2021). Prioritizing the mental health and well-being of healthcare workers: an urgent global public health priority. *Frontiers in public health*, 9, 679397.
- StataCorp, L. (2021). Mata reference manual. *College Station, TX, StataCorp LLC*.
- Theorell, T., Hammarström, A., Aronsson, G., Träskman Bendz, L., Grape, T., Hogstedt, C., Marteinsdottir, I., Skoog, I., & Hall, C. (2015). A systematic review including meta-analysis of work environment and depressive symptoms. *BMC Public Health*, 15(1), 1-14.
- Vaidya, V., Gabriel, M. H., Gangan, N., & Borse, M. (2014). Characteristics of prescription and nonprescription sleep medication users in the United States. *Population health management*, 17(6), 345-350.
- Virtanen, M., Ferrie, J. E., Gimeno, D., Vahtera, J., Elovainio, M., Singh-Manoux, A., Marmot, M. G., & Kivimäki, M. (2009). Long working hours and sleep disturbances: the Whitehall II prospective cohort study. *Sleep*, 32(6), 737-745.

- Virtanen, M., Ferrie, J. E., Singh-Manoux, A., Shipley, M. J., Stansfeld, S. A., Marmot, M. G., Ahola, K., Vahtera, J., & Kivimäki, M. (2011). Long working hours and symptoms of anxiety and depression: a 5-year follow-up of the Whitehall II study. *Psychological medicine, 41*(12), 2485-2494.
- Virtanen, M., Heikkilä, K., Jokela, M., Ferrie, J. E., Batty, G. D., Vahtera, J., & Kivimäki, M. (2012). Long working hours and coronary heart disease: a systematic review and meta-analysis. *American journal of epidemiology, 176*(7), 586-596.
- Virtanen, M., Jokela, M., Madsen, I. E., Hanson, L. L. M., Lallukka, T., Nyberg, S. T., Alfredsson, L., Batty, G. D., Bjorner, J. B., & Borritz, M. (2018). Long working hours and depressive symptoms: systematic review and meta-analysis of published studies and unpublished individual participant data. *Scandinavian journal of work, environment & health, 44*(3), 239-250.
- Virtanen, M., Jokela, M., Nyberg, S. T., Madsen, I. E., Lallukka, T., Ahola, K., Alfredsson, L., Batty, G. D., Bjorner, J. B., & Borritz, M. (2015). Long working hours and alcohol use: systematic review and meta-analysis of published studies and unpublished individual participant data. *BMJ, 350*.
- Virtanen, M., & Kivimäki, M. (2012). Saved by the bell: does working too much increase the likelihood of depression? *Expert review of neurotherapeutics, 12*(5), 497-499.
- Votaw, V. R., Geyer, R., Rieselbach, M. M., & McHugh, R. K. (2019). The epidemiology of benzodiazepine misuse: a systematic review. *Drug and alcohol dependence, 200*, 95-114.

- Wang, W., & Griswold, M. E. (2017). Natural interpretations in Tobit regression models using marginal estimation methods. *Statistical methods in medical research*, 26(6), 2622-2632.
- Watanabe, K., Imamura, K., & Kawakami, N. (2016). Working hours and the onset of depressive disorder: a systematic review and meta-analysis. *Occupational and environmental medicine*, 73(12), 877-884.
- Weir, R. C., Toyoji, M., McKee, M., Li, V., & Wang, C. C. (2018). Assessing the impact of electronic health record interventions on hepatitis b screening and vaccination. *Journal of Health Care for the Poor and Underserved*, 29(4), 1587-1605.
- Weissman, M. M., Greenwald, S., Niño-Murcia, G., & Dement, W. C. (1997). The morbidity of insomnia uncomplicated by psychiatric disorders. *General hospital psychiatry*, 19(4), 245-250.
- Williams, J. (2001). *Unbending gender: Why family and work conflict and what to do about it*. Oxford University Press.
- Wisetborisut, A., Angkurawaranon, C., Jiraporncharoen, W., Uaphanthasath, R., & Wiwatanadate, P. (2014). Shift work and burnout among health care workers. *Occupational Medicine*, 64(4), 279-286.
- Wong, K., Chan, A. H., & Ngan, S. (2019). The effect of long working hours and overtime on occupational health: a meta-analysis of evidence from 1998 to 2018. *International journal of environmental research and public health*, 16(12), 2102.
- Wooden, M., Warren, D., & Drago, R. (2009). Working time mismatch and subjective well-being. *British Journal of Industrial Relations*, 47(1), 147-179.

- World Economic Forum (2010). The new discipline of workforce wellness: Enhancing corporate performance by tackling chronic disease. Geneva.
- Yang, H., Schnall, P. L., Jauregui, M., Su, T.-C., & Baker, D. (2006). Work hours and self-reported hypertension among working people in California. *Hypertension*, 48(4), 744-750.
- You, W., & Henneberg, M. (2022). Large household reduces dementia mortality: A cross-sectional data analysis of 183 populations. *Plos One*, 17(3), e0263309.

## APPENDICES



## APPENDIX CHAPTER 1:

### Appendix 1. Medications with sedative properties (Comparisons eAnswers., 2021).

#### Drug Facts And Comparisons

[Acetaminophen and Codeine Oral](#)

[Acetaminophen, Caffeine, and Dihydrocodeine Oral](#)

[Alfentanil Injection](#)

[Alosetron Oral](#)

[ALPRAZolam Oral](#)

[Amitriptyline Oral](#)

[Ampicillin and Sulbactam Injection](#)

[ARIPiprazole Injection](#)

[ARIPiprazole Oral](#)

[Azelastine and Fluticasone Intranasal](#)

[Benzonatate Oral](#)

[Brexpiprazole Oral](#)

[Brivaracetam Injection](#)

[Brivaracetam Oral](#)

[Brompheniramine, Pseudoephedrine, and Dextromethorphan Oral](#)

[Buprenorphine Implant](#)

[Buprenorphine Injection](#)

[Buprenorphine Oral](#)

[Buprenorphine Transdermal](#)

[Butalbital, Acetaminophen, and Caffeine Oral](#)

[Cannabidiol Oral](#)

[Capecitabine Oral](#)

[Carbinoxamine Oral](#)

[Cenobamate Oral](#)

[Cetirizine Injection](#)

[Cetirizine Oral](#)

[Chlorpheniramine and Phenylephrine Oral](#)

[Chlorpheniramine, Phenylephrine, and Dextromethorphan Oral](#)

[Chlorpheniramine, Pseudoephedrine, and Codeine Oral](#)

[Chlorpheniramine, Pseudoephedrine, and Dextromethorphan Oral](#)

[CloBAZam Oral](#)

[Clonidine and Clorthalidone Oral](#)

[CloNIDine Injection](#)

[CloNIDine Oral](#)

[CloNIDine Transdermal](#)

[CloZAPine Oral](#)

[Codeine and Chlorpheniramine Oral](#)

[Codeine Oral](#)

[Cyproheptadine Oral](#)

[Deutetrabenazine Oral](#)

[Dexchlorpheniramine Oral](#)

[DiphenhydrAMINE Injection](#)

[DiphenhydrAMINE Oral](#)

[Diphenoxylate and Atropine Oral](#)

[Doxepin Oral](#)

[Eluxadoline Oral](#)

[Esketamine Intranasal](#)

[Fenfluramine Oral](#)

[FentaNYL Buccal](#)

[FentaNYL Injection](#)

[FentaNYL Intranasal](#)

FentaNYL Sublingual  
FentaNYL Transdermal  
Flibanserin Oral  
Flucytosine Oral  
Flurazepam Oral  
Gabapentin Enacarbil Oral  
Guaifenesin and Codeine Oral  
GuanFACINE Oral  
Haloperidol Injection  
Haloperidol Oral  
Hydrocodone and Acetaminophen Oral  
Hydrocodone and Chlorpheniramine Oral  
Hydrocodone and Homatropine Oral  
Hydrocodone and Pseudoephedrine Oral  
HYDRQcodone Oral  
HYDROmorphine Injection  
HYDROmorphine Oral  
HYDROmorphine Rectal  
Isocarboxazid Oral  
LevETIRAcetam Injection  
LevETIRAcetam Oral  
Lithium Oral  
Lofexidine Oral  
Loratadine Oral  
LORazepam Injection  
LORazepam Oral  
Loxapine Inhalation  
Loxapine Oral  
Lumateperone Oral  
Mecamylamine Oral  
Meperidine Injection  
Meperidine Oral  
Methadone Injection  
Methadone Oral  
Methocarbamol Injection  
Methocarbamol Oral  
Methyldopa and Hydrochlorothiazide Oral  
Methyldopa Oral  
Methyldopate Injection  
Methylphenidate Oral  
Methylphenidate Transdermal  
MetyraPONE Oral  
MetyroSINE Oral  
Midazolam Injection  
Midazolam Intranasal  
Midazolam Oral  
Mitotane Oral  
Morphine and Naltrexone Oral  
Morphine Injection  
Morphine Oral  
Morphine Rectal  
Nabilone Oral  
Nadolol and Bendroflumethiazide Oral  
Nadolol Oral  
Nalbuphine Injection  
Naltrexone Injection  
Naltrexone Oral  
Olanzapine and Samidorphan Oral  
OLANZapine Injection  
OLANZapine Oral  
Oliceridine Injection  
Ondansetron Injection

Ondansetron Oral  
Oxaprozin Oral  
Oxycodone and Acetaminophen Oral  
Oxycodone and Aspirin Oral  
OxyMORphone Injection  
OxyMORphone Oral  
Paliperidone Injection  
Paliperidone Oral  
Papaverine Injection  
Pentazocine and Naloxone Oral  
Pentazocine Injection  
Perphenazine Oral  
PHENobarbital Oral  
Pimozide Oral  
Pregabalin Oral  
Promethazine and Codeine Oral  
Promethazine and Phenylephrine Oral  
Promethazine Injection  
Promethazine Oral  
Promethazine Rectal  
Promethazine, Phenylephrine, and Codeine Oral  
RisperiDONE Injection  
RisperiDONE Oral  
SUFentanil Injection  
SUFentanil Oral  
SUMatriptan Injection  
SUMatriptan Intranasal  
SUMatriptan Oral  
Suvorexant Oral  
Tapentadol Oral  
Tetrabenazine Oral  
TiZANidine Oral  
TraMADol Oral  
Triazolam Oral  
Valbenazine Oral  
Vigabatrin Oral  
Vilazodone Oral  
Viloxazine Oral

N/B: Facts and Comparisons at the time had a search feature that enabled a whole database search for medications that have a side effect of interest.

**Appendix 2a. Heckman Analysis for the Use of Prescription Sleep Aids and Medications with Sedative Properties:  $\geq 36$  h/week and  $\geq 46$  h/week**

| Variables  | $\geq 36$ h/week               |                               |   |                               | $\geq 46$ h/week               |                               |   |                               |
|--|--------------------------------|-------------------------------|---|-------------------------------|--------------------------------|-------------------------------|---|-------------------------------|
|  | Use of Prescription Sleep Aids |                               | Use of Medications with Sedative Properties |                               | Use of Prescription Sleep Aids |                               | Use of Medications with Sedative Properties |                               |
| Work hours/week                                      | Sleep Aids                     | select                        | Sedative Properties                         | select                        | Sleep Aids                     | select                        | Sedative Properties                         | select                        |
| h/week   | 0.135<br>(-0.362, 0.633)       | 0.007<br>(-0.036, 0.050)      | 0.415<br>(-0.902, 1.731)                    | 0.018<br>(-0.010, 0.046)      | 0.051<br>(-0.426, 0.529)       | 0.009<br>(-0.026, 0.044)      | 0.347<br>(-0.813, 1.507)                    | 0.015<br>(-0.008, 0.038)      |
| 27-64 years  | 3.101<br>(-10.214, 16.415)     | 0.356***<br>(0.291, 0.422)    | 3.905<br>(-4.788, 12.598)                   | 0.150***<br>(0.115, 0.186)    | 3.132<br>(-10.285, 16.549)     | 0.357***<br>(0.291, 0.422)    | 4.119<br>(-5.103, 13.342)                   | 0.151***<br>(0.115, 0.187)    |
| $\geq 65$ years                                      | 2.513<br>(-8.970, 13.996)      | 0.306***<br>(0.218, 0.394)    | 5.868<br>(-7.904, 19.641)                   | 0.244***<br>(0.191, 0.296)    | 2.520<br>(-9.025, 14.065)      | 0.305***<br>(0.217, 0.394)    | 6.150<br>(-8.323, 20.622)                   | 0.242***<br>(0.190, 0.295)    |
| <b>Race (White - REF)</b>                            |                                |                               |   |                               |                                |                               |   |                               |
| Black  | -2.996<br>(-16.358, 10.366)    | -0.360***<br>(-0.404, -0.316) | -0.729<br>(-1.692, 0.234)                   | -0.011<br>(-0.035, 0.014)     | -3.015<br>(-16.467, 10.436)    | -0.360***<br>(-0.404, -0.316) | -0.732<br>(-1.735, 0.271)                   | -0.010<br>(-0.035, 0.015)     |
| Other  | -1.854<br>(-8.623, 4.914)      | -0.183***<br>(-0.238, -0.128) | -2.083<br>(-6.375, 2.208)                   | -0.074***<br>(-0.108, -0.041) | -1.861<br>(-8.668, 4.945)      | -0.183***<br>(-0.238, -0.127) | -2.164<br>(-6.661, 2.334)                   | -0.074***<br>(-0.108, -0.040) |
| <b>Sex (Male - REF)</b>                              |                                |                               |   |                               |                                |                               |   |                               |
| Female   | 0.329<br>(-1.606, 2.263)       | 0.052***<br>(0.022, 0.082)    | 0.140<br>(-0.609, 0.889)                    | 0.008<br>(-0.011, 0.027)      | 0.331<br>(-1.641, 2.302)       | 0.053***<br>(0.023, 0.083)    | 0.164<br>(-0.654, 0.983)                    | 0.009<br>(-0.011, 0.028)      |
| <b>Marital status (Married - REF)</b>                |                                |                               |   |                               |                                |                               |   |                               |
| Divorced   | 0.469<br>(-1.628, 2.566)       | 0.057**<br>(0.021, 0.092)     | 1.057<br>(-1.001, 3.115)                    | 0.036**<br>(0.012, 0.060)     | 0.475<br>(-1.636, 2.587)       | 0.057**<br>(0.021, 0.092)     | 1.100<br>(-1.072, 3.273)                    | 0.036**<br>(0.012, 0.060)     |
| Unmarried  | -0.011<br>(-0.596, 0.574)      | 0.012<br>(-0.027, 0.051)      | -1.182<br>(-4.237, 1.874)                   | -0.053***<br>(-0.078, -0.028) | -0.009<br>(-0.602, 0.585)      | 0.012<br>(-0.027, 0.051)      | -1.247<br>(-4.469, 1.975)                   | -0.053***<br>(-0.078, -0.028) |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                                |                               |   |                               |                                |                               |   |                               |
| 4-6 Members  | -1.601<br>(-8.643, 5.442)      | -0.191***<br>(-0.225, -0.157) | -3.066<br>(-10.065, 3.934)                  | -0.125***<br>(-0.146, -0.103) | -1.611<br>(-8.704, 5.483)      | -0.191***<br>(-0.225, -0.157) | -3.225<br>(-10.618, 4.168)                  | -0.125***<br>(-0.146, -0.103) |
| > 6 Members  | -2.678<br>(-16.781, 11.424)    | -0.375***<br>(-0.521, -0.230) | -5.962<br>(-19.564, 7.641)                  | -0.237***<br>(-0.313, -0.161) | -2.696<br>(-16.887, 11.495)    | -0.375***<br>(-0.521, -0.229) | -6.247<br>(-20.565, 8.072)                  | -0.236***<br>(-0.312, -0.160) |

| <b>Education (Less than College - REF)</b>  |                           |                             |                            |                               |                           |                             |                            |                               |
|---|---------------------------|-----------------------------|----------------------------|-------------------------------|---------------------------|-----------------------------|----------------------------|-------------------------------|
| Some college or more                        | 0.662<br>(-3.150, 4.474)  | 0.103***<br>(0.072, 0.135)  | -0.024<br>(-0.634, 0.586)  | -0.001<br>(-0.021, 0.019)     | 0.668<br>(-3.160, 4.496)  | 0.103***<br>(0.072, 0.134)  | -0.033<br>(-0.682, 0.615)  | -0.002<br>(-0.021, 0.018)     |
| <b>Occupation (Natural resources - REF)</b> |                           |                             |                            |                               |                           |                             |                            |                               |
| Hospitality services                        | 0.308<br>(-0.633, 1.249)  | 0.013<br>(-0.069, 0.096)    | 0.869<br>(-1.883, 3.621)   | 0.041<br>(-0.009, 0.091)      | 0.291<br>(-0.644, 1.226)  | 0.012<br>(-0.070, 0.095)    | 0.872<br>(-1.937, 3.681)   | 0.039<br>(-0.011, 0.089)      |
| Trade                                       | 0.675<br>(-1.493, 2.843)  | 0.056<br>(-0.016, 0.127)    | 1.322<br>(-1.767, 4.412)   | 0.050*<br>(0.005, 0.094)      | 0.671<br>(-1.512, 2.854)  | 0.056<br>(-0.016, 0.127)    | 1.376<br>(-1.870, 4.621)   | 0.049*<br>(0.005, 0.093)      |
| Professional services                       | 1.217<br>(-3.467, 5.901)  | 0.126***<br>(0.063, 0.190)  | 1.782<br>(-2.704, 6.268)   | 0.077***<br>(0.038, 0.117)    | 1.226<br>(-3.504, 5.956)  | 0.127***<br>(0.063, 0.190)  | 1.887<br>(-2.870, 6.644)   | 0.078***<br>(0.038, 0.117)    |
| Manufacturing                               | 0.987<br>(-2.774, 4.747)  | 0.100**<br>(0.030, 0.171)   | 1.311<br>(-2.458, 5.081)   | 0.063**<br>(0.019, 0.107)     | 1.000<br>(-2.809, 4.809)  | 0.101**<br>(0.031, 0.171)   | 1.426<br>(-2.636, 5.487)   | 0.064**<br>(0.020, 0.108)     |
| Other                                       | 0.851<br>(-2.873, 4.575)  | 0.100**<br>(0.035, 0.164)   | 1.532<br>(-2.757, 5.820)   | 0.074***<br>(0.034, 0.114)    | 0.862<br>(-2.912, 4.636)  | 0.100**<br>(0.036, 0.165)   | 1.650<br>(-2.941, 6.240)   | 0.075***<br>(0.035, 0.115)    |
| <b>Year (2010 - REF)</b>                    |                           |                             |                            |                               |                           |                             |                            |                               |
| 2011  | 0.129<br>(-0.821, 1.078)  | -0.020<br>(-0.082, 0.042)   | -1.959<br>(-7.907, 3.989)  | -0.106***<br>(-0.146, -0.066) | 0.130<br>(-0.825, 1.085)  | -0.020<br>(-0.082, 0.042)   | -2.089<br>(-8.366, 4.188)  | -0.106***<br>(-0.146, -0.066) |
| 2012  | -0.160<br>(-2.906, 2.586) | -0.073*<br>(-0.135, -0.011) | -3.415<br>(-12.609, 5.780) | -0.165***<br>(-0.205, -0.125) | -0.164<br>(-2.927, 2.598) | -0.073*<br>(-0.135, -0.011) | -3.618<br>(-13.318, 6.082) | -0.165***<br>(-0.205, -0.125) |
| 2013  | 0.090<br>(-1.481, 1.661)  | -0.040<br>(-0.103, 0.023)   | -1.971<br>(-7.587, 3.646)  | -0.100***<br>(-0.141, -0.060) | 0.086<br>(-1.491, 1.664)  | -0.039<br>(-0.103, 0.024)   | -2.091<br>(-8.011, 3.830)  | -0.100***<br>(-0.140, -0.059) |
| 2014  | -0.177<br>(-1.296, 0.943) | -0.026<br>(-0.088, 0.036)   | -2.302<br>(-8.629, 4.025)  | -0.113***<br>(-0.153, -0.073) | -0.178<br>(-1.307, 0.951) | -0.026<br>(-0.088, 0.036)   | -2.445<br>(-9.131, 4.242)  | -0.113***<br>(-0.153, -0.073) |
| 2015  | -0.400<br>(-2.595, 1.796) | -0.058<br>(-0.120, 0.005)   | -2.000<br>(-7.452, 3.451)  | -0.097***<br>(-0.137, -0.057) | -0.404<br>(-2.617, 1.809) | -0.058<br>(-0.120, 0.005)   | -2.125<br>(-7.892, 3.642)  | -0.097***<br>(-0.137, -0.057) |
| 2016  | -0.243<br>(-2.175, 1.689) | -0.050<br>(-0.113, 0.013)   | -2.228<br>(-8.607, 4.151)  | -0.114***<br>(-0.155, -0.073) | -0.243<br>(-2.186, 1.699) | -0.050<br>(-0.113, 0.013)   | -2.368<br>(-9.099, 4.364)  | -0.114***<br>(-0.154, -0.073) |
| 2017  | 0.371<br>(-0.250, 0.992)  | -0.003<br>(-0.066, 0.060)   | -1.997<br>(-8.361, 4.367)  | -0.114***<br>(-0.155, -0.072) | 0.373<br>(-0.254, 0.999)  | -0.003<br>(-0.066, 0.060)   | -2.143<br>(-8.873, 4.588)  | -0.114***<br>(-0.155, -0.072) |
| 2018  | 0.316                     | 0.003                       | -1.756                     | -0.104***                     | 0.318                     | 0.003                       | -1.892                     | -0.104***                     |

|  |                   |                 |                    |                  |                   |                 |                    |                  |
|--|-------------------|-----------------|--------------------|------------------|-------------------|-----------------|--------------------|------------------|
| 2019                                     | (-0.305, 0.938)   | (-0.060, 0.066) | (-7.600, 4.088)    | (-0.146, -0.063) | (-0.309, 0.944)   | (-0.060, 0.066) | (-8.078, 4.295)    | (-0.146, -0.063) |
|  | 0.695             | 0.091**         | -0.214             | -0.021           | 0.696             | 0.091**         | -0.249             | -0.022           |
|  | (-2.701, 4.090)   | (0.028, 0.155)  | (-1.952, 1.523)    | (-0.064, 0.022)  | (-2.719, 4.111)   | (0.027, 0.155)  | (-2.099, 1.601)    | (-0.065, 0.021)  |
| <b>Region (Northeast - REF)</b>          |                   |                 |                    |                  |                   |                 |                    |                  |
| North central, Midwest                   | 0.071             |                 | 0.150              |                  | 0.074             |                 | 0.151              |                  |
|  | (-0.236, 0.378)   |                 | (-0.511, 0.811)    |                  | (-0.235, 0.383)   |                 | (-0.548, 0.850)    |                  |
| South                                    | 0.030             |                 | 0.075              |                  | 0.034             |                 | 0.077              |                  |
|  | (-0.262, 0.323)   |                 | (-0.543, 0.693)    |                  | (-0.261, 0.329)   |                 | (-0.576, 0.730)    |                  |
| West                                     | -0.232            |                 | -0.123             |                  | -0.229            |                 | -0.121             |                  |
|  | (-0.545, 0.081)   |                 | (-0.786, 0.541)    |                  | (-0.545, 0.086)   |                 | (-0.823, 0.580)    |                  |
| <b>Income (Poor to low income - REF)</b> |                   |                 |                    |                  |                   |                 |                    |                  |
| Middle to high income                    | 0.049             | 0.020           | -0.091             | -0.004           | 0.059             | 0.020           | -0.085             | -0.004           |
|  | (-0.773, 0.871)   | (-0.021, 0.060) | (-0.901, 0.718)    | (-0.029, 0.021)  | (-0.767, 0.885)   | (-0.021, 0.060) | (-0.931, 0.762)    | (-0.029, 0.021)  |
| <b>Receipt of food stamp (No - REF)</b>  |                   |                 |                    |                  |                   |                 |                    |                  |
| YES                                      | 0.568             | 0.045           | 2.230              | 0.086***         | 0.557             | 0.045           | 2.314              | 0.085***         |
|  | (-1.160, 2.296)   | (-0.011, 0.101) | (-2.634, 7.094)    | (0.051, 0.120)   | (-1.168, 2.281)   | (-0.011, 0.101) | (-2.767, 7.395)    | (0.050, 0.119)   |
| <b>Health status (Excellent - REF)</b>   |                   |                 |                    |                  |                   |                 |                    |                  |
| Good                                     | 1.489             | 0.179***        | 4.617              | 0.183***         | 1.500             | 0.179***        | 4.858              | 0.183***         |
|  | (-5.162, 8.139)   | (0.135, 0.224)  | (-5.889, 15.123)   | (0.157, 0.210)   | (-5.206, 8.205)   | (0.135, 0.224)  | (-6.244, 15.960)   | (0.157, 0.210)   |
| Fair                                     | 3.124             | 0.318***        | 10.311             | 0.409***         | 3.143             | 0.318***        | 10.831             | 0.410***         |
|  | (-8.612, 14.860)  | (0.261, 0.375)  | (-12.394, 33.017)  | (0.373, 0.446)   | (-8.687, 14.973)  | (0.261, 0.375)  | (-13.156, 34.817)  | (0.373, 0.446)   |
| Poor                                     | 4.403             | 0.470***        | 15.086             | 0.585***         | 4.433             | 0.470***        | 15.806             | 0.585***         |
|  | (-12.837, 21.643) | (0.361, 0.579)  | (-16.544, 46.715)  | (0.507, 0.663)   | (-12.938, 21.804) | (0.361, 0.579)  | (-17.600, 49.212)  | (0.507, 0.663)   |
| Lambda                                   | 8.386             |                 | 30.596             |                  | 8.461             |                 | 32.377             |                  |
|  | (-34.820, 51.593) |                 | (-47.237, 108.430) |                  | (-35.067, 51.988) |                 | (-49.835, 114.589) |                  |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference; Instrumenting on Region

**Appendix 2b. Heckman Analysis for the Use of Prescription Sleep Aids and Medications with Sedative Properties:  $\geq 56$  h/week and  $\geq 66$  h/week**

| Variables  | $\geq 56$ h/week               |                               |   |                               | $\geq 66$ h/week               |                               |   |                               |
|--|--------------------------------|-------------------------------|---|-------------------------------|--------------------------------|-------------------------------|---|-------------------------------|
|  | Use of Prescription Sleep Aids |                               | Use of Medications with Sedative Properties |                               | Use of Prescription Sleep Aids |                               | Use of Medications with Sedative Properties |                               |
| Work hours/week                                      | Sleep Aids                     | select                        | Sedative Properties                         | select                        | Sleep Aids                     | select                        | Sedative Properties                         | select                        |
| h/week   | 0.826<br>(-1.573, 3.225)       | 0.060*<br>(0.006, 0.114)      | 1.649<br>(-1.833, 5.131)                    | 0.056**<br>(0.020, 0.092)     | 0.451<br>(-1.062, 1.964)       | 0.037<br>(-0.061, 0.136)      | 0.693<br>(-1.651, 3.038)                    | 0.020<br>(-0.045, 0.085)      |
| <b>Age (18-26years - REF)</b>                        |                                |                               |   |                               |                                |                               |   |                               |
| 27-64 years  | 3.270<br>(-10.825, 17.366)     | 0.356***<br>(0.290, 0.422)    | 4.181<br>(-5.098, 13.460)                   | 0.151***<br>(0.115, 0.186)    | 2.874<br>(-9.241, 14.989)      | 0.357***<br>(0.291, 0.423)    | 4.046<br>(-5.016, 13.108)                   | 0.152***<br>(0.116, 0.187)    |
| $\geq 65$ years                                      | 2.617<br>(-9.475, 14.710)      | 0.304***<br>(0.216, 0.392)    | 6.227<br>(-8.296, 20.749)                   | 0.241***<br>(0.189, 0.294)    | 2.289<br>(-8.117, 12.695)      | 0.305***<br>(0.217, 0.393)    | 6.011<br>(-8.159, 20.182)                   | 0.242***<br>(0.190, 0.295)    |
| <b>Race (White - REF)</b>                            |                                |                               |   |                               |                                |                               |   |                               |
| Black  | -3.178<br>(-17.353, 10.997)    | -0.360***<br>(-0.404, -0.316) | -0.747<br>(-1.781, 0.286)                   | -0.010<br>(-0.035, 0.014)     | -2.756<br>(-14.897, 9.386)     | -0.360***<br>(-0.404, -0.316) | -0.734<br>(-1.723, 0.256)                   | -0.010<br>(-0.035, 0.014)     |
| Other  | -1.958<br>(-9.136, 5.220)      | -0.183***<br>(-0.238, -0.128) | -2.207<br>(-6.757, 2.344)                   | -0.074***<br>(-0.108, -0.040) | -1.736<br>(-7.888, 4.415)      | -0.183***<br>(-0.238, -0.128) | -2.132<br>(-6.557, 2.292)                   | -0.074***<br>(-0.108, -0.040) |
| <b>Sex (Male - REF)</b>                              |                                |                               |   |                               |                                |                               |   |                               |
| Female   | 0.396<br>(-1.764, 2.556)       | 0.055***<br>(0.025, 0.085)    | 0.216<br>(-0.660, 1.093)                    | 0.010<br>(-0.010, 0.029)      | 0.298<br>(-1.474, 2.070)       | 0.052***<br>(0.022, 0.082)    | 0.139<br>(-0.618, 0.895)                    | 0.008<br>(-0.012, 0.027)      |
| <b>Marital status (Married - REF)</b>                |                                |                               |   |                               |                                |                               |   |                               |
| Divorced   | 0.491<br>(-1.720, 2.702)       | 0.056**<br>(0.021, 0.092)     | 1.108<br>(-1.072, 3.288)                    | 0.035**<br>(0.012, 0.059)     | 0.435<br>(-1.472, 2.342)       | 0.057**<br>(0.021, 0.092)     | 1.081<br>(-1.046, 3.209)                    | 0.036**<br>(0.012, 0.060)     |
| Unmarried  | -0.001<br>(-0.639, 0.637)      | 0.013<br>(-0.026, 0.051)      | -1.265<br>(-4.503, 1.973)                   | -0.053***<br>(-0.078, -0.028) | -0.015<br>(-0.552, 0.522)      | 0.012<br>(-0.026, 0.051)      | -1.220<br>(-4.383, 1.943)                   | -0.053***<br>(-0.078, -0.028) |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                                |                               |   |                               |                                |                               |   |                               |
| 4-6 Members  | -1.692<br>(-9.150, 5.767)      | -0.190***<br>(-0.224, -0.156) | -3.283<br>(-10.727, 4.162)                  | -0.124***<br>(-0.146, -0.103) | -1.474<br>(-7.869, 4.921)      | -0.191***<br>(-0.225, -0.156) | -3.153<br>(-10.389, 4.082)                  | -0.125***<br>(-0.146, -0.103) |
| > 6 Members  | -2.837                         | -0.374***                     | -6.359                                      | -0.236***                     | -2.420                         | -0.375***                     | -6.119                                      | -0.236***                     |

|   |                           |                             |                            |                               |                           |                                 |                            |                               |
|---|---------------------------|-----------------------------|----------------------------|-------------------------------|---------------------------|---------------------------------|----------------------------|-------------------------------|
|   | (-17.758,<br>12.084)      | (-0.520, -0.228)            | (-20.792, 8.074)           | (-0.312, -0.160)              | (-15.230, 10.390)         | (-0.521, -<br>0.229)            | (-20.149, 7.912)           | (-0.312, -0.161)              |
| <b>Education (Less than College - REF)</b>  |                           |                             |                            |                               |                           |                                 |                            |                               |
| Some<br>college or<br>more                  | 0.706<br>(-3.314, 4.725)  | 0.103***<br>(0.071, 0.134)  | -0.035<br>(-0.696, 0.626)  | -0.001<br>(-0.021, 0.018)     | 0.600<br>(-2.875, 4.075)  | 0.104***<br>(0.072, 0.135)      | -0.013<br>(-0.641, 0.615)  | -0.001<br>(-0.021, 0.019)     |
| <b>Occupation (Natural resources - REF)</b> |                           |                             |                            |                               |                           |                                 |                            |                               |
| Hospitality<br>services                     | 0.350<br>(-0.696, 1.397)  | 0.015<br>(-0.067, 0.097)    | 0.946<br>(-1.977, 3.869)   | 0.040<br>(-0.010, 0.090)      | 0.284<br>(-0.561, 1.129)  | 0.013<br>(-0.070, 0.095)        | 0.844<br>(-1.876, 3.565)   | 0.038<br>(-0.012, 0.088)      |
| Trade                                       | 0.741<br>(-1.663, 3.145)  | 0.058<br>(-0.013, 0.130)    | 1.461<br>(-1.919, 4.841)   | 0.051*<br>(0.007, 0.095)      | 0.633<br>(-1.342, 2.609)  | 0.056<br>(-0.016, 0.127)        | 1.341<br>(-1.803, 4.485)   | 0.049*<br>(0.004, 0.093)      |
| Professiona<br>l services                   | 1.322<br>(-3.748, 6.392)  | 0.129***<br>(0.065, 0.192)  | 1.974<br>(-2.909, 6.857)   | 0.079***<br>(0.039, 0.119)    | 1.133<br>(-3.129, 5.396)  | 0.126***<br>(0.063, 0.190)      | 1.832<br>(-2.782, 6.447)   | 0.077***<br>(0.037, 0.117)    |
| Manufactur<br>ing                           | 1.090<br>(-3.039, 5.218)  | 0.104**<br>(0.034, 0.175)   | 1.534<br>(-2.701, 5.768)   | 0.067**<br>(0.023, 0.111)     | 0.933<br>(-2.522, 4.388)  | 0.102**<br>(0.031, 0.172)       | 1.396<br>(-2.574, 5.367)   | 0.064**<br>(0.020, 0.108)     |
| Other                                       | 0.945<br>(-3.119, 5.008)  | 0.103**<br>(0.038, 0.167)   | 1.735<br>(-2.979, 6.449)   | 0.076***<br>(0.036, 0.116)    | 0.784<br>(-2.612, 4.180)  | 0.100**<br>(0.036, 0.165)       | 1.594<br>(-2.853, 6.041)   | 0.074***<br>(0.034, 0.114)    |
| <b>Year (2010 - REF)</b>                    |                           |                             |                            |                               |                           |                                 |                            |                               |
| 2011  | 0.123<br>(-0.889, 1.136)  | -0.020<br>(-0.083, 0.042)   | -2.146<br>(-8.491, 4.199)  | -0.106***<br>(-0.147, -0.066) | 0.138<br>(-0.730, 1.007)  | -0.020<br>(-0.083, 0.042)       | -2.035<br>(-8.186, 4.116)  | -0.106***<br>(-0.146, -0.066) |
| 2012  | -0.191<br>(-3.094, 2.711) | -0.073*<br>(-0.135, -0.010) | -3.693<br>(-13.464, 6.079) | -0.165***<br>(-0.204, -0.125) | -0.112<br>(-2.608, 2.383) | -0.073*<br>(-0.135, -<br>0.011) | -3.528<br>(-13.026, 5.970) | -0.165***<br>(-0.205, -0.125) |
| 2013  | 0.075<br>(-1.578, 1.727)  | -0.039<br>(-0.102, 0.024)   | -2.131<br>(-8.088, 3.827)  | -0.100***<br>(-0.140, -0.059) | 0.115<br>(-1.309, 1.539)  | -0.040<br>(-0.103, 0.024)       | -2.035<br>(-7.832, 3.761)  | -0.100***<br>(-0.140, -0.059) |
| 2014  | -0.193<br>(-1.388, 1.003) | -0.026<br>(-0.088, 0.036)   | -2.501<br>(-9.246, 4.244)  | -0.113***<br>(-0.153, -0.073) | -0.163<br>(-1.183, 0.857) | -0.026<br>(-0.088, 0.036)       | -2.379<br>(-8.916, 4.158)  | -0.113***<br>(-0.153, -0.073) |
| 2015  | -0.424<br>(-2.755, 1.907) | -0.058<br>(-0.120, 0.005)   | -2.174<br>(-7.995, 3.647)  | -0.097***<br>(-0.137, -0.057) | -0.366<br>(-2.361, 1.630) | -0.058<br>(-0.120, 0.005)       | -2.065<br>(-7.694, 3.564)  | -0.097***<br>(-0.137, -0.057) |
| 2016  | -0.259<br>(-2.295, 1.778) | -0.050<br>(-0.113, 0.014)   | -2.415<br>(-9.189, 4.359)  | -0.114***<br>(-0.154, -0.073) | -0.209<br>(-1.956, 1.539) | -0.050<br>(-0.113, 0.013)       | -2.298<br>(-8.872, 4.276)  | -0.114***<br>(-0.154, -0.073) |
| 2017  | 0.382<br>(-0.281, 1.044)  | -0.003<br>(-0.066, 0.061)   | -2.187<br>(-8.954, 4.579)  | -0.113***<br>(-0.155, -0.072) | 0.378<br>(-0.185, 0.941)  | -0.003<br>(-0.066, 0.060)       | -2.072<br>(-8.642, 4.499)  | -0.113***<br>(-0.155, -0.072) |
| 2018  | 0.323<br>(-0.343, 0.989)  | 0.003<br>(-0.060, 0.066)    | -1.934<br>(-8.159, 4.291)  | -0.104***<br>(-0.146, -0.063) | 0.317<br>(-0.248, 0.883)  | 0.003<br>(-0.060, 0.066)        | -1.827<br>(-7.866, 4.212)  | -0.104***<br>(-0.146, -0.063) |
| 2019  | 0.746                     | 0.091**                     | -0.261                     | -0.022                        | 0.636                     | 0.092**                         | -0.231                     | -0.021                        |



|  |                   |                 |                    |                 |                   |                 |                    |                 |
|--|-------------------|-----------------|--------------------|-----------------|-------------------|-----------------|--------------------|-----------------|
|  | (-2.855, 4.346)   | (0.028, 0.155)  | (-2.142, 1.620)    | (-0.065, 0.021) | (-2.454, 3.725)   | (0.028, 0.156)  | (-2.030, 1.568)    | (-0.064, 0.022) |
| <b>Region (Northeast - REF)</b>          |                   |                 |                    |                 |                   |                 |                    |                 |
| North central,                           |                   |                 |                    |                 |                   |                 |                    |                 |
| Midwest                                  | 0.064             |                 | 0.143              |                 | 0.072             |                 | 0.151              |                 |
|  | (-0.264, 0.393)   |                 | (-0.571, 0.857)    |                 | (-0.207, 0.350)   |                 | (-0.531, 0.833)    |                 |
| South                                    | 0.030             |                 | 0.075              |                 | 0.033             |                 | 0.076              |                 |
|  | (-0.283, 0.343)   |                 | (-0.592, 0.742)    |                 | (-0.233, 0.299)   |                 | (-0.561, 0.713)    |                 |
| West                                     | -0.223            |                 | -0.120             |                 | -0.227            |                 | -0.122             |                 |
|  | (-0.558, 0.112)   |                 | (-0.837, 0.596)    |                 | (-0.511, 0.057)   |                 | (-0.806, 0.563)    |                 |
| <b>Income (Poor to low income - REF)</b> |                   |                 |                    |                 |                   |                 |                    |                 |
| Middle to high income                    | 0.049             | 0.019           | -0.100             | -0.004          | 0.048             | 0.020           | -0.058             | -0.003          |
|  | (-0.795, 0.892)   | (-0.021, 0.059) | (-0.966, 0.765)    | (-0.029, 0.021) | (-0.715, 0.811)   | (-0.020, 0.060) | (-0.866, 0.750)    | (-0.028, 0.022) |
| <b>Receipt of food stamp (No - REF)</b>  |                   |                 |                    |                 |                   |                 |                    |                 |
| YES                                      | 0.563             | 0.044           | 2.339              | 0.084***        | 0.516             | 0.044           | 2.258              | 0.085***        |
|  | (-1.230, 2.355)   | (-0.012, 0.100) | (-2.754, 7.433)    | (0.050, 0.119)  | (-1.027, 2.060)   | (-0.012, 0.100) | (-2.700, 7.216)    | (0.050, 0.119)  |
| <b>Health status (Excellent - REF)</b>   |                   |                 |                    |                 |                   |                 |                    |                 |
| Good                                     | 1.604             | 0.180***        | 4.963              | 0.184***        | 1.374             | 0.179***        | 4.749              | 0.183***        |
|  | (-5.487, 8.695)   | (0.136, 0.224)  | (-6.246, 16.172)   | (0.157, 0.210)  | (-4.674, 7.421)   | (0.135, 0.224)  | (-6.104, 15.603)   | (0.156, 0.210)  |
| Fair                                     | 3.302             | 0.319***        | 11.036             | 0.409***        | 2.917             | 0.318***        | 10.591             | 0.409***        |
|  | (-9.168, 15.773)  | (0.262, 0.376)  | (-13.140, 35.211)  | (0.373, 0.446)  | (-7.744, 13.579)  | (0.261, 0.375)  | (-12.856, 34.039)  | (0.373, 0.445)  |
| Poor                                     | 4.655             | 0.470***        | 16.073             | 0.585***        | 4.105             | 0.470***        | 15.485             | 0.585***        |
|  | (-13.634, 22.943) | (0.361, 0.579)  | (-17.579, 49.726)  | (0.507, 0.663)  | (-11.563, 19.773) | (0.361, 0.579)  | (-17.200, 48.170)  | (0.507, 0.663)  |
| Lambda                                   | 8.986             |                 | 33.075             |                 | 7.625             |                 | 31.591             |                 |
|  | (-36.846, 54.817) |                 | (-49.800, 115.951) |                 | (-31.646, 46.896) |                 | (-48.865, 112.048) |                 |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference; Instrumenting on Region

Appendix 3a. Regressions for Use of Prescription Sleep Aids:  $\geq 36$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Prescription Sleep Aid = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 36$ h/week                                     | 0.005<br>(-0.014, 0.024)      | 0.007<br>(-0.031, 0.045)      | 1.01<br>(0.923, 1.106)     | 0.001<br>(-0.005, 0.006)      | 0.084<br>(-0.115, 0.283)  |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.104***<br>(0.080, 0.128)    | 0.305***<br>(0.255, 0.356)    | 2.206***<br>(1.897, 2.565) | 0.034***<br>(0.030, 0.039)    | 0.522**<br>(0.188, 0.856)   |
| $\geq 65$ years                                      | 0.069***<br>(0.033, 0.105)    | 0.253***<br>(0.181, 0.325)    | 1.995***<br>(1.644, 2.420) | 0.029***<br>(0.020, 0.037)    | 0.293<br>(-0.132, 0.718)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.127***<br>(-0.144, -0.109) | -0.313***<br>(-0.349, -0.277) | 0.456***<br>(0.413, 0.505) | -0.037***<br>(-0.040, -0.033) | -0.407***<br>(-0.628, -0.186)                                       |
| Other  | -0.087***<br>(-0.109, -0.064) | -0.171***<br>(-0.218, -0.123) | 0.687***<br>(0.609, 0.775) | -0.021***<br>(-0.026, -0.015) | -0.546***<br>(-0.811, -0.281)                                       |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | 0.012<br>(-0.001, 0.025)      | 0.043**<br>(0.017, 0.070)     | 1.108**<br>(1.041, 1.180)  | 0.006**<br>(0.002, 0.009)     | -0.042<br>(-0.177, 0.093)   |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.029***<br>(0.012, 0.045)    | 0.053**<br>(0.021, 0.085)     | 1.127**<br>(1.047, 1.212)  | 0.007**<br>(0.003, 0.011)     | 0.068<br>(-0.090, 0.226)  |
| Unmarried  | 0.002<br>(-0.015, 0.019)      | 0.009<br>(-0.026, 0.043)      | 1.034<br>(0.954, 1.121)    | 0.002<br>(-0.003, 0.006)      | -0.098<br>(-0.275, 0.078)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.072***<br>(-0.086, -0.058) | -0.171***<br>(-0.200, -0.142) | 0.670***<br>(0.623, 0.720) | -0.021***<br>(-0.025, -0.018) | -0.237**<br>(-0.397, -0.077)  |
| > 6 Members  | -0.101***<br>(-0.151, -0.051) | -0.308***<br>(-0.419, -0.198) | 0.437***<br>(0.311, 0.615) | -0.037***<br>(-0.048, -0.027) | 0.043<br>(-0.714, 0.800)  |
| <b>Education (Less than College - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | 0.026***<br>(0.013, 0.040)    | 0.087***<br>(0.059, 0.114)    | 1.237***<br>(1.158, 1.321) | 0.012***<br>(0.008, 0.015)    | -0.074<br>(-0.217, 0.069)   |
| <b>Occupation (Natural resources - REF)</b>          |                               |                               |                            |                               |   |
| Hospitality services                                 | 0.021<br>(-0.013, 0.055)      | 0.018<br>(-0.052, 0.088)      | 1.033<br>(0.866, 1.232)    | 0.002<br>(-0.007, 0.010)      | 0.214<br>(-0.173, 0.600)  |
| Trade  | 0.033*<br>(0.003, 0.063)      | 0.057<br>(-0.004, 0.118)      | 1.114<br>(0.956, 1.297)    | 0.005<br>(-0.002, 0.013)      | 0.277<br>(-0.057, 0.610)  |
| Professional services                                | 0.061***<br>(0.034, 0.088)    | 0.120***<br>(0.066, 0.175)    | 1.301***<br>(1.137, 1.490) | 0.014***<br>(0.008, 0.021)    | 0.317*<br>(0.023, 0.611)  |
| Manufacturing  | 0.046**<br>(0.016, 0.076)     | 0.091**<br>(0.030, 0.151)     | 1.221**<br>(1.051, 1.418)  | 0.011**<br>(0.003, 0.018)     | 0.27<br>(-0.058, 0.597)   |
| Other  | 0.039**                       | 0.091**                       | 1.231**                    | 0.011**                       | 0.139   |

|  |                            |                            |                            |                             |                             |
|--|----------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|
|  | (0.012, 0.067)             | (0.036, 0.146)             | (1.073, 1.412)             | (0.004, 0.018)              | (-0.161, 0.439)             |
| <b>Year (2010 - REF)</b>                 |                            |                            |                            |                             |                             |
| 2011                                     | 0.011<br>(-0.017, 0.038)   | -0.008<br>(-0.064, 0.047)  | 0.964<br>(0.847, 1.097)    | -0.002<br>(-0.010, 0.005)   | 0.271<br>(-0.009, 0.551)    |
| 2012                                     | -0.004<br>(-0.031, 0.023)  | -0.051<br>(-0.106, 0.003)  | 0.856*<br>(0.751, 0.975)   | -0.009*<br>(-0.016, -0.001) | 0.359*<br>(0.076, 0.642)    |
| 2013                                     | 0.009<br>(-0.019, 0.036)   | -0.021<br>(-0.077, 0.035)  | 0.922<br>(0.808, 1.051)    | -0.005<br>(-0.012, 0.003)   | 0.370*<br>(0.085, 0.656)    |
| 2014                                     | -0.005<br>(-0.033, 0.022)  | -0.02<br>(-0.075, 0.036)   | 0.953<br>(0.837, 1.084)    | -0.003<br>(-0.010, 0.005)   | 0.006<br>(-0.274, 0.287)    |
| 2015                                     | -0.019<br>(-0.046, 0.008)  | -0.049<br>(-0.104, 0.006)  | 0.885<br>(0.777, 1.009)    | -0.007<br>(-0.014, 0.001)   | 0.01<br>(-0.273, 0.293)     |
| 2016                                     | -0.012<br>(-0.039, 0.016)  | -0.041<br>(-0.097, 0.015)  | 0.896<br>(0.785, 1.022)    | -0.006<br>(-0.014, 0.001)   | 0.112<br>(-0.174, 0.399)    |
| 2017                                     | 0.022<br>(-0.007, 0.050)   | 0.011<br>(-0.046, 0.068)   | 0.985<br>(0.864, 1.123)    | -0.001<br>(-0.009, 0.007)   | 0.393**<br>(0.109, 0.677)   |
| 2018                                     | 0.018<br>(-0.011, 0.046)   | 0.013<br>(-0.044, 0.070)   | 0.996<br>(0.874, 1.134)    | 0<br>(-0.008, 0.008)        | 0.293*<br>(0.010, 0.576)    |
| 2019                                     | 0.033*<br>(0.003, 0.063)   | 0.083**<br>(0.024, 0.141)  | 1.192**<br>(1.047, 1.358)  | 0.011**<br>(0.003, 0.020)   | 0.048<br>(-0.234, 0.330)    |
| <b>Region (Northeast - REF)</b>          |                            |                            |                            |                             |                             |
| North<br>central,<br>Midwest             | 0.028**<br>(0.008, 0.049)  | 0.059**<br>(0.018, 0.101)  | 1.151**<br>(1.045, 1.268)  | 0.008**<br>(0.003, 0.014)   | 0.072<br>(-0.139, 0.282)    |
| South                                    | 0.015<br>(-0.004, 0.035)   | 0.034<br>(-0.005, 0.073)   | 1.09<br>(0.994, 1.196)     | 0.005<br>(-0.000, 0.010)    | 0.033<br>(-0.169, 0.234)    |
| West                                     | -0.01<br>(-0.031, 0.010)   | 0.004<br>(-0.037, 0.045)   | 1.036<br>(0.939, 1.143)    | 0.002<br>(-0.003, 0.007)    | -0.232*<br>(-0.447, -0.016) |
| <b>Income (Poor to low income - REF)</b> |                            |                            |                            |                             |                             |
| Middle to<br>high<br>income              | 0.002<br>(-0.015, 0.019)   | 0.015<br>(-0.021, 0.050)   | 1.043<br>(0.958, 1.135)    | 0.002<br>(-0.002, 0.007)    | -0.091<br>(-0.280, 0.097)   |
| <b>Receipt of food stamp (No - REF)</b>  |                            |                            |                            |                             |                             |
| YES                                      | 0.029*<br>(0.005, 0.052)   | 0.047<br>(-0.004, 0.098)   | 1.093<br>(0.971, 1.230)    | 0.005<br>(-0.002, 0.013)    | 0.25<br>(-0.015, 0.515)     |
| <b>Health status (Excellent - REF)</b>   |                            |                            |                            |                             |                             |
| Good                                     | 0.058***<br>(0.040, 0.075) | 0.154***<br>(0.118, 0.191) | 1.466***<br>(1.331, 1.615) | 0.019***<br>(0.015, 0.023)  | 0.202<br>(-0.011, 0.416)    |
| Fair                                     | 0.156***<br>(0.132, 0.181) | 0.307***<br>(0.257, 0.358) | 1.949***<br>(1.728, 2.198) | 0.037***<br>(0.030, 0.044)  | 0.852***<br>(0.588, 1.116)  |
| Poor                                     | 0.258***<br>(0.204, 0.313) | 0.470***<br>(0.359, 0.581) | 2.611***<br>(2.104, 3.241) | 0.061***<br>(0.043, 0.080)  | 1.069***<br>(0.606, 1.532)  |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 3b. Regressions for the Use of Prescription Medications with Sedative Properties:  $\geq 36$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Sedative Medications = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 36$ h/week                                     | 0.017<br>(-0.034, 0.068)      | 0.022<br>(-0.014, 0.057)      | 1.028<br>(0.983, 1.076)    | 0.006<br>(-0.004, 0.016)      | 0.02<br>(-0.104, 0.144)   |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.294***<br>(0.230, 0.359)    | 0.235***<br>(0.192, 0.278)    | 1.290***<br>(1.215, 1.369) | 0.054***<br>(0.042, 0.067)    | 0.521***<br>(0.350, 0.693)  |
| $\geq 65$ years                                      | 0.382***<br>(0.285, 0.479)    | 0.339***<br>(0.271, 0.406)    | 1.500***<br>(1.377, 1.635) | 0.089***<br>(0.070, 0.108)    | 0.499***<br>(0.262, 0.737)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.201***<br>(-0.248, -0.154) | -0.088***<br>(-0.121, -0.056) | 0.974<br>(0.934, 1.016)    | -0.006<br>(-0.015, 0.003)     | -0.502***<br>(-0.615, -0.388)                                     |
| Other  | -0.215***<br>(-0.277, -0.153) | -0.138***<br>(-0.181, -0.095) | 0.893***<br>(0.844, 0.945) | -0.025***<br>(-0.037, -0.013) | -0.452***<br>(-0.609, -0.294)                                     |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | -0.007<br>(-0.042, 0.029)     | 0.003<br>(-0.022, 0.029)      | 1.013<br>(0.981, 1.045)    | 0.003<br>(-0.004, 0.010)      | -0.038<br>(-0.125, 0.049)   |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.158***<br>(0.114, 0.203)    | 0.086***<br>(0.055, 0.118)    | 1.061**<br>(1.020, 1.103)  | 0.013**<br>(0.004, 0.022)     | 0.301***<br>(0.196, 0.406)  |
| Unmarried  | -0.054*<br>(-0.100, -0.008)   | -0.061***<br>(-0.092, -0.029) | 0.919***<br>(0.882, 0.957) | -0.019***<br>(-0.028, -0.010) | -0.022<br>(-0.136, 0.091)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.233***<br>(-0.271, -0.194) | -0.184***<br>(-0.211, -0.158) | 0.816***<br>(0.788, 0.845) | -0.045***<br>(-0.052, -0.037) | -0.331***<br>(-0.428, -0.235)                                     |
| > 6 Members  | -0.436***<br>(-0.571, -0.300) | -0.344***<br>(-0.433, -0.255) | 0.681***<br>(0.600, 0.774) | -0.082***<br>(-0.107, -0.056) | -0.709***<br>(-1.076, -0.343)                                     |
| <b>Education (Less than College - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | 0.001<br>(-0.036, 0.038)      | 0<br>(-0.026, 0.025)          | 0.999<br>(0.968, 1.032)    | 0<br>(-0.007, 0.007)          | -0.001<br>(-0.089, 0.088)   |
| <b>Occupation (Natural resources - REF)</b>          |                               |                               |                            |                               |   |
| Hospitality services                                 | 0.045<br>(-0.047, 0.137)      | 0.045<br>(-0.019, 0.109)      | 1.069<br>(0.984, 1.161)    | 0.014<br>(-0.003, 0.032)      | -0.023<br>(-0.252, 0.206)   |
| Trade  | 0.138***<br>(0.056, 0.219)    | 0.089**<br>(0.032, 0.145)     | 1.083*<br>(1.007, 1.165)   | 0.017*<br>(0.002, 0.033)      | 0.236*<br>(0.034, 0.438)  |
| Professional services                                | 0.118**<br>(0.045, 0.191)     | 0.102***<br>(0.052, 0.153)    | 1.135***<br>(1.063, 1.211) | 0.028***<br>(0.014, 0.042)    | 0.09<br>(-0.091, 0.271)   |

|  |                              |                               |                            |                               |                            |
|--|------------------------------|-------------------------------|----------------------------|-------------------------------|----------------------------|
| Manufacturing                            | 0.045<br>(-0.037, 0.126)     | 0.060*<br>(0.004, 0.116)      | 1.103**<br>(1.026, 1.186)  | 0.021**<br>(0.006, 0.037)     | -0.067<br>(-0.268, 0.134)  |
| Other                                    | 0.06<br>(-0.014, 0.134)      | 0.075**<br>(0.024, 0.126)     | 1.130***<br>(1.057, 1.207) | 0.027***<br>(0.012, 0.041)    | -0.078<br>(-0.262, 0.105)  |
| <b>Year (2010 - REF)</b>                 |                              |                               |                            |                               |                            |
| 2011                                     | -0.001<br>(-0.076, 0.074)    | -0.076**<br>(-0.130, -0.023)  | 0.841***<br>(0.787, 0.898) | -0.039***<br>(-0.054, -0.024) | 0.324***<br>(0.146, 0.501) |
| 2012                                     | -0.112**<br>(-0.186, -0.039) | -0.165***<br>(-0.217, -0.113) | 0.764***<br>(0.716, 0.815) | -0.060***<br>(-0.075, -0.046) | 0.16<br>(-0.017, 0.338)    |
| 2013                                     | -0.036<br>(-0.111, 0.040)    | -0.087**<br>(-0.140, -0.033)  | 0.851***<br>(0.796, 0.908) | -0.037***<br>(-0.052, -0.022) | 0.178*<br>(0.000, 0.357)   |
| 2014                                     | -0.058<br>(-0.133, 0.016)    | -0.106***<br>(-0.159, -0.054) | 0.834***<br>(0.781, 0.891) | -0.041***<br>(-0.056, -0.026) | 0.133<br>(-0.044, 0.310)   |
| 2015                                     | -0.065<br>(-0.139, 0.009)    | -0.097***<br>(-0.149, -0.044) | 0.855***<br>(0.801, 0.912) | -0.036***<br>(-0.050, -0.021) | 0.084<br>(-0.092, 0.260)   |
| 2016                                     | -0.035<br>(-0.110, 0.041)    | -0.098***<br>(-0.151, -0.044) | 0.830***<br>(0.777, 0.887) | -0.042***<br>(-0.057, -0.027) | 0.226*<br>(0.047, 0.405)   |
| 2017                                     | 0.046<br>(-0.031, 0.123)     | -0.066*<br>(-0.120, -0.011)   | 0.831***<br>(0.777, 0.889) | -0.042***<br>(-0.057, -0.027) | 0.450***<br>(0.267, 0.632) |
| 2018                                     | 0.068<br>(-0.009, 0.145)     | -0.05<br>(-0.105, 0.005)      | 0.844***<br>(0.789, 0.903) | -0.038***<br>(-0.054, -0.023) | 0.482***<br>(0.299, 0.665) |
| 2019                                     | 0.061<br>(-0.020, 0.141)     | 0.01<br>(-0.047, 0.068)       | 0.965<br>(0.900, 1.035)    | -0.008<br>(-0.024, 0.008)     | 0.237*<br>(0.050, 0.424)   |
| <b>Region (Northeast - REF)</b>          |                              |                               |                            |                               |                            |
| North central, Midwest                   | 0.075**<br>(0.019, 0.131)    | 0.050*<br>(0.011, 0.090)      | 1.044<br>(0.993, 1.097)    | 0.01<br>(-0.002, 0.021)       | 0.146*<br>(0.009, 0.283)   |
| South                                    | 0.039<br>(-0.013, 0.092)     | 0.029<br>(-0.008, 0.066)      | 1.029<br>(0.982, 1.078)    | 0.006<br>(-0.004, 0.017)      | 0.075<br>(-0.053, 0.202)   |
| West                                     | -0.060*<br>(-0.116, -0.004)  | -0.03<br>(-0.069, 0.009)      | 0.983<br>(0.935, 1.033)    | -0.004<br>(-0.015, 0.007)     | -0.128<br>(-0.265, 0.010)  |
| <b>Income (Poor to low income - REF)</b> |                              |                               |                            |                               |                            |
| Middle to high income                    | -0.003<br>(-0.049, 0.043)    | -0.003<br>(-0.036, 0.029)     | 0.993<br>(0.953, 1.035)    | -0.002<br>(-0.011, 0.008)     | 0.008<br>(-0.105, 0.121)   |
| <b>Receipt of food stamp (No - REF)</b>  |                              |                               |                            |                               |                            |
| YES                                      | 0.218***<br>(0.154, 0.282)   | 0.154***<br>(0.107, 0.201)    | 1.149***<br>(1.086, 1.216) | 0.031***<br>(0.018, 0.044)    | 0.369***<br>(0.216, 0.522) |
| <b>Health status (Excellent - REF)</b>   |                              |                               |                            |                               |                            |
| Good                                     | 0.308***<br>(0.260, 0.356)   | 0.258***<br>(0.226, 0.290)    | 1.355***<br>(1.295, 1.417) | 0.064***<br>(0.055, 0.073)    | 0.504***<br>(0.374, 0.633) |
| Fair                                     | 0.932***<br>(0.865, 0.998)   | 0.678***<br>(0.630, 0.727)    | 1.957***<br>(1.844, 2.076) | 0.150***<br>(0.137, 0.163)    | 1.404***<br>(1.241, 1.566) |
| Poor                                     | 1.842***<br>(1.695, 1.990)   | 1.200***<br>(1.072, 1.329)    | 2.593***<br>(2.288, 2.938) | 0.219***<br>(0.189, 0.249)    | 2.694***<br>(2.386, 3.002) |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 4a. Regressions for the Use of Prescription Sleep Aids:  $\geq 46$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Prescription Sleep Aid = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 46$ h/week                                     | 0.002<br>(-0.014, 0.017)      | 0.006<br>(-0.025, 0.037)      | 1.015<br>(0.943, 1.092)    | 0.001<br>(-0.003, 0.005)      | -0.013<br>(-0.173, 0.147)   |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.104***<br>(0.081, 0.128)    | 0.306***<br>(0.255, 0.356)    | 2.206***<br>(1.897, 2.564) | 0.034***<br>(0.030, 0.039)    | 0.530**<br>(0.196, 0.864)   |
| $\geq 65$ years                                      | 0.069***<br>(0.033, 0.105)    | 0.252***<br>(0.181, 0.324)    | 1.993***<br>(1.642, 2.418) | 0.028***<br>(0.020, 0.037)    | 0.285<br>(-0.139, 0.709)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.126***<br>(-0.144, -0.109) | -0.313***<br>(-0.349, -0.277) | 0.457***<br>(0.413, 0.505) | -0.037***<br>(-0.040, -0.033) | -0.405***<br>(-0.627, -0.184)                                       |
| Other  | -0.086***<br>(-0.109, -0.063) | -0.171***<br>(-0.218, -0.123) | 0.687***<br>(0.610, 0.775) | -0.021***<br>(-0.026, -0.015) | -0.544***<br>(-0.809, -0.279)                                       |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | 0.012<br>(-0.001, 0.025)      | 0.044**<br>(0.017, 0.070)     | 1.109**<br>(1.042, 1.181)  | 0.006**<br>(0.002, 0.009)     | -0.048<br>(-0.184, 0.089)   |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.029***<br>(0.012, 0.045)    | 0.053**<br>(0.021, 0.085)     | 1.127**<br>(1.047, 1.212)  | 0.007**<br>(0.003, 0.011)     | 0.071<br>(-0.087, 0.229)  |
| Unmarried  | 0.002<br>(-0.015, 0.019)      | 0.009<br>(-0.026, 0.043)      | 1.034<br>(0.954, 1.121)    | 0.002<br>(-0.003, 0.006)      | -0.097<br>(-0.274, 0.079)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.072***<br>(-0.086, -0.058) | -0.171***<br>(-0.200, -0.142) | 0.670***<br>(0.623, 0.720) | -0.021***<br>(-0.025, -0.018) | -0.235**<br>(-0.395, -0.075)  |
| > 6 Members  | -0.101***<br>(-0.151, -0.051) | -0.308***<br>(-0.419, -0.198) | 0.437***<br>(0.311, 0.615) | -0.037***<br>(-0.048, -0.027) | 0.046<br>(-0.711, 0.803)  |
| <b>Education (Less than College - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | 0.026***<br>(0.013, 0.040)    | 0.087***<br>(0.059, 0.114)    | 1.236***<br>(1.158, 1.320) | 0.012***<br>(0.008, 0.015)    | -0.072<br>(-0.215, 0.072)   |
| <b>Occupation (Natural resources - REF)</b>          |                               |                               |                            |                               |   |
| Hospitality services                                 | 0.02<br>(-0.014, 0.054)       | 0.017<br>(-0.052, 0.087)      | 1.032<br>(0.865, 1.231)    | 0.002<br>(-0.007, 0.010)      | 0.2<br>(-0.185, 0.585)  |
| Trade  | 0.033*<br>(0.003, 0.063)      | 0.057<br>(-0.004, 0.118)      | 1.114<br>(0.956, 1.298)    | 0.005<br>(-0.002, 0.013)      | 0.27<br>(-0.063, 0.603)   |
| Professional services                                | 0.061***                      | 0.121***                      | 1.302***                   | 0.014***                      | 0.316*  |

|  |                 |                 |                |                  |                  |
|--|-----------------|-----------------|----------------|------------------|------------------|
|  | (0.034, 0.088)  | (0.066, 0.175)  | (1.137, 1.491) | (0.008, 0.021)   | (0.022, 0.610)   |
| Manufacturing                            | 0.046**         | 0.091**         | 1.222**        | 0.011**          | 0.273            |
|  | (0.016, 0.076)  | (0.031, 0.152)  | (1.052, 1.419) | (0.003, 0.018)   | (-0.054, 0.601)  |
| Other                                    | 0.039**         | 0.092**         | 1.232**        | 0.011**          | 0.139            |
|  | (0.012, 0.067)  | (0.037, 0.147)  | (1.074, 1.414) | (0.004, 0.018)   | (-0.161, 0.440)  |
| <b>Year (2010 - REF)</b>                 |                 |                 |                |                  |                  |
| 2011                                     | 0.011           | -0.008          | 0.964          | -0.002           | 0.273            |
|  | (-0.017, 0.038) | (-0.064, 0.048) | (0.847, 1.097) | (-0.010, 0.005)  | (-0.007, 0.553)  |
| 2012                                     | -0.004          | -0.051          | 0.856*         | -0.009*          | 0.359*           |
|  | (-0.031, 0.023) | (-0.106, 0.004) | (0.751, 0.975) | (-0.016, -0.001) | (0.076, 0.642)   |
| 2013                                     | 0.009           | -0.021          | 0.922          | -0.005           | 0.368*           |
|  | (-0.019, 0.036) | (-0.077, 0.035) | (0.808, 1.051) | (-0.012, 0.003)  | (0.083, 0.654)   |
| 2014                                     | -0.005          | -0.02           | 0.953          | -0.003           | 0.006            |
|  | (-0.033, 0.022) | (-0.075, 0.036) | (0.837, 1.084) | (-0.010, 0.005)  | (-0.274, 0.287)  |
| 2015                                     | -0.019          | -0.049          | 0.885          | -0.007           | 0.009            |
|  | (-0.046, 0.008) | (-0.104, 0.006) | (0.777, 1.009) | (-0.014, 0.001)  | (-0.274, 0.292)  |
| 2016                                     | -0.012          | -0.041          | 0.896          | -0.006           | 0.114            |
|  | (-0.039, 0.016) | (-0.096, 0.015) | (0.786, 1.022) | (-0.014, 0.001)  | (-0.173, 0.400)  |
| 2017                                     | 0.022           | 0.011           | 0.985          | -0.001           | 0.394**          |
|  | (-0.007, 0.050) | (-0.046, 0.068) | (0.864, 1.122) | (-0.009, 0.007)  | (0.110, 0.678)   |
| 2018                                     | 0.018           | 0.013           | 0.996          | 0                | 0.295*           |
|  | (-0.011, 0.046) | (-0.044, 0.070) | (0.874, 1.134) | (-0.008, 0.007)  | (0.011, 0.578)   |
| 2019                                     | 0.033*          | 0.082**         | 1.192**        | 0.011**          | 0.044            |
|  | (0.003, 0.063)  | (0.024, 0.141)  | (1.046, 1.358) | (0.003, 0.020)   | (-0.238, 0.326)  |
| <b>Region (Northeast - REF)</b>          |                 |                 |                |                  |                  |
| North central, Midwest                   | 0.028**         | 0.060**         | 1.151**        | 0.008**          | 0.075            |
|  | (0.008, 0.049)  | (0.018, 0.101)  | (1.045, 1.268) | (0.003, 0.014)   | (-0.136, 0.285)  |
| South                                    | 0.016           | 0.034           | 1.091          | 0.005            | 0.037            |
|  | (-0.004, 0.035) | (-0.004, 0.073) | (0.995, 1.196) | (-0.000, 0.010)  | (-0.165, 0.238)  |
| West                                     | -0.01           | 0.004           | 1.037          | 0.002            | -0.229*          |
|  | (-0.031, 0.011) | (-0.037, 0.045) | (0.940, 1.144) | (-0.003, 0.007)  | (-0.444, -0.014) |
| <b>Income (Poor to low income - REF)</b> |                 |                 |                |                  |                  |
| Middle to high income                    | 0.002           | 0.015           | 1.042          | 0.002            | -0.082           |
|  | (-0.015, 0.019) | (-0.020, 0.051) | (0.958, 1.134) | (-0.002, 0.007)  | (-0.270, 0.105)  |
| <b>Receipt of food stamp (No - REF)</b>  |                 |                 |                |                  |                  |
| YES                                      | 0.028*          | 0.047           | 1.093          | 0.005            | 0.239            |
|  | (0.005, 0.052)  | (-0.004, 0.098) | (0.971, 1.230) | (-0.002, 0.012)  | (-0.024, 0.503)  |
| <b>Health status (Excellent - REF)</b>   |                 |                 |                |                  |                  |
| Good                                     | 0.058***        | 0.154***        | 1.467***       | 0.019***         | 0.201            |
|  | (0.040, 0.075)  | (0.118, 0.191)  | (1.331, 1.616) | (0.015, 0.023)   | (-0.013, 0.415)  |
| Fair                                     | 0.156***        | 0.307***        | 1.949***       | 0.037***         | 0.850***         |
|  | (0.132, 0.181)  | (0.257, 0.358)  | (1.728, 2.199) | (0.030, 0.044)   | (0.585, 1.114)   |
| Poor                                     | 0.258***        | 0.470***        | 2.611***       | 0.061***         | 1.069***         |
|  | (0.204, 0.313)  | (0.359, 0.581)  | (2.104, 3.241) | (0.043, 0.080)   | (0.606, 1.532)   |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 4b. Regressions for the Use of Medications with Sedative Properties:  $\geq 46$  h/week

| Variable                                    | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Sedative Medications = YES) |
|---|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                      |                               |                               |                            |                               |   |
| ≥ 46 h/week                                 | 0.015<br>(-0.027, 0.058)      | 0.016<br>(-0.014, 0.047)      | 1.023<br>(0.985, 1.063)    | 0.005<br>(-0.003, 0.014)      | 0<br>(-0.104, 0.104)  |
| <b>Age (18-26years - REF)</b>               |                               |                               |                            |                               |   |
| 27-64 years                                 | 0.295***<br>(0.231, 0.359)    | 0.236***<br>(0.193, 0.279)    | 1.291***<br>(1.217, 1.371) | 0.055***<br>(0.042, 0.067)    | 0.523***<br>(0.352, 0.694)  |
| ≥ 65 years                                  | 0.381***<br>(0.284, 0.478)    | 0.337***<br>(0.269, 0.404)    | 1.498***<br>(1.374, 1.632) | 0.088***<br>(0.069, 0.107)    | 0.498***<br>(0.261, 0.735)  |
| <b>Race (White - REF)</b>                   |                               |                               |                            |                               |   |
| Black                                       | -0.200***<br>(-0.248, -0.153) | -0.088***<br>(-0.120, -0.056) | 0.975<br>(0.935, 1.016)    | -0.006<br>(-0.015, 0.004)     | -0.501***<br>(-0.615, -0.388)                                     |
| Other                                       | -0.214***<br>(-0.277, -0.152) | -0.137***<br>(-0.180, -0.094) | 0.894***<br>(0.845, 0.945) | -0.025***<br>(-0.037, -0.012) | -0.451***<br>(-0.608, -0.294)                                     |
| <b>Sex (Male - REF)</b>                     |                               |                               |                            |                               |   |
| Female                                      | -0.006<br>(-0.042, 0.030)     | 0.004<br>(-0.021, 0.029)      | 1.014<br>(0.982, 1.047)    | 0.003<br>(-0.004, 0.010)      | -0.039<br>(-0.127, 0.049)   |
| <b>Marital status (Married - REF)</b>       |                               |                               |                            |                               |   |
| Divorced                                    | 0.158***<br>(0.114, 0.203)    | 0.086***<br>(0.055, 0.118)    | 1.061**<br>(1.020, 1.102)  | 0.013**<br>(0.004, 0.022)     | 0.301***<br>(0.196, 0.406)  |
| Unmarried                                   | -0.054*<br>(-0.100, -0.008)   | -0.060***<br>(-0.092, -0.028) | 0.919***<br>(0.882, 0.957) | -0.019***<br>(-0.028, -0.010) | -0.022<br>(-0.136, 0.091)   |
| <b>Family size (≤ 3 Members REF)</b>        |                               |                               |                            |                               |   |
| 4-6 Members                                 | -0.233***<br>(-0.271, -0.194) | -0.184***<br>(-0.211, -0.158) | 0.816***<br>(0.788, 0.845) | -0.045***<br>(-0.052, -0.037) | -0.331***<br>(-0.428, -0.235)                                     |
| > 6 Members                                 | -0.435***<br>(-0.570, -0.299) | -0.343***<br>(-0.432, -0.254) | 0.682***<br>(0.600, 0.775) | -0.081***<br>(-0.106, -0.056) | -0.708***<br>(-1.075, -0.342)                                     |
| <b>Education (Less than College - REF)</b>  |                               |                               |                            |                               |   |
| Some college or more                        | 0.001<br>(-0.036, 0.037)      | -0.001<br>(-0.027, 0.025)     | 0.999<br>(0.967, 1.032)    | 0<br>(-0.007, 0.007)          | -0.001<br>(-0.089, 0.088)   |
| <b>Occupation (Natural resources - REF)</b> |                               |                               |                            |                               |   |
| Hospitality services                        | 0.043<br>(-0.049, 0.135)      | 0.042<br>(-0.021, 0.106)      | 1.066<br>(0.982, 1.157)    | 0.014<br>(-0.004, 0.032)      | -0.026<br>(-0.254, 0.202)   |
| Trade                                       | 0.137***<br>(0.056, 0.219)    | 0.088**<br>(0.032, 0.145)     | 1.083*<br>(1.007, 1.165)   | 0.017*<br>(0.002, 0.033)      | 0.234*<br>(0.032, 0.436)  |
| Professional services                       | 0.118**                       | 0.102***                      | 1.136***                   | 0.028***                      | 0.089   |



|  |                  |                  |                |                  |                 |
|--|------------------|------------------|----------------|------------------|-----------------|
|  | (0.046, 0.191)   | (0.052, 0.153)   | (1.064, 1.212) | (0.014, 0.042)   | (-0.092, 0.270) |
| Manufacturing                            | 0.046            | 0.062*           | 1.106**        | 0.022**          | -0.066          |
|  | (-0.035, 0.128)  | (0.006, 0.118)   | (1.028, 1.189) | (0.006, 0.038)   | (-0.267, 0.135) |
| Other                                    | 0.061            | 0.076**          | 1.131***       | 0.027***         | -0.078          |
|  | (-0.012, 0.135)  | (0.025, 0.127)   | (1.059, 1.209) | (0.013, 0.041)   | (-0.262, 0.105) |
| <b>Year (2010 - REF)</b>                 |                  |                  |                |                  |                 |
| 2011                                     | -0.001           | -0.076**         | 0.841***       | -0.039***        | 0.324***        |
|  | (-0.076, 0.074)  | (-0.129, -0.023) | (0.788, 0.898) | (-0.054, -0.024) | (0.146, 0.502)  |
| 2012                                     | -0.112**         | -0.165***        | 0.764***       | -0.060***        | 0.161           |
|  | (-0.186, -0.038) | (-0.216, -0.113) | (0.716, 0.815) | (-0.075, -0.046) | (-0.017, 0.338) |
| 2013                                     | -0.035           | -0.086**         | 0.851***       | -0.037***        | 0.178           |
|  | (-0.111, 0.040)  | (-0.140, -0.033) | (0.797, 0.909) | (-0.052, -0.022) | (-0.000, 0.356) |
| 2014                                     | -0.058           | -0.106***        | 0.834***       | -0.041***        | 0.133           |
|  | (-0.133, 0.016)  | (-0.159, -0.054) | (0.781, 0.890) | (-0.056, -0.026) | (-0.044, 0.310) |
| 2015                                     | -0.065           | -0.097***        | 0.854***       | -0.036***        | 0.084           |
|  | (-0.139, 0.009)  | (-0.149, -0.044) | (0.801, 0.912) | (-0.050, -0.021) | (-0.092, 0.260) |
| 2016                                     | -0.035           | -0.098***        | 0.830***       | -0.042***        | 0.227*          |
|  | (-0.110, 0.041)  | (-0.151, -0.044) | (0.777, 0.887) | (-0.057, -0.027) | (0.048, 0.406)  |
| 2017                                     | 0.046            | -0.066*          | 0.831***       | -0.042***        | 0.450***        |
|  | (-0.031, 0.123)  | (-0.120, -0.011) | (0.776, 0.888) | (-0.057, -0.027) | (0.267, 0.632)  |
| 2018                                     | 0.068            | -0.05            | 0.844***       | -0.039***        | 0.482***        |
|  | (-0.010, 0.145)  | (-0.105, 0.004)  | (0.788, 0.902) | (-0.054, -0.023) | (0.299, 0.664)  |
| 2019                                     | 0.061            | 0.01             | 0.965          | -0.008           | 0.237*          |
|  | (-0.020, 0.141)  | (-0.047, 0.068)  | (0.900, 1.034) | (-0.024, 0.008)  | (0.050, 0.423)  |
| <b>Region (Northeast - REF)</b>          |                  |                  |                |                  |                 |
| North central, Midwest                   | 0.075**          | 0.051*           | 1.044          | 0.01             | 0.147*          |
|  | (0.019, 0.132)   | (0.011, 0.090)   | (0.994, 1.098) | (-0.001, 0.021)  | (0.010, 0.284)  |
| South                                    | 0.04             | 0.03             | 1.031          | 0.007            | 0.076           |
|  | (-0.012, 0.093)  | (-0.006, 0.067)  | (0.984, 1.080) | (-0.004, 0.017)  | (-0.052, 0.203) |
| West                                     | -0.059*          | -0.029           | 0.984          | -0.003           | -0.127          |
|  | (-0.115, -0.003) | (-0.068, 0.010)  | (0.937, 1.035) | (-0.014, 0.007)  | (-0.264, 0.010) |
| <b>Income (Poor to low income - REF)</b> |                  |                  |                |                  |                 |
| Middle to high income                    | -0.003           | -0.003           | 0.994          | -0.001           | 0.01            |
|  | (-0.049, 0.043)  | (-0.035, 0.030)  | (0.954, 1.035) | (-0.010, 0.008)  | (-0.102, 0.122) |
| <b>Receipt of food stamp (No - REF)</b>  |                  |                  |                |                  |                 |
| YES                                      | 0.217***         | 0.153***         | 1.148***       | 0.031***         | 0.367***        |
|  | (0.153, 0.281)   | (0.106, 0.200)   | (1.085, 1.214) | (0.018, 0.044)   | (0.214, 0.520)  |
| <b>Health status (Excellent - REF)</b>   |                  |                  |                |                  |                 |
| Good                                     | 0.308***         | 0.258***         | 1.355***       | 0.064***         | 0.503***        |
|  | (0.260, 0.356)   | (0.227, 0.290)   | (1.296, 1.417) | (0.055, 0.073)   | (0.374, 0.633)  |
| Fair                                     | 0.932***         | 0.678***         | 1.957***       | 0.150***         | 1.403***        |
|  | (0.865, 0.998)   | (0.630, 0.727)   | (1.844, 2.076) | (0.137, 0.163)   | (1.241, 1.566)  |
| Poor                                     | 1.842***         | 1.200***         | 2.593***       | 0.219***         | 2.694***        |
|  | (1.695, 1.990)   | (1.072, 1.329)   | (2.288, 2.938) | (0.189, 0.249)   | (2.386, 3.002)  |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

#### Appendix 5a. Regressions for the Use of Prescription Sleep Aid: $\geq 66$ h/week

| Variables                                   | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Prescription Sleep Aid = YES) |
|---|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                      |                               |                               |                            |                               |   |
| ≥ 66 h/week                                 | 0.026<br>(-0.019, 0.070)      | 0.039<br>(-0.050, 0.128)      | 1.077<br>(0.879, 1.320)    | 0.004<br>(-0.008, 0.017)      | 0.21<br>(-0.232, 0.652)   |
| <b>Age (18-26years - REF)</b>               |                               |                               |                            |                               |   |
| 27-64 years                                 | 0.104***<br>(0.081, 0.128)    | 0.306***<br>(0.255, 0.356)    | 2.207***<br>(1.898, 2.566) | 0.034***<br>(0.030, 0.039)    | 0.527**<br>(0.194, 0.861)   |
| ≥ 65 years                                  | 0.069***<br>(0.033, 0.104)    | 0.252***<br>(0.180, 0.324)    | 1.992***<br>(1.642, 2.417) | 0.028***<br>(0.020, 0.037)    | 0.277<br>(-0.148, 0.701)  |
| <b>Race (White - REF)</b>                   |                               |                               |                            |                               |   |
| Black                                       | -0.127***<br>(-0.144, -0.109) | -0.313***<br>(-0.349, -0.277) | 0.456***<br>(0.413, 0.505) | -0.037***<br>(-0.040, -0.033) | -0.402***<br>(-0.624, -0.181)                                       |
| Other                                       | -0.086***<br>(-0.109, -0.064) | -0.171***<br>(-0.218, -0.123) | 0.687***<br>(0.609, 0.775) | -0.021***<br>(-0.026, -0.015) | -0.547***<br>(-0.812, -0.282)                                       |
| <b>Sex (Male - REF)</b>                     |                               |                               |                            |                               |   |
| Female                                      | 0.012<br>(-0.001, 0.025)      | 0.044**<br>(0.017, 0.070)     | 1.109**<br>(1.042, 1.181)  | 0.006**<br>(0.002, 0.009)     | -0.042<br>(-0.177, 0.094)   |
| <b>Marital status (Married - REF)</b>       |                               |                               |                            |                               |   |
| Divorced                                    | 0.029***<br>(0.012, 0.045)    | 0.053**<br>(0.021, 0.085)     | 1.127**<br>(1.047, 1.212)  | 0.007**<br>(0.003, 0.011)     | 0.07<br>(-0.088, 0.229)   |
| Unmarried                                   | 0.002<br>(-0.015, 0.019)      | 0.009<br>(-0.025, 0.043)      | 1.034<br>(0.954, 1.121)    | 0.002<br>(-0.003, 0.007)      | -0.095<br>(-0.272, 0.081)   |
| <b>Family size (≤ 3 Members REF)</b>        |                               |                               |                            |                               |   |
| 4-6 Members                                 | -0.072***<br>(-0.086, -0.058) | -0.171***<br>(-0.200, -0.142) | 0.670***<br>(0.623, 0.721) | -0.021***<br>(-0.025, -0.018) | -0.235**<br>(-0.395, -0.075)  |
| > 6 Members                                 | -0.101***<br>(-0.151, -0.051) | -0.308***<br>(-0.419, -0.198) | 0.437***<br>(0.311, 0.615) | -0.037***<br>(-0.048, -0.027) | 0.053<br>(-0.705, 0.810)  |
| <b>Education (Less than College - REF)</b>  |                               |                               |                            |                               |   |
| Some college or more                        | 0.027***<br>(0.013, 0.040)    | 0.087***<br>(0.060, 0.115)    | 1.237***<br>(1.159, 1.321) | 0.012***<br>(0.008, 0.016)    | -0.071<br>(-0.214, 0.072)   |
| <b>Occupation (Natural resources - REF)</b> |                               |                               |                            |                               |   |
| Hospitality services                        | 0.02<br>(-0.013, 0.054)       | 0.018<br>(-0.052, 0.087)      | 1.033<br>(0.866, 1.232)    | 0.002<br>(-0.007, 0.010)      | 0.201<br>(-0.184, 0.586)  |
| Trade                                       | 0.033*<br>(0.003, 0.063)      | 0.057<br>(-0.004, 0.118)      | 1.115<br>(0.957, 1.298)    | 0.006<br>(-0.002, 0.013)      | 0.27<br>(-0.063, 0.603)   |
| Professional services                       | 0.061***<br>(0.034, 0.088)    | 0.121***<br>(0.066, 0.175)    | 1.302***<br>(1.137, 1.491) | 0.014***<br>(0.008, 0.021)    | 0.314*<br>(0.020, 0.608)  |

|  |                 |                 |                |                  |                  |
|--|-----------------|-----------------|----------------|------------------|------------------|
| Manufacturing                            | 0.047**         | 0.092**         | 1.224**        | 0.011**          | 0.274            |
|  | (0.017, 0.077)  | (0.031, 0.152)  | (1.053, 1.421) | (0.003, 0.018)   | (-0.054, 0.601)  |
| Other                                    | 0.040**         | 0.092**         | 1.232**        | 0.011**          | 0.135            |
|  | (0.012, 0.067)  | (0.036, 0.147)  | (1.074, 1.414) | (0.004, 0.018)   | (-0.165, 0.435)  |
| <b>Year (2010 - REF)</b>                 |                 |                 |                |                  |                  |
| 2011                                     | 0.011           | -0.008          | 0.963          | -0.002           | 0.269            |
|  | (-0.017, 0.038) | (-0.064, 0.047) | (0.847, 1.096) | (-0.010, 0.005)  | (-0.011, 0.549)  |
| 2012                                     | -0.004          | -0.051          | 0.856*         | -0.009*          | 0.360*           |
|  | (-0.031, 0.023) | (-0.106, 0.003) | (0.751, 0.975) | (-0.016, -0.001) | (0.077, 0.643)   |
| 2013                                     | 0.009           | -0.021          | 0.922          | -0.005           | 0.369*           |
|  | (-0.019, 0.036) | (-0.077, 0.035) | (0.808, 1.051) | (-0.012, 0.003)  | (0.084, 0.655)   |
| 2014                                     | -0.005          | -0.02           | 0.953          | -0.003           | 0.004            |
|  | (-0.033, 0.022) | (-0.075, 0.036) | (0.837, 1.084) | (-0.010, 0.005)  | (-0.277, 0.284)  |
| 2015                                     | -0.019          | -0.049          | 0.886          | -0.007           | 0.006            |
|  | (-0.046, 0.009) | (-0.104, 0.006) | (0.778, 1.009) | (-0.014, 0.001)  | (-0.277, 0.289)  |
| 2016                                     | -0.011          | -0.04           | 0.897          | -0.006           | 0.112            |
|  | (-0.039, 0.016) | (-0.096, 0.016) | (0.786, 1.023) | (-0.014, 0.001)  | (-0.174, 0.399)  |
| 2017                                     | 0.022           | 0.011           | 0.985          | -0.001           | 0.396**          |
|  | (-0.007, 0.050) | (-0.046, 0.068) | (0.865, 1.123) | (-0.009, 0.007)  | (0.112, 0.680)   |
| 2018                                     | 0.018           | 0.013           | 0.996          | 0                | 0.296*           |
|  | (-0.011, 0.046) | (-0.044, 0.070) | (0.875, 1.135) | (-0.008, 0.008)  | (0.012, 0.579)   |
| 2019                                     | 0.033*          | 0.083**         | 1.192**        | 0.011**          | 0.047            |
|  | (0.003, 0.063)  | (0.024, 0.141)  | (1.047, 1.358) | (0.003, 0.020)   | (-0.235, 0.329)  |
| <b>Region (Northeast - REF)</b>          |                 |                 |                |                  |                  |
| North central, Midwest                   | 0.028**         | 0.060**         | 1.151**        | 0.008**          | 0.072            |
|  | (0.008, 0.049)  | (0.018, 0.101)  | (1.045, 1.269) | (0.003, 0.014)   | (-0.138, 0.283)  |
| South                                    | 0.015           | 0.034           | 1.091          | 0.005            | 0.035            |
|  | (-0.004, 0.035) | (-0.005, 0.073) | (0.994, 1.196) | (-0.000, 0.010)  | (-0.166, 0.236)  |
| West                                     | -0.01           | 0.004           | 1.036          | 0.002            | -0.227*          |
|  | (-0.031, 0.011) | (-0.037, 0.045) | (0.939, 1.143) | (-0.003, 0.007)  | (-0.441, -0.012) |
| <b>Income (Poor to low income - REF)</b> |                 |                 |                |                  |                  |
| Middle to high income                    | 0.002           | 0.015           | 1.043          | 0.002            | -0.083           |
|  | (-0.015, 0.019) | (-0.020, 0.051) | (0.959, 1.135) | (-0.002, 0.007)  | (-0.270, 0.104)  |
| <b>Receipt of food stamp (No - REF)</b>  |                 |                 |                |                  |                  |
| YES                                      | 0.028*          | 0.046           | 1.092          | 0.005            | 0.233            |
|  | (0.005, 0.052)  | (-0.005, 0.097) | (0.970, 1.229) | (-0.002, 0.012)  | (-0.031, 0.497)  |
| <b>Health status (Excellent - REF)</b>   |                 |                 |                |                  |                  |
| Good                                     | 0.058***        | 0.154***        | 1.466***       | 0.019***         | 0.204            |
|  | (0.040, 0.075)  | (0.118, 0.191)  | (1.331, 1.616) | (0.015, 0.023)   | (-0.010, 0.417)  |
| Fair                                     | 0.156***        | 0.307***        | 1.948***       | 0.037***         | 0.853***         |
|  | (0.132, 0.181)  | (0.256, 0.358)  | (1.727, 2.197) | (0.030, 0.044)   | (0.589, 1.117)   |
| Poor                                     | 0.258***        | 0.470***        | 2.610***       | 0.061***         | 1.073***         |
|  | (0.204, 0.313)  | (0.359, 0.581)  | (2.103, 3.239) | (0.043, 0.080)   | (0.610, 1.537)   |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 5b. Regressions for the Use of Medications with Sedative Properties:  $\geq 66$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Sedative Medications = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 66$ h/week                                     | 0.11<br>(-0.011, 0.230)       | 0.058<br>(-0.028, 0.144)      | 1.034<br>(0.930, 1.149)    | 0.007<br>(-0.016, 0.031)      | 0.244<br>(-0.043, 0.531)  |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.295***<br>(0.231, 0.359)    | 0.237***<br>(0.193, 0.280)    | 1.293***<br>(1.218, 1.372) | 0.055***<br>(0.043, 0.067)    | 0.522***<br>(0.351, 0.693)  |
| $\geq 65$ years                                      | 0.380***<br>(0.283, 0.477)    | 0.336***<br>(0.269, 0.404)    | 1.498***<br>(1.374, 1.633) | 0.088***<br>(0.069, 0.107)    | 0.494***<br>(0.257, 0.731)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.201***<br>(-0.248, -0.154) | -0.088***<br>(-0.121, -0.056) | 0.974<br>(0.934, 1.016)    | -0.006<br>(-0.015, 0.003)     | -0.502***<br>(-0.615, -0.388)                                     |
| Other  | -0.215***<br>(-0.277, -0.153) | -0.138***<br>(-0.180, -0.095) | 0.893***<br>(0.844, 0.945) | -0.025***<br>(-0.037, -0.013) | -0.452***<br>(-0.609, -0.295)                                     |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | -0.005<br>(-0.041, 0.030)     | 0.004<br>(-0.022, 0.029)      | 1.012<br>(0.980, 1.045)    | 0.003<br>(-0.004, 0.010)      | -0.034<br>(-0.121, 0.053)   |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.158***<br>(0.114, 0.203)    | 0.086***<br>(0.055, 0.118)    | 1.061**<br>(1.020, 1.103)  | 0.013**<br>(0.004, 0.022)     | 0.301***<br>(0.196, 0.406)  |
| Unmarried  | -0.054*<br>(-0.099, -0.008)   | -0.060***<br>(-0.092, -0.028) | 0.919***<br>(0.882, 0.957) | -0.019***<br>(-0.028, -0.010) | -0.021<br>(-0.134, 0.093)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.232***<br>(-0.271, -0.194) | -0.184***<br>(-0.211, -0.157) | 0.816***<br>(0.788, 0.845) | -0.045***<br>(-0.052, -0.037) | -0.330***<br>(-0.427, -0.233)                                     |
| > 6 Members  | -0.435***<br>(-0.571, -0.300) | -0.343***<br>(-0.432, -0.254) | 0.682***<br>(0.600, 0.774) | -0.081***<br>(-0.107, -0.056) | -0.707***<br>(-1.074, -0.341)                                     |
| <b>Education (Less than College - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | 0.002<br>(-0.035, 0.038)      | 0<br>(-0.026, 0.026)          | 1<br>(0.968, 1.033)        | 0<br>(-0.007, 0.007)          | 0.001<br>(-0.087, 0.090)  |
| <b>Occupation (Natural resources - REF)</b>          |                               |                               |                            |                               |   |
| Hospitality services                                 | 0.044<br>(-0.047, 0.136)      | 0.043<br>(-0.021, 0.106)      | 1.065<br>(0.981, 1.156)    | 0.014<br>(-0.004, 0.031)      | -0.019<br>(-0.247, 0.209)   |

|  |                              |                               |                            |                               |                            |
|--|------------------------------|-------------------------------|----------------------------|-------------------------------|----------------------------|
| Trade                                    | 0.139***<br>(0.057, 0.220)   | 0.088**<br>(0.032, 0.145)     | 1.082*<br>(1.006, 1.164)   | 0.017*<br>(0.001, 0.033)      | 0.240*<br>(0.038, 0.442)   |
| Professional services                    | 0.119**<br>(0.046, 0.192)    | 0.102***<br>(0.052, 0.153)    | 1.134***<br>(1.063, 1.211) | 0.028***<br>(0.014, 0.042)    | 0.094<br>(-0.087, 0.275)   |
| Manufacturing                            | 0.048<br>(-0.033, 0.130)     | 0.063*<br>(0.006, 0.119)      | 1.105**<br>(1.028, 1.189)  | 0.022**<br>(0.006, 0.038)     | -0.058<br>(-0.259, 0.143)  |
| Other                                    | 0.062<br>(-0.012, 0.136)     | 0.076**<br>(0.025, 0.127)     | 1.130***<br>(1.058, 1.207) | 0.027***<br>(0.012, 0.041)    | -0.074<br>(-0.257, 0.110)  |
| <b>Year (2010 - REF)</b>                 |                              |                               |                            |                               |                            |
| 2011                                     | -0.002<br>(-0.077, 0.073)    | -0.077**<br>(-0.130, -0.023)  | 0.841***<br>(0.787, 0.898) | -0.039***<br>(-0.054, -0.024) | 0.323***<br>(0.145, 0.501) |
| 2012                                     | -0.112**<br>(-0.186, -0.038) | -0.165***<br>(-0.216, -0.113) | 0.764***<br>(0.716, 0.815) | -0.060***<br>(-0.075, -0.046) | 0.161<br>(-0.016, 0.338)   |
| 2013                                     | -0.035<br>(-0.110, 0.040)    | -0.086**<br>(-0.140, -0.033)  | 0.851***<br>(0.797, 0.909) | -0.037***<br>(-0.052, -0.022) | 0.180*<br>(0.002, 0.358)   |
| 2014                                     | -0.058<br>(-0.133, 0.016)    | -0.106***<br>(-0.159, -0.053) | 0.834***<br>(0.781, 0.891) | -0.041***<br>(-0.056, -0.026) | 0.134<br>(-0.044, 0.311)   |
| 2015                                     | -0.065<br>(-0.139, 0.010)    | -0.096***<br>(-0.149, -0.044) | 0.855***<br>(0.801, 0.912) | -0.036***<br>(-0.050, -0.021) | 0.085<br>(-0.091, 0.261)   |
| 2016                                     | -0.034<br>(-0.109, 0.042)    | -0.097***<br>(-0.150, -0.044) | 0.831***<br>(0.777, 0.887) | -0.042***<br>(-0.057, -0.027) | 0.229*<br>(0.049, 0.408)   |
| 2017                                     | 0.047<br>(-0.030, 0.123)     | -0.065*<br>(-0.120, -0.011)   | 0.831***<br>(0.777, 0.889) | -0.042***<br>(-0.057, -0.027) | 0.451***<br>(0.269, 0.634) |
| 2018                                     | 0.068<br>(-0.009, 0.145)     | -0.05<br>(-0.105, 0.005)      | 0.844***<br>(0.789, 0.903) | -0.038***<br>(-0.054, -0.023) | 0.483***<br>(0.300, 0.666) |
| 2019                                     | 0.061<br>(-0.019, 0.142)     | 0.01<br>(-0.047, 0.068)       | 0.965<br>(0.900, 1.035)    | -0.008<br>(-0.024, 0.008)     | 0.238*<br>(0.051, 0.424)   |
| <b>Region (Northeast - REF)</b>          |                              |                               |                            |                               |                            |
| North central, Midwest                   | 0.075**<br>(0.019, 0.132)    | 0.051*<br>(0.012, 0.091)      | 1.045<br>(0.994, 1.098)    | 0.01<br>(-0.001, 0.021)       | 0.146*<br>(0.010, 0.283)   |
| South                                    | 0.04<br>(-0.013, 0.092)      | 0.03<br>(-0.007, 0.067)       | 1.031<br>(0.984, 1.080)    | 0.007<br>(-0.004, 0.017)      | 0.075<br>(-0.053, 0.203)   |
| West                                     | -0.059*<br>(-0.115, -0.003)  | -0.029<br>(-0.068, 0.010)     | 0.984<br>(0.936, 1.034)    | -0.004<br>(-0.014, 0.007)     | -0.127<br>(-0.264, 0.010)  |
| <b>Income (Poor to low income - REF)</b> |                              |                               |                            |                               |                            |
| Middle to high income                    | -0.002<br>(-0.048, 0.044)    | -0.002<br>(-0.034, 0.031)     | 0.995<br>(0.956, 1.037)    | -0.001<br>(-0.010, 0.008)     | 0.008<br>(-0.104, 0.120)   |
| <b>Receipt of food stamp (No - REF)</b>  |                              |                               |                            |                               |                            |
| YES                                      | 0.217***<br>(0.153, 0.281)   | 0.153***<br>(0.105, 0.200)    | 1.147***<br>(1.084, 1.214) | 0.031***<br>(0.018, 0.044)    | 0.365***<br>(0.212, 0.518) |
| <b>Health status (Excellent - REF)</b>   |                              |                               |                            |                               |                            |
| Good                                     | 0.308***<br>(0.260, 0.356)   | 0.258***<br>(0.226, 0.290)    | 1.355***<br>(1.295, 1.417) | 0.064***<br>(0.055, 0.073)    | 0.505***<br>(0.375, 0.634) |
| Fair                                     | 0.931***<br>(0.864, 0.998)   | 0.678***<br>(0.629, 0.726)    | 1.956***<br>(1.843, 2.075) | 0.150***<br>(0.137, 0.163)    | 1.403***<br>(1.241, 1.566) |
| Poor                                     | 1.842***<br>(1.694, 1.989)   | 1.200***<br>(1.071, 1.329)    | 2.592***<br>(2.287, 2.937) | 0.219***<br>(0.189, 0.249)    | 2.694***<br>(2.386, 3.002) |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

## Weighted Results

### Appendix 6. Bivariate Analysis

| Variables                           | Use of Prescription Sleep Aid |       |       | Use of Prescription Medications with Sedative Properties |       |       |
|-------------------------------------|-------------------------------|-------|-------|--|-------|-------|
|                                     | Yes                           |       |       | Yes  |       |       |
|                                     | %                             | X     | P     | %  | X     | P     |
| <b>Part 1: Work Hour Categories</b> |                               |       |       |  |       |       |
| <b>≥ 36 hours/week</b>              |                               |       |       |  |       |       |
| 30-35                               | 10.9                          | 4.5   | 0.245 | 11.4   | 1.6   | 0.256 |
| ≥ 36                                | 89.1                          |       |       | 88.6   |       |       |
| <b>≥ 46 hours/week</b>              |                               |       |       |  |       |       |
| 31-45                               | 77.5                          | 2.8   | 0.394 | 78.3   | 0.0   | 0.899 |
| ≥ 46                                | 22.5                          |       |       | 21.7   |       |       |
| <b>≥ 56 hours/week</b>              |                               |       |       |  |       |       |
| 31-55                               | 91.6                          | 7.5   | 0.016 | 92.3   | 7.8   | 0.030 |
| ≥ 56                                | 8.4                           |       |       | 7.7  |       |       |
| <b>≥ 66 hours/week</b>              |                               |       |       |  |       |       |
| 31-65                               | 97.3                          | 8.8   | 0.058 | 97.7   | 3.0   | 0.270 |
| ≥ 66                                | 2.7                           |       |       | 2.3  |       |       |
| <b>Part 2: Predisposing Factors</b> |                               |       |       |  |       |       |
| <b>Age (years)</b>                  |                               |       |       |  |       |       |
| 18-26                               | 4.1                           | 174.5 | 0.001 | 7.2  | 208.5 | 0.000 |
| 27-64                               | 90.2                          |       |       | 86.3   |       |       |
| ≥ 65                                | 5.6                           |       |       | 6.4  |       |       |
| <b>Race</b>                         |                               |       |       |  |       |       |
| White                               | 90.3                          | 199.0 | 0.000 | 84.2   | 30.9  | 0.000 |
| Black                               | 5.4                           |       |       | 10.1   |       |       |
| Other                               | 4.3                           |       |       | 5.7  |       |       |
| <b>Sex</b>                          |                               |       |       |  |       |       |
| Male                                | 41.9                          | 16.5  | 0.024 | 44.2   | 1.9   | 0.282 |
| Female                              | 58.1                          |       |       | 55.8   |       |       |
| <b>Marital Status</b>               |                               |       |       |  |       |       |
| Married                             | 55.5                          | 92.9  | 0.000 | 57.3   | 156.9 | 0.000 |
| Divorced                            | 24.8                          |       |       | 22.0   |       |       |
| Unmarried                           | 19.7                          |       |       | 20.7   |       |       |
| <b>Family Size</b>                  |                               |       |       |  |       |       |
| 0 - 3 members                       | 80.3                          | 136.8 | 0.000 | 76.2   | 148.2 | 0.000 |
| 4 - 6 members                       | 19.3                          |       |       | 23.1   |       |       |
| > 6 members                         | 0.4                           |       |       | 0.7  |       |       |
| <b>Education Level</b>              |                               |       |       |  |       |       |
| Less than College                   | 28.5                          | 29.4  | 0.008 | 32.5   | 12.1  | 0.001 |
| Some College or more                | 71.6                          |       |       | 67.5   |       |       |

Appendix 6. (Continued)

| Variables                       | Use of Prescription Sleep Aid |       | Use of Prescription Medications with Sedative Properties |      |       |       |
|---------------------------------|-------------------------------|-------|--|------|-------|-------|
|                                 | Yes                           |       |  |      | Yes   |       |
|                                 | %                             | X     | P  | %    | X     | P     |
| <b>Occupation</b>               |                               |       |  |      |       |       |
| Natural resources               | 5.6                           | 62.3  | 0.006  | 6.6  | 8.8   | 0.351 |
| Hospitality services            | 5.0                           |       |  | 5.8  |       |       |
| Trade                           | 10.0                          |       |  | 11.2 |       |       |
| Professional services           | 43.9                          |       |  | 39.6 |       |       |
| Manufacturing                   | 10.7                          |       |  | 11.1 |       |       |
| Other                           | 24.7                          |       |  | 25.8 |       |       |
| <b>Year</b>                     |                               |       |  |      |       |       |
| 2010                            | 10.3                          | 45.2  | 0.014  | 10.4 | 68.9  | 0.013 |
| 2011                            | 11.1                          |       |  | 10.5 |       |       |
| 2012                            | 9.7                           |       |  | 10.0 |       |       |
| 2013                            | 9.5                           |       |  | 9.9  |       |       |
| 2014                            | 11.4                          |       |  | 10.9 |       |       |
| 2015                            | 9.8                           |       |  | 10.7 |       |       |
| 2016                            | 10.5                          |       |  | 10.1 |       |       |
| 2017                            | 9.4                           |       |  | 9.4  |       |       |
| 2018                            | 8.3                           |       |  | 9.1  |       |       |
| 2019                            | 10.2                          |       |  | 8.9  |       |       |
| <b>Part 3: Enabling Factors</b> |                               |       |  |      |       |       |
| <b>Region</b>                   |                               |       |  |      |       |       |
| Northeast                       | 13.5                          | 21.9  | 0.073  | 14.5 | 26.5  | 0.001 |
| North Central/Midwest           | 26.2                          |       |  | 24.6 |       |       |
| South                           | 38.7                          |       |  | 40.0 |       |       |
| West                            | 21.6                          |       |  | 20.8 |       |       |
| <b>Income</b>                   |                               |       |  |      |       |       |
| Poor to low income              | 14.1                          | 10.9  | 0.055  | 16.0 | 3.6   | 0.031 |
| Middle to high income           | 86.0                          |       |  | 84.1 |       |       |
| No                              | 94.6                          | 2.2   | 0.539  | 93.6 | 30.4  | 0.003 |
| Yes                             | 5.4                           |       |  | 6.4  |       |       |
| <b>Part 4: Need Factors</b>     |                               |       |  |      |       |       |
| <b>Health Status</b>            |                               |       |  |      |       |       |
| Excellent                       | 10.4                          | 151.7 | 0.001  | 11.5 | 788.4 | 0.000 |
| Good                            | 75.5                          |       |  | 74.3 |       |       |
| Fair                            | 12.2                          |       |  | 12.4 |       |       |
| Poor                            | 2.0                           |       |  | 1.7  |       |       |

Appendix 7a. Regressions for the Use of Prescription Sleep Aids:  $\geq 56$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Prescription Sleep Aid = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 56$ h/week                                     | 0.062*<br>(0.017, 0.106)      | 0.090*<br>(0.013, 0.167)      | 1.165*<br>(1.014, 1.338)   | 0.010*<br>(0.000, 0.020)      | 0.483**<br>(0.200, 0.767)   |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.115***<br>(0.082, 0.148)    | 0.334***<br>(0.228, 0.439)    | 2.322***<br>(1.933, 2.791) | 0.039***<br>(0.033, 0.045)    | 0.355<br>(-0.022, 0.732)  |
| $\geq 65$ years                                      | 0.068**<br>(0.039, 0.097)     | 0.256***<br>(0.191, 0.320)    | 1.990***<br>(1.564, 2.531) | 0.030***<br>(0.019, 0.040)    | 0.064<br>(-0.208, 0.336)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.136***<br>(-0.149, -0.123) | -0.331***<br>(-0.388, -0.275) | 0.442***<br>(0.392, 0.498) | -0.039***<br>(-0.044, -0.035) | -0.389*<br>(-0.767, -0.011)   |
| Other  | -0.107***<br>(-0.141, -0.073) | -0.219**<br>(-0.330, -0.108)  | 0.618***<br>(0.532, 0.717) | -0.027***<br>(-0.033, -0.020) | -0.671***<br>(-0.837, -0.506)                                       |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | 0.014<br>(-0.023, 0.050)      | 0.032<br>(-0.014, 0.077)      | 1.068<br>(0.991, 1.150)    | 0.004<br>(-0.001, 0.009)      | 0.058<br>(-0.403, 0.518)  |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.044***<br>(0.036, 0.052)    | 0.084***<br>(0.060, 0.107)    | 1.204***<br>(1.102, 1.316) | 0.012***<br>(0.006, 0.018)    | 0.047<br>(-0.183, 0.277)  |
| Unmarried  | 0<br>(-0.015, 0.014)          | 0.022<br>(-0.020, 0.065)      | 1.072<br>(0.974, 1.179)    | 0.004<br>(-0.002, 0.010)      | -0.21<br>(-0.487, 0.068)  |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.067***<br>(-0.070, -0.065) | -0.157***<br>(-0.174, -0.140) | 0.697***<br>(0.639, 0.761) | -0.021***<br>(-0.025, -0.016) | -0.161<br>(-0.380, 0.058)   |
| > 6 Members  | -0.103**<br>(-0.166, -0.041)  | -0.284***<br>(-0.356, -0.211) | 0.486***<br>(0.321, 0.734) | -0.036***<br>(-0.051, -0.021) | 0.034<br>(-1.205, 1.273)  |
| <b>Education (Less than college - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | 0.013<br>(-0.001, 0.028)      | 0.054**<br>(0.025, 0.084)     | 1.137**<br>(1.051, 1.231)  | 0.008**<br>(0.003, 0.013)     | -0.081**<br>(-0.115, -0.046)  |
| <b>Occupation (Natural resources - REF)</b>          |                               |                               |                            |                               |   |
| Hospitality services                                 | 0.04                          | 0.066                         | 1.166                      | 0.008                         | 0.175   |



|  |                 |                  |                |                  |                 |
|--|-----------------|------------------|----------------|------------------|-----------------|
|  | (-0.006, 0.087) | (-0.130, 0.262)  | (0.948, 1.433) | (-0.003, 0.020)  | (-0.185, 0.535) |
| Trade                                    | 0.033           | 0.06             | 1.121          | 0.006            | 0.238           |
|  | (-0.076, 0.142) | (-0.181, 0.301)  | (0.927, 1.355) | (-0.004, 0.016)  | (-0.248, 0.725) |
| Professional services                    | 0.074*          | 0.153*           | 1.406***       | 0.020***         | 0.221           |
|  | (0.008, 0.139)  | (0.008, 0.298)   | (1.195, 1.653) | (0.011, 0.029)   | (-0.178, 0.620) |
| Manufacturing                            | 0.042           | 0.088            | 1.216*         | 0.011*           | 0.188*          |
|  | (-0.003, 0.087) | (-0.044, 0.221)  | (1.015, 1.456) | (0.001, 0.021)   | (0.043, 0.333)  |
| Other                                    | 0.034*          | 0.085            | 1.222*         | 0.011*           | 0.032           |
|  | (0.002, 0.066)  | (-0.048, 0.217)  | (1.037, 1.440) | (0.002, 0.020)   | (-0.178, 0.242) |
| <b>Year (2010 - REF)</b>                 |                 |                  |                |                  |                 |
| 2011                                     | 0.022           | 0.014            | 1.013          | 0.001            | 0.274           |
|  | (-0.044, 0.089) | (-0.064, 0.091)  | (0.869, 1.180) | (-0.009, 0.011)  | (-0.370, 0.919) |
| 2012                                     | 0.007           | -0.039***        | 0.875          | -0.008           | 0.475*          |
|  | (-0.009, 0.023) | (-0.042, -0.036) | (0.744, 1.028) | (-0.018, 0.002)  | (0.159, 0.790)  |
| 2013                                     | 0.005           | -0.033           | 0.903          | -0.006           | 0.351           |
|  | (-0.071, 0.081) | (-0.105, 0.040)  | (0.769, 1.060) | (-0.016, 0.004)  | (-0.485, 1.187) |
| 2014                                     | 0.008           | -0.002           | 0.986          | -0.001           | 0.123           |
|  | (-0.037, 0.053) | (-0.019, 0.014)  | (0.845, 1.152) | (-0.011, 0.009)  | (-0.752, 0.997) |
| 2015                                     | -0.009          | -0.046*          | 0.885          | -0.008           | 0.21            |
|  | (-0.054, 0.036) | (-0.089, -0.003) | (0.759, 1.032) | (-0.017, 0.002)  | (-0.702, 1.123) |
| 2016                                     | 0.003           | -0.004           | 0.977          | -0.002           | 0.169           |
|  | (-0.036, 0.043) | (-0.024, 0.016)  | (0.835, 1.142) | (-0.012, 0.009)  | (-0.632, 0.970) |
| 2017                                     | 0.007           | -0.030**         | 0.9            | -0.007           | 0.395***        |
|  | (-0.003, 0.017) | (-0.047, -0.013) | (0.770, 1.051) | (-0.016, 0.003)  | (0.274, 0.517)  |
| 2018                                     | -0.018          | -0.076**         | 0.814**        | -0.012**         | 0.276           |
|  | (-0.064, 0.029) | (-0.122, -0.031) | (0.699, 0.947) | (-0.022, -0.003) | (-0.218, 0.771) |
| 2019                                     | 0.026           | 0.066            | 1.138          | 0.009            | 0.075           |
|  | (-0.027, 0.079) | (-0.003, 0.134)  | (0.977, 1.327) | (-0.002, 0.019)  | (-0.401, 0.550) |
| <b>Region (Northeast - REF)</b>          |                 |                  |                |                  |                 |
| North central, Midwest                   | 0.052***        | 0.102***         | 1.248***       | 0.013***         | 0.210**         |
|  | (0.041, 0.062)  | (0.075, 0.128)   | (1.110, 1.403) | (0.006, 0.020)   | (0.089, 0.332)  |
| South                                    | 0.041***        | 0.077***         | 1.183**        | 0.010**          | 0.216           |
|  | (0.029, 0.054)  | (0.056, 0.099)   | (1.059, 1.323) | (0.004, 0.016)   | (-0.032, 0.465) |
| West                                     | 0.026           | 0.063            | 1.162*         | 0.009*           | 0.054           |
|  | (-0.039, 0.091) | (-0.057, 0.183)  | (1.033, 1.307) | (0.002, 0.015)   | (-0.300, 0.408) |
| <b>Income (Poor to low income - REF)</b> |                 |                  |                |                  |                 |
| Middle to high income                    | -0.003          | -0.003           | 0.997          | 0                | -0.048          |

|   |                 |                 |                |                 |                 |
|---|-----------------|-----------------|----------------|-----------------|-----------------|
|   | (-0.018, 0.012) | (-0.063, 0.057) | (0.900, 1.104) | (-0.007, 0.006) | (-0.323, 0.226) |
| <b>Receipt of food stamp (No - REF)</b> |                 |                 |                |                 |                 |
| YES                                     | 0.036           | 0.042           | 1.068          | 0.004           | 0.368**         |
|   | (-0.038, 0.109) | (-0.100, 0.185) | (0.921, 1.239) | (-0.005, 0.014) | (0.178, 0.558)  |
| <b>Health status (Excellent - REF)</b>  |                 |                 |                |                 |                 |
| Good                                    | 0.067***        | 0.164***        | 1.469***       | 0.021***        | 0.255*          |
|   | (0.057, 0.077)  | (0.140, 0.187)  | (1.303, 1.656) | (0.015, 0.026)  | (0.087, 0.424)  |
| Fair                                    | 0.189***        | 0.321***        | 1.906***       | 0.039***        | 1.085***        |
|   | (0.158, 0.220)  | (0.252, 0.391)  | (1.646, 2.207) | (0.030, 0.048)  | (0.746, 1.424)  |
| Poor                                    | 0.290**         | 0.490***        | 2.589***       | 0.066***        | 1.226***        |
|   | (0.162, 0.418)  | (0.313, 0.666)  | (1.999, 3.353) | (0.042, 0.090)  | (1.012, 1.439)  |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF=reference

Appendix 7b. Regressions for the Use of Prescription Medications with Sedative Properties:  $\geq 56$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Sedative Medications = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 56$ h/week                                     | 0.157*<br>(0.025, 0.289)      | 0.098*<br>(0.027, 0.170)      | 1.079*<br>(1.004, 1.160)   | 0.017*<br>(0.001, 0.033)      | 0.301*<br>(0.020, 0.582)  |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.256***<br>(0.221, 0.291)    | 0.223***<br>(0.180, 0.266)    | 1.287***<br>(1.194, 1.388) | 0.054***<br>(0.039, 0.070)    | 0.373*<br>(0.060, 0.687)  |
| $\geq 65$ years                                      | 0.328*<br>(0.111, 0.546)      | 0.314**<br>(0.167, 0.460)     | 1.466***<br>(1.317, 1.632) | 0.084***<br>(0.060, 0.107)    | 0.345<br>(-0.061, 0.751)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.263***<br>(-0.288, -0.237) | -0.117***<br>(-0.133, -0.102) | 0.963<br>(0.916, 1.012)    | -0.008<br>(-0.019, 0.003)     | -0.648***<br>(-0.698, -0.598)                                     |
| Other  | -0.289***<br>(-0.325, -0.253) | -0.188***<br>(-0.225, -0.151) | 0.855***<br>(0.797, 0.916) | -0.034***<br>(-0.049, -0.019) | -0.589***<br>(-0.659, -0.518)                                     |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | 0.007<br>(-0.015, 0.029)      | 0.005<br>(-0.015, 0.025)      | 1.007<br>(0.969, 1.047)    | 0.002<br>(-0.007, 0.010)      | 0.003<br>(-0.101, 0.108)  |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.174***<br>(0.145, 0.202)    | 0.097***<br>(0.080, 0.113)    | 1.068**<br>(1.018, 1.120)  | 0.015**<br>(0.004, 0.026)     | 0.325***<br>(0.259, 0.392)  |
| Unmarried  | -0.057<br>(-0.196, 0.082)     | -0.068<br>(-0.160, 0.023)     | 0.907***<br>(0.863, 0.954) | -0.021***<br>(-0.032, -0.010) | -0.001<br>(-0.236, 0.233)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.217***<br>(-0.246, -0.188) | -0.177***<br>(-0.200, -0.154) | 0.823***<br>(0.787, 0.859) | -0.043***<br>(-0.052, -0.033) | -0.294***<br>(-0.350, -0.237)                                     |
| > 6 Members  | -0.457***<br>(-0.591, -0.324) | -0.367**<br>(-0.527, -0.207)  | 0.659***<br>(0.562, 0.772) | -0.088***<br>(-0.119, -0.057) | -0.662***<br>(-0.883, -0.441)                                     |
| <b>Education (Less than college - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | -0.023<br>(-0.057, 0.010)     | -0.015<br>(-0.035, 0.006)     | 0.986<br>(0.947, 1.026)    | -0.003<br>(-0.012, 0.006)     | -0.027<br>(-0.113, 0.059)   |

| <b>Occupation (Natural resources - REF)</b> |                  |                  |                |                  |                 |
|---|------------------|------------------|----------------|------------------|-----------------|
| Hospitality services                        | 0.02             | 0.051            | 1.109*         | 0.023*           | -0.181          |
|   | (-0.345, 0.385)  | (-0.191, 0.294)  | (1.001, 1.229) | (0.000, 0.045)   | (-0.843, 0.481) |
| Trade                                       | 0.051            | 0.049            | 1.073          | 0.015            | -0.009          |
|   | (-0.340, 0.442)  | (-0.169, 0.267)  | (0.979, 1.175) | (-0.005, 0.035)  | (-0.946, 0.928) |
| Professional services                       | 0.047            | 0.069            | 1.123**        | 0.025**          | -0.119          |
|   | (-0.260, 0.355)  | (-0.088, 0.227)  | (1.035, 1.218) | (0.008, 0.043)   | (-0.889, 0.650) |
| Manufacturing                               | -0.032           | 0.023            | 1.087          | 0.018            | -0.281          |
|   | (-0.356, 0.292)  | (-0.179, 0.224)  | (0.994, 1.190) | (-0.001, 0.038)  | (-0.905, 0.343) |
| Other                                       | -0.016           | 0.04             | 1.118**        | 0.025**          | -0.303          |
|   | (-0.286, 0.253)  | (-0.119, 0.199)  | (1.030, 1.214) | (0.007, 0.042)   | (-0.940, 0.335) |
| <b>Year (2010 - REF)</b>                    |                  |                  |                |                  |                 |
| 2011  | 0.025            | -0.031           | 0.907*         | -0.022*          | 0.277***        |
|   | (-0.033, 0.083)  | (-0.080, 0.017)  | (0.838, 0.982) | (-0.040, -0.004) | (0.247, 0.308)  |
| 2012  | -0.079*          | -0.115***        | 0.831***       | -0.042***        | 0.130***        |
|   | (-0.133, -0.026) | (-0.153, -0.078) | (0.766, 0.900) | (-0.060, -0.024) | (0.087, 0.173)  |
| 2013  | -0.002           | -0.060*          | 0.878**        | -0.029**         | 0.234*          |
|   | (-0.086, 0.082)  | (-0.104, -0.016) | (0.809, 0.952) | (-0.048, -0.011) | (0.012, 0.455)  |
| 2014  | 0.022            | -0.051           | 0.879**        | -0.029**         | 0.289           |
|   | (-0.164, 0.208)  | (-0.122, 0.020)  | (0.811, 0.953) | (-0.047, -0.011) | (-0.231, 0.809) |
| 2015  | -0.021           | -0.057           | 0.897**        | -0.025**         | 0.15            |
|   | (-0.171, 0.129)  | (-0.145, 0.031)  | (0.829, 0.970) | (-0.043, -0.007) | (-0.164, 0.464) |
| 2016  | -0.035           | -0.079           | 0.868***       | -0.032***        | 0.162**         |
|   | (-0.121, 0.051)  | (-0.158, 0.000)  | (0.802, 0.940) | (-0.050, -0.014) | (0.097, 0.228)  |
| 2017  | -0.016           | -0.095**         | 0.826***       | -0.043***        | 0.304*          |
|   | (-0.046, 0.014)  | (-0.130, -0.059) | (0.761, 0.896) | (-0.061, -0.025) | (0.081, 0.528)  |
| 2018  | 0.017            | -0.084*          | 0.822***       | -0.044***        | 0.407***        |
|   | (-0.065, 0.099)  | (-0.154, -0.014) | (0.759, 0.890) | (-0.062, -0.026) | (0.339, 0.475)  |
| 2019  | 0.005            | -0.015           | 0.956          | -0.01            | 0.12            |
|   | (-0.130, 0.140)  | (-0.111, 0.081)  | (0.881, 1.038) | (-0.029, 0.009)  | (-0.136, 0.376) |
| <b>Region (Northeast - REF)</b>             |                  |                  |                |                  |                 |
| North central, Midwest                      | 0.129***         | 0.093***         | 1.089**        | 0.019**          | 0.230***        |
|   | (0.101, 0.158)   | (0.071, 0.115)   | (1.025, 1.157) | (0.005, 0.032)   | (0.211, 0.249)  |
| South                                       | 0.124***         | 0.091***         | 1.088**        | 0.019**          | 0.226***        |
|   | (0.088, 0.160)   | (0.069, 0.113)   | (1.027, 1.153) | (0.006, 0.031)   | (0.143, 0.309)  |
| West  | 0.008            | 0.022            | 1.036          | 0.008            | -0.013          |
|   | (-0.056, 0.072)  | (-0.011, 0.055)  | (0.974, 1.101) | (-0.006, 0.021)  | (-0.156, 0.131) |

| <b>Income (Poor to low income - REF)</b> |                 |                 |                |                 |                 |
|--|-----------------|-----------------|----------------|-----------------|-----------------|
| Middle to high income                    | -0.02           | -0.019          | 0.976          | -0.005          | -0.016          |
|  | (-0.043, 0.003) | (-0.047, 0.010) | (0.927, 1.028) | (-0.017, 0.006) | (-0.054, 0.022) |
| <b>Receipt of food stamp (No - REF)</b>  |                 |                 |                |                 |                 |
| YES                                      | 0.308***        | 0.213***        | 1.203***       | 0.042***        | 0.483**         |
|  | (0.211, 0.406)  | (0.169, 0.256)  | (1.120, 1.293) | (0.025, 0.059)  | (0.200, 0.766)  |
| <b>Health status (Excellent - REF)</b>   |                 |                 |                |                 |                 |
| Good                                     | 0.361***        | 0.303***        | 1.412***       | 0.073***        | 0.590***        |
|  | (0.314, 0.408)  | (0.253, 0.354)  | (1.336, 1.493) | (0.062, 0.084)  | (0.497, 0.683)  |
| Fair                                     | 1.075***        | 0.783***        | 2.113***       | 0.168***        | 1.586***        |
|  | (1.009, 1.140)  | (0.731, 0.834)  | (1.962, 2.275) | (0.151, 0.184)  | (1.378, 1.795)  |
| Poor                                     | 1.856***        | 1.224***        | 2.638***       | 0.223***        | 2.696***        |
|  | (1.791, 1.922)  | (1.181, 1.266)  | (2.264, 3.075) | (0.186, 0.260)  | (2.422, 2.970)  |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 8a. Regressions for Use of Prescription Sleep Aids:  $\geq 36$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Prescription Sleep Aid = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 36$ h/week                                     | 0.029<br>(-0.036, 0.095)      | 0.023<br>(-0.001, 0.048)      | 1.036<br>(0.966, 1.111)    | 0.002<br>(-0.002, 0.006)      | 0.047<br>(-0.028, 0.122)  |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.115***<br>(0.082, 0.148)    | 0.334***<br>(0.229, 0.439)    | 2.323***<br>(1.716, 3.145) | 0.039***<br>(0.029, 0.049)    | 0.347<br>(-0.020, 0.714)  |
| $\geq 65$ years                                      | 0.071**<br>(0.040, 0.101)     | 0.261***<br>(0.198, 0.324)    | 1.999***<br>(1.594, 2.508) | 0.030***<br>(0.024, 0.036)    | 0.159<br>(-0.030, 0.348)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.136***<br>(-0.148, -0.123) | -0.331***<br>(-0.388, -0.274) | 0.442***<br>(0.371, 0.527) | -0.039***<br>(-0.046, -0.033) | -0.381*<br>(-0.750, -0.012)   |
| Other  | -0.107***<br>(-0.143, -0.072) | -0.219**<br>(-0.332, -0.105)  | 0.617**<br>(0.458, 0.833)  | -0.027**<br>(-0.041, -0.013)  | -0.658***<br>(-0.793, -0.522)                                       |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | 0.011<br>(-0.026, 0.048)      | 0.027<br>(-0.019, 0.073)      | 1.058<br>(0.980, 1.143)    | 0.004<br>(-0.001, 0.008)      | 0.034<br>(-0.430, 0.498)  |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.045***<br>(0.037, 0.053)    | 0.084***<br>(0.060, 0.109)    | 1.207***<br>(1.132, 1.286) | 0.012***<br>(0.008, 0.016)    | 0.035<br>(-0.176, 0.246)  |
| Unmarried  | 0<br>(-0.014, 0.013)          | 0.022<br>(-0.020, 0.063)      | 1.07<br>(0.952, 1.203)     | 0.004<br>(-0.003, 0.011)      | -0.217<br>(-0.501, 0.066)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.067***<br>(-0.070, -0.064) | -0.157***<br>(-0.175, -0.139) | 0.697***<br>(0.657, 0.740) | -0.021***<br>(-0.024, -0.017) | -0.173<br>(-0.383, 0.037)   |
| > 6 Members  | -0.104**<br>(-0.170, -0.038)  | -0.285***<br>(-0.362, -0.208) | 0.486***<br>(0.446, 0.529) | -0.036***<br>(-0.040, -0.032) | -0.019<br>(-1.348, 1.309)   |
| <b>Education (Less than college - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | 0.013<br>(-0.000, 0.027)      | 0.055**<br>(0.026, 0.084)     | 1.138**<br>(1.060, 1.222)  | 0.008**<br>(0.004, 0.012)     | -0.076***<br>(-0.099, -0.053)                                       |
| <b>Occupation (Natural resources - REF)</b>          |                               |                               |                            |                               |   |

|  |                 |                  |                |                  |                 |
|--|-----------------|------------------|----------------|------------------|-----------------|
| Hospitality services                     | 0.04            | 0.064            | 1.164          | 0.008            | 0.143           |
|  | (-0.003, 0.084) | (-0.129, 0.257)  | (0.713, 1.898) | (-0.018, 0.035)  | (-0.214, 0.500) |
| Trade                                    | 0.031           | 0.057            | 1.115          | 0.006            | 0.208           |
|  | (-0.078, 0.140) | (-0.184, 0.298)  | (0.652, 1.906) | (-0.023, 0.035)  | (-0.276, 0.692) |
| Professional services                    | 0.071*          | 0.149*           | 1.398*         | 0.020*           | 0.165           |
|  | (0.010, 0.133)  | (0.009, 0.289)   | (1.020, 1.914) | (0.003, 0.036)   | (-0.173, 0.503) |
| Manufacturing                            | 0.037           | 0.08             | 1.201          | 0.01             | 0.113           |
|  | (-0.006, 0.080) | (-0.049, 0.210)  | (0.873, 1.653) | (-0.006, 0.027)  | (-0.127, 0.354) |
| Other                                    | 0.031*          | 0.079            | 1.213          | 0.011            | -0.035          |
|  | (0.001, 0.061)  | (-0.049, 0.208)  | (0.877, 1.678) | (-0.006, 0.028)  | (-0.280, 0.210) |
| <b>Year (2010 - REF)</b>                 |                 |                  |                |                  |                 |
| 2011                                     | 0.023           | 0.014            | 1.014          | 0.001            | 0.28            |
|  | (-0.045, 0.091) | (-0.065, 0.094)  | (0.894, 1.151) | (-0.007, 0.009)  | (-0.363, 0.922) |
| 2012                                     | 0.008           | -0.039***        | 0.875***       | -0.008***        | 0.484**         |
|  | (-0.011, 0.026) | (-0.039, -0.038) | (0.858, 0.892) | (-0.009, -0.007) | (0.200, 0.768)  |
| 2013                                     | 0.005           | -0.033           | 0.901          | -0.006           | 0.358           |
|  | (-0.073, 0.083) | (-0.108, 0.042)  | (0.800, 1.016) | (-0.014, 0.001)  | (-0.486, 1.201) |
| 2014                                     | 0.009           | -0.002           | 0.987          | -0.001           | 0.136           |
|  | (-0.039, 0.056) | (-0.016, 0.013)  | (0.911, 1.068) | (-0.006, 0.004)  | (-0.751, 1.024) |
| 2015                                     | -0.009          | -0.047*          | 0.885          | -0.008           | 0.198           |
|  | (-0.055, 0.036) | (-0.090, -0.003) | (0.756, 1.036) | (-0.017, 0.002)  | (-0.723, 1.120) |
| 2016                                     | 0.003           | -0.005           | 0.975          | -0.002           | 0.159           |
|  | (-0.037, 0.043) | (-0.025, 0.015)  | (0.880, 1.080) | (-0.008, 0.005)  | (-0.632, 0.950) |
| 2017                                     | 0.007           | -0.031**         | 0.898**        | -0.007**         | 0.379***        |
|  | (-0.004, 0.017) | (-0.049, -0.013) | (0.862, 0.937) | (-0.009, -0.004) | (0.288, 0.470)  |
| 2018                                     | -0.018          | -0.077**         | 0.813**        | -0.012**         | 0.265           |
|  | (-0.064, 0.029) | (-0.122, -0.032) | (0.746, 0.885) | (-0.017, -0.007) | (-0.202, 0.732) |
| 2019                                     | 0.026           | 0.065            | 1.138*         | 0.009*           | 0.077           |
|  | (-0.027, 0.079) | (-0.002, 0.133)  | (1.008, 1.285) | (0.000, 0.018)   | (-0.373, 0.528) |
| <b>Region (Northeast - REF)</b>          |                 |                  |                |                  |                 |
| North central, Midwest                   | 0.052***        | 0.103***         | 1.250***       | 0.013***         | 0.219**         |
|  | (0.043, 0.061)  | (0.077, 0.129)   | (1.180, 1.324) | (0.010, 0.017)   | (0.119, 0.319)  |
| South                                    | 0.041***        | 0.078***         | 1.183***       | 0.010***         | 0.221           |
|  | (0.028, 0.053)  | (0.055, 0.100)   | (1.117, 1.254) | (0.007, 0.013)   | (-0.020, 0.462) |
| West                                     | 0.025           | 0.063            | 1.162          | 0.009            | 0.047           |
|  | (-0.040, 0.090) | (-0.058, 0.183)  | (0.904, 1.494) | (-0.006, 0.024)  | (-0.299, 0.392) |
| <b>Income (Poor to low income - REF)</b> |                 |                  |                |                  |                 |

|   |                            |                            |                            |                            |                            |
|---|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| Middle to high income                   | -0.003<br>(-0.017, 0.011)  | -0.003<br>(-0.062, 0.056)  | 0.998<br>(0.869, 1.145)    | 0<br>(-0.009, 0.008)       | -0.055<br>(-0.320, 0.210)  |
| <b>Receipt of food stamp (No - REF)</b> |                            |                            |                            |                            |                            |
| YES                                     | 0.037<br>(-0.039, 0.114)   | 0.045<br>(-0.102, 0.193)   | 1.072<br>(0.758, 1.516)    | 0.004<br>(-0.018, 0.027)   | 0.417**<br>(0.218, 0.617)  |
| <b>Health status (Excellent - REF)</b>  |                            |                            |                            |                            |                            |
| Good                                    | 0.066***<br>(0.057, 0.075) | 0.162***<br>(0.141, 0.183) | 1.465***<br>(1.382, 1.552) | 0.020***<br>(0.018, 0.023) | 0.232*<br>(0.070, 0.394)   |
| Fair                                    | 0.189***<br>(0.158, 0.219) | 0.320***<br>(0.253, 0.388) | 1.903***<br>(1.593, 2.275) | 0.039***<br>(0.026, 0.051) | 1.071**<br>(0.717, 1.424)  |
| Poor                                    | 0.290**<br>(0.161, 0.420)  | 0.488***<br>(0.309, 0.667) | 2.587***<br>(1.921, 3.486) | 0.066**<br>(0.035, 0.097)  | 1.182***<br>(0.985, 1.379) |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference



Appendix 8b. Regressions for the Use of Prescription Medications with Sedative Properties:  $\geq 36$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Sedative Medications = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 36$ h/week                                     | 0.071<br>(-0.073, 0.214)      | 0.056<br>(0.078, 0.114)       | 1.062<br>(0.966, 1.168)    | 0.013<br>(-0.007, 0.034)      | 0.092<br>(-0.174, 0.359)  |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.255***<br>(0.221, 0.290)    | 0.222***<br>(0.180, 0.263)    | 1.284***<br>(1.170, 1.409) | 0.054***<br>(0.034, 0.073)    | 0.374*<br>(0.070, 0.679)  |
| $\geq 65$ years                                      | 0.336*<br>(0.120, 0.552)      | 0.321**<br>(0.176, 0.466)     | 1.474***<br>(1.277, 1.702) | 0.085***<br>(0.054, 0.116)    | 0.369<br>(-0.057, 0.796)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.262***<br>(-0.288, -0.236) | -0.117***<br>(-0.133, -0.102) | 0.963**<br>(0.945, 0.981)  | -0.008**<br>(-0.013, -0.004)  | -0.646***<br>(-0.694, -0.598)                                     |
| Other  | -0.290***<br>(-0.325, -0.254) | -0.188***<br>(-0.226, -0.151) | 0.854***<br>(0.825, 0.885) | -0.034***<br>(-0.042, -0.027) | -0.589***<br>(-0.662, -0.516)                                     |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | 0<br>(-0.023, 0.023)          | 0.002<br>(-0.024, 0.027)      | 1.005<br>(0.957, 1.055)    | 0.001<br>(-0.010, 0.012)      | -0.014<br>(-0.115, 0.086)   |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.176***<br>(0.147, 0.204)    | 0.098***<br>(0.081, 0.115)    | 1.069***<br>(1.045, 1.093) | 0.015***<br>(0.010, 0.020)    | 0.327***<br>(0.261, 0.393)  |
| Unmarried  | -0.058<br>(-0.197, 0.082)     | -0.068<br>(-0.160, 0.024)     | 0.908*<br>(0.827, 0.996)   | -0.021*<br>(-0.042, -0.001)   | -0.005<br>(-0.241, 0.232)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.217***<br>(-0.245, -0.189) | -0.177***<br>(-0.198, -0.156) | 0.823***<br>(0.815, 0.831) | -0.043***<br>(-0.045, -0.041) | -0.295***<br>(-0.349, -0.241)                                     |
| > 6 Members  | -0.459***<br>(-0.582, -0.336) | -0.369**<br>(-0.525, -0.213)  | 0.658**<br>(0.533, 0.811)  | -0.088**<br>(-0.129, -0.047)  | -0.668***<br>(-0.881, -0.456)                                     |
| <b>Education (Less than college - REF)</b>           |                               |                               |                            |                               |   |

|   |                             |                               |                            |                               |                            |
|---|-----------------------------|-------------------------------|----------------------------|-------------------------------|----------------------------|
| Some college or more                        | -0.023<br>(-0.055, 0.008)   | -0.015<br>(-0.035, 0.005)     | 0.985<br>(0.959, 1.012)    | -0.003<br>(-0.009, 0.003)     | -0.026<br>(-0.112, 0.059)  |
| <b>Occupation (Natural resources - REF)</b> |                             |                               |                            |                               |                            |
| Hospitality services                        | 0.022<br>(-0.341, 0.385)    | 0.055<br>(-0.180, 0.291)      | 1.116<br>(0.906, 1.374)    | 0.024<br>(-0.022, 0.070)      | -0.183<br>(-0.883, 0.516)  |
| Trade                                       | 0.048<br>(-0.342, 0.437)    | 0.049<br>(-0.168, 0.265)      | 1.072<br>(0.941, 1.222)    | 0.015<br>(-0.013, 0.044)      | -0.014<br>(-0.963, 0.935)  |
| Professional services                       | 0.042<br>(-0.257, 0.342)    | 0.067<br>(-0.085, 0.219)      | 1.121*<br>(1.028, 1.223)   | 0.025*<br>(0.006, 0.044)      | -0.129<br>(-0.892, 0.634)  |
| Manufacturing                               | -0.045<br>(-0.357, 0.267)   | 0.014<br>(-0.184, 0.212)      | 1.079<br>(0.909, 1.281)    | 0.017<br>(-0.021, 0.054)      | -0.306<br>(-0.887, 0.276)  |
| Other                                       | -0.024<br>(-0.286, 0.238)   | 0.036<br>(-0.119, 0.190)      | 1.114<br>(0.995, 1.247)    | 0.024<br>(-0.001, 0.048)      | -0.316<br>(-0.938, 0.305)  |
| <b>Year (2010 - REF)</b>                    |                             |                               |                            |                               |                            |
| 2011  | 0.027<br>(-0.034, 0.088)    | -0.03<br>(-0.081, 0.020)      | 0.908*<br>(0.848, 0.972)   | -0.022*<br>(-0.038, -0.006)   | 0.280***<br>(0.245, 0.315) |
| 2012  | -0.079*<br>(-0.136, -0.021) | -0.115***<br>(-0.154, -0.075) | 0.831***<br>(0.785, 0.879) | -0.042***<br>(-0.055, -0.029) | 0.130***<br>(0.086, 0.174) |
| 2013  | -0.003<br>(-0.091, 0.085)   | -0.060*<br>(-0.105, -0.016)   | 0.878**<br>(0.825, 0.934)  | -0.030**<br>(-0.043, -0.016)  | 0.233*<br>(0.002, 0.463)   |
| 2014  | 0.023<br>(-0.167, 0.213)    | -0.05<br>(-0.124, 0.023)      | 0.880***<br>(0.870, 0.890) | -0.029***<br>(-0.032, -0.026) | 0.289<br>(-0.235, 0.814)   |
| 2015  | -0.02<br>(-0.170, 0.130)    | -0.057<br>(-0.144, 0.030)     | 0.897*<br>(0.825, 0.975)   | -0.025*<br>(-0.044, -0.006)   | 0.149<br>(-0.163, 0.461)   |
| 2016  | -0.036<br>(-0.122, 0.050)   | -0.080*<br>(-0.158, -0.001)   | 0.868*<br>(0.780, 0.966)   | -0.032*<br>(-0.056, -0.008)   | 0.160**<br>(0.096, 0.223)  |
| 2017  | -0.017<br>(-0.047, 0.014)   | -0.095**<br>(-0.132, -0.058)  | 0.826**<br>(0.753, 0.905)  | -0.043**<br>(-0.064, -0.022)  | 0.304*<br>(0.081, 0.526)   |
| 2018  | 0.017<br>(-0.066, 0.099)    | -0.084*<br>(-0.154, -0.014)   | 0.822**<br>(0.735, 0.920)  | -0.044**<br>(-0.069, -0.019)  | 0.405***<br>(0.342, 0.468) |
| 2019  | 0.006<br>(-0.129, 0.141)    | -0.014<br>(-0.111, 0.082)     | 0.957<br>(0.861, 1.064)    | -0.01<br>(-0.034, 0.014)      | 0.12<br>(-0.135, 0.375)    |
| <b>Region (Northeast - REF)</b>             |                             |                               |                            |                               |                            |
| North central, Midwest                      | 0.130***<br>(0.098, 0.162)  | 0.093***<br>(0.067, 0.118)    | 1.088**<br>(1.052, 1.125)  | 0.019**<br>(0.011, 0.026)     | 0.235***<br>(0.216, 0.253) |
| South                                       | 0.122***                    | 0.089***                      | 1.086***                   | 0.018***                      | 0.224**                    |

|  |                 |                 |                |                 |                 |
|--|-----------------|-----------------|----------------|-----------------|-----------------|
|  | (0.085, 0.159)  | (0.069, 0.109)  | (1.073, 1.098) | (0.016, 0.021)  | (0.132, 0.316)  |
| West                                     | 0.006           | 0.02            | 1.033*         | 0.007*          | -0.017          |
|  | (-0.057, 0.069) | (-0.014, 0.054) | (1.006, 1.061) | (0.001, 0.013)  | (-0.152, 0.118) |
| <b>Income (Poor to low income - REF)</b> |                 |                 |                |                 |                 |
| Middle to high income                    | -0.022          | -0.022          | 0.972          | -0.006          | -0.016          |
|  | (-0.044, 0.001) | (-0.055, 0.012) | (0.922, 1.026) | (-0.018, 0.006) | (-0.069, 0.037) |
| <b>Receipt of food stamp (No - REF)</b>  |                 |                 |                |                 |                 |
| YES                                      | 0.314***        | 0.218***        | 1.209**        | 0.043**         | 0.497**         |
|  | (0.213, 0.414)  | (0.172, 0.264)  | (1.125, 1.299) | (0.027, 0.060)  | (0.235, 0.759)  |
| <b>Health status (Excellent - REF)</b>   |                 |                 |                |                 |                 |
| Good                                     | 0.359***        | 0.302***        | 1.411***       | 0.073***        | 0.584***        |
|  | (0.308, 0.410)  | (0.249, 0.355)  | (1.335, 1.492) | (0.062, 0.084)  | (0.487, 0.682)  |
| Fair                                     | 1.074***        | 0.783***        | 2.113***       | 0.168***        | 1.583***        |
|  | (1.006, 1.143)  | (0.729, 0.837)  | (2.033, 2.196) | (0.160, 0.176)  | (1.375, 1.791)  |
| Poor                                     | 1.856***        | 1.224***        | 2.639***       | 0.223***        | 2.692***        |
|  | (1.790, 1.922)  | (1.183, 1.265)  | (2.475, 2.815) | (0.206, 0.240)  | (2.418, 2.965)  |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 9a. Regressions for the Use of Prescription Sleep Aids:  $\geq 46$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Prescription Sleep Aid = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 46$ h/week                                     | 0.006<br>(-0.020, 0.031)      | 0.013<br>(-0.035, 0.061)      | 1.033<br>(0.911, 1.171)    | 0.002<br>(-0.006, 0.010)      | -0.005<br>(-0.056, 0.047)   |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.116***<br>(0.084, 0.148)    | 0.335***<br>(0.231, 0.438)    | 2.324***<br>(1.722, 3.136) | 0.039***<br>(0.030, 0.049)    | 0.372*<br>(0.011, 0.734)  |
| $\geq 65$ years                                      | 0.069**<br>(0.039, 0.100)     | 0.259***<br>(0.197, 0.321)    | 1.993***<br>(1.596, 2.489) | 0.030***<br>(0.024, 0.036)    | 0.13<br>(-0.040, 0.301)   |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.135***<br>(-0.148, -0.123) | -0.330***<br>(-0.386, -0.274) | 0.442***<br>(0.372, 0.526) | -0.039***<br>(-0.046, -0.033) | -0.376*<br>(-0.747, -0.006)   |
| Other  | -0.107***<br>(-0.142, -0.072) | -0.218**<br>(-0.332, -0.105)  | 0.618**<br>(0.458, 0.835)  | -0.026**<br>(-0.041, -0.012)  | -0.653***<br>(-0.797, -0.508)                                       |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | 0.01<br>(-0.029, 0.049)       | 0.028<br>(-0.022, 0.077)      | 1.061<br>(0.975, 1.154)    | 0.004<br>(-0.002, 0.009)      | 0.021<br>(-0.461, 0.504)  |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.045***<br>(0.037, 0.053)    | 0.084***<br>(0.061, 0.107)    | 1.206***<br>(1.134, 1.283) | 0.012***<br>(0.008, 0.016)    | 0.046<br>(-0.173, 0.264)  |
| Unmarried  | -0.001<br>(-0.015, 0.014)     | 0.022<br>(-0.020, 0.064)      | 1.07<br>(0.951, 1.205)     | 0.004<br>(-0.003, 0.011)      | -0.216<br>(-0.477, 0.045)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.068***<br>(-0.070, -0.065) | -0.157***<br>(-0.175, -0.140) | 0.697***<br>(0.658, 0.739) | -0.021***<br>(-0.024, -0.017) | -0.169<br>(-0.378, 0.040)   |
| > 6 Members  | -0.103*<br>(-0.170, -0.036)   | -0.284***<br>(-0.363, -0.205) | 0.486***<br>(0.444, 0.532) | -0.036***<br>(-0.040, -0.032) | -0.003<br>(-1.315, 1.308)   |
| <b>Education (Less than college - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | 0.014*<br>(0.001, 0.027)      | 0.055**<br>(0.028, 0.081)     | 1.137**<br>(1.064, 1.215)  | 0.008**<br>(0.004, 0.012)     | -0.073**<br>(-0.099, -0.047)  |

| <b>Occupation (Natural resources - REF)</b> |                 |                  |                |                  |                 |
|---|-----------------|------------------|----------------|------------------|-----------------|
| Hospitality services                        | 0.038           | 0.061            | 1.16           | 0.008            | 0.101           |
|   | (-0.007, 0.082) | (-0.133, 0.256)  | (0.707, 1.902) | (-0.019, 0.035)  | (-0.306, 0.507) |
| Trade                                       | 0.03            | 0.056            | 1.115          | 0.006            | 0.196           |
|   | (-0.078, 0.138) | (-0.184, 0.297)  | (0.654, 1.902) | (-0.023, 0.034)  | (-0.277, 0.668) |
| Professional services                       | 0.071*          | 0.149*           | 1.399*         | 0.020*           | 0.165           |
|   | (0.009, 0.134)  | (0.008, 0.290)   | (1.020, 1.919) | (0.003, 0.036)   | (-0.168, 0.498) |
| Manufacturing                               | 0.038           | 0.082            | 1.205          | 0.01             | 0.126           |
|   | (-0.004, 0.081) | (-0.046, 0.211)  | (0.881, 1.649) | (-0.006, 0.027)  | (-0.099, 0.352) |
| Other                                       | 0.031*          | 0.08             | 1.216          | 0.011            | -0.031          |
|   | (0.002, 0.061)  | (-0.048, 0.209)  | (0.880, 1.679) | (-0.006, 0.028)  | (-0.273, 0.211) |
| <b>Year (2010 - REF)</b>                    |                 |                  |                |                  |                 |
| 2011  | 0.023           | 0.014            | 1.014          | 0.001            | 0.284           |
|   | (-0.045, 0.091) | (-0.065, 0.093)  | (0.894, 1.149) | (-0.007, 0.009)  | (-0.371, 0.938) |
| 2012  | 0.007           | -0.039***        | 0.875***       | -0.008***        | 0.476*          |
|   | (-0.011, 0.026) | (-0.040, -0.038) | (0.857, 0.893) | (-0.009, -0.007) | (0.169, 0.782)  |
| 2013  | 0.005           | -0.033           | 0.902          | -0.006           | 0.351           |
|   | (-0.074, 0.083) | (-0.109, 0.043)  | (0.799, 1.017) | (-0.014, 0.001)  | (-0.509, 1.210) |
| 2014  | 0.008           | -0.002           | 0.986          | -0.001           | 0.136           |
|   | (-0.039, 0.056) | (-0.017, 0.013)  | (0.909, 1.070) | (-0.006, 0.004)  | (-0.758, 1.029) |
| 2015  | -0.009          | -0.047*          | 0.885          | -0.008           | 0.198           |
|   | (-0.055, 0.036) | (-0.090, -0.004) | (0.755, 1.036) | (-0.017, 0.002)  | (-0.765, 1.160) |
| 2016  | 0.003           | -0.004           | 0.976          | -0.002           | 0.159           |
|   | (-0.036, 0.043) | (-0.024, 0.015)  | (0.880, 1.081) | (-0.008, 0.005)  | (-0.671, 0.989) |
| 2017  | 0.007           | -0.031**         | 0.898**        | -0.007***        | 0.380***        |
|   | (-0.004, 0.017) | (-0.049, -0.013) | (0.863, 0.935) | (-0.009, -0.004) | (0.294, 0.466)  |
| 2018  | -0.018          | -0.077**         | 0.812**        | -0.012**         | 0.268           |
|   | (-0.064, 0.028) | (-0.122, -0.032) | (0.747, 0.884) | (-0.017, -0.008) | (-0.219, 0.756) |
| 2019  | 0.026           | 0.065            | 1.138*         | 0.009            | 0.063           |
|   | (-0.028, 0.079) | (-0.004, 0.134)  | (1.006, 1.286) | (-0.000, 0.018)  | (-0.434, 0.559) |
| <b>Region (Northeast - REF)</b>             |                 |                  |                |                  |                 |
| North central, Midwest                      | 0.053***        | 0.104***         | 1.251***       | 0.013***         | 0.230**         |
|   | (0.043, 0.062)  | (0.077, 0.130)   | (1.180, 1.326) | (0.010, 0.017)   | (0.134, 0.325)  |
| South                                       | 0.042***        | 0.078***         | 1.185***       | 0.010***         | 0.234           |
|   | (0.030, 0.054)  | (0.057, 0.100)   | (1.122, 1.252) | (0.007, 0.013)   | (-0.004, 0.471) |
| West  | 0.026           | 0.064            | 1.164          | 0.009            | 0.057           |
|   | (-0.038, 0.091) | (-0.055, 0.183)  | (0.909, 1.491) | (-0.006, 0.024)  | (-0.304, 0.418) |

| <b>Income (Poor to low income - REF)</b> |                 |                 |                |                 |                 |
|--|-----------------|-----------------|----------------|-----------------|-----------------|
| Middle to high income                    | -0.002          | -0.002          | 0.998          | 0               | -0.034          |
|  | (-0.017, 0.014) | (-0.064, 0.061) | (0.862, 1.156) | (-0.009, 0.009) | (-0.304, 0.235) |
| <b>Receipt of food stamp (No - REF)</b>  |                 |                 |                |                 |                 |
| YES                                      | 0.036           | 0.044           | 1.07           | 0.004           | 0.382**         |
|  | (-0.039, 0.112) | (-0.102, 0.189) | (0.760, 1.506) | (-0.018, 0.027) | (0.174, 0.590)  |
| <b>Health status (Excellent - REF)</b>   |                 |                 |                |                 |                 |
| Good                                     | 0.066***        | 0.162***        | 1.466***       | 0.020***        | 0.223*          |
|  | (0.057, 0.076)  | (0.142, 0.182)  | (1.388, 1.548) | (0.018, 0.023)  | (0.043, 0.403)  |
| Fair                                     | 0.189***        | 0.321***        | 1.905***       | 0.039***        | 1.058**         |
|  | (0.160, 0.218)  | (0.255, 0.386)  | (1.602, 2.266) | (0.027, 0.051)  | (0.673, 1.443)  |
| Poor                                     | 0.290**         | 0.488***        | 2.589***       | 0.066**         | 1.183***        |
|  | (0.159, 0.421)  | (0.307, 0.670)  | (1.915, 3.500) | (0.034, 0.098)  | (0.963, 1.402)  |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 9b. Regressions for the Use of Medications with Sedative Properties:  $\geq 46$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Sedative Properties Medications = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|--|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |  |
| $\geq 46$ h/week                                     | -0.005<br>(-0.117, 0.108)     | 0.002<br>(-0.060, 0.065)      | 1.009<br>(0.970, 1.049)    | 0.002<br>(-0.007, 0.011)      | -0.028<br>(-0.303, 0.247)  |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |  |
| 27-64 years  | 0.260***<br>(0.229, 0.291)    | 0.225***<br>(0.179, 0.271)    | 1.289**<br>(1.166, 1.424)  | 0.054**<br>(0.034, 0.075)     | 0.381*<br>(0.080, 0.683)   |
| $\geq 65$ years                                      | 0.332*<br>(0.115, 0.550)      | 0.317**<br>(0.173, 0.461)     | 1.468***<br>(1.276, 1.689) | 0.084***<br>(0.054, 0.114)    | 0.364<br>(-0.054, 0.782)   |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |  |
| Black  | -0.262***<br>(-0.291, -0.233) | -0.117***<br>(-0.134, -0.100) | 0.963**<br>(0.945, 0.982)  | -0.008**<br>(-0.013, -0.004)  | -0.646***<br>(-0.699, -0.593)  |
| Other  | -0.289***<br>(-0.328, -0.251) | -0.188***<br>(-0.227, -0.150) | 0.855***<br>(0.825, 0.885) | -0.034***<br>(-0.042, -0.027) | -0.589***<br>(-0.667, -0.511)  |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |  |
| Female   | -0.003<br>(-0.025, 0.018)     | -0.001<br>(-0.017, 0.016)     | 1.003<br>(0.964, 1.043)    | 0.001<br>(-0.008, 0.009)      | -0.022<br>(-0.132, 0.088)  |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |  |
| Divorced   | 0.175***<br>(0.147, 0.204)    | 0.098***<br>(0.081, 0.115)    | 1.069***<br>(1.045, 1.093) | 0.015***<br>(0.010, 0.020)    | 0.328***<br>(0.265, 0.391)   |
| Unmarried  | -0.059<br>(-0.197, 0.080)     | -0.069<br>(-0.160, 0.022)     | 0.907*<br>(0.827, 0.994)   | -0.021*<br>(-0.042, -0.001)   | -0.006<br>(-0.243, 0.230)  |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |  |
| 4-6 Members  | -0.218***<br>(-0.247, -0.188) | -0.178***<br>(-0.200, -0.155) | 0.822***<br>(0.813, 0.832) | -0.043***<br>(-0.045, -0.041) | -0.295***<br>(-0.349, -0.242)  |
| > 6 Members  | -0.457***<br>(-0.580, -0.335) | -0.367**<br>(-0.521, -0.213)  | 0.659**<br>(0.537, 0.809)  | -0.088**<br>(-0.128, -0.048)  | -0.664***<br>(-0.874, -0.454)  |
| <b>Education (Less than college - REF)</b>           |                               |                               |                            |                               |  |

|   |                             |                               |                            |                               |                            |
|---|-----------------------------|-------------------------------|----------------------------|-------------------------------|----------------------------|
| Some college or more                        | -0.022<br>(-0.057, 0.012)   | -0.014<br>(-0.035, 0.007)     | 0.986<br>(0.960, 1.012)    | -0.003<br>(-0.009, 0.003)     | -0.023<br>(-0.114, 0.067)  |
| <b>Occupation (Natural resources - REF)</b> |                             |                               |                            |                               |                            |
| Hospitality services                        | 0.012<br>(-0.355, 0.379)    | 0.047<br>(-0.199, 0.292)      | 1.106<br>(0.883, 1.384)    | 0.022<br>(-0.027, 0.071)      | -0.201<br>(-0.876, 0.475)  |
| Trade                                       | 0.044<br>(-0.350, 0.438)    | 0.045<br>(-0.175, 0.266)      | 1.069<br>(0.933, 1.225)    | 0.015<br>(-0.015, 0.044)      | -0.022<br>(-0.966, 0.922)  |
| Professional services                       | 0.04<br>(-0.268, 0.348)     | 0.065<br>(-0.093, 0.223)      | 1.120*<br>(1.020, 1.229)   | 0.025*<br>(0.005, 0.045)      | -0.134<br>(-0.904, 0.637)  |
| Manufacturing                               | -0.042<br>(-0.362, 0.277)   | 0.017<br>(-0.183, 0.217)      | 1.082<br>(0.914, 1.283)    | 0.017<br>(-0.020, 0.054)      | -0.303<br>(-0.911, 0.305)  |
| Other                                       | -0.024<br>(-0.297, 0.248)   | 0.036<br>(-0.125, 0.197)      | 1.115<br>(0.992, 1.253)    | 0.024<br>(-0.001, 0.049)      | -0.319<br>(-0.962, 0.324)  |
| <b>Year (2010 - REF)</b>                    |                             |                               |                            |                               |                            |
| 2011  | 0.027<br>(-0.034, 0.088)    | -0.03<br>(-0.081, 0.020)      | 0.908*<br>(0.849, 0.971)   | -0.022*<br>(-0.037, -0.007)   | 0.281***<br>(0.248, 0.313) |
| 2012  | -0.079*<br>(-0.137, -0.021) | -0.115***<br>(-0.155, -0.076) | 0.831***<br>(0.785, 0.879) | -0.042***<br>(-0.055, -0.029) | 0.130***<br>(0.086, 0.173) |
| 2013  | -0.003<br>(-0.092, 0.085)   | -0.061*<br>(-0.105, -0.016)   | 0.877**<br>(0.825, 0.934)  | -0.030**<br>(-0.043, -0.016)  | 0.231*<br>(0.004, 0.458)   |
| 2014  | 0.022<br>(-0.170, 0.215)    | -0.051<br>(-0.125, 0.024)     | 0.879***<br>(0.869, 0.890) | -0.029***<br>(-0.032, -0.026) | 0.289<br>(-0.240, 0.818)   |
| 2015  | -0.02<br>(-0.174, 0.133)    | -0.057<br>(-0.146, 0.032)     | 0.897*<br>(0.824, 0.975)   | -0.025*<br>(-0.044, -0.006)   | 0.149<br>(-0.171, 0.469)   |
| 2016  | -0.036<br>(-0.123, 0.051)   | -0.079*<br>(-0.158, -0.000)   | 0.868*<br>(0.779, 0.967)   | -0.032*<br>(-0.056, -0.008)   | 0.160**<br>(0.096, 0.224)  |
| 2017  | -0.017<br>(-0.049, 0.015)   | -0.095**<br>(-0.133, -0.058)  | 0.825**<br>(0.753, 0.904)  | -0.043**<br>(-0.064, -0.023)  | 0.303*<br>(0.084, 0.522)   |
| 2018  | 0.016<br>(-0.068, 0.100)    | -0.085*<br>(-0.156, -0.014)   | 0.822**<br>(0.734, 0.919)  | -0.044**<br>(-0.069, -0.019)  | 0.403***<br>(0.342, 0.464) |
| 2019  | 0.004<br>(-0.131, 0.139)    | -0.015<br>(-0.111, 0.080)     | 0.956<br>(0.862, 1.061)    | -0.01<br>(-0.034, 0.014)      | 0.117<br>(-0.136, 0.371)   |
| <b>Region (Northeast - REF)</b>             |                             |                               |                            |                               |                            |
| North central, Midwest                      | 0.133***<br>(0.102, 0.164)  | 0.095***<br>(0.072, 0.118)    | 1.090***<br>(1.059, 1.122) | 0.019***<br>(0.013, 0.026)    | 0.241***<br>(0.215, 0.267) |



|  |                            |                            |                            |                            |                            |
|--|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| South                                    | 0.125***<br>(0.089, 0.162) | 0.092***<br>(0.069, 0.115) | 1.089***<br>(1.071, 1.107) | 0.019***<br>(0.015, 0.023) | 0.230***<br>(0.145, 0.314) |
| West                                     | 0.009<br>(-0.055, 0.074)   | 0.023<br>(-0.010, 0.056)   | 1.036*<br>(1.010, 1.063)   | 0.008*<br>(0.002, 0.013)   | -0.012<br>(-0.165, 0.140)  |
| <b>Income (Poor to low income - REF)</b> |                            |                            |                            |                            |                            |
| Middle to high income                    | -0.015<br>(-0.039, 0.008)  | -0.016<br>(-0.041, 0.008)  | 0.978<br>(0.939, 1.018)    | -0.005<br>(-0.014, 0.004)  | -0.005<br>(-0.062, 0.053)  |
| <b>Receipt of food stamp (No - REF)</b>  |                            |                            |                            |                            |                            |
| YES                                      | 0.309***<br>(0.209, 0.409) | 0.213***<br>(0.169, 0.258) | 1.204**<br>(1.116, 1.299)  | 0.042**<br>(0.025, 0.060)  | 0.488**<br>(0.200, 0.776)  |
| <b>Health status (Excellent - REF)</b>   |                            |                            |                            |                            |                            |
| Good                                     | 0.359***<br>(0.311, 0.407) | 0.302***<br>(0.251, 0.353) | 1.411***<br>(1.337, 1.490) | 0.073***<br>(0.062, 0.084) | 0.582***<br>(0.495, 0.669) |
| Fair                                     | 1.073***<br>(1.008, 1.138) | 0.782***<br>(0.731, 0.833) | 2.112***<br>(2.033, 2.194) | 0.168***<br>(0.160, 0.176) | 1.578***<br>(1.382, 1.774) |
| Poor                                     | 1.855***<br>(1.790, 1.920) | 1.223***<br>(1.181, 1.265) | 2.637***<br>(2.471, 2.814) | 0.223***<br>(0.206, 0.240) | 2.691***<br>(2.418, 2.964) |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 10a. Regressions for the Use of Prescription Sleep Aid:  $\geq 66$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Prescription Sleep Aid = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|---|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |   |
| $\geq 66$ h/week                                     | 0.091<br>(-0.085, 0.266)      | 0.143<br>(-0.119, 0.406)      | 1.28<br>(0.781, 2.098)     | 0.017<br>(-0.020, 0.054)      | 0.566*<br>(0.104, 1.028)  |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |   |
| 27-64 years  | 0.116***<br>(0.084, 0.147)    | 0.335***<br>(0.256, 0.414)    | 2.328***<br>(1.853, 2.924) | 0.039***<br>(0.031, 0.047)    | 0.366*<br>(0.006, 0.726)  |
| $\geq 65$ years                                      | 0.069<br>(-0.006, 0.144)      | 0.256***<br>(0.112, 0.400)    | 1.993***<br>(1.401, 2.835) | 0.030***<br>(0.013, 0.047)    | 0.077<br>(-0.136, 0.291)  |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |   |
| Black  | -0.136***<br>(-0.164, -0.108) | -0.331***<br>(-0.395, -0.266) | 0.442***<br>(0.370, 0.527) | -0.039***<br>(-0.046, -0.033) | -0.377<br>(-0.758, 0.004)   |
| Other  | -0.107***<br>(-0.138, -0.077) | -0.219***<br>(-0.305, -0.133) | 0.618***<br>(0.493, 0.774) | -0.027***<br>(-0.037, -0.016) | -0.667***<br>(-0.845, -0.489)                                       |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |   |
| Female   | 0.012<br>(-0.015, 0.039)      | 0.03<br>(-0.025, 0.085)       | 1.064<br>(0.940, 1.205)    | 0.004<br>(-0.004, 0.012)      | 0.042<br>(-0.432, 0.515)  |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |   |
| Divorced   | 0.044*<br>(0.006, 0.083)      | 0.084*<br>(0.016, 0.152)      | 1.204*<br>(1.040, 1.395)   | 0.012*<br>(0.002, 0.022)      | 0.048<br>(-0.196, 0.293)  |
| Unmarried  | 0<br>(-0.031, 0.030)          | 0.022<br>(-0.042, 0.087)      | 1.072<br>(0.923, 1.246)    | 0.004<br>(-0.005, 0.014)      | -0.209<br>(-0.502, 0.083)   |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |   |
| 4-6 Members  | -0.067***<br>(-0.095, -0.039) | -0.157***<br>(-0.215, -0.099) | 0.698***<br>(0.607, 0.801) | -0.021***<br>(-0.028, -0.013) | -0.169<br>(-0.395, 0.058)   |
| > 6 Members  | -0.104**<br>(-0.166, -0.041)  | -0.284**<br>(-0.499, -0.070)  | 0.485*<br>(0.257, 0.917)   | -0.036**<br>(-0.059, -0.013)  | 0.017<br>(-1.261, 1.296)  |
| <b>Education (Less than college - REF)</b>           |                               |                               |                            |                               |   |
| Some college or more                                 | 0.014<br>(-0.017, 0.046)      | 0.056<br>(-0.003, 0.115)      | 1.141<br>(1.000, 1.302)    | 0.008*<br>(0.000, 0.016)      | -0.068**<br>(-0.094, -0.043)  |
| <b>Occupation (Natural resources - REF)</b>          |                               |                               |                            |                               |   |

|  |                 |                 |                |                  |                 |
|--|-----------------|-----------------|----------------|------------------|-----------------|
| Hospitality services                     | 0.038           | 0.062           | 1.161          | 0.008            | 0.103           |
|  | (-0.032, 0.109) | (-0.087, 0.211) | (0.815, 1.654) | (-0.011, 0.027)  | (-0.314, 0.521) |
| Trade                                    | 0.031           | 0.056           | 1.116          | 0.006            | 0.178           |
|  | (-0.042, 0.104) | (-0.095, 0.208) | (0.780, 1.599) | (-0.013, 0.025)  | (-0.260, 0.615) |
| Professional services                    | 0.072*          | 0.149*          | 1.399*         | 0.020*           | 0.161           |
|  | (0.013, 0.130)  | (0.025, 0.273)  | (1.043, 1.876) | (0.004, 0.035)   | (-0.187, 0.508) |
| Manufacturing                            | 0.04            | 0.084           | 1.21           | 0.011            | 0.129           |
|  | (-0.022, 0.103) | (-0.047, 0.216) | (0.888, 1.650) | (-0.006, 0.027)  | (-0.044, 0.301) |
| Other                                    | 0.032           | 0.08            | 1.215          | 0.011            | -0.045          |
|  | (-0.027, 0.091) | (-0.047, 0.207) | (0.898, 1.644) | (-0.005, 0.027)  | (-0.326, 0.236) |
| <b>Year (2010 - REF)</b>                 |                 |                 |                |                  |                 |
| 2011                                     | 0.022           | 0.012           | 1.011          | 0.001            | 0.262           |
|  | (-0.022, 0.066) | (-0.067, 0.091) | (0.851, 1.202) | (-0.011, 0.012)  | (-0.417, 0.941) |
| 2012                                     | 0.007           | -0.04           | 0.874          | -0.008           | 0.460*          |
|  | (-0.042, 0.056) | (-0.136, 0.055) | (0.700, 1.090) | (-0.022, 0.005)  | (0.115, 0.804)  |
| 2013                                     | 0.005           | -0.033          | 0.903          | -0.006           | 0.352           |
|  | (-0.037, 0.047) | (-0.118, 0.053) | (0.745, 1.094) | (-0.018, 0.006)  | (-0.510, 1.215) |
| 2014                                     | 0.008           | -0.002          | 0.987          | -0.001           | 0.126           |
|  | (-0.034, 0.051) | (-0.088, 0.084) | (0.812, 1.200) | (-0.013, 0.012)  | (-0.775, 1.027) |
| 2015                                     | -0.009          | -0.046          | 0.886          | -0.007           | 0.192           |
|  | (-0.056, 0.038) | (-0.138, 0.045) | (0.721, 1.090) | (-0.020, 0.005)  | (-0.749, 1.133) |
| 2016                                     | 0.004           | -0.004          | 0.978          | -0.001           | 0.155           |
|  | (-0.037, 0.044) | (-0.091, 0.083) | (0.803, 1.190) | (-0.014, 0.011)  | (-0.663, 0.973) |
| 2017                                     | 0.007           | -0.03           | 0.9            | -0.007           | 0.382***        |
|  | (-0.034, 0.048) | (-0.114, 0.054) | (0.741, 1.094) | (-0.019, 0.006)  | (0.300, 0.463)  |
| 2018                                     | -0.018          | -0.076          | 0.814*         | -0.012*          | 0.267           |
|  | (-0.059, 0.024) | (-0.164, 0.011) | (0.665, 0.996) | (-0.024, -0.000) | (-0.246, 0.780) |
| 2019                                     | 0.026           | 0.066           | 1.14           | 0.009            | 0.07            |
|  | (-0.023, 0.075) | (-0.034, 0.166) | (0.918, 1.416) | (-0.006, 0.024)  | (-0.429, 0.570) |
| <b>Region (Northeast - REF)</b>          |                 |                 |                |                  |                 |
| North central, Midwest                   | 0.052*          | 0.102*          | 1.250*         | 0.013*           | 0.214*          |
|  | (0.004, 0.101)  | (0.007, 0.197)  | (1.006, 1.552) | (0.001, 0.026)   | (0.076, 0.352)  |
| South                                    | 0.041           | 0.078           | 1.183          | 0.01             | 0.229           |
|  | (-0.003, 0.085) | (-0.011, 0.166) | (0.965, 1.451) | (-0.002, 0.021)  | (-0.007, 0.464) |
| West                                     | 0.026           | 0.063           | 1.162          | 0.009            | 0.062           |
|  | (-0.017, 0.068) | (-0.027, 0.154) | (0.938, 1.438) | (-0.003, 0.021)  | (-0.300, 0.425) |
| <b>Income (Poor to low income - REF)</b> |                 |                 |                |                  |                 |

|   |                            |                            |                            |                            |                            |
|---|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| Middle to high income                   | -0.002<br>(-0.034, 0.030)  | -0.001<br>(-0.069, 0.067)  | 1<br>(0.856, 1.167)        | 0<br>(-0.010, 0.010)       | -0.034<br>(-0.295, 0.226)  |
| <b>Receipt of food stamp (No - REF)</b> |                            |                            |                            |                            |                            |
| YES                                     | 0.036<br>(-0.016, 0.087)   | 0.042<br>(-0.057, 0.140)   | 1.068<br>(0.859, 1.327)    | 0.004<br>(-0.010, 0.018)   | 0.361**<br>(0.150, 0.573)  |
| <b>Health status (Excellent - REF)</b>  |                            |                            |                            |                            |                            |
| Good                                    | 0.067***<br>(0.036, 0.098) | 0.163***<br>(0.088, 0.238) | 1.466***<br>(1.209, 1.779) | 0.021***<br>(0.011, 0.030) | 0.239*<br>(0.069, 0.410)   |
| Fair                                    | 0.188***<br>(0.126, 0.250) | 0.320***<br>(0.219, 0.421) | 1.900***<br>(1.530, 2.361) | 0.039***<br>(0.026, 0.052) | 1.074***<br>(0.744, 1.404) |
| Poor                                    | 0.290***<br>(0.137, 0.443) | 0.489***<br>(0.287, 0.692) | 2.588***<br>(1.781, 3.761) | 0.066***<br>(0.033, 0.099) | 1.210***<br>(0.982, 1.437) |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

Appendix 10b. Regressions for the Use of Medications with Sedative Properties:  $\geq 66$  h/week

| Variables  | Linear Regression             | Tobit Marginal Effect         | Adjusted Logit Odds Ratio  | Logit Marginal Effect         | Linear Regression Among Users (Use of Sedative Properties Medications = YES) |
|--|-------------------------------|-------------------------------|----------------------------|-------------------------------|--|
| <b>Work hours/week</b>                               |                               |                               |                            |                               |  |
| $\geq 66$ h/week                                     | 0.093<br>(-0.156, 0.342)      | 0.063<br>(-0.108, 0.233)      | 1.056<br>(0.889, 1.255)    | 0.012<br>(-0.027, 0.051)      | 0.162*<br>(0.012, 0.312)   |
| <b>Age (18-26years - REF)</b>                        |                               |                               |                            |                               |  |
| 27-64 years  | 0.259***<br>(0.108, 0.411)    | 0.225***<br>(0.147, 0.303)    | 1.289***<br>(1.194, 1.391) | 0.054***<br>(0.039, 0.070)    | 0.379*<br>(0.059, 0.698)   |
| $\geq 65$ years                                      | 0.331**<br>(0.123, 0.540)     | 0.316***<br>(0.197, 0.435)    | 1.468***<br>(1.302, 1.655) | 0.084***<br>(0.057, 0.111)    | 0.358<br>(-0.055, 0.772)   |
| <b>Race (White - REF)</b>                            |                               |                               |                            |                               |  |
| Black  | -0.262***<br>(-0.330, -0.194) | -0.117***<br>(-0.164, -0.070) | 0.963<br>(0.911, 1.018)    | -0.008<br>(-0.021, 0.004)     | -0.646***<br>(-0.696, -0.597)  |
| Other  | -0.289***<br>(-0.368, -0.210) | -0.188***<br>(-0.253, -0.123) | 0.854***<br>(0.787, 0.927) | -0.034***<br>(-0.052, -0.017) | -0.589***<br>(-0.660, -0.517)  |
| <b>Sex (Male - REF)</b>                              |                               |                               |                            |                               |  |
| Female   | 0<br>(-0.058, 0.057)          | 0.001<br>(-0.039, 0.040)      | 1.003<br>(0.960, 1.049)    | 0.001<br>(-0.009, 0.011)      | -0.014<br>(-0.108, 0.080)  |
| <b>Marital status (Married - REF)</b>                |                               |                               |                            |                               |  |
| Divorced   | 0.175***<br>(0.080, 0.270)    | 0.097**<br>(0.035, 0.159)     | 1.068*<br>(1.005, 1.136)   | 0.015*<br>(0.001, 0.029)      | 0.327***<br>(0.262, 0.392)   |
| Unmarried  | -0.058<br>(-0.137, 0.021)     | -0.069**<br>(-0.119, -0.019)  | 0.907***<br>(0.858, 0.958) | -0.021***<br>(-0.034, -0.009) | -0.005<br>(-0.243, 0.234)  |
| <b>Family size (<math>\leq 3</math> Members REF)</b> |                               |                               |                            |                               |  |
| 4-6 Members  | -0.217***<br>(-0.279, -0.156) | -0.177***<br>(-0.224, -0.130) | 0.823***<br>(0.779, 0.868) | -0.043***<br>(-0.055, -0.031) | -0.295***<br>(-0.350, -0.239)  |
| > 6 Members  | -0.458***<br>(-0.599, -0.316) | -0.367***<br>(-0.493, -0.241) | 0.659***<br>(0.552, 0.786) | -0.088***<br>(-0.123, -0.054) | -0.663***<br>(-0.876, -0.450)  |
| <b>Education (Less than college - REF)</b>           |                               |                               |                            |                               |  |
| Some college or more                                 | -0.022<br>(-0.105, 0.061)     | -0.014<br>(-0.065, 0.038)     | 0.987<br>(0.939, 1.036)    | -0.003<br>(-0.014, 0.008)     | -0.023<br>(-0.103, 0.057)  |
| <b>Occupation (Natural resources - REF)</b>          |                               |                               |                            |                               |  |
| Hospitality services                                 | 0.013<br>(-0.212, 0.239)      | 0.047<br>(-0.079, 0.173)      | 1.106<br>(0.983, 1.244)    | 0.022<br>(-0.004, 0.048)      | -0.195<br>(-0.849, 0.458)  |
| Trade  | 0.045                         | 0.046                         | 1.069                      | 0.015                         | -0.019   |

|  |                 |                  |                |                  |                 |
|--|-----------------|------------------|----------------|------------------|-----------------|
|  | (-0.174, 0.264) | (-0.079, 0.171)  | (0.955, 1.198) | (-0.010, 0.039)  | (-0.945, 0.907) |
| Professional services                    | 0.041           | 0.066            | 1.120*         | 0.025*           | -0.13           |
|  | (-0.147, 0.229) | (-0.037, 0.169)  | (1.020, 1.230) | (0.005, 0.045)   | (-0.889, 0.630) |
| Manufacturing                            | -0.04           | 0.018            | 1.083          | 0.017            | -0.297          |
|  | (-0.244, 0.164) | (-0.101, 0.137)  | (0.967, 1.214) | (-0.007, 0.042)  | (-0.899, 0.306) |
| Other                                    | -0.023          | 0.036            | 1.115*         | 0.024*           | -0.315          |
|  | (-0.201, 0.155) | (-0.064, 0.136)  | (1.013, 1.226) | (0.003, 0.045)   | (-0.943, 0.313) |
| <b>Year (2010 - REF)</b>                 |                 |                  |                |                  |                 |
| 2011                                     | 0.026           | -0.031           | 0.907*         | -0.022*          | 0.278***        |
|  | (-0.075, 0.127) | (-0.098, 0.036)  | (0.838, 0.981) | (-0.040, -0.004) | (0.242, 0.315)  |
| 2012                                     | -0.079          | -0.115**         | 0.830***       | -0.042***        | 0.129***        |
|  | (-0.201, 0.042) | (-0.198, -0.033) | (0.757, 0.911) | (-0.062, -0.021) | (0.086, 0.172)  |
| 2013                                     | -0.003          | -0.06            | 0.878**        | -0.030**         | 0.232           |
|  | (-0.124, 0.118) | (-0.142, 0.021)  | (0.804, 0.958) | (-0.049, -0.010) | (-0.004, 0.469) |
| 2014                                     | 0.022           | -0.051           | 0.880**        | -0.029**         | 0.29            |
|  | (-0.131, 0.175) | (-0.142, 0.040)  | (0.804, 0.962) | (-0.049, -0.009) | (-0.240, 0.819) |
| 2015                                     | -0.02           | -0.057           | 0.897*         | -0.025*          | 0.149           |
|  | (-0.143, 0.103) | (-0.140, 0.027)  | (0.819, 0.982) | (-0.045, -0.004) | (-0.168, 0.467) |
| 2016                                     | -0.035          | -0.079*          | 0.868**        | -0.032***        | 0.161**         |
|  | (-0.145, 0.075) | (-0.156, -0.002) | (0.799, 0.944) | (-0.051, -0.013) | (0.096, 0.226)  |
| 2017                                     | -0.016          | -0.095*          | 0.826***       | -0.043***        | 0.304*          |
|  | (-0.128, 0.096) | (-0.171, -0.018) | (0.756, 0.901) | (-0.062, -0.023) | (0.086, 0.522)  |
| 2018                                     | 0.017           | -0.085*          | 0.822***       | -0.044***        | 0.403***        |
|  | (-0.105, 0.138) | (-0.168, -0.002) | (0.752, 0.898) | (-0.064, -0.024) | (0.342, 0.465)  |
| 2019                                     | 0.005           | -0.015           | 0.956          | -0.01            | 0.119           |
|  | (-0.114, 0.124) | (-0.098, 0.068)  | (0.871, 1.050) | (-0.032, 0.011)  | (-0.137, 0.375) |
| <b>Region (Northeast - REF)</b>          |                 |                  |                |                  |                 |
| North central, Midwest                   | 0.132*          | 0.095*           | 1.090*         | 0.019*           | 0.239***        |
|  | (0.022, 0.242)  | (0.018, 0.172)   | (1.002, 1.186) | (0.001, 0.038)   | (0.207, 0.270)  |
| South                                    | 0.125*          | 0.092*           | 1.089*         | 0.019*           | 0.229***        |
|  | (0.019, 0.230)  | (0.017, 0.166)   | (1.003, 1.182) | (0.001, 0.037)   | (0.146, 0.312)  |
| West                                     | 0.009           | 0.023            | 1.036          | 0.008            | -0.012          |
|  | (-0.087, 0.105) | (-0.047, 0.092)  | (0.957, 1.121) | (-0.010, 0.025)  | (-0.160, 0.136) |
| <b>Income (Poor to low income - REF)</b> |                 |                  |                |                  |                 |
| Middle to high income                    | -0.016          | -0.017           | 0.978          | -0.005           | -0.008          |
|  | (-0.092, 0.059) | (-0.069, 0.036)  | (0.922, 1.038) | (-0.018, 0.008)  | (-0.040, 0.023) |
| <b>Receipt of food stamp (No - REF)</b>  |                 |                  |                |                  |                 |
| YES                                      | 0.309***        | 0.213***         | 1.204***       | 0.042***         | 0.486**         |
|  | (0.158, 0.460)  | (0.123, 0.303)   | (1.114, 1.301) | (0.024, 0.060)   | (0.200, 0.772)  |
| <b>Health status (Excellent - REF)</b>   |                 |                  |                |                  |                 |
| Good                                     | 0.359***        | 0.302***         | 1.411***       | 0.073***         | 0.585***        |

|      |                |                |                |                |                |
|------|----------------|----------------|----------------|----------------|----------------|
|      | (0.293, 0.426) | (0.246, 0.359) | (1.323, 1.505) | (0.060, 0.086) | (0.485, 0.685) |
| Fair | 1.073***       | 0.781***       | 2.111***       | 0.168***       | 1.581***       |
|      | (0.937, 1.209) | (0.688, 0.875) | (1.946, 2.290) | (0.150, 0.186) | (1.367, 1.795) |
| Poor | 1.855***       | 1.223***       | 2.637***       | 0.223***       | 2.694***       |
|      | (1.450, 2.261) | (0.962, 1.485) | (2.157, 3.223) | (0.174, 0.272) | (2.414, 2.975) |

Note: Confidence Interval in parentheses; \*\*\* p<0.001, \*\* p<0.01, \* p<0.05; REF = reference

## APPENDIX CHAPTER 2:

Appendix 1: Medications with sedative properties (Comparisons eAnswers., 2021).

### Drug Facts And Comparisons

Acetaminophen and Codeine Oral

Acetaminophen, Caffeine, and Dihydrocodeine Oral

Alfentanil Injection

Alosetron Oral

ALPRAZolam Oral

Amitriptyline Oral

Ampicillin and Sulbactam Injection

ARIPiprazole Injection

ARIPiprazole Oral

Azelastine and Fluticasone Intranasal

Benzonatate Oral

Brexpiprazole Oral

Brivaracetam Injection

Brivaracetam Oral

Brompheniramine, Pseudoephedrine, and Dextromethorphan Oral

Buprenorphine Implant

Buprenorphine Injection

Buprenorphine Oral

Buprenorphine Transdermal

Butalbital, Acetaminophen, and Caffeine Oral

Cannabidiol Oral

Capecitabine Oral

Carbinoxamine Oral

Cenobamate Oral

Cetirizine Injection

Cetirizine Oral

Chlorpheniramine and Phenylephrine Oral

Chlorpheniramine, Phenylephrine, and Dextromethorphan Oral

Chlorpheniramine, Pseudoephedrine, and Codeine Oral

Chlorpheniramine, Pseudoephedrine, and Dextromethorphan Oral

CloBAZam Oral

Clonidine and Clorthalidone Oral

CloNIDine Injection

CloNIDine Oral

CloNIDine Transdermal

CloZAPine Oral

Codeine and Chlorpheniramine Oral

Codeine Oral

Cyproheptadine Oral

Deutetrabenazine Oral

Dexchlorpheniramine Oral

DiphenhydrAMINE Injection

DiphenhydrAMINE Oral

Diphenoxylate and Atropine Oral

Doxepin Oral

Eluxadoline Oral

Esketamine Intranasal

Fenfluramine Oral

FentaNYL Buccal

FentaNYL Injection

FentaNYL Intranasal



[FentaNYL Sublingual](#)  
[FentaNYL Transdermal](#)  
[Flibanserin Oral](#)  
[Flucytosine Oral](#)  
[Flurazepam Oral](#)  
[Gabapentin Enacarbil Oral](#)  
[Guaifenesin and Codeine Oral](#)  
[GuanFACINE Oral](#)  
[Haloperidol Injection](#)  
[Haloperidol Oral](#)  
[Hydrocodone and Acetaminophen Oral](#)  
[Hydrocodone and Chlorpheniramine Oral](#)  
[Hydrocodone and Homatropine Oral](#)  
[Hydrocodone and Pseudoephedrine Oral](#)  
[HYDRQcodone Oral](#)  
[HYDROmorphine Injection](#)  
[HYDROmorphine Oral](#)  
[HYDROmorphine Rectal](#)  
[Isocarboxazid Oral](#)  
[LevETIRAcetam Injection](#)  
[LevETIRAcetam Oral](#)  
[Lithium Oral](#)  
[Lofexidine Oral](#)  
[Loratadine Oral](#)  
[LORazepam Injection](#)  
[LORazepam Oral](#)  
[Loxapine Inhalation](#)  
[Loxapine Oral](#)  
[Lumateperone Oral](#)  
[Mecamylamine Oral](#)  
[Meperidine Injection](#)  
[Meperidine Oral](#)  
[Methadone Injection](#)  
[Methadone Oral](#)  
[Methocarbamol Injection](#)  
[Methocarbamol Oral](#)  
[Methyldopa and Hydrochlorothiazide Oral](#)  
[Methyldopa Oral](#)  
[Methyldopate Injection](#)  
[Methylphenidate Oral](#)  
[Methylphenidate Transdermal](#)  
[MetyraPONE Oral](#)  
[MetyroSINE Oral](#)  
[Midazolam Injection](#)  
[Midazolam Intranasal](#)  
[Midazolam Oral](#)  
[Mitotane Oral](#)  
[Morphine and Naltrexone Oral](#)  
[Morphine Injection](#)  
[Morphine Oral](#)  
[Morphine Rectal](#)  
[Nabilone Oral](#)  
[Nadolol and Bendroflumethiazide Oral](#)  
[Nadolol Oral](#)  
[Nalbuphine Injection](#)  
[Naltrexone Injection](#)  
[Naltrexone Oral](#)  
[Olanzapine and Samidorphan Oral](#)  
[OLANZapine Injection](#)  
[OLANZapine Oral](#)  
[Oliceridine Injection](#)  
[Ondansetron Injection](#)

Ondansetron Oral  
Oxaprozin Oral  
Oxycodone and Acetaminophen Oral  
Oxycodone and Aspirin Oral  
OxyMORphone Injection  
OxyMORphone Oral  
Paliperidone Injection  
Paliperidone Oral  
Papaverine Injection  
Pentazocine and Naloxone Oral  
Pentazocine Injection  
Perphenazine Oral  
PHENobarbital Oral  
Pimozide Oral  
Pregabalin Oral  
Promethazine and Codeine Oral  
Promethazine and Phenylephrine Oral  
Promethazine Injection  
Promethazine Oral  
Promethazine Rectal  
Promethazine, Phenylephrine, and Codeine Oral  
RisperiDONE Injection  
RisperiDONE Oral  
SUFentanil Injection  
SUFentanil Oral  
SUMatriptan Injection  
SUMatriptan Intranasal  
SUMatriptan Oral  
Suvorexant Oral  
Tapentadol Oral  
Tetrabenazine Oral  
TiZANidine Oral  
TraMADol Oral  
Triazolam Oral  
Valbenazine Oral  
Vigabatrin Oral  
Vilazodone Oral  
Viloxazine Oral

Appendix 2. Distribution of Psychological Distress in the Panel.

| <b>Psychological Distress</b> | <b>Year 1</b> | <b>Year 2</b> | <b>Total</b>  |
|-------------------------------|---------------|---------------|---------------|
| No                            | 27,562        | 25,110        | 52,672        |
| Yes                           | 0             | 2,452         | 2,452         |
| Yes %                         | 0             | 8.9           | 4             |
| <b>Total</b>                  | <b>27,562</b> | <b>27,562</b> | <b>55,124</b> |

*Note:* Balanced sample of the 27,562 individuals with the two years; No psychological distress at baseline (year 1).

Appendix 3. Transition frequencies in and out of work hour groups.

|  | Weekly Working Hours |        |       |       |       |
|--|----------------------|--------|-------|-------|-------|
|  | 30-35                | 36-40  | 41-50 | 51-60 | ≥ 61  |
| <b>Overall Sample</b>                    |                      |        |       |       |       |
| Observations in year 1                   | 2,903                | 16,695 | 5,221 | 1,990 | 753   |
| Frequency in                             | 172                  | 495    | 274   | 126   | 46    |
| Frequency out                            | 237                  | 399    | 281   | 141   | 55    |
| Frequency in and out                     | 409                  | 894    | 555   | 267   | 101   |
| Observations in year 2                   | 2,838                | 16,791 | 5,214 | 1,975 | 744   |
| Rate of movement as % year 1 observation | 14.1%                | 5.4%   | 10.6% | 13.4% | 13.4% |
| <b>All Men</b>                           |                      |        |       |       |       |
| Observations in year 1                   | 1,185                | 9,132  | 3,367 | 1,415 | 565   |
| Frequency in                             | 73                   | 298    | 180   | 101   | 35    |
| Frequency out                            | 123                  | 235    | 190   | 97    | 42    |
| Frequency in and out                     | 196                  | 533    | 370   | 198   | 77    |
| Observations in year 2                   | 1,135                | 9,195  | 3,357 | 1,419 | 558   |
| Rate of movement as % year 1 observation | 16.5%                | 5.8%   | 11.0% | 14.0% | 13.6% |
| <b>All Women</b>                         |                      |        |       |       |       |
| Observations in year 1                   | 1,718                | 7,563  | 1,854 | 575   | 188   |
| Frequency in                             | 99                   | 197    | 94    | 25    | 11    |
| Frequency out                            | 114                  | 164    | 91    | 44    | 13    |
| Frequency in and out                     | 213                  | 361    | 185   | 69    | 24    |
| Observations in year 2                   | 1,703                | 7,596  | 1,857 | 556   | 186   |
| Rate of movement as % year 1 observation | 12.4%                | 4.8%   | 10.0% | 12.0% | 12.8% |

*Note:* Frequency in: number of observations that entered a work hour subgroup after year 1; Frequency out: number of observations that exited the work hour subgroup after year 1.

Appendix 4. Transition frequencies out of each work hour group.

| Weekly Working Hours       | Overall       |               | All Men      |              | All Women    |              |
|----------------------------|---------------|---------------|--------------|--------------|--------------|--------------|
|                            | Year 1        | Year 2        | Year 1       | Year 2       | Year 1       | Year 2       |
| <b>30-35 h/week select</b> |               |               |              |              |              |              |
| 30-35 h/week               | 2,903         | 2,666         | 1,185        | 1,062        | 1,718        | 1,604        |
| 36-40 h/week               | 0             | 195           | 0            | 100          | 0            | 95           |
| 41-50 h/week               | 0             | 31            | 0            | 19           | 0            | 12           |
| 51-60 h/week               | 0             | 9             | 0            | 3            | 0            | 6            |
| ≥ 61 h/week                | 0             | 2             | 0            | 1            | 0            | 1            |
| <b>Total</b>               | <b>2,903</b>  | <b>2,903</b>  | <b>1,185</b> | <b>1,185</b> | <b>1,718</b> | <b>1,718</b> |
| <b>36-40 h/week select</b> |               |               |              |              |              |              |
| 30-35 h/week               | 0             | 138           | 0            | 55           | 0            | 83           |
| 36-40 h/week               | 16,695        | 16,296        | 9,132        | 8,897        | 7,563        | 7,399        |
| 41-50 h/week               | 0             | 186           | 0            | 119          | 0            | 67           |
| 51-60 h/week               | 0             | 62            | 0            | 53           | 0            | 9            |
| ≥ 61 h/week                | 0             | 13            | 0            | 8            | 0            | 5            |
| <b>Total</b>               | <b>16,695</b> | <b>16,695</b> | <b>9,132</b> | <b>9,132</b> | <b>7,563</b> | <b>7,563</b> |
| <b>41-50 h/week select</b> |               |               |              |              |              |              |
| 30-35 h/week               | 0             | 24            | 0            | 15           | 0            | 9            |
| 36-40 h/week               | 0             | 200           | 0            | 129          | 0            | 71           |
| 41-50 h/week               | 5,221         | 4,940         | 3,367        | 3,177        | 1,854        | 1,763        |
| 51-60 h/week               | 0             | 40            | 0            | 31           | 0            | 9            |
| ≥ 61 h/week                | 0             | 17            | 0            | 15           | 0            | 2            |
| <b>Total</b>               | <b>5,221</b>  | <b>5,221</b>  | <b>3,367</b> | <b>3,367</b> | <b>1,854</b> | <b>1,854</b> |
| <b>51-60 h/week select</b> |               |               |              |              |              |              |
| 30-35 h/week               | 0             | 6             | 0            | 3            | 0            | 3            |
| 36-40 h/week               | 0             | 79            | 0            | 51           | 0            | 28           |
| 41-50 h/week               | 0             | 42            | 0            | 32           | 0            | 10           |
| 51-60 h/week               | 1,990         | 1,849         | 1,415        | 1,318        | 575          | 531          |
| ≥ 61 h/week                | 0             | 14            | 0            | 11           | 0            | 3            |
| <b>Total</b>               | <b>1,990</b>  | <b>1,990</b>  | <b>1,415</b> | <b>1,415</b> | <b>575</b>   | <b>575</b>   |
| <b>≥ 61 h/week select</b>  |               |               |              |              |              |              |
| 30-35 h/week               | 0             | 4             | 0            | 0            | 0            | 4            |
| 36-40 h/week               | 0             | 21            | 0            | 18           | 0            | 3            |
| 41-50 h/week               | 0             | 15            | 0            | 10           | 0            | 5            |
| 51-60 h/week               | 0             | 15            | 0            | 14           | 0            | 1            |
| ≥ 61 h/week                | 753           | 698           | 565          | 523          | 188          | 175          |
| <b>Total</b>               | <b>753</b>    | <b>753</b>    | <b>565</b>   | <b>565</b>   | <b>188</b>   | <b>188</b>   |

## Appendix 5. Covariate Transitions

| Covariate Transitions             |                           |                 |                  |                         |                           |   |
|-----------------------------------|---------------------------|-----------------|------------------|-------------------------|---------------------------|---|
|                                   | Observations<br>in year 1 | Frequency<br>in | Frequency<br>out | Frequency<br>in and out | Observations<br>in year 2 | Rate of<br>movement as<br>% year 2<br>observation |
| Overall Sample - Age              |                           |                 |                  |                         |                           |   |
| 18-64yrs                          | 26,556                    | 0               | 228              | 228                     | 26,328                    | 0.9%  |
| ≥ 65yrs                           | 1,006                     | 228             | 0                | 228                     | 1,234                     | 18.5%   |
| All Men Sample - Age              |                           |                 |                  |                         |                           |   |
| 18-64yrs                          | 15,060                    | 0               | 132              | 132                     | 14,928                    | 0.9%  |
| ≥ 65yrs                           | 604                       | 132             | 0                | 132                     | 736                       | 17.9%   |
| All Women Sample - Age            |                           |                 |                  |                         |                           |   |
| 18-64yrs                          | 11,496                    | 0               | 96               | 96                      | 11,400                    | 0.8%  |
| ≥ 65yrs                           | 402                       | 96              | 0                | 96                      | 498                       | 19.3%   |
| Overall Sample - Marital Status   |                           |                 |                  |                         |                           |   |
| Married                           | 16,221                    | 443             | 186              | 629                     | 16,478                    | 3.9%  |
| Not married                       | 11,341                    | 186             | 443              | 629                     | 11,084                    | 5.5%  |
| All Men Sample - Marital Status   |                           |                 |                  |                         |                           |   |
| Married                           | 9,959                     | 264             | 90               | 354                     | 10,133                    | 3.6%  |
| Not married                       | 5,705                     | 90              | 264              | 354                     | 5,531                     | 6.2%  |
| All Women Sample - Marital Status |                           |                 |                  |                         |                           |   |
| Married                           | 6,262                     | 179             | 96               | 275                     | 6,345                     | 4.4%  |
| Not married                       | 5,636                     | 96              | 179              | 275                     | 5,553                     | 4.9%  |
| Overall Sample - Family size      |                           |                 |                  |                         |                           |   |
| ≤ 2<br>Members                    | 11,677                    | 724             | 461              | 1,185                   | 11,940                    | 10.1%   |
| > 2<br>Members                    | 15,885                    | 461             | 724              | 1,185                   | 15,622                    | 7.5%  |
| All Men Sample - Family size      |                           |                 |                  |                         |                           |   |
| ≤ 2<br>Members                    | 6,424                     | 363             | 276              | 639                     | 6,511                     | 9.9%  |
| > 2<br>Members                    | 9,240                     | 276             | 363              | 639                     | 9,153                     | 6.9%  |
| All Women Sample - Family size    |                           |                 |                  |                         |                           |   |
| ≤ 2<br>Members                    | 5,253                     | 361             | 185              | 546                     | 5,429                     | 10.4%   |
| > 2<br>Members                    | 6,645                     | 185             | 361              | 546                     | 6,469                     | 8.2%  |

Appendix 5. Covariate Transitions (Continued).

| Covariate Transitions                            |                           |                 |                  |                         |                           |   |
|--|---------------------------|-----------------|------------------|-------------------------|---------------------------|---|
|  | Observations<br>in year 1 | Frequency<br>in | Frequency<br>out | Frequency<br>in and out | Observations<br>in year 2 | Rate of<br>movement as<br>% year 2<br>observation |
| <b>Overall Sample - Occupation</b>               |                           |                 |                  |                         |                           |   |
| Professional<br>and<br>management<br>occupations | 10,796                    | 237             | 206              | 443                     | 10,827                    | 4.1%  |
| Other  | 16,766                    | 206             | 237              | 443                     | 16,735                    | 2.6%  |
| <b>All Men Sample - Occupation</b>               |                           |                 |                  |                         |                           |   |
| Professional<br>and<br>management<br>occupations | 5,454                     | 132             | 113              | 245                     | 5,473                     | 4.5%  |
| Other  | 10,210                    | 113             | 132              | 245                     | 10,191                    | 2.4%  |
| <b>All Women Sample - Occupation</b>             |                           |                 |                  |                         |                           |   |
| Professional<br>and<br>management<br>occupations | 5,342                     | 105             | 93               | 198                     | 5,354                     | 3.7%  |
| Other  | 6,556                     | 93              | 105              | 198                     | 6,544                     | 3.0%  |
| <b>Overall Sample - Income</b>                   |                           |                 |                  |                         |                           |   |
| Poor to low<br>income                            | 5876                      | 1425            | 1830             | 3255                    | 5471                      | 55.4%   |
| Middle to<br>high income                         | 21686                     | 1830            | 1425             | 3255                    | 22091                     | 15.0%   |
| <b>All Men Sample - Income</b>                   |                           |                 |                  |                         |                           |   |
| Poor to low<br>income                            | 3,306                     | 835             | 1,089            | 1,924                   | 3,052                     | 58.2%   |
| Middle to<br>high income                         | 12,358                    | 1,089           | 835              | 1,924                   | 12,612                    | 15.6%   |
| <b>All Women Sample - Income</b>                 |                           |                 |                  |                         |                           |   |
| Poor to low<br>income                            | 2,570                     | 590             | 741              | 1,331                   | 2,419                     | 51.8%   |
| Middle to<br>high income                         | 9,328                     | 741             | 590              | 1,331                   | 9,479                     | 14.3%   |
| <b>Overall Sample - Receipt of Food Stamps</b>   |                           |                 |                  |                         |                           |   |
| no   | 25,682                    | 705             | 601              | 1,306                   | 25,786                    | 5.1%  |
| yes  | 1,880                     | 601             | 705              | 1,306                   | 1,776                     | 69.5%   |
| <b>All Men Sample - Receipt of Food Stamps</b>   |                           |                 |                  |                         |                           |   |
| no   | 14,667                    | 378             | 332              | 710                     | 14,713                    | 4.8%  |
| yes  | 997                       | 332             | 378              | 710                     | 951                       | 71.2%   |
| <b>All Women Sample - Receipt of Food Stamps</b> |                           |                 |                  |                         |                           |   |
| no   | 11,015                    | 327             | 269              | 596                     | 11,073                    | 5.4%  |
| yes  | 883                       | 269             | 327              | 596                     | 825                       | 67.5%   |

*Note:* Frequency in: number of observations that entered a work hour subgroup after year 1; Frequency out: number of observations that exited the work hour subgroup after year 1.

Appendix 6: Association between long working hours and onset of psychological distress: fixed effects estimate without covariates.

| <b>Variable</b>                              | <b>Overall</b>            | <b>All Men</b>            | <b>All Women</b>          |
|--|---------------------------|---------------------------|---------------------------|
| <b>Work hours/week (30-35hrs/week - REF)</b> |                           |                           |                           |
| 36-40 h/week                                 | 0.037**<br>(0.001, 0.073) | 0.050*<br>(-0.004, 0.105) | 0.026<br>(-0.021, 0.073)  |
| 41-50 h/week                                 | 0.032<br>(-0.011, 0.075)  | 0.049<br>(-0.012, 0.110)  | 0.014<br>(-0.052, 0.080)  |
| 51-60 h/week                                 | 0.037<br>(-0.014, 0.088)  | 0.073**<br>(0.003, 0.143) | -0.035<br>(-0.120, 0.049) |
| ≥ 61 h/week                                  | 0.073**<br>(0.011, 0.135) | 0.083**<br>(0.001, 0.164) | 0.090<br>(-0.018, 0.198)  |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Results – Weighted

Appendix 7. Association Between long working Hours and the onset of psychological distress: Fixed effects regression analysis – Weighted.

| <b>Variables</b>                             | <b>Overall</b>               | <b>All Men</b>            | <b>All Women</b>              |
|--|------------------------------|---------------------------|-------------------------------|
| <b>Work hours/week (30-35hrs/week - REF)</b> |                              |                           |                               |
| 36-40 h/week                                 | 0.060**<br>(0.008, 0.111)    | 0.077*<br>(-0.007, 0.161) | 0.040<br>(-0.019, 0.100)      |
| 41-50 h/week                                 | 0.061**<br>(0.001, 0.121)    | 0.071<br>(-0.017, 0.158)  | 0.059<br>(-0.034, 0.153)      |
| 51-60 h/week                                 | 0.047<br>(-0.020, 0.113)     | 0.088*<br>(-0.010, 0.187) | -0.054<br>(-0.152, 0.043)     |
| ≥ 61 h/week                                  | 0.097**<br>(0.016, 0.178)    | 0.098*<br>(-0.007, 0.203) | 0.155*<br>(-0.018, 0.329)     |
| <b>Age (18-64years - REF)</b>                |                              |                           |                               |
| ≥ 65 years                                   | 0.044***<br>(0.017, 0.070)   | 0.046**<br>(0.010, 0.081) | 0.041**<br>(0.003, 0.079)     |
| <b>Marital Status (Married - REF)</b>        |                              |                           |                               |
| Not married                                  | -0.015<br>(-0.042, 0.013)    | -0.025<br>(-0.061, 0.012) | -0.000<br>(-0.041, 0.041)     |
| <b>Family Size (≤ 2 Members REF)</b>         |                              |                           |                               |
| > 2 Members                                  | -0.013<br>(-0.034, 0.009)    | 0.013<br>(-0.014, 0.040)  | -0.046***<br>(-0.080, -0.011) |
| <b>Occupation (Natural resources - REF)</b>  |                              |                           |                               |
| Professional services                        | -0.040**<br>(-0.075, -0.005) | -0.032<br>(-0.072, 0.008) | -0.062*<br>(-0.124, 0.000)    |
| <b>Income (Poor to low income - REF)</b>     |                              |                           |                               |
| Middle to high income                        | 0.016**<br>(0.003, 0.029)    | 0.011<br>(-0.005, 0.027)  | 0.024**<br>(0.001, 0.048)     |
| <b>Receipt of food stamp (No - REF)</b>      |                              |                           |                               |
| Yes  | 0.024*<br>(-0.004, 0.052)    | 0.041**<br>(0.009, 0.073) | 0.001<br>(-0.047, 0.048)      |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference



Appendix 8. Association between long working hours and onset of psychological distress: fixed effects estimate without covariates- Weighted.

| <b>Variables</b>                             | <b>Overall</b>            | <b>All Men</b>            | <b>All Women</b>          |
|--|---------------------------|---------------------------|---------------------------|
| <b>Work hours/week (30-35hrs/week - REF)</b> |                           |                           |                           |
| 36-40 h/week                                 | 0.062**<br>(0.011, 0.114) | 0.077*<br>(-0.008, 0.161) | 0.049<br>(-0.012, 0.111)  |
| 41-50 h/week                                 | 0.065**<br>(0.004, 0.126) | 0.071<br>(-0.016, 0.159)  | 0.071<br>(-0.026, 0.167)  |
| 51-60 h/week                                 | 0.051<br>(-0.016, 0.117)  | 0.088*<br>(-0.011, 0.187) | -0.036<br>(-0.134, 0.063) |
| ≥ 61 h/week                                  | 0.101**<br>(0.020, 0.183) | 0.099*<br>(-0.006, 0.204) | 0.156*<br>(-0.021, 0.333) |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

## APPENDIX CHAPTER 3:

Appendix 1: Medications with sedative properties (Comparisons eAnswers., 2021).

### Drug Facts And Comparisons

Acetaminophen and Codeine Oral

Acetaminophen, Caffeine, and Dihydrocodeine Oral

Alfentanil Injection

Alosetron Oral

ALPRAZolam Oral

Amitriptyline Oral

Ampicillin and Sulbactam Injection

ARIPiprazole Injection

ARIPiprazole Oral

Azelastine and Fluticasone Intranasal

Benzonatate Oral

Brexpiprazole Oral

Brivaracetam Injection

Brivaracetam Oral

Brompheniramine, Pseudoephedrine, and Dextromethorphan Oral

Buprenorphine Implant

Buprenorphine Injection

Buprenorphine Oral

Buprenorphine Transdermal

Butalbital, Acetaminophen, and Caffeine Oral

Cannabidiol Oral

Capecitabine Oral

Carbinoxamine Oral

Cenobamate Oral

Cetirizine Injection

Cetirizine Oral

Chlorpheniramine and Phenylephrine Oral

Chlorpheniramine, Phenylephrine, and Dextromethorphan Oral

Chlorpheniramine, Pseudoephedrine, and Codeine Oral

Chlorpheniramine, Pseudoephedrine, and Dextromethorphan Oral

CloBAZam Oral

Clonidine and Clorthalidone Oral

CloNIDine Injection

CloNIDine Oral

CloNIDine Transdermal

CloZAPine Oral

Codeine and Chlorpheniramine Oral

Codeine Oral

Cyproheptadine Oral

Deutetrabenazine Oral

Dexchlorpheniramine Oral

DiphenhydrAMINE Injection

DiphenhydrAMINE Oral

Diphenoxylate and Atropine Oral

Doxepin Oral

Eluxadoline Oral

Esketamine Intranasal

Fenfluramine Oral

FentaNYL Buccal

FentaNYL Injection

FentaNYL Intranasal

FentaNYL Sublingual  
FentaNYL Transdermal  
Flibanserin Oral  
Flucytosine Oral  
Flurazepam Oral  
Gabapentin Enacarbil Oral  
Guaifenesin and Codeine Oral  
GuanFACINE Oral  
Haloperidol Injection  
Haloperidol Oral  
Hydrocodone and Acetaminophen Oral  
Hydrocodone and Chlorpheniramine Oral  
Hydrocodone and Homatropine Oral  
Hydrocodone and Pseudoephedrine Oral  
HYDRQcodone Oral  
HYDROmorphine Injection  
HYDROmorphine Oral  
HYDROmorphine Rectal  
Isocarboxazid Oral  
LevETIRAcetam Injection  
LevETIRAcetam Oral  
Lithium Oral  
Lofexidine Oral  
Loratadine Oral  
LORazepam Injection  
LORazepam Oral  
Loxapine Inhalation  
Loxapine Oral  
Lumateperone Oral  
Mecamylamine Oral  
Meperidine Injection  
Meperidine Oral  
Methadone Injection  
Methadone Oral  
Methocarbamol Injection  
Methocarbamol Oral  
Methyldopa and Hydrochlorothiazide Oral  
Methyldopa Oral  
Methyldopate Injection  
Methylphenidate Oral  
Methylphenidate Transdermal  
MetyraPONE Oral  
MetyroSINE Oral  
Midazolam Injection  
Midazolam Intranasal  
Midazolam Oral  
Mitotane Oral  
Morphine and Naltrexone Oral  
Morphine Injection  
Morphine Oral  
Morphine Rectal  
Nabilone Oral  
Nadolol and Bendroflumethiazide Oral  
Nadolol Oral  
Nalbuphine Injection  
Naltrexone Injection  
Naltrexone Oral  
Olanzapine and Samidorphan Oral  
OLANZapine Injection  
OLANZapine Oral  
Oliceridine Injection  
Ondansetron Injection

Ondansetron Oral  
Oxaprozin Oral  
Oxycodone and Acetaminophen Oral  
Oxycodone and Aspirin Oral  
OxyMORphone Injection  
OxyMORphone Oral  
Paliperidone Injection  
Paliperidone Oral  
Papaverine Injection  
Pentazocine and Naloxone Oral  
Pentazocine Injection  
Perphenazine Oral  
PHENobarbital Oral  
Pimozide Oral  
Pregabalin Oral  
Promethazine and Codeine Oral  
Promethazine and Phenylephrine Oral  
Promethazine Injection  
Promethazine Oral  
Promethazine Rectal  
Promethazine, Phenylephrine, and Codeine Oral  
RisperiDONE Injection  
RisperiDONE Oral  
SUFentanil Injection  
SUFentanil Oral  
SUMatriptan Injection  
SUMatriptan Intranasal  
SUMatriptan Oral  
Suvorexant Oral  
Tapentadol Oral  
Tetrabenazine Oral  
TiZANidine Oral  
TraMADol Oral  
Triazolam Oral  
Valbenazine Oral  
Vigabatrin Oral  
Vilazodone Oral  
Viloxazine Oral

Appendix 2a. Fixed Effect Linear Model – Overall Sample.

| Fixed Effect Linear Model – Overall Sample  |                               |                              |                               |                           |
|---|-------------------------------|------------------------------|-------------------------------|---------------------------|
| Variables                                   | Office Based                  | Emergency Room               | Hospital Discharges           | Dental                    |
| <b>Work hours/week (30-35 h/week - REF)</b> |                               |                              |                               |                           |
| 36-40 h/week                                | 0<br>(-0.029, 0.028)          | -0.007<br>(-0.031, 0.017)    | 0.012<br>(-0.003, 0.027)      | 0.004<br>(-0.022, 0.031)  |
| 41-50 h/week                                | 0<br>(-0.036, 0.035)          | -0.021<br>(-0.050, 0.009)    | 0.006<br>(-0.014, 0.025)      | 0.01<br>(-0.024, 0.044)   |
| 51-55 h/week                                | -0.012<br>(-0.061, 0.036)     | -0.013<br>(-0.051, 0.024)    | 0.009<br>(-0.014, 0.032)      | -0.024<br>(-0.069, 0.021) |
| ≥ 61 h/week                                 | -0.001<br>(-0.069, 0.068)     | -0.014<br>(-0.067, 0.038)    | 0.003<br>(-0.029, 0.034)      | 0<br>(-0.060, 0.061)      |
| <b>Age (18-26years - REF)</b>               |                               |                              |                               |                           |
| 27-64 years                                 | -0.032*<br>(-0.067, 0.002)    | -0.025*<br>(-0.050, 0.001)   | 0.01<br>(-0.007, 0.026)       | -0.013<br>(-0.046, 0.020) |
| ≥ 65 years                                  | -0.061**<br>(-0.116, -0.006)  | -0.034<br>(-0.079, 0.011)    | 0.028*<br>(-0.003, 0.059)     | -0.037<br>(-0.096, 0.022) |
| <b>Marital Status (Married - REF)</b>       |                               |                              |                               |                           |
| Unmarried                                   | 0.022<br>(-0.010, 0.053)      | -0.005<br>(-0.031, 0.022)    | -0.014<br>(-0.031, 0.003)     | -0.005<br>(-0.036, 0.025) |
| <b>Family Size (≤ 2 Members REF)</b>        |                               |                              |                               |                           |
| > 2 Members                                 | -0.037***<br>(-0.055, -0.019) | -0.020**<br>(-0.035, -0.005) | -0.009<br>(-0.022, 0.003)     | -0.011<br>(-0.030, 0.007) |
| <b>Occupation (Non-professional - REF)</b>  |                               |                              |                               |                           |
| Professional services                       | -0.016<br>(-0.045, 0.012)     | 0.000<br>(-0.023, 0.023)     | -0.01<br>(-0.024, 0.005)      | 0.013<br>(-0.015, 0.041)  |
| <b>Income (Poor to low income - REF)</b>    |                               |                              |                               |                           |
| middle to high income                       | 0.000<br>(-0.014, 0.014)      | 0.002<br>(-0.009, 0.013)     | -0.015***<br>(-0.023, -0.008) | 0.013**<br>(0.000, 0.026) |
| <b>Receipt of food stamp (No - REF)</b>     |                               |                              |                               |                           |
| Yes   | -0.011<br>(-0.031, 0.010)     | 0.001<br>(-0.017, 0.019)     | 0.013**<br>(0.000, 0.025)     | -0.006<br>(-0.024, 0.013) |
| <b>Limitation (No - REF)</b>                |                               |                              |                               |                           |
| Yes   | 0.043***<br>(0.029, 0.057)    | 0.036***<br>(0.021, 0.052)   | 0.034***<br>(0.023, 0.045)    | 0.020**<br>(0.004, 0.036) |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 2b. Fixed Logit Marginal Analysis – Overall sample.

| Fixed Logit Marginal Analysis – Overall sample |                               |                              |                               |                           |
|--|-------------------------------|------------------------------|-------------------------------|---------------------------|
| Variables                                      | Office Based                  | Emergency Room               | Hospital Discharges           | Dental                    |
| <b>Work hours/week (30-35 h/week - REF)</b>    |                               |                              |                               |                           |
| 36-40 h/week                                   | 0.000<br>(-0.046, 0.046)      | -0.014<br>(-0.068, 0.041)    | 0.065<br>(-0.025, 0.155)      | 0.008<br>(-0.044, 0.060)  |
| 41-50 h/week                                   | 0.005<br>(-0.056, 0.065)      | -0.048<br>(-0.118, 0.022)    | 0.021<br>(-0.092, 0.133)      | 0.021<br>(-0.045, 0.086)  |
| 51-55 h/week                                   | -0.016<br>(-0.094, 0.061)     | -0.032<br>(-0.132, 0.069)    | 0.049<br>(-0.124, 0.223)      | -0.050<br>(-0.137, 0.038) |
| ≥ 61 h/week                                    | -0.002<br>(-0.101, 0.098)     | -0.029<br>(-0.156, 0.099)    | 0.002<br>(-0.215, 0.218)      | 0.003<br>(-0.115, 0.120)  |
| <b>Age (18-26years - REF)</b>                  |                               |                              |                               |                           |
| 27-64 years                                    | -0.052*<br>(-0.105, 0.002)    | -0.071*<br>(-0.144, 0.002)   | 0.068<br>(-0.048, 0.184)      | -0.023<br>(-0.081, 0.036) |
| ≥ 65 years                                     | -0.116**<br>(-0.226, -0.006)  | -0.101<br>(-0.235, 0.033)    | 0.178*<br>(-0.027, 0.383)     | -0.067<br>(-0.178, 0.043) |
| <b>Marital Status (Married - REF)</b>          |                               |                              |                               |                           |
| Unmarried                                      | 0.041<br>(-0.016, 0.097)      | -0.013<br>(-0.077, 0.051)    | -0.093*<br>(-0.197, 0.010)    | -0.009<br>(-0.067, 0.049) |
| <b>Family Size (≤ 2 Members REF)</b>           |                               |                              |                               |                           |
| > 2 Members                                    | -0.067***<br>(-0.100, -0.034) | -0.049**<br>(-0.086, -0.011) | -0.034<br>(-0.083, 0.015)     | -0.021<br>(-0.055, 0.012) |
| <b>Occupation (Non-professional - REF)</b>     |                               |                              |                               |                           |
| Professional services                          | -0.027<br>(-0.076, 0.021)     | -0.005<br>(-0.065, 0.054)    | -0.071<br>(-0.168, 0.025)     | 0.025<br>(-0.026, 0.076)  |
| <b>Income (Poor to low income - REF)</b>       |                               |                              |                               |                           |
| middle to high income                          | -0.000<br>(-0.024, 0.023)     | 0.004<br>(-0.025, 0.033)     | -0.096***<br>(-0.142, -0.049) | 0.027**<br>(0.001, 0.054) |
| <b>Receipt of food stamp (No - REF)</b>        |                               |                              |                               |                           |
| Yes  | -0.017<br>(-0.053, 0.018)     | -0.000<br>(-0.042, 0.041)    | 0.053*<br>(-0.009, 0.114)     | -0.012<br>(-0.053, 0.030) |
| <b>Limitation (No - REF)</b>                   |                               |                              |                               |                           |
| Yes  | 0.104***<br>(0.069, 0.140)    | 0.074***<br>(0.042, 0.107)   | 0.136***<br>(0.092, 0.181)    | 0.037**<br>(0.007, 0.067) |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 3. Fixed Effects Regressions for Outpatient Physicians.

| <b>Fixed Effects Regressions for Outpatient Physicians</b> |                            |                            |                            |
|--|----------------------------|----------------------------|----------------------------|
| <b>Variables</b>   | <b>Logit Model</b>         | <b>Marginal Effect</b>     | <b>Linear Model</b>        |
| <b>Work hours/week (30-35 h/week - REF)</b>                |                            |                            |                            |
| 36-40 h/week   | 1.173<br>(0.673, 2.044)    | 0.037<br>(-0.092, 0.167)   | 0.004<br>(-0.006, 0.015)   |
| 41-50 h/week   | 0.864<br>(0.393, 1.900)    | -0.034<br>(-0.214, 0.146)  | -0.002<br>(-0.014, 0.010)  |
| 51-55 h/week   | 0.893<br>(0.304, 2.618)    | -0.026<br>(-0.273, 0.221)  | -0.001<br>(-0.017, 0.015)  |
| ≥ 61 h/week  | 5.038*<br>(0.954, 26.595)  | 0.333**<br>(0.046, 0.621)  | 0.023**<br>(0.001, 0.044)  |
| <b>Age (18-26years - REF)</b>                              |                            |                            |                            |
| 27-64 years  | 1.486<br>(0.432, 5.115)    | 0.091<br>(-0.195, 0.378)   | 0.002<br>(-0.005, 0.009)   |
| ≥ 65 years   | 0.897<br>(0.165, 4.883)    | -0.024<br>(-0.402, 0.353)  | -0.003<br>(-0.020, 0.015)  |
| <b>Marital Status (Married - REF)</b>                      |                            |                            |                            |
| Unmarried  | 0.703<br>(0.256, 1.930)    | -0.081<br>(-0.310, 0.148)  | -0.001<br>(-0.009, 0.006)  |
| <b>Family Size (≤ 2 Members REF)</b>                       |                            |                            |                            |
| > 2 Members  | 0.947<br>(0.482, 1.858)    | -0.013<br>(-0.168, 0.142)  | 0.000<br>(-0.004, 0.004)   |
| <b>Occupation (Non-professional - REF)</b>                 |                            |                            |                            |
| Professional services                                      | 0.696<br>(0.352, 1.379)    | -0.083<br>(-0.240, 0.074)  | -0.005<br>(-0.014, 0.005)  |
| <b>Income (Poor to low income - REF)</b>                   |                            |                            |                            |
| middle to high income                                      | 0.643*<br>(0.393, 1.051)   | -0.102*<br>(-0.215, 0.011) | -0.003*<br>(-0.006, 0.000) |
| <b>Receipt of food stamp (No - REF)</b>                    |                            |                            |                            |
| Yes  | 1.353<br>(0.814, 2.249)    | 0.070<br>(-0.047, 0.187)   | 0.004<br>(-0.003, 0.010)   |
| <b>Limitation (No - REF)</b>                               |                            |                            |                            |
| Yes  | 2.534***<br>(1.564, 4.106) | 0.217***<br>(0.111, 0.324) | 0.009***<br>(0.004, 0.014) |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 4a. Fixed Effect Logit Model – Among Medication Users.

| <b>Fixed Effect Logit Model – Among Medication Users</b> |                            |                            |                            |                           |
|--|----------------------------|----------------------------|----------------------------|---------------------------|
| <b>Variables</b>   | <b>Office Based</b>        | <b>Emergency Room</b>      | <b>Hospital Discharges</b> | <b>Dental</b>             |
| <b>Work hours/week (30-35 h/week - REF)</b>              |                            |                            |                            |                           |
| 36-40 h/week   | 1.012<br>(0.681, 1.504)    | 0.875<br>(0.636, 1.206)    | 1.288<br>(0.820, 2.023)    | 1.134<br>(0.814, 1.579)   |
| 41-50 h/week   | 0.892<br>(0.524, 1.517)    | 0.814<br>(0.538, 1.234)    | 1.157<br>(0.654, 2.046)    | 1.255<br>(0.827, 1.903)   |
| 51-55 h/week   | 1.001<br>(0.465, 2.152)    | 0.925<br>(0.495, 1.728)    | 0.954<br>(0.365, 2.491)    | 1.703<br>(0.854, 3.399)   |
| ≥ 61 h/week  | 1.173<br>(0.465, 2.958)    | 0.755<br>(0.346, 1.647)    | 0.595<br>(0.169, 2.095)    | 1.393<br>(0.596, 3.254)   |
| <b>Age (18-26years - REF)</b>                            |                            |                            |                            |                           |
| 27-64 years  | 0.734<br>(0.449, 1.197)    | 0.540**<br>(0.337, 0.864)  | 1.186<br>(0.626, 2.248)    | 0.768<br>(0.497, 1.187)   |
| ≥ 65 years   | 0.444<br>(0.118, 1.674)    | 0.613<br>(0.271, 1.386)    | 2.616<br>(0.759, 9.011)    | 0.877<br>(0.393, 1.959)   |
| <b>Marital Status (Married - REF)</b>                    |                            |                            |                            |                           |
| Unmarried  | 1.355<br>(0.842, 2.178)    | 1.147<br>(0.785, 1.677)    | 0.819<br>(0.465, 1.442)    | 1.007<br>(0.681, 1.487)   |
| <b>Family Size (≤ 2 Members REF)</b>                     |                            |                            |                            |                           |
| > 2 Members  | 0.937<br>(0.695, 1.263)    | 0.894<br>(0.717, 1.115)    | 0.907<br>(0.696, 1.181)    | 1.044<br>(0.830, 1.314)   |
| <b>Occupation (Non-professional - REF)</b>               |                            |                            |                            |                           |
| Professional services                                    | 0.996<br>(0.630, 1.574)    | 0.989<br>(0.698, 1.402)    | 1.030<br>(0.616, 1.723)    | 0.870<br>(0.613, 1.236)   |
| <b>Income (Poor to low income - REF)</b>                 |                            |                            |                            |                           |
| middle to high income                                    | 1.002<br>(0.809, 1.240)    | 0.898<br>(0.753, 1.071)    | 0.716***<br>(0.558, 0.920) | 1.220**<br>(1.009, 1.476) |
| <b>Receipt of food stamp (No - REF)</b>                  |                            |                            |                            |                           |
| Yes  | 0.941<br>(0.701, 1.263)    | 1.028<br>(0.808, 1.309)    | 1.451**<br>(1.038, 2.029)  | 0.984<br>(0.750, 1.290)   |
| <b>Limitation (No - REF)</b>                             |                            |                            |                            |                           |
| Yes  | 2.057***<br>(1.589, 2.665) | 1.477***<br>(1.248, 1.749) | 1.819***<br>(1.453, 2.277) | 1.133<br>(0.951, 1.349)   |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference



## Appendix 4b: Fixed Effect Linear Model – Among Medication Users

| Fixed Effect Linear Model – Among Medication Users |                            |                               |                               |                           |
|--|----------------------------|-------------------------------|-------------------------------|---------------------------|
| Variables  | Office Based               | Emergency Room                | Hospital Discharges           | Dental                    |
| <b>Work hours/week (30-35 h/week - REF)</b>        |                            |                               |                               |                           |
| 36-40 h/week                                       | 0.007<br>(-0.041, 0.055)   | -0.027<br>(-0.086, 0.032)     | 0.021<br>(-0.022, 0.064)      | 0.026<br>(-0.032, 0.083)  |
| 41-50 h/week                                       | -0.005<br>(-0.061, 0.051)  | -0.038<br>(-0.109, 0.033)     | 0.014<br>(-0.039, 0.067)      | 0.044<br>(-0.027, 0.116)  |
| 51-55 h/week                                       | 0.005<br>(-0.083, 0.092)   | -0.018<br>(-0.124, 0.088)     | -0.001<br>(-0.072, 0.070)     | 0.077<br>(-0.021, 0.176)  |
| ≥ 61 h/week  | 0.028<br>(-0.095, 0.150)   | -0.060<br>(-0.204, 0.083)     | -0.037<br>(-0.131, 0.057)     | 0.046<br>(-0.086, 0.179)  |
| <b>Age (18-26years - REF)</b>                      |                            |                               |                               |                           |
| 27-64 years  | -0.038<br>(-0.107, 0.032)  | -0.098***<br>(-0.171, -0.024) | 0.015<br>(-0.039, 0.068)      | -0.048<br>(-0.125, 0.030) |
| ≥ 65 years   | -0.059<br>(-0.145, 0.026)  | -0.081<br>(-0.196, 0.035)     | 0.063<br>(-0.017, 0.143)      | -0.031<br>(-0.149, 0.086) |
| <b>Marital Status (Married - REF)</b>              |                            |                               |                               |                           |
| Unmarried  | 0.031<br>(-0.017, 0.079)   | 0.022<br>(-0.038, 0.082)      | -0.018<br>(-0.059, 0.024)     | 0.002<br>(-0.056, 0.060)  |
| <b>Family Size (≤ 2 Members REF)</b>               |                            |                               |                               |                           |
| > 2 Members  | -0.006<br>(-0.032, 0.021)  | -0.018<br>(-0.055, 0.018)     | -0.015<br>(-0.045, 0.015)     | 0.006<br>(-0.029, 0.040)  |
| <b>Occupation (Non-professional - REF)</b>         |                            |                               |                               |                           |
| Professional services                              | -0.003<br>(-0.047, 0.041)  | 0.000<br>(-0.058, 0.058)      | 0.005<br>(-0.034, 0.045)      | -0.025<br>(-0.083, 0.032) |
| <b>Income (Poor to low income - REF)</b>           |                            |                               |                               |                           |
| middle to high income                              | 0.001<br>(-0.025, 0.026)   | -0.017<br>(-0.047, 0.014)     | -0.031***<br>(-0.053, -0.009) | 0.030**<br>(0.001, 0.058) |
| <b>Receipt of food stamp (No - REF)</b>            |                            |                               |                               |                           |
| Yes  | -0.008<br>(-0.045, 0.030)  | 0.012<br>(-0.033, 0.058)      | 0.040**<br>(0.006, 0.073)     | -0.004<br>(-0.045, 0.036) |
| <b>Limitation (No - REF)</b>                       |                            |                               |                               |                           |
| Yes  | 0.052***<br>(0.034, 0.070) | 0.062***<br>(0.036, 0.089)    | 0.057***<br>(0.037, 0.078)    | 0.019<br>(-0.007, 0.045)  |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 4c. Fixed Marginal Analysis – Among Medication Users.

| <b>Fixed Marginal Analysis – Among Medication Users</b> |                            |                               |                              |                           |
|---|----------------------------|-------------------------------|------------------------------|---------------------------|
| <b>Variables</b>  | <b>Office Based</b>        | <b>Emergency Room</b>         | <b>Hospital Discharges</b>   | <b>Dental</b>             |
| <b>Work hours/week (30-35 h/week - REF)</b>             |                            |                               |                              |                           |
| 36-40 h/week  | 0.003<br>(-0.094, 0.099)   | -0.030<br>(-0.101, 0.041)     | 0.060<br>(-0.046, 0.167)     | 0.031<br>(-0.051, 0.114)  |
| 41-50 h/week  | -0.028<br>(-0.156, 0.100)  | -0.046<br>(-0.135, 0.044)     | 0.035<br>(-0.101, 0.171)     | 0.056<br>(-0.047, 0.160)  |
| 51-55 h/week  | 0.000<br>(-0.186, 0.186)   | -0.018<br>(-0.158, 0.123)     | -0.011<br>(-0.242, 0.219)    | 0.131<br>(-0.034, 0.296)  |
| ≥ 61 h/week   | 0.039<br>(-0.187, 0.265)   | -0.062<br>(-0.226, 0.103)     | -0.123<br>(-0.409, 0.164)    | 0.082<br>(-0.126, 0.291)  |
| <b>Age (18-26years - REF)</b>                           |                            |                               |                              |                           |
| 27-64 years   | -0.076<br>(-0.195, 0.044)  | -0.145***<br>(-0.248, -0.042) | 0.041<br>(-0.113, 0.195)     | -0.065<br>(-0.173, 0.043) |
| ≥ 65 years  | -0.194<br>(-0.489, 0.101)  | -0.117<br>(-0.302, 0.068)     | 0.219*<br>(-0.029, 0.466)    | -0.032<br>(-0.230, 0.166) |
| <b>Marital Status (Married - REF)</b>                   |                            |                               |                              |                           |
| Unmarried   | 0.074<br>(-0.043, 0.191)   | 0.030<br>(-0.056, 0.116)      | -0.048<br>(-0.183, 0.088)    | 0.002<br>(-0.095, 0.098)  |
| <b>Family Size (≤ 2 Members REF)</b>                    |                            |                               |                              |                           |
| > 2 Members   | -0.016<br>(-0.088, 0.057)  | -0.025<br>(-0.073, 0.023)     | -0.023<br>(-0.087, 0.040)    | 0.011<br>(-0.046, 0.068)  |
| <b>Occupation (Non-professional - REF)</b>              |                            |                               |                              |                           |
| Professional services                                   | -0.001<br>(-0.112, 0.110)  | -0.002<br>(-0.079, 0.074)     | 0.007<br>(-0.115, 0.129)     | -0.034<br>(-0.122, 0.053) |
| <b>Income (Poor to low income - REF)</b>                |                            |                               |                              |                           |
| middle to high income                                   | 0.000<br>(-0.052, 0.052)   | -0.024<br>(-0.063, 0.015)     | -0.079**<br>(-0.139, -0.018) | 0.049**<br>(0.002, 0.097) |
| <b>Receipt of food stamp (No - REF)</b>                 |                            |                               |                              |                           |
| Yes   | -0.015<br>(-0.086, 0.056)  | 0.006<br>(-0.048, 0.060)      | 0.087**<br>(0.011, 0.164)    | -0.004<br>(-0.071, 0.063) |
| <b>Limitation (No - REF)</b>                            |                            |                               |                              |                           |
| Yes   | 0.175***<br>(0.114, 0.236) | 0.089***<br>(0.046, 0.132)    | 0.142***<br>(0.088, 0.195)   | 0.031<br>(-0.012, 0.074)  |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 5. Fixed Effect Logit Model for Outpatient Physicians – Among Medication Users.

| <b>Fixed Effect Logit Model for Outpatient Physicians – Among Medication Users</b> |                            |                              |                              |
|--|----------------------------|------------------------------|------------------------------|
| <b>Variables</b>   | <b>Logit Model</b>         | <b>Marginal Effect</b>       | <b>Linear Model</b>          |
| <b>Work hours/week (30-35 h/week - REF)</b>  |                            |                              |                              |
| 36-40 h/week   | 1.284<br>(0.604, 2.727)    | 0.051<br>(-0.107, 0.209)     | 0.012<br>(-0.016, 0.040)     |
| 41-50 h/week   | 0.784<br>(0.269, 2.280)    | -0.048<br>(-0.257, 0.160)    | -0.008<br>(-0.039, 0.022)    |
| 51-55 h/week   | 0.769<br>(0.153, 3.876)    | -0.052<br>(-0.363, 0.260)    | -0.012<br>(-0.057, 0.033)    |
| ≥ 61 h/week  | 6.411<br>(0.651, 63.149)   | 0.366*<br>(-0.019, 0.751)    | 0.065*<br>(-0.002, 0.132)    |
| <b>Age (18-26years - REF)</b>  |                            |                              |                              |
| 27-64 years  | 1.592<br>(0.292, 8.699)    | 0.093<br>(-0.260, 0.446)     | 0.003<br>(-0.019, 0.026)     |
| ≥ 65 years   | 3.185<br>(0.168, 60.260)   | 0.239<br>(-0.376, 0.854)     | 0.012<br>(-0.022, 0.047)     |
| <b>Marital Status (Married - REF)</b>  |                            |                              |                              |
| Unmarried  | 0.227*<br>(0.040, 1.273)   | -0.308*<br>(-0.622, 0.006)   | -0.013<br>(-0.029, 0.004)    |
| <b>Family Size (≤ 2 Members REF)</b>   |                            |                              |                              |
| > 2 Members  | 0.963<br>(0.389, 2.382)    | -0.008<br>(-0.191, 0.176)    | -0.002<br>(-0.011, 0.008)    |
| <b>Occupation (Non-professional - REF)</b>   |                            |                              |                              |
| Professional services  | 0.738<br>(0.293, 1.859)    | -0.062<br>(-0.250, 0.125)    | -0.012<br>(-0.037, 0.012)    |
| <b>Income (Poor to low income - REF)</b>   |                            |                              |                              |
| middle to high income  | 0.522**<br>(0.272, 1.000)  | -0.132**<br>(-0.262, -0.003) | -0.010**<br>(-0.019, -0.001) |
| <b>Receipt of food stamp (No - REF)</b>  |                            |                              |                              |
| Yes  | 1.718<br>(0.886, 3.330)    | 0.111<br>(-0.027, 0.250)     | 0.015*<br>(-0.003, 0.034)    |
| <b>Limitation (No - REF)</b>   |                            |                              |                              |
| Yes  | 2.607***<br>(1.358, 5.007) | 0.199***<br>(0.051, 0.348)   | 0.013***<br>(0.005, 0.021)   |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

## Weighted Results

Appendix 6a. Fixed Effect Logit Model – Overall Sample (Weighted)

| Fixed Effect Logit Model – Overall Sample   |                            |                            |                            |                            |                           |
|---|----------------------------|----------------------------|----------------------------|----------------------------|---------------------------|
| Variables                                   | Outpatient                 | Office Based               | Emergency Room             | Hospital Discharges        | Dental                    |
| <b>Work hours/week (30-35 h/week - REF)</b> |                            |                            |                            |                            |                           |
| 36-40 h/week                                | 1.379*<br>(0.971, 1.958)   | 1.095<br>(0.852, 1.407)    | 0.941<br>(0.706, 1.253)    | 1.369<br>(0.872, 2.150)    | 1.006<br>(0.777, 1.303)   |
| 41-45 h/week                                | 1.709**<br>(1.022, 2.857)  | 0.912<br>(0.623, 1.336)    | 0.808<br>(0.512, 1.275)    | 1.465<br>(0.730, 2.939)    | 1.029<br>(0.711, 1.488)   |
| 46-50 h/week                                | 1.342<br>(0.821, 2.195)    | 1.290<br>(0.883, 1.886)    | 0.748<br>(0.484, 1.158)    | 1.538<br>(0.767, 3.086)    | 0.989<br>(0.677, 1.444)   |
| 51-55 h/week                                | 2.062*<br>(0.945, 4.499)   | 0.796<br>(0.458, 1.385)    | 0.963<br>(0.474, 1.955)    | 2.304<br>(0.605, 8.770)    | 0.656<br>(0.366, 1.177)   |
| 56-60 h/week                                | 1.247<br>(0.625, 2.488)    | 1.132<br>(0.692, 1.853)    | 0.750<br>(0.412, 1.363)    | 1.296<br>(0.379, 4.431)    | 0.786<br>(0.478, 1.293)   |
| ≥ 61 h/week                                 | 1.632<br>(0.734, 3.628)    | 0.868<br>(0.521, 1.443)    | 1.039<br>(0.536, 2.014)    | 1.140<br>(0.431, 3.013)    | 0.917<br>(0.523, 1.607)   |
| <b>Age (18-26years - REF)</b>               |                            |                            |                            |                            |                           |
| 27-64 years                                 | 1.389<br>(0.896, 2.154)    | 0.786*<br>(0.592, 1.044)   | 0.736<br>(0.504, 1.075)    | 1.161<br>(0.619, 2.177)    | 0.860<br>(0.640, 1.156)   |
| ≥ 65 years                                  | 1.900*<br>(0.966, 3.737)   | 0.716<br>(0.377, 1.361)    | 0.573<br>(0.273, 1.202)    | 1.164<br>(0.420, 3.227)    | 0.962<br>(0.539, 1.717)   |
| <b>Marital Status (Married - REF)</b>       |                            |                            |                            |                            |                           |
| Unmarried                                   | 0.878<br>(0.581, 1.326)    | 1.372**<br>(1.021, 1.843)  | 1.008<br>(0.715, 1.422)    | 0.727<br>(0.425, 1.242)    | 0.888<br>(0.664, 1.187)   |
| <b>Family Size (≤ 2 Members REF)</b>        |                            |                            |                            |                            |                           |
| > 2 Members                                 | 0.750***<br>(0.603, 0.931) | 0.759***<br>(0.636, 0.906) | 0.878<br>(0.718, 1.074)    | 0.909<br>(0.699, 1.182)    | 0.919<br>(0.775, 1.091)   |
| <b>Occupation (Non-professional - REF)</b>  |                            |                            |                            |                            |                           |
| Professional services                       | 0.798<br>(0.558, 1.142)    | 0.881<br>(0.685, 1.133)    | 0.968<br>(0.710, 1.320)    | 0.935<br>(0.559, 1.565)    | 1.310**<br>(1.021, 1.679) |
| <b>Income (Poor to low income - REF)</b>    |                            |                            |                            |                            |                           |
| middle to high income                       | 0.846*<br>(0.704, 1.017)   | 1.074<br>(0.946, 1.219)    | 1.077<br>(0.925, 1.255)    | 0.728**<br>(0.566, 0.937)  | 1.101<br>(0.961, 1.261)   |
| <b>Receipt of food stamp (No - REF)</b>     |                            |                            |                            |                            |                           |
| Yes   | 1.259<br>(0.956, 1.659)    | 1.040<br>(0.856, 1.265)    | 0.986<br>(0.787, 1.235)    | 1.428*<br>(0.981, 2.078)   | 0.882<br>(0.708, 1.099)   |
| <b>Limitation (No - REF)</b>                |                            |                            |                            |                            |                           |
| Yes   | 1.426***<br>(1.220, 1.668) | 1.482***<br>(1.236, 1.777) | 1.454***<br>(1.234, 1.714) | 1.898***<br>(1.509, 2.387) | 1.149*<br>(0.988, 1.336)  |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 6b. Fixed Marginal Analysis – Overall Sample (Weighted).

| <b>Fixed Marginal Analysis – Overall Sample</b> |                   |                     |                       |                            |                 |
|---|-------------------|---------------------|-----------------------|----------------------------|-----------------|
| <b>Variables</b>                                | <b>Outpatient</b> | <b>Office Based</b> | <b>Emergency Room</b> | <b>Hospital Discharges</b> | <b>Dental</b>   |
| <b>Work hours/week (30-35 h/week - REF)</b>     |                   |                     |                       |                            |                 |
| 36-40 h/week                                    | 0.078*            | 0.022               | -0.015                | 0.076                      | 0.002           |
|   | (-0.006, 0.163)   | (-0.039, 0.083)     | (-0.084, 0.055)       | (-0.033, 0.186)            | (-0.063, 0.066) |
| 41-45 h/week                                    | 0.129**           | -0.022              | -0.051                | 0.093                      | 0.007           |
|   | (0.010, 0.249)    | (-0.113, 0.069)     | (-0.159, 0.056)       | (-0.075, 0.260)            | (-0.085, 0.099) |
| 46-50 h/week                                    | 0.072             | 0.062               | -0.069                | 0.104                      | -0.003          |
|   | (-0.047, 0.191)   | (-0.031, 0.155)     | (-0.171, 0.032)       | (-0.062, 0.271)            | (-0.097, 0.091) |
| 51-55 h/week                                    | 0.172*            | -0.054              | -0.009                | 0.198                      | -0.103          |
|   | (-0.001, 0.344)   | (-0.184, 0.075)     | (-0.182, 0.163)       | (-0.096, 0.491)            | (-0.243, 0.036) |
| 56-60 h/week                                    | 0.054             | 0.030               | -0.069                | 0.063                      | -0.060          |
|   | (-0.114, 0.222)   | (-0.090, 0.151)     | (-0.208, 0.070)       | (-0.236, 0.362)            | (-0.182, 0.063) |
| ≥ 61 h/week                                     | 0.119             | -0.034              | 0.009                 | 0.032                      | -0.022          |
|   | (-0.069, 0.306)   | (-0.155, 0.087)     | (-0.153, 0.172)       | (-0.205, 0.268)            | (-0.161, 0.118) |
| <b>Age (18-26years - REF)</b>                   |                   |                     |                       |                            |                 |
| 27-64 years                                     | 0.080             | -0.059*             | -0.075                | 0.036                      | -0.037          |
|   | (-0.025, 0.186)   | (-0.128, 0.010)     | (-0.166, 0.016)       | (-0.116, 0.188)            | (-0.111, 0.036) |
| ≥ 65 years                                      | 0.154**           | -0.081              | -0.134                | 0.037                      | -0.010          |
|   | (0.001, 0.307)    | (-0.236, 0.073)     | (-0.302, 0.035)       | (-0.209, 0.283)            | (-0.153, 0.134) |
| <b>Marital Status (Married - REF)</b>           |                   |                     |                       |                            |                 |
| Unmarried                                       | -0.031            | 0.078**             | 0.002                 | -0.077                     | -0.029          |
|   | (-0.131, 0.069)   | (0.005, 0.151)      | (-0.081, 0.085)       | (-0.207, 0.053)            | (-0.101, 0.042) |
| <b>Family Size (≤ 2 Members REF)</b>            |                   |                     |                       |                            |                 |
| > 2 Members                                     | -0.069**          | -0.068***           | -0.031                | -0.023                     | -0.021          |
|   | (-0.123, -0.016)  | (-0.110, -0.025)    | (-0.079, 0.017)       | (-0.086, 0.041)            | (-0.063, 0.022) |
| <b>Occupation (Non-professional - REF)</b>      |                   |                     |                       |                            |                 |
| Professional services                           | -0.054            | -0.031              | -0.008                | -0.016                     | 0.067**         |
|   | (-0.141, 0.033)   | (-0.092, 0.030)     | (-0.082, 0.067)       | (-0.141, 0.108)            | (0.005, 0.129)  |
| <b>Income (Poor to low income - REF)</b>        |                   |                     |                       |                            |                 |
| middle to high income                           | -0.040*           | 0.017               | 0.018                 | -0.076**                   | 0.024           |
|   | (-0.084, 0.005)   | (-0.014, 0.048)     | (-0.019, 0.055)       | (-0.137, -0.014)           | (-0.010, 0.057) |
| <b>Receipt of food stamp (No - REF)</b>         |                   |                     |                       |                            |                 |
| Yes   | 0.054*            | 0.010               | -0.003                | 0.085*                     | -0.031          |
|   | (-0.010, 0.118)   | (-0.038, 0.057)     | (-0.057, 0.051)       | (-0.003, 0.172)            | (-0.086, 0.023) |
| <b>Limitation (No - REF)</b>                    |                   |                     |                       |                            |                 |
| Yes   | 0.084***          | 0.096***            | 0.092***              | 0.153***                   | 0.034*          |

(0.047, 0.121) (0.052, 0.140) (0.050, 0.133) (0.098, 0.208) (-0.003, 0.072)

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 7a. Fixed Effect Logit Model – Among Medication Users (Weighted)

| <b>Fixed Effect Logit Model – Among Medication Users</b> |                   |                     |                       |                            |                |
|--|-------------------|---------------------|-----------------------|----------------------------|----------------|
| <b>Variables</b>   | <b>Outpatient</b> | <b>Office Based</b> | <b>Emergency Room</b> | <b>Hospital Discharges</b> | <b>Dental</b>  |
| <b>Work hours/week (30-35 h/week - REF)</b>              |                   |                     |                       |                            |                |
| 36-40 h/week   | 2.547*            | 1.088               | 0.902                 | 1.308                      | 1.113          |
|  | (0.837, 7.749)    | (0.669, 1.768)      | (0.604, 1.348)        | (0.752, 2.275)             | (0.742, 1.669) |
| 41-45 h/week   | 2.847             | 1.046               | 0.703                 | 1.568                      | 1.543          |
|  | (0.410, 19.777)   | (0.462, 2.365)      | (0.390, 1.268)        | (0.677, 3.634)             | (0.854, 2.785) |
| 46-50 h/week   | 1.709             | 1.221               | 0.622                 | 1.141                      | 1.170          |
|  | (0.209, 13.995)   | (0.549, 2.716)      | (0.336, 1.154)        | (0.509, 2.562)             | (0.630, 2.170) |
| 51-55 h/week   | 28.917***         | 0.782               | 0.855                 | 0.519                      | 0.842          |
|  | (3.851, 217.141)  | (0.220, 2.778)      | (0.302, 2.420)        | (0.097, 2.789)             | (0.288, 2.456) |
| 56-60 h/week   | 20.830**          | 1.499               | 0.876                 | 0.950                      | 2.294          |
|  | (1.139, 381.100)  | (0.482, 4.667)      | (0.381, 2.011)        | (0.283, 3.191)             | (0.815, 6.457) |
| ≥ 61 h/week  | 49.085***         | 0.800               | 1.030                 | 0.677                      | 1.340          |
|  | (3.623, 664.958)  | (0.273, 2.344)      | (0.398, 2.663)        | (0.185, 2.472)             | (0.458, 3.923) |
| <b>Age (18-26years - REF)</b>                            |                   |                     |                       |                            |                |
| 27-64 years  | 0.219             | 0.750               | 0.606*                | 1.392                      | 0.707          |
|  | (0.004, 10.771)   | (0.425, 1.323)      | (0.346, 1.061)        | (0.623, 3.110)             | (0.405, 1.235) |
| ≥ 65 years   | 0.534             | 0.630               | 0.547                 | 2.446                      | 1.032          |
|  | (0.007, 38.470)   | (0.132, 3.003)      | (0.210, 1.428)        | (0.605, 9.886)             | (0.379, 2.812) |
| <b>Marital Status (Married - REF)</b>                    |                   |                     |                       |                            |                |
| Unmarried  | 0.866             | 1.453               | 1.356                 | 0.905                      | 0.833          |
|  | (0.206, 3.640)    | (0.842, 2.508)      | (0.843, 2.184)        | (0.453, 1.811)             | (0.519, 1.334) |
| <b>Family Size (≤ 2 Members - REF)</b>                   |                   |                     |                       |                            |                |
| > 2 Members  | 1.018             | 0.947               | 1.053                 | 1.051                      | 0.983          |
|  | (0.434, 2.389)    | (0.641, 1.400)      | (0.799, 1.387)        | (0.742, 1.488)             | (0.736, 1.312) |
| <b>Occupation (Non-professional - REF)</b>               |                   |                     |                       |                            |                |
| Professional services                                    | 13.119***         | 0.882               | 0.917                 | 1.190                      | 0.926          |
|  | (1.888, 91.140)   | (0.516, 1.507)      | (0.603, 1.396)        | (0.614, 2.306)             | (0.610, 1.404) |
| <b>Income (Poor to low income - REF)</b>                 |                   |                     |                       |                            |                |
| middle to high income                                    | 0.608             | 0.967               | 0.958                 | 0.789                      | 1.136          |
|  | (0.260, 1.423)    | (0.737, 1.267)      | (0.769, 1.194)        | (0.568, 1.096)             | (0.897, 1.439) |
| <b>Receipt of food stamp (No - REF)</b>                  |                   |                     |                       |                            |                |
| Yes  | 0.347**           | 1.025               | 1.105                 | 1.471                      | 0.943          |
|  | (0.129, 0.936)    | (0.701, 1.499)      | (0.814, 1.499)        | (0.918, 2.357)             | (0.675, 1.319) |
| <b>Limitation (No - REF)</b>                             |                   |                     |                       |                            |                |
| Yes  | 1.486             | 2.117***            | 1.580***              | 2.103***                   | 1.096          |
|  | (0.893, 2.472)    | (1.517, 2.953)      | (1.278, 1.953)        | (1.599, 2.766)             | (0.883, 1.361) |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 7b. Fixed Marginal Analysis – Among Medication Users (Weighted)

| Fixed Marginal Analysis – Among Medication Users |                            |                            |                            |                            |                           |
|--|----------------------------|----------------------------|----------------------------|----------------------------|---------------------------|
| Variables  | Outpatient                 | Office Based               | Emergency Room             | Hospital Discharges        | Dental                    |
| <b>Work hours/week (30-35 h/week - REF)</b>      |                            |                            |                            |                            |                           |
| 36-40 h/week                                     | 0.154*<br>(-0.027, 0.335)  | 0.020<br>(-0.098, 0.138)   | -0.025<br>(-0.119, 0.070)  | 0.060<br>(-0.059, 0.178)   | 0.026<br>(-0.074, 0.126)  |
| 41-45 h/week                                     | 0.171<br>(-0.133, 0.476)   | 0.011<br>(-0.187, 0.209)   | -0.082<br>(-0.213, 0.049)  | 0.098<br>(-0.073, 0.268)   | 0.107<br>(-0.039, 0.254)  |
| 46-50 h/week                                     | 0.089<br>(-0.262, 0.441)   | 0.048<br>(-0.146, 0.243)   | -0.109<br>(-0.242, 0.025)  | 0.030<br>(-0.150, 0.210)   | 0.038<br>(-0.114, 0.191)  |
| 51-55 h/week                                     | 0.457***<br>(0.124, 0.790) | -0.059<br>(-0.356, 0.238)  | -0.037<br>(-0.281, 0.206)  | -0.155<br>(-0.553, 0.243)  | -0.041<br>(-0.293, 0.210) |
| 56-60 h/week                                     | 0.430**<br>(0.082, 0.778)  | 0.098<br>(-0.174, 0.370)   | -0.032<br>(-0.227, 0.164)  | -0.012<br>(-0.292, 0.269)  | 0.203*<br>(-0.038, 0.443) |
| ≥ 61 h/week                                      | 0.489**<br>(0.091, 0.887)  | -0.053<br>(-0.306, 0.200)  | 0.007<br>(-0.222, 0.236)   | -0.092<br>(-0.402, 0.218)  | 0.072<br>(-0.195, 0.339)  |
| <b>Age (18-26years - REF)</b>                    |                            |                            |                            |                            |                           |
| 27-64 years                                      | -0.209<br>(-0.770, 0.352)  | -0.070<br>(-0.207, 0.068)  | -0.121*<br>(-0.251, 0.010) | 0.075<br>(-0.102, 0.253)   | -0.085<br>(-0.222, 0.051) |
| ≥ 65 years                                       | -0.078<br>(-0.641, 0.485)  | -0.112<br>(-0.485, 0.261)  | -0.144<br>(-0.362, 0.074)  | 0.189<br>(-0.064, 0.441)   | 0.008<br>(-0.238, 0.253)  |
| <b>Marital Status (Married - REF)</b>            |                            |                            |                            |                            |                           |
| Unmarried  | -0.022<br>(-0.245, 0.200)  | 0.091<br>(-0.042, 0.224)   | 0.072<br>(-0.044, 0.187)   | -0.022<br>(-0.176, 0.132)  | -0.045<br>(-0.159, 0.070) |
| <b>Family Size (≤ 2 Members - REF)</b>           |                            |                            |                            |                            |                           |
| > 2 Members                                      | 0.003<br>(-0.128, 0.133)   | -0.013<br>(-0.108, 0.081)  | 0.012<br>(-0.052, 0.077)   | 0.011<br>(-0.065, 0.086)   | -0.004<br>(-0.075, 0.067) |
| <b>Occupation (Non-professional - REF)</b>       |                            |                            |                            |                            |                           |
| Professional services                            | 0.491**<br>(0.090, 0.893)  | -0.030<br>(-0.160, 0.099)  | -0.020<br>(-0.117, 0.077)  | 0.038<br>(-0.103, 0.179)   | -0.019<br>(-0.121, 0.083) |
| <b>Income (Poor to low income - REF)</b>         |                            |                            |                            |                            |                           |
| middle to high income                            | -0.075<br>(-0.205, 0.055)  | -0.008<br>(-0.074, 0.057)  | -0.010<br>(-0.061, 0.041)  | -0.051<br>(-0.124, 0.023)  | 0.031<br>(-0.027, 0.089)  |
| <b>Receipt of food stamp (No - REF)</b>          |                            |                            |                            |                            |                           |
| Yes  | -0.171*<br>(-0.345, 0.004) | 0.006<br>(-0.086, 0.098)   | 0.023<br>(-0.049, 0.096)   | 0.080<br>(-0.016, 0.176)   | -0.014<br>(-0.096, 0.068) |
| <b>Limitation (No - REF)</b>                     |                            |                            |                            |                            |                           |
| Yes  | 0.060<br>(-0.021, 0.142)   | 0.180***<br>(0.103, 0.258) | 0.109***<br>(0.055, 0.163) | 0.156***<br>(0.080, 0.231) | 0.022<br>(-0.031, 0.076)  |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference

Appendix 8. Fixed Effect Logit Model for Outpatient Physicians (Weighted)

| <b>Fixed Effect Logit Model for Outpatient Physicians</b> |                               |                              |                               |                              |
|---|-------------------------------|------------------------------|-------------------------------|------------------------------|
| <b>Variables</b>  | <b>Overall Sample</b>         |                              | <b>Among Medication Users</b> |                              |
|   | <b>Logit Model</b>            | <b>Marginal Effect</b>       | <b>Logit Model</b>            | <b>Marginal Effect</b>       |
| <b>Work hours/week (30-35 h/week - REF)</b>               |                               |                              |                               |                              |
| 36-40 h/week  | 1.047<br>(0.521, 2.102)       | 0.009<br>(-0.133, 0.151)     | 1.169<br>(0.511, 2.672)       | 0.030<br>(-0.129, 0.188)     |
| 41-45 h/week  | 0.859<br>(0.268, 2.759)       | -0.030<br>(-0.256, 0.196)    | 0.520<br>(0.108, 2.502)       | -0.124<br>(-0.413, 0.166)    |
| 46-50 h/week  | 0.789<br>(0.276, 2.254)       | -0.046<br>(-0.242, 0.150)    | 0.378<br>(0.063, 2.265)       | -0.180<br>(-0.501, 0.141)    |
| 51-55 h/week  | 1.571<br>(0.288, 8.577)       | 0.096<br>(-0.275, 0.467)     | 1.959<br>(0.216, 17.779)      | 0.126<br>(-0.273, 0.524)     |
| 56-60 h/week  | 0.521<br>(0.143, 1.895)       | -0.118<br>(-0.335, 0.099)    | 0.290<br>(0.049, 1.697)       | -0.224<br>(-0.513, 0.065)    |
| ≥ 61 h/week   | 16.048***<br>(2.490, 103.423) | 0.525***<br>(0.254, 0.796)   | 13.705*<br>(0.956, 196.438)   | 0.390**<br>(0.011, 0.770)    |
| <b>Age (18-26years - REF)</b>                             |                               |                              |                               |                              |
| 27-64 years   | 1.344<br>(0.237, 7.637)       | 0.058<br>(-0.291, 0.406)     | 4.352<br>(0.553, 34.233)      | 0.267<br>(-0.141, 0.675)     |
| ≥ 65 years  | 1.039<br>(0.111, 9.753)       | 0.007<br>(-0.415, 0.430)     | 6.007<br>(0.220, 164.409)     | 0.329<br>(-0.301, 0.959)     |
| <b>Marital Status (Married - REF)</b>                     |                               |                              |                               |                              |
| Unmarried   | 0.426<br>(0.136, 1.332)       | -0.168<br>(-0.375, 0.039)    | 0.182**<br>(0.038, 0.865)     | -0.330**<br>(-0.614, -0.047) |
| <b>Family Size (≤ 2 Members - REF)</b>                    |                               |                              |                               |                              |
| > 2 Members   | 1.390<br>(0.585, 3.299)       | 0.066<br>(-0.116, 0.249)     | 1.296<br>(0.396, 4.240)       | 0.048<br>(-0.170, 0.265)     |
| <b>Occupation (Non-professional - REF)</b>                |                               |                              |                               |                              |
| Professional services                                     | 0.471*<br>(0.194, 1.145)      | -0.155<br>(-0.342, 0.032)    | 0.536<br>(0.135, 2.128)       | -0.115<br>(-0.377, 0.147)    |
| <b>Income (Poor to low income - REF)</b>                  |                               |                              |                               |                              |
| middle to high income                                     | 0.527**<br>(0.295, 0.940)     | -0.130**<br>(-0.250, -0.010) | 0.406**<br>(0.188, 0.876)     | -0.161**<br>(-0.297, -0.025) |
| <b>Receipt of food stamp (No - REF)</b>                   |                               |                              |                               |                              |
| Yes   | 1.353<br>(0.679, 2.697)       | 0.062<br>(-0.081, 0.205)     | 1.662<br>(0.635, 4.347)       | 0.093<br>(-0.080, 0.266)     |
| <b>Limitation (No - REF)</b>                              |                               |                              |                               |                              |
| Yes   | 2.252***<br>(1.260, 4.023)    | 0.168**<br>(0.028, 0.309)    | 2.707**<br>(1.235, 5.930)     | 0.184**<br>(0.039, 0.330)    |

Note: Robust confidence interval in parentheses; \*\*\* p<0.01, \*\* p<0.05, \* p<0.1; REF=reference



## CURRICULUM VITAE

Emmanuel Ezekekwa, BPharm, MHA, PhD(c)

euezek01@louisville.edu | 202-817-9737

### EDUCATION

---

- 2023 (May) Ph.D., Public Health Sciences, Health Management & Systems Sciences.  
Dissertation: “The Association of Long Working Hours and the Use of Prescription Sedatives among U.S. Workers”
- 2017 Master of Health Administration  
**Western Kentucky University**
- 2010 Bachelor of Pharmacy (Hons)  
**University of Nigeria**

### RESEARCH AND TEACHING EXPERIENCE

---

- 2018 – 2022 Graduate Research Assistant  
**Center for Health Organization Transformation (CHOT),  
School of Public Health & Information Sciences, University of  
Louisville, Louisville, KY**
- 2016 – 2017 Graduate Research Assistant  
**Department of Public Health, Western Kentucky University,  
Bowling Green Kentucky**
- Summer, 2017 Administrative Intern  
**Commonwealth Regional Specialty Hospital, Bowling Green  
Kentucky**

## SELECTED PUBLICATIONS

---

- Ezekekwa**, E., Salunkhe, S. S., Jennings, J. A. C., & Kelly Pryor, B. N. (2022). Community-Based and System-Level Interventions for Improving Food Security and Nutritious Food Consumption: A Systematic Review. *Journal of Hunger & Environmental Nutrition*, 17(2), 149-169.
- Thornewill, J., Antimisiaris, D., **Ezekekwa**, E., & Esterhay, R. (2022). Transformational strategies for optimizing use of medications and related therapies through us pharmacists and pharmacies: Findings from a national study. *Journal of the American Pharmacists Association*, 62(2), 450-460.
- Karimi, S.M., Salunkhe, S.S., White, K.B., Little, B.B., McKinney, W.P., Mitra, R., Chen, Y., Adkins, E.R., Barclay, J.A., **Ezekekwa**, E. and He, C.X. (2021). Prevalence of unmasked and improperly masked behavior in indoor public areas during the COVID-19 pandemic: Analysis of a stratified random sample from Louisville, Kentucky. *Plos one*, 16(7), e0248324.
- Karimi, S.M., Salunkhe, S.S., White, K.B., Little, B.M., McKinney, W.P.M., Mitra, R.M., Chen, Y.M., Adkins, E.R., Barclay, J.A., **Ezekekwa**, E.M. and He, C.X. (2021). Stratified Random Sampling Methodology for Observing Community Mask Use within Indoor Settings: Results from Louisville, Kentucky during the COVID-19 Pandemic. *BioRxiv*.
- Karimi, S. M., Salunkhe, S. S., White, K. B., Alzahrani, S. A., Little, B., McKinney, W. P., ... **Ezekekwa**, E. & Moyer, S. (2021). Facial Mask Use and COVID-19 Protection Measures in Jefferson County, Kentucky: Results from an Observational Survey, November 5– 11, 2020. *The University of Louisville Journal of Respiratory Infections*, 5(1), 7.
- Thornewill, J., Antimisiaris, D., **Ezekekwa**, E., & Esterhay, R. (2021) The RAPID Alliance Report.
- Kabir, U. Y., Askew, A., Jiang, Y., Bhuyan, S. S., **Ezekekwa**, E., & Dobalian, A. (2020). Moderate psychological distress as a barrier to breast cancer screening among women. *Journal of Hospital Administration*, 9(4).
- Kabir, U. Y., **Ezekekwa**, E., Bhuyan, S. S., Mahmood, A., & Dobalian, A. (2020). Trends and best practices in health care cybersecurity insurance policy. *Journal of Healthcare Risk Management*, 40(2), 10-14.
- Patel,N., **Ezekekwa**, E. U. (2019). Ketamine-Opioid report. *CHI St. Joseph Report* Lexington, KY, November 2019.

Mkanta, W. N., Reece, M. C., Alamri, A. D., **Ezekekwa**, E. U., Potluri, A., & Chumbler, N. R. (2018). A 3-State Analysis of Black–White Disparities in Diabetes Hospitalizations Among Medicaid Beneficiaries. *Health Services Research and Managerial Epidemiology*, 5, 2333392818783513.

Mkanta, W. N., Eustace, R. W., Reece, M. C., Alamri, A. D., Davis, T., **Ezekekwa**, E. U., & Potluri, A. (2018). From images to voices: A photo analysis of medical and social support needs of people living with HIV/AIDS in Tanzania. *Journal of Global Health Reports*, 2.

Mkanta, W. N., Chumbler, N. R., Yang, K., Saigal, R., Abdollahi, M., Mejia de Grubb, M. C., & **Ezekekwa**, E. U. (2017). An Examination of the Likelihood of Home Discharge After General Hospitalizations among Medicaid Recipients. *INQUIRY: The Journal of Health Care Organization, Provision, and Financing*, 54, 0046958017711783.

#### WORKING PAPERS

**Ezekekwa**, E., Kabir, U., & Thornewill, J. Hospital ownership and responses to readmissions: A resource dependence theory framework.

William N. Mkanta, **Ezekekwa**, Emmanuel, Michelle C. Reece, Neale R. Chumbler. Cross Sectional Analysis of The Risk Factor for Repeated Hospitalizations Among Female Medicaid Beneficiaries.

Kabir, U., **Ezekekwa**, E., Meredith R., Dobalian, A. Adherence Level of Recommended Preventative Services for Individuals with Psychological Distress. Preventing Chronic Diseases.

Mkanta, W., Reece M., Alamri A., Pope D., **Ezekekwa E.** Cross-Sectional Analysis of the Risk Factors for Multiple Hospitalizations among Female Medicaid Beneficiaries. *Journal of Healthcare Management*.

#### SELECTED PRESENTATIONS

**Ezekekwa**, E., Thornewill, J., Antimisiaris, D., & Esterhay, R. (2022). Transformational strategies for optimizing use of medications and related therapies through us pharmacists and pharmacies: Findings from a national study.

**Ezekekwa**, E., Brandon Mitchell and Judah Thornewill. Robust Public Health Preparedness for Preventing the Next Pandemic. Presentation at RSPP Workshop on Pandemics, Location and Mobility. Ponta Delgada, Azores, Portugal (November 1 – 2, 2021).

- Ezekekwa, E.**, Seyed, K and Johnson, C. Evaluating the Use of Sedatives in the US. Working Population. Poster presentation at AcademyHealth Annual Research Meeting, June 14-17, 2021.
- Ezekekwa, E.**, Seyed, K and Johnson, C. Examining the Use of Sedative Drugs in the US. Working Population. Poster presentation at Virtual ISPOR 2021 (May 17-20, 2021). International Society for Pharmacoeconomics and Outcomes.
- Ezekekwa, E.**, Salunkhe, S., and Jennings, J.C. “Systematic Review of Effective Community-Based and System Level Interventions for Improving Food Security.” Poster presentation accepted at AcademyHealth Annual Research Meeting, Boston, MA, June 2020. (Conference held virtually in response to COVID-19).
- Ezekekwa, E.**, Salunkhe, S., and Jennings, J.C. “Systematic Review of Effective Community-Based and System Level Interventions for Improving Food Security.” Poster presentation at Kentucky Public Health Association Annual Conference, April 2020. (Conference held virtually in response to COVID-19).
- Ezekekwa, E.** (2020, October). Hospital ownership and responses to readmissions: A resource dependence theory framework. Oral presentation at *APHA's 2020 VIRTUAL Annual Meeting and Expo (Oct. 24-28)*. American Public Health Association.
- Jennings, J.C **Ezekekwa, E.U.**, and Ali, M. Advancements in Social Determinants of Health CHOT IAB Meeting. Malvern, PA. April 2019
- William N. Mkanta, Michelle C. Reece, **Ezekekwa Emmanuel U.**, Niyati thaku, Abdulgafir Ibrahim, Darnez Pope, Rukhaiya khaton, Abeer Alamri. Mental Health and Healthcare Access among Refugees. Oral and Poster Presentation at Refugees Health Summit, April 18th, 2019 Bowling Green, KY, USA.
- Ali, M. & **Ezekekwa, E.U.** (October 2018) - Interventions to Reduce 30-Day Hospital Readmission Rates for Adults with Heart Failure: A Systematic Literature Review. Poster presentation at NSF Center for Health Organization’s 2019 Spring Industry Advisory Board meeting on October 11th and 12th, Mclean, Virginia, USA.
- Ezekekwa, E. U.** (March 26, 2017) - An Examination of the likelihood of Home Discharge after General Hospitalizations among Medicaid Recipients. Oral presentation at the 47th WKU Student Research Conference, Bowling Green, KY, USA.
- Ezekekwa, E.U** (November 2016) - Evaluation of Long-term care in Kentucky: Oral presentation at: Department of Public Health; Bowling Green, KY, USA.

**Ezekekwu, E.U** (March 2016). Pattern of Marijuana and Illicit Drug consumption. Poster presentation at 46th WKU Student Research Conference, Bowling Green, KY, USA.

#### ACADEMIC SERVICE

---

2019 – 2022                    **University of Louisville School of Public Health and Information Sciences Department of Health Management and Systems Sciences**

- MSHA and EMSHA Survey development and
- 

2021 (Spring)                Advisor (Christopher E Johnson, PhD)

- PHMS 383 Healthcare Quality Management, PHMS 282 Healthcare Information & Technology Management, PHMS 305 Healthcare Organizational Leadership & Governance.

2016 – 2017                    **Western Kentucky University**

Advisor (William Mkanta, PhD)

Course: International Comparisons of Health Care Systems (Spring and Fall, 2017)

HCA 347                        Course: Health Care Quality and Accountability (Spring and Fall, 2016)

#### ACADEMIC SERVICE

---

2020 – Present                Reviewer, Academy of Management Annual Conference

2020 – Present                Reviewer, APHA 2020 Annual Meeting and Expo

2020 – Present                Reviewer, Journal of Primary Care & Community Health

2020 – Present                Reviewer, CDC Chronic Diseases

2020 – Present                Reviewer, Journal of Healthcare Management

2019 – Present                Reviewer, The Journal of Health Care Organization, Provision, and Financing

2019 – Present      Reviewer, AcademyHealth Annual Conference

## SKILLS

---

- Microsoft office including Excel, PowerPoint and Forms
- Project management
- Statistical Software: STATA, SPSS, and R
- SQL
- Qualtrics
- Endnote
- Review Manager
- Quality Improvement
- Data analytics and informatics
- Risk Management
- Time-efficient, systematic working methodology.

## CERTIFICATIONS

---

2018 – Present      CITI Programs: Responsible Conduct of Research and  
Basic/Refresher Course - Human Subjects Research

2013                      Certificate in Key accounts management

2013                      Certificate in customer centric interactions

## SCHOLARSHIPS AND AWARDS

---

2023                      Dean's Award, Graduation, University of Louisville

2023                      Winner of the 2022 Integrated Public Use Microdata Series  
(IPUMS) Research Award competition.

2021                      First Place Recipient, 2020 Jules Delambre Student Paper Contest  
by the Anthropologists and Sociologists of Kentucky

2020                      Recipient, School of Public Health and Information Sciences  
Scholarship for attendance at the APHA 2020 Annual Meeting and  
Expo

2018 – 2021              Recipient, Presidential Diversity Fellow, University of Louisville,

|           |   |
|-----------|---|
| 2018-2021 | Recipient, University of Louisville School of Public Health and Information Sciences Full Tuition Scholarship |
| 2017      | Dean's Award, Graduation, WKU   |
| 2017      | Recipient, American College of Healthcare Executives (ACHE) student scholarship                               |
| 2016-2017 | Recipient, College of Health and Human Services (CHHS) Scholarship, WKU                                       |

### PROFESSIONAL MEMBERSHIPS AND ACTIVITIES

- Member, AcademyHealth
- Member, Academy of Management
- Member, International Society for Pharmacoeconomics and Outcomes
- Member, American Public Health Association.
- Member, National Association of Health Services Executives
- Member, American College of Healthcare Executives
- Past Senate President (2016-2017), African Student Union, WKU Chapter
- Past Deputy secretary General (2007-2008), Pharmaceutical Association of Nigerian Students

### INDUSTRY EXPERIENCE

|                |  |
|----------------|--|
| 2022 – Present | Director of Research<br><b>Intrinsic Global, Inc.</b> , Louisville, KY                                       |
| 2021 – 2022    | Data Analyst and Co-Investigator<br><b>Intrinsic Global, Inc. (formerly Rapid Alliance)</b> , Louisville, KY |
| Spring, 2017   | Job Shadowing<br><b>Wayne County Hospital</b> , Monticello Kentucky  |
| 2014 -2015     | National Sales Manager<br><b>Tricare Pharmaceutical Ltd</b> , Lagos.   |
| 2013 -2014     | Specialist Medical Representative (Territory Coordinator)<br><b>MERCK &amp; CO</b> , Nigeria.                |
| 2012 – 2013    | Medical Representative<br><b>Sanofi</b>  |

2011-2012                      Pharmacy Manager  
**Jurdynal Health Pharmacy, Lagos.**

**VOLUNTEER SERVICE**

---

2021                              LouVax (Louisville metropolis vaccination)

2020 - 2021                      Louisville Metro Mask Survey Observation Project

2016                              Salvation Army, Bowling Green, Kentucky, USA

2016                              International Refugee Center, Bowling Green, Kentucky, USA

2010                              Red Cross, Nigeria.

**References Available Upon Request**