




Contents lists available at ScienceDirect

## Seminars in Cell and Developmental Biology

journal homepage: [www.elsevier.com/locate/semcdb](http://www.elsevier.com/locate/semcdb)

Review

The Calvin-Benson-Bassham cycle in  $C_4$  and Crassulacean acid metabolism speciesMartha Ludwig<sup>a,\*</sup>, James Hartwell<sup>b,2</sup>, Christine A. Raines<sup>c,3</sup>, Andrew J. Simkin<sup>c,d,4</sup><sup>a</sup> School of Molecular Sciences, University of Western Australia, Perth, Western Australia, Australia<sup>b</sup> Department of Biochemistry and Systems Biology, Institute of Systems, Molecular and Integrative Biology, University of Liverpool, Liverpool L69 7ZB, UK<sup>c</sup> University of Essex, Wivenhoe Park, Colchester CO4 3SQ, UK<sup>d</sup> School of Biosciences, University of Kent, Canterbury CT2 7NJ, UK

## ARTICLE INFO

## Keywords:

Calvin-Benson-Bassham cycle

 $C_4$ 

CAM

CO<sub>2</sub> assimilation

Crop improvement

Photosynthesis

## ABSTRACT

The Calvin-Benson-Bassham (CBB) cycle is the ancestral CO<sub>2</sub> assimilation pathway and is found in all photosynthetic organisms. Biochemical extensions to the CBB cycle have evolved that allow the resulting pathways to act as CO<sub>2</sub> concentrating mechanisms, either spatially in the case of  $C_4$  photosynthesis or temporally in the case of Crassulacean acid metabolism (CAM). While the biochemical steps in the  $C_4$  and CAM pathways are known, questions remain on their integration and regulation with CBB cycle activity. The application of omic and transgenic technologies is providing a more complete understanding of the biochemistry of  $C_4$  and CAM species and will also provide insight into the CBB cycle in these plants. As the global population increases, new solutions are required to increase crop yields and meet demands for food and other bioproducts. Previous work in  $C_3$  species has shown that increasing carbon assimilation through genetic manipulation of the CBB cycle can increase biomass and yield. There may also be options to improve photosynthesis in species using  $C_4$  photosynthesis and CAM through manipulation of the CBB cycle in these plants. This is an underexplored strategy and requires more basic knowledge of CBB cycle operation in these species to enable approaches for increased productivity.

## 1. Introduction

All photosynthetic organisms use the Calvin-Benson-Bassham (CBB) cycle to fix atmospheric CO<sub>2</sub> into organic compounds needed for

development and growth. The steps in this cycle and the enzymes involved were elucidated starting in the 1950 s by Calvin, Benson, Bassham, and colleagues [1,2]. As the first stable product of the CBB cycle is the 3-carbon compound, 3-phosphoglycerate (3-PGA), this

**Abbreviations:** Ald, fructose-1,6-bisphosphate aldolase; BS, bundle sheath; CA, carbonic anhydrase; CAM, Crassulacean acid metabolism; CBB, Calvin-Benson-Bassham; CCM, carbon concentrating mechanism;  $C_i$ , internal partial pressures; DHAP, dihydroxyacetone phosphate; E-4 P, erythrose-4-phosphate; FB, fructose-1,6-bisphosphatase; Fru-1, 6-BP, fructose 1,6-bisphosphate; Fru-6 P, fructose 6-phosphate; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; h, hour; HPLC-MS, high performance liquid chromatography-mass spectrometry; LD, light-dark cycle; LL, constant light conditions; LSU, large subunit; OAA, oxaloacetate; MDH, malate dehydrogenase; min, minute; NAD-ME, NAD-malic enzyme; NADP-ME, NADP-malic enzyme; PCK, PEP carboxykinase; 3-PGA, 3-phosphoglycerate; PEP, phosphoenolpyruvate; PEPC, phosphoenolpyruvate carboxylase; PGK, phosphoglycerate kinase; PNUe, photosynthetic nitrogen-use efficiency; PPDK, pyruvate phosphate dikinase; PRK, phosphoribulokinase; PSII, Photosystem II; PWUE, photosynthetic water-use efficiency; R-5 P, ribose-5-phosphate; RAF1, Rubisco activation factor 1; RBCS, Rubisco small subunit; RCA, Rubisco activase; RNA-Seq, RNA sequencing; RPI, ribose 5-phosphate isomerase; Rubisco, ribulose 1,5-bisphosphate carboxylase/oxygenase; RuBP, ribulose 1,5-bisphosphate; SB, sedoheptulose-1,7-bisphosphatase; Sed-1, 7-BP, sedoheptulose 1,7-bisphosphate; Sed-7 P, sedoheptulose 7-phosphate; SSU, small subunit; TK, transketolase; TP, triose phosphate; TPI, triose-P isomerase; TPT, triose phosphate:phosphate translocator; WT, wild type; Xyl-5 P, xylulose 5-phosphate.

\* Corresponding author.

E-mail address: [martha.ludwig@uwa.edu.au](mailto:martha.ludwig@uwa.edu.au) (M. Ludwig).<sup>1</sup> 0000-0002-0324-7602<sup>2</sup> 0000-0001-50000-223X<sup>3</sup> 0000-0001-7997-7823<sup>4</sup> 0000-0001-5056-1306<https://doi.org/10.1016/j.semcdb.2023.07.013>

Received 7 March 2023; Received in revised form 3 July 2023; Accepted 25 July 2023

1084-9521/© 2023 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).


pathway is also referred to as the C<sub>3</sub> cycle. Plants utilizing only this pathway to assimilate CO<sub>2</sub> are often referred to as C<sub>3</sub> plants and make up over 85% of plant species.

There are 13 reactions in the CBB cycle, which are catalyzed by 11 different enzymes (Fig. 1), and the cycle can be divided into three catalytic phases: carboxylation, reduction, and regeneration [1–3]. The carboxylation and reduction phases form the linear, assimilatory part of the C<sub>3</sub> cycle catalyzed by the enzymes phosphoribulokinase (PRK), ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco), phosphoglycerate kinase (PGK), and glyceraldehyde 3-phosphate dehydrogenase (GAPDH). The reactions catalyzed by PRK and PGK use ATP resulting from the light reactions of photosynthesis while NADPH produced through light harvesting is used by GAPDH. The products resulting from these reactions are triose phosphates (TPs), which can be exported from the chloroplast to the cytosol via the triose phosphate:phosphate translocator (TPT) for sucrose and isoprenoid synthesis, or they can enter the branched, regeneration phase of the cycle (Fig. 1). This regenerative phase results in the formation of the 5-carbon CO<sub>2</sub> acceptor molecule ribulose 1,5-bisphosphate (RuBP), and also produces a number of different carbon compounds which exit the cycle and form the basis of the biosynthetic pathways for starch (fructose 6-phosphate; Fru-6 P), thiamine, nucleotides (ribose 5-phosphate; R-5 P) and shikimate (erythrose 4-phosphate; E-4 P), with the latter being vital for the production of key metabolites including aromatic amino acids and phenylpropanoids (Fig. 1). Although the majority (five-sixths) of the TPs produced in the CBB cycle remains within the cycle to regenerate RuBP, one-sixth of the carbon compounds exit the cycle. The relative amount of carbon going to the different biosynthetic pathways is not fixed, and is likely to change during development, under different environmental conditions, and in response to changes in the rate of photosynthesis. However, a key factor that must remain constant is the retention of five

out of every six carbons within the cycle, which prevents the cycle from becoming depleted in the metabolic intermediates it needs to continue to function [1–3].

Rubisco, the enzyme catalyzing the carboxylation of RuBP, has several limitations. It is a relatively inefficient enzyme with a low turnover number, and it also catalyzes the oxygenation of RuBP, with the substrates for the carboxylation and oxygenation reactions competing for the same catalytic site [4]. The oxygenation reaction directs the flow of carbon through the photorespiratory pathway (Fig. 1), and this can result in losses of between 25% and 30% of the carbon fixed [5,6]. Environmental variables, such as high temperature and drought, can result in an increase in the oxygenase activity relative to the carboxylation reaction [7]. Therefore, promoting the Rubisco carboxylation reaction, which would simultaneously minimize the oxygenation reaction, has the potential to increase carbon assimilation significantly and would represent a step change in photosynthesis (up to 100% depending on temperature [8]).

Although all plants use the CBB cycle some species have evolved additional metabolic pathways that capture atmospheric CO<sub>2</sub> using a different carboxylase enzyme that has no oxygenase activity. These pathways act as carbon concentrating mechanisms (CCMs) that limit the oxygenase reaction of Rubisco and photorespiration. Our knowledge of CBB cycle operation, regulation, and metabolic integration in these plants is limited but understanding these processes is critical as it offers opportunities to inform strategies for photosynthetic enhancement and crop improvement.



**Fig. 1.** Schematic representation of the Calvin-Benson-Bassham (CBB) Cycle. Fructose-1,6-bisphosphatase (FB: EC.3.1.3.11), fructose-1,6-bisphosphate aldolase (Ald: EC.4.1.2.13), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), glycine decarboxylase (GDC), triosephosphate isomerase (TPI: EC 5.3.1.1), phosphoglycerate kinase (PGK: EC.2.7.2.3), phosphoribulokinase (PRK: EC.2.7.1.19), ribulose-1,5-bisphosphate carboxylase/oxygenase (R: EC.4.1.1.39), sedoheptulose-1,7-bisphosphatase (SB: EC.3.1.3.37), transketolase (TK: EC.2.2.1.1), 1,3-bisphosphoglyceric acid (1,3-BPGA), dihydroxyacetone phosphate (DHAP), erythrose-4-phosphate (E-4 P), fructose 1,6-bisphosphate (Fru-1,6-BP), fructose 6-phosphate (Fru-6 P), glyceraldehyde 3-phosphate (GA-3 P), 3-phosphoglycerate (3-PGA), ribulose-5-phosphate (Ru-5 P), ribulose biphosphate (RuBP), sedoheptulose 7-phosphate (Sed-7 P), sedoheptulose 1,7-bisphosphate (Sed-1,7-BP), xylulose 5-phosphate (Xyl-5 P) (see Simkin [127]).

## 2. Angiosperm carbon concentrating mechanisms favour carboxylation, minimise photorespiration, and have evolved in diverse lineages

Within angiosperms, two distinct metabolic adaptations of photosynthetic  $\text{CO}_2$  fixation have evolved that limit the oxygenation of RuBP and loss of fixed carbon through photorespiration (Fig. 2). The adaptations are known as  $\text{C}_4$  photosynthesis, where the first stable compound synthesized is a 4-carbon acid, oxaloacetate (OAA), and Crassulacean acid metabolism (CAM), which also generates OAA as the initial product of primary atmospheric  $\text{CO}_2$  fixation [9]. Both pathways are metabolic additions to the CBB cycle and increase the concentration of  $\text{CO}_2$  around Rubisco, thereby reducing the oxygenation reaction and flux through the photorespiratory pathway.


The  $\text{C}_4$  and CAM pathways concentrate  $\text{CO}_2$  around Rubisco through either a spatial separation of primary and secondary  $\text{CO}_2$  fixation across two distinct, but connected, photosynthetic cell types within the leaf ( $\text{C}_4$ ; Fig. 2), or a temporal separation of primary and secondary  $\text{CO}_2$  fixation within each photosynthetic mesophyll cell (CAM; Fig. 2). Both systems catalyze primary atmospheric  $\text{CO}_2$  fixation using phosphoenolpyruvate (PEP) carboxylase (PEPC), a cytosolic enzyme that does not recognize oxygen and has a higher affinity for  $\text{CO}_2$  (specifically  $\text{HCO}_3^-$ ) than Rubisco, making it a superior carboxylase [10].

### 2.1. $\text{C}_4$ photosynthesis

A  $\text{C}_4$  photosynthetic syndrome has evolved more than 60 independent times in the angiosperms, in both monocot and eudicot taxa [11]. While species operating a  $\text{C}_4$  pathway in a single cell type are known [12], most  $\text{C}_4$  species demonstrate Kranz anatomy in which leaf mesophyll cells surround bundle sheath (BS) cells, which surround the


vascular tissue. In Kranz  $\text{C}_4$  species, carbonic anhydrase (CA) converts  $\text{CO}_2$  entering the cytosol of leaf mesophyll cells to  $\text{HCO}_3^-$ , which is then used by PEPC to produce the 4-carbon OAA from PEP in the light period, and malate and/or aspartate move through plasmodesmata connections to BS cells where they are decarboxylated (Fig. 2). This decarboxylase activity results in high  $\text{CO}_2$  concentrations in the vicinity of Rubisco, which is found only in BS chloroplasts in  $\text{C}_4$  plants [13], and the  $\text{CO}_2$  is fixed by the CBB cycle [14]. The 3-carbon compound resulting from the decarboxylation reaction is used to regenerate PEP in the mesophyll cells through the activity of pyruvate phosphate dikinase (PPDK; Fig. 2). Under standard growth conditions, the production of carbohydrates is split between the two leaf cell types in  $\text{C}_4$  species with starch synthesized predominantly in the BS and mesophyll cells being the primary site of sucrose production although the BS also has capacity for sucrose synthesis in many  $\text{C}_4$  species (Fig. 3) [15–17].

Historically,  $\text{C}_4$  species were categorized into one of three subtypes based on the decarboxylase with the highest activity in their leaf tissue [14]. In NADP-malic enzyme (NADP-ME) species, decarboxylation occurs in the BS chloroplasts whereas in plants using NAD-malic enzyme (NAD-ME) as the major decarboxylase,  $\text{CO}_2$  is released in the BS mitochondria. In the third subtype, PEP carboxykinase (PCK),  $\text{CO}_2$  is released in the BS cytosol through the activity of PCK while an NAD-ME isoform is also active in BS mitochondria. Currently, it is recognised that for most  $\text{C}_4$  species, more than one type of  $\text{C}_4$  acid moves into the BS where more than one of the above decarboxylases are active [18,19]. Due to the high affinity of PEPC for  $\text{HCO}_3^-$ , the sequestration of Rubisco to BS chloroplasts, and Kranz anatomy where BS cell connections to intracellular airspaces are limited, the concentration of  $\text{CO}_2$  around Rubisco is high. This combination of biochemistry and anatomy results in a  $\text{C}_4$   $\text{CO}_2$  pump that limits Rubisco oxygenase activity and photorespiration and allows  $\text{C}_4$  plants to outcompete  $\text{C}_3$  species in hot, dry, and/or high light



**Fig. 2.** Schematic representation of the  $\text{CO}_2$  fluxes in  $\text{C}_3$ ,  $\text{C}_4$ , and Crassulacean acid metabolism leaves.  $\text{C}_3$  plants convert atmospheric  $\text{CO}_2$  into 3-phosphoglycerate (3-PGA) via ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco; Fig. 1). Sedoheptulose-1,7-bisphosphatase (SB) catalyzes a step in the regeneration phase of the Calvin-Benson-Bassham (CBB) cycle. In the leaves of  $\text{C}_4$  plants,  $\text{CO}_2$  is converted into the 4-carbon oxaloacetate (OAA) by the enzyme phosphoenolpyruvate (PEP) carboxylase (PEPC).  $\text{C}_4$  plants are categorized into three subtypes, based on the  $\text{C}_4$  acid decarboxylation enzyme (see text for detail). The NADP-malic enzyme (NADP-ME) type is shown in the figure (see also Fig. 3). The  $\text{CO}_2$  released through the decarboxylation event in the bundle sheath is fixed by Rubisco (Fig. 1). Crassulacean acid metabolism (CAM) plants fix  $\text{CO}_2$  at night (grey area), with PEPC converting  $\text{CO}_2$  into OAA, which is stored as the 4-carbon intermediate malic acid. During the day (green area), malate is decarboxylated and the  $\text{CO}_2$  released is fixed by Rubisco (Fig. 1). CAM plants are represented by two subtypes, 1, the NAD(P)-ME type, and 2, the PEP carboxykinase type. Carbonic anhydrase (CA), NAD(P)-malate dehydrogenase (NAD(P)-MDH); pyruvate phosphate dikinase (PPDK). Stars show locations

of enzymes targeted for manipulation to increase photosynthesis (see Tables 1–2). Adapted from Yamori et al. [168] and Ainsworth et al. [169].



**Fig. 3.** Schematic representation of NADP-malic enzyme-type  $C_4$  photosynthesis. Atmospheric  $CO_2$  and phosphoenolpyruvate (PEP) are converted into the 4-carbon oxaloacetate (OAA) in the mesophyll cell cytosol. In the mesophyll cell chloroplast, OAA is converted into malate (Mal), which is then translocated to a bundle sheath cell and decarboxylated in the chloroplast. The  $CO_2$  released enters the Calvin-Benson-Bassham (CBB) cycle and the 3-carbon pyruvate (PYR) diffuses into a mesophyll cell where it is regenerated into PEP. 3-phosphoglycerate (3-PGA) made in the CBB cycle is shuttled to a mesophyll chloroplast from the bundle sheath chloroplast where it can be either reduced to triose phosphate (TP) and returned to the CBB cycle in the bundle sheath chloroplast or converted to PEP via phosphoglycerate mutase and enolase. Sucrose synthesis occurs mainly in the mesophyll cell cytosol while starch is made in bundle sheath chloroplasts. Adapted from von Caemmerer and Furbank [68].

environments.

## 2.2. Crassulacean acid metabolism

In CAM species (Fig. 2), primary  $CO_2$  fixation by PEPC occurs in the cytosol of each chloroplast-containing mesophyll cell and happens in the dark period [20]. Stomata open at night and PEPC fixes both atmospheric and internal respiratory  $CO_2$  with PEP, forming OAA initially. OAA is rapidly converted to malate by malate dehydrogenase (MDH). Malate is stored in the mesophyll vacuole as malic acid throughout the night and reaches high concentrations by dawn. This nocturnal primary atmospheric  $CO_2$  fixation during CAM has been defined as phase I [21].

Phases II, III and IV of CAM occur in the light period [21]. Phase II represents a brief period of 1 to 2 h of atmospheric  $CO_2$  fixation after sunrise when stomata open and PEPC and Rubisco are active simultaneously. Phase III follows and represents a period of stomatal closure in the light period when  $CO_2$  from malate decarboxylation is refixed by Rubisco. Phase IV defines a period that can occur when all malate has been consumed and stomata reopen, allowing direct atmospheric  $CO_2$  fixation by Rubisco later in the light period [21].

In terms of the timing of the biochemical steps of the CAM pathway (Fig. 2), once the sun has risen, malic acid leaves the vacuole and is decarboxylated by either NAD(P)-ME or the combination of MDH, working in the opposite direction to the night, and PCK [20]. The mechanism of malate decarboxylation in the light depends on the specific CAM species, with the adaptation being found in at least 37 plant families and having evolved independently multiple times in several families [22]. In addition, the extent of CAM expression is dependent on the species and, in some species, dependent on the environmental conditions. CAM species have therefore been categorized as displaying either constitutive/obligate CAM or facultative/inducible CAM [23]. Obligate CAM species develop CAM even when well watered and continue to rely on CAM as their main mode of primary  $CO_2$  fixation throughout their lives. Facultative CAM species use  $C_3$  when well

watered and induce CAM in response to drought stress. A range of other environmental signals has also been reported to induce CAM in specific species, including high light intensity and a decrease in day length. Importantly, facultative CAM species are able to return to  $C_3$  when rewatered [23].

The  $CO_2$  released through malate decarboxylation in the light period results in high internal partial pressures of  $CO_2$  ( $C_i$ ) within the leaf air spaces due to diffusion of  $CO_2$  from the mesophyll cells. This rise in  $C_i$  is believed to promote stomatal closure and efficient secondary fixation of  $CO_2$  to sugars via the CBB cycle (phase III of CAM). It has been hypothesized that CAM species close their stomatal pores for much of the hot and dry light period in response to this build up of internal  $CO_2$  from malate decarboxylation [24].

The high  $CO_2$  concentration that is established in CAM photosynthetic tissues in phase III is widely assumed to favour the carboxylase activity of Rubisco over the oxygenation reaction. However, the extent to which photorespiration is suppressed in CAM species during phase III requires further detailed studies across a diverse range of independent CAM origins. This is important because stomatal closure in the light period also prevents oxygen, generated in chloroplasts through water splitting, escaping from the leaf. Thus, oxygen concentrations inside the leaf rise simultaneously with  $C_i$ . Data have been reported for the concentration of  $CO_2$  and  $O_2$  in the leaves of CAM species during phase III, and the results revealed that the concentration of  $O_2$  inside CAM tissues varied from 21% to 80% [25,26]. In turn, the ratio between the partial pressure of  $O_2$  and  $CO_2$  inside leaves of CAM species varied from 285 in *Hoya carnosus* to 81 in *Sedum praealtum* [26]. These ratios support the conclusion that  $CO_2$  will be enriched over  $O_2$  behind closed stomata in the light in phase III as the equivalent ratio for ambient air is 525 and that for the air in a  $C_3$  leaf is 840. Thus, the highest ratio reported for a CAM species (285 in *H. carnosus*) was three times better, in terms of favouring Rubisco carboxylation over oxygenation, relative to the air in a  $C_3$  leaf. Lüttge [25] used these published values for the concentrations of  $CO_2$  and  $O_2$  inside CAM tissues in phase III to calculate theoretical

values for the ratio between the reaction rate of Rubisco with O<sub>2</sub> and CO<sub>2</sub>. These calculations were performed using two possible equations and a range of published kinetic constants for Rubisco. The resulting calculated values led Lüttge [25] to conclude that the level of photorespiration in CAM leaves or stems behind closed stomata in the light period may be either very similar to that in C<sub>3</sub> leaves, at best half that in C<sub>3</sub> leaves, or that photorespiration may be suppressed very strongly in phase III of CAM.

### 2.3. Increased photosynthetic and nitrogen- and water-use efficiencies result from C<sub>4</sub> and Crassulacean acid metabolism carbon concentrating mechanisms

In plants that use C<sub>4</sub> photosynthesis, the concentration of CO<sub>2</sub> in the vicinity of Rubisco within the chloroplast, where the CBB cycle functions, is increased, thereby favouring carboxylation over oxygenation. This facet results in an increase in the amount of CO<sub>2</sub> being fixed per photon of sunlight absorbed by C<sub>4</sub> plants when compared to C<sub>3</sub> species [27,28]. Moreover, increasing the CO<sub>2</sub> concentration around Rubisco allows it to operate at close to its maximum catalytic rate, which leads to C<sub>4</sub> species being able to function optimally with lower amounts of Rubisco than C<sub>3</sub> plants [29]. Given that Rubisco accounts for a high proportion of total protein, and thus nitrogen, in C<sub>3</sub> leaves, the lower requirement for Rubisco in C<sub>4</sub> plants leads to increased photosynthetic nitrogen-use efficiency (PNUE; reviewed in Long, [29]). It is less clear that CAM plants achieve higher PNUE relative to C<sub>3</sub> species, but this may, at least in part, be due to a lack of appropriate studies in CAM species [30–32]. Overall, this means that CO<sub>2</sub> assimilation rates are greater per unit leaf nitrogen in the leaves of C<sub>4</sub> plants compared to those of C<sub>3</sub> plants [33]. In addition, C<sub>4</sub> biochemistry allows C<sub>4</sub> plants to achieve rates of CO<sub>2</sub> fixation sufficient to satisfy the C<sub>4</sub> pump at intercellular CO<sub>2</sub> partial pressures much lower than those found in C<sub>3</sub> leaves [34]. Thus, C<sub>4</sub> plants can maintain high rates of photosynthesis with lower stomatal conductance relative to a C<sub>3</sub> plant achieving the same rate, which in turn means that the transpirational loss of water during atmospheric CO<sub>2</sub> uptake for photosynthesis is lower in C<sub>4</sub> species than in C<sub>3</sub> plants [20,35]. Consequently, C<sub>4</sub> species demonstrate higher photosynthetic water-use efficiency (PWUE) than C<sub>3</sub> plants, and CAM plants have substantially greater PWUE than both C<sub>3</sub> and C<sub>4</sub> species because they open their stomata during the cooler and more humid dark period and close them for much of the hot and dry light period [20,36]. Under optimal conditions, these advantages allow C<sub>4</sub> plants to fix more carbon per unit of light, per unit of nitrogen, and per unit of water transpired, when compared to C<sub>3</sub> plants [20,30,33]. While these relationships are often less well characterized for CAM species, CAM plants do fix more CO<sub>2</sub> per unit of water transpired relative to C<sub>3</sub> and C<sub>4</sub> plants, and may display superior fixation of carbon per unit of light and per unit of nitrogen, although more data are required for CAM species for these latter two use efficiencies.

## 3. Biochemistry of the Calvin-Benson-Bassham cycle in C<sub>4</sub> photosynthesis and Crassulacean acid metabolism

While the enzymes catalyzing the steps of the CBB cycle are the same in all organisms, their properties, concentrations, and/or cellular locations differ between photosynthetic types and notably between plants using the same photosynthetic pathway.

### 3.1. C<sub>4</sub> photosynthesis: Calvin-Benson-Bassham cycle reactions are shared between mesophyll and bundle-sheath cells

In early work involving the isolation of mesophyll and BS cells and/or chloroplasts from plants representing all three C<sub>4</sub> subtypes, the enzymes catalyzing the reduction phase of the CBB cycle, PGK and GAPDH, as well as triose-P isomerase (TPI), were found in both cell types, while Rubisco, PRK, and chloroplast fructose-1,6-bisphosphatase (FB) showed

BS specificity (Fig. 1) [37–39]. While these results should be considered with the knowledge that most enzymes of C<sub>4</sub> photosynthesis are encoded by gene families [40], they have been supported by more recent proteomic studies that identified CBB cycle-associated isoforms in maize through mass spectrometry-based technologies [41,42]. Analyses of the maize leaf proteome data also indicated preferential localization of sedoheptulose-1,7-bisphosphatase (SB), fructose-1,6-bisphosphate aldolase (Ald), and transketolase (TK) (Fig. 1) to the BS [41,42].

Rubisco has been the most examined and best characterized of all the CBB cycle enzymes within datasets for C<sub>4</sub> species [43–46]. As described above, the C<sub>4</sub> CCM concentrates CO<sub>2</sub> around Rubisco in BS chloroplasts, which results in C<sub>4</sub> plant leaves containing less Rubisco than those of C<sub>3</sub> species [47]. Consistent with the activity of a CCM, C<sub>4</sub> plant Rubisco isoforms demonstrate lower CO<sub>2</sub> specificities compared to those of C<sub>3</sub> species, but with the trade-off of greater  $k_{cat}$  values ([48,49] and references therein). Studies in which C<sub>3</sub> plant Rubisco small (SSU) or large (LSU) subunits were replaced with those of C<sub>4</sub> plants have indicated elements controlling Rubisco kinetics are located on both subunits of the enzymes in both C<sub>3</sub> and C<sub>4</sub> species (Section 5.1.1) [50–52].

Comparisons of the amount or activity of other CBB cycle enzymes between C<sub>3</sub> and C<sub>4</sub> species are limited. Whole leaf extracts of several NADP-ME-, NAD-ME-, and PCK-type C<sub>4</sub> grasses showed PRK activity, which is predominantly in the BS (see above), is similar to that of wheat whole leaf extracts [39]. Whole leaf TPI activity, which is split between mesophyll and BS cells (see above), in these same C<sub>4</sub> grasses was comparable to or greater than that of wheat leaf extracts [39]. By contrast, the activity of chloroplast GAPDH, which is also split between the two cell types (see above), in species representing the three C<sub>4</sub> subtypes was lower than that of wheat [53]. As for studies reporting CBB cycle enzyme location, the presence of multiple enzyme isoforms needs to be considered when interpreting the results of these whole leaf enzyme activity studies.

Chloroplast dimorphism is typically seen between mesophyll and BS cells of NADP-ME-type C<sub>4</sub> species although the extent of the differences between the plastids is variable among species and likely dependent on development and environmental factors [54–63]. Under standard growth conditions, when dimorphic chloroplasts are present, the BS organelles exhibit few grana stacks and significant reduction in Photosystem II (PSII) activity [64], which contribute to maintaining low O<sub>2</sub> levels around Rubisco due to the absence of PSII H<sub>2</sub>O-splitting complexes. However, limited linear electron flow and only about half the NADPH requirement of the CBB cycle being met through the NADP-ME decarboxylation reaction in the BS chloroplasts [65] would inhibit RuBP regeneration. This photochemical imbalance is mitigated through a 3-PGA/TP shuttle with approximately half the 3-PGA generated by Rubisco transported into mesophyll chloroplasts (Fig. 3) [14], which are characterised by thylakoid grana with PSII activity [64]. In mesophyll chloroplasts, 3-PGA is converted to TPs, of which at least two-thirds [66] must be translocated to BS cells for entry into the CBB cycle and regeneration of RuBP as well as starch synthesis (Fig. 3). The TPs remaining in the mesophyll are used predominantly for sucrose synthesis (Fig. 3) [17] while some BS-generated 3-PGA may also be converted to PEP in the mesophyll, thereby supporting PEPC activity (Fig. 3) [67].

Although BS chloroplasts of NADP-ME- and PCK-type C<sub>4</sub> species contain grana stacks and PSII activity [64], as noted above, the enzymes of the CBB cycle reduction phase are also found in both mesophyll and BS cells of these plants, indicating they also contain a functional 3-PGA/TP shuttle [14,39,68]. The separation of the CBB cycle reactions between mesophyll and BS cells along with the likelihood of mixed decarboxylation pathways operating in all C<sub>4</sub> species allow flexibility in balancing the photochemical load and carbohydrate synthesis between the two cell types in response to environmental changes and/or during leaf and plant development (reviewed in [17,18,69–71]). However, details on the flux through the individual C<sub>4</sub> acid transfer cycles in a single species, including its regulation, involvement of a 3-PGA/TP

shuttle, and mesophyll/BS partitioning of sucrose and starch synthesis, under different growth conditions and stages of development are limited.

### 3.2. Crassulacean acid metabolism: the temporal integration of two carboxylases

Although a relatively limited number of studies have explored the properties of the enzymes of the CBB cycle in CAM species, and/or studied the operation of the CBB cycle as a whole in the photosynthetic tissues of CAM plants, some insights have been gleaned from studies of *in vitro* enzyme activities in extracts from *Kalanchoë* leaves and instantaneous online carbon isotope discrimination. Early work established that several enzymes within the CBB cycle are activated by light in a redox-dependent manner, and thus share this characteristic in common with their counterparts in  $C_4$  and  $C_3$  species [72,73]. Specifically, Gupta and Anderson [72] reported that PRK, NADP-GAPDH and SB were light-activated in *K. blossfeldiana* cv. Tetra Vulcan, but FB was not, whereas Hutcheson and Buchanan [73] demonstrated that FB and SB from leaves of *K. daigremontiana* were subject to light-activation by reduced thioredoxin-f.

Additional understanding of the CBB cycle in a CAM species (pineapple; *Ananas comosus*) was provided by applying high performance liquid chromatography-mass spectrometry (HPLC-MS) with mixed-mode stationary phases to identify and quantify the notoriously difficult CBB cycle intermediates. The concentrations of Fru-6 P, glucose 6-phosphate, dihydroxyacetone phosphate (DHAP), 3-PGA, PEP, and fructose 1,6-bisphosphate (Fru-1,6-BP) were all measured at levels comparable to those in the  $C_3$  model species *Arabidopsis thaliana*, plus R-5 P, xylose 5-phosphate (Xyl-5 P), and E-4 P were below the level of detection, which was also the case for *A. thaliana* [74]. However, *A. thaliana* and three other  $C_3$  species (barley, peppermint, and pea) all had detectable levels of RuBP at the level of  $< 5 \mu\text{mol m}^{-2}$ , whereas RuBP was below the level of detection in pineapple [74]. Despite the absence of a large body of literature studying the characteristics of the CBB cycle enzymes and associated metabolite levels in a diverse range of CAM species, these studies do support the general conclusion that the CBB cycle in CAM species is operating behind closed stomata in the light period in much the same way as it is known to be operating in  $C_3$  species behind open stomata.

Most work on CAM species in relation to the CBB cycle has focused on Rubisco, which like the  $C_4$  homologues have lower  $\text{CO}_2$  specificity and greater  $k_{\text{cat}}$  values relative to the Rubiscos of  $C_3$  species [48], as well as the interplay between PEPC and Rubisco activities throughout the light period. In particular, studies in species of *Kalanchoë* that perform constitutive/obligate CAM, and bromeliads such as the facultative CAM species *Guzmania monostachia*, demonstrated that Rubisco activation during the light period is delayed relative to the norm for  $C_3$  species, where Rubisco is activated rapidly at dawn [75,76]. Findings demonstrated that Rubisco activation state remains low during malate decarboxylation in phase III of CAM, but then Rubisco activase (RCA) protein abundance and Rubisco activation state rise in anticipation of phase IV of CAM when stomatal pores can re-open later in the light period and atmospheric  $\text{CO}_2$  is assimilated directly via the CBB cycle [76]. Furthermore, online carbon isotope discrimination data for the leaves of *K. daigremontiana* sampled at different time points across the light period revealed that during phase II of CAM, a short period of stomatal opening and atmospheric  $\text{CO}_2$  fixation for 1–2 h after dawn, the instantaneous carbon isotope discrimination transitioned from a signal indicative of PEPC fixing most of the  $\text{CO}_2$  ( $-6\text{‰}$  to  $-7\text{‰}$ ) to a Rubisco signal ( $-16\text{‰}$ ) [76,77]. Likewise, in the bromeliad *Tillandsia utriculata*, discrimination transitioned from PEPC in early phase II to Rubisco late in phase II [78] and a similar progression was measured in *Clusia rosea*, which displays constitutive CAM. By contrast the facultative CAM species *C. minor* had an instantaneous discrimination signal indicative of Rubisco dominating in both phase II and phase IV in the light period [79]. In phase IV, when

stomata re-open for the final few hours of the light period, instantaneous discrimination in *K. daigremontiana* declined from  $-13\text{‰}$  to  $-11\text{‰}$ , indicating a transition from Rubisco to PEPC as dusk was approached [76]. Equivalent data for phase IV in *T. utriculata* also showed that the instantaneous discrimination transitioned from a Rubisco signal early in phase IV to a more PEPC-influenced signal late in phase IV [78]. More recently, experiments using short periods of darkness applied during phase IV of CAM with the orchid *Phalaenopsis* cv. Sacramento revealed that PEPC was the major carboxylase in the last few hours of phase IV as  $\text{CO}_2$  fixation did not decline in response to short 5-min blackouts applied at 2 h and 0.5 h before dark [80]. By contrast, *K. blossfeldiana* cv. Saja leaves displayed a sharp decline in  $\text{CO}_2$  fixation in response to 5-min blackouts applied throughout phase IV, indicating that Rubisco remained the major carboxylase right through to dusk [80]. However, it is important to note that the 24 h pattern of  $\text{CO}_2$  exchange for *Phalaenopsis* cv. Sacramento in this study did not show the classic decline in  $\text{CO}_2$  fixation to zero or below zero at dawn and dusk that mark the start of phase II and end of phase IV in other CAM species that display the four phases of gas exchange. Instead, phase IV in *Phalaenopsis* cv. Sacramento appeared to be a prolonged start to phase I, with the rate of  $\text{CO}_2$  assimilation continuing without a dip through the light to dark transition, and phase II also had the characteristics of a continuation of phase I into the early hours of the light period, with no dip in  $\text{CO}_2$  fixation at the dark to light transition [80]. Hence, the physiology of *Phalaenopsis* cv. Sacramento reported by van Tongerlo et al. [80] was abnormal at dawn and dusk relative to all previous reports of CAM gas exchange patterns in other species; consequently, the response to short periods of darkness applied in phase IV may not be broadly applicable across diverse CAM species.

Such data from both *Kalanchoë* species, the orchid *Phalaenopsis* cv. Sacramento, and the bromeliads *G. monostachia* and *T. utriculata*, which represent distantly related, independent origins of the CAM adaptation, emphasize that the circadian clock most likely regulates this fine balancing between PEPC and Rubisco activity at different times across the light period in a wide diversity of CAM origins. An important role for the clock is likely because PEPC and Rubisco are clock controlled via PEPC kinase and RCA, respectively. However, it is also clear that the specific timing of the transitioning and balancing between PEPC and Rubisco differs between CAM species, which in turn emphasizes the need to study the carboxylase interplay over the light period in a much greater diversity of CAM plants. Furthermore, far greater attention should be paid to studying these aspects of CAM under natural dawn and dusk transitions in light, temperature and humidity, as many lab experiments use square wave light/ dark cycles that do not reflect the gradual transitions that occur in the field.

Addressing the role of the circadian clock in balancing the two carboxylases during CAM directly, Davies and Griffiths [81] studied the facultative CAM species, *Mesembryanthemum crystallinum* (common iceplant) under constant light (LL) and temperature conditions. LL conditions are a key experimental method to establish whether or not the endogenous circadian clock is regulating the observed temporal changes. Again, instantaneous carbon isotope discrimination measurements on CAM-induced leaves measured under circadian free-running LL conditions demonstrated that there was a transition from a Rubisco-mediated  $C_3$  discrimination signal in the early hours of the subjective night, to a  $C_4$  discrimination due to PEPC late in the subjective dark [81]. This study also demonstrated that *in vitro* Rubisco activation state reached a 24 h minimum at the start of the dark period under light-dark (LD) cycles, but then rose for much of the dark period, perhaps pre-empting dawn. The activation state also reached a minimum shortly after subjective dawn under LL conditions, coincident with the timing of maximal malate decarboxylation [81]. Under LL, Rubisco activation state reached its peak level in the subjective dark period, overlapping with the timing of the Rubisco-dominated isotope discrimination signal in the first hours of the subjective dark, and prior to the transition to a PEPC-dominated  $C_4$  discrimination signal later in

the subjective dark. Furthermore, immunoblot analysis showed that two isoforms of RCA were detected in C<sub>3</sub> leaves of *M. crystallinum* when measured every 24 h across 3 days of LL conditions, whereas the larger RCA isoform was barely detectable in the CAM-induced leaves under LL. These findings, spanning multiple independent origins of CAM across monocots and eudicots, emphasize that there is a delicate interplay between the regulation and balancing of the two competing carboxylases in CAM leaves, with circadian control important for the optimal timing of peak PEPC activity in the dark, and Rubisco in the light.

#### 4. Insights into Calvin-Benson-Basham cycle operation in C<sub>4</sub> photosynthesis and Crassulacean acid metabolism from omics

In this era when high-throughput omics approaches have been leveraged to delve more deeply and globally into a wide diversity of biological questions, it is not surprising that advances are being made possible in the fields of C<sub>4</sub> and CAM research through applying genome-, transcriptome-, proteome- and metabolome-wide approaches. Early insights into genome content, organization, and evolution were achieved through the publication of the sorghum and maize genomes for C<sub>4</sub> [82, 83], and the genomes of two orchid (Orchidaceae) species, *Phalaenopsis equestris* and *Dendrobium officinale*, and *Ananas comosus* for independent origins of CAM in monocots, plus *K. fedtschenkoi* for CAM in a eudicot [84–86]. Since these initial breakthroughs, complete genome sequences have continued to be published for species representing independent origins of both C<sub>4</sub> and CAM [87]. Further novel insights into each pathway have also been gained from several detailed large-scale transcriptome, proteome, and/or metabolome datasets generated from photosynthetic tissues of a range of C<sub>4</sub> and CAM species to answer specific questions on C<sub>4</sub> photosynthesis and CAM [88–96]. The availability of such ‘big data’ for a wide diversity of C<sub>4</sub> and CAM species offers unprecedented opportunities to elucidate the complete genetic blueprint for each pathway, and for a range of independent evolutionary origins. In contrast to most previous work, it is now possible to globally explore the regulation and functioning of the genes, transcripts, proteins, and metabolites of the CBB cycle in a wide diversity of C<sub>4</sub> and CAM species. As C<sub>4</sub> and CAM species use the CBB cycle for secondary CO<sub>2</sub> fixation to provide the building blocks supporting wider metabolism, growth, and development, it is clear that understanding the regulation and operation of the CBB cycle and its integration with the C<sub>4</sub> cycle and CAM are vitally important for achieving a complete understanding of both pathways. Such work is now possible through analyses of existing published datasets.

##### 4.1. C<sub>4</sub> photosynthesis

In addition to solidifying mesophyll and/or BS cell locations of CBB enzymes (Section 3.1) [41,42], proteome analyses have also given insight into CBB enzyme expression patterns during the differentiation of mesophyll and BS chloroplasts in maize [55]. A developmental gradient exists in grass leaves with cells at the leaf base containing proplastids and those near the tip fully differentiated mesophyll and BS chloroplasts [97]. Using maize leaf and BS-strand sections along this developmental gradient for proteome analysis, Majeran et al. [55] showed the differential enrichment of CBB cycle enzymes seen in mesophyll and BS chloroplasts at the leaf tip begins at about 4 cm from the leaf base, at the source and sink transition zone. Plastids at the base of the leaf were found to lack CBB cycle enzymes although mesophyll and BS chloroplasts could be distinguished in electron micrographs of sections between 2 and 3 cm from the leaf base due to differences in grana stacking (Section 3.1) [55]. Interestingly, the abundance of C<sub>4</sub> cycle enzymes (Section 2.1; Figs. 2, 3) also begins to increase in leaf cells located about 4 cm from the base, indicating the activities of the enzymes in this cycle are closely linked with those of the CBB cycle [55]. While the above work has increased our knowledge of CBB cycle operation during leaf development in a NADP-ME-type C<sub>4</sub> species,

corresponding information for NAD-ME- and PCK-type species is lacking. A systems biology study reported Rubisco protein amounts did not differ between mature and immature phytomers of *Setaria italica*, another NADP-ME-type C<sub>4</sub> species [98]; however, no information was reported for other CBB enzymes. Consequently, it is also not known if C<sub>4</sub> species of the same decarboxylation subtype exhibit similar developmental patterns.

Variation in CBB cycle operation among different C<sub>4</sub> species of the NADP-ME subtype has been detected at the level of CBB cycle intermediates in recent metabolite profiling studies, with the diversity attributed to differing levels of 3-PGA/TP shuttle activity in the four species (Section 3.1; [65,99] and Clapero et al. [100], this special issue). Principal component analyses in these studies also showed that RuBP was the driver separating the C<sub>3</sub> and C<sub>4</sub> species examined; C<sub>4</sub> taxa have lower RuBP levels, enabled by the presence of a CCM [65,99,100]. Metabolite profiles of C<sub>3</sub>, C<sub>3</sub>-C<sub>4</sub> (species showing leaf anatomy and/or photosynthetic biochemistry intermediate to C<sub>3</sub> and C<sub>4</sub> plants), C<sub>4</sub>-like (species lacking optimization of C<sub>4</sub> anatomy and/or biochemistry), and C<sub>4</sub> species of *Flaveria* have shown the C<sub>4</sub> species have higher levels of 3-PGA relative to the non-C<sub>4</sub> congeners [101]. Few differences in the levels of CBB cycle intermediates were found between the C<sub>3</sub> and C<sub>3</sub>-C<sub>4</sub> intermediate *Flaveria* species, whereas RuBP, Fru-6 P; Fru-1,6-BP, sedoheptulose 7-phosphate (Sed-7 P), and sedoheptulose 1,7-bisphosphate (Sed-1,7-BP) (Fig. 1) levels showed an overall trend toward lower levels in the C<sub>4</sub>-like species relative to the C<sub>3</sub>-C<sub>4</sub> species and lower again in the true C<sub>4</sub> species [101]. These modifications are consistent with the integration of the CBB cycle in the progressive evolution of a full C<sub>4</sub> syndrome [102].

At the transcript level, RNA-Seq datasets have been generated from whole leaf tissue of closely related members of the Cleomaceae, *Flaveria*, Paniceae, and *Moricandia* for comparative transcriptomic studies [103–106]. Not surprisingly, relative to the C<sub>3</sub> comparators, a decreased abundance of transcripts encoding CBB cycle enzymes was found in all the C<sub>4</sub> species [104–106]. Bräutigam et al. [105] suggested this down-regulation of expression should be considered in strategies to engineer a C<sub>4</sub> pathway into C<sub>3</sub> crop plants as it, along with reduced expression of proteins in the photorespiratory pathway and protein translation, may be important for achieving the high PNUE demonstrated by C<sub>4</sub> plants (Section 2.3). The genus *Moricandia* contains C<sub>3</sub> and C<sub>3</sub>-C<sub>4</sub> photosynthetic intermediate species but no true C<sub>4</sub> species [107]. A decreasing trend in the abundance of transcripts coding for most of the CBB cycle enzymes was found for two C<sub>3</sub>-C<sub>4</sub> intermediate *Moricandia* species relative to the C<sub>3</sub> congener *M. moricandioides*, and differences between the intermediate species were apparent [103]. This decrease in transcripts coding for CBB cycle enzymes in *Moricandia* C<sub>3</sub>-C<sub>4</sub> intermediates, however, was not paralleled in the C<sub>3</sub>-C<sub>4</sub> intermediate *F. ramosissima*, where only transcripts encoding Rubisco were decreased relative to the levels of the *Flaveria* C<sub>3</sub> congener [104]. In addition, an increase in transcripts encoding components of the C<sub>4</sub> cycle accompanied the decrease in transcripts coding for the CBB cycle enzymes in *Flaveria* C<sub>3</sub>-C<sub>4</sub> and C<sub>4</sub> species but not in *Moricandia* photosynthetic intermediates [103].

Predictably, transcript profiles along leaf developmental gradients from the NADP-ME-type C<sub>4</sub> grasses maize, sugarcane, and the non-model *Antaenanthia lanata*; the NAD-ME-type C<sub>4</sub> eudicot *Gynandropsis gynandra*; and C<sub>3</sub> and proto-Kranz (incipient C<sub>3</sub>-C<sub>4</sub> photosynthetic intermediate) comparators showed investment in transcripts encoding CBB cycle enzymes was high in mature regions of the leaf but lower in C<sub>4</sub> than in C<sub>3</sub> or proto-Kranz species [71,108–112]. In the case of maize for which corresponding proteome information is available [55], a high correlation was seen between CBB cycle enzyme transcript and protein profiles across the gradient [71,109]. Interestingly, *G. gynandra* exhibited correlated delays in the accumulation of mRNAs encoding CBB cycle enzymes and leaf cell differentiation relative to its C<sub>3</sub> congener *Tarenaya hassleriana* [110]. A co-expression study using leaf developmental transcriptomes of sorghum, *Setaria viridis*, maize, and rice found

Ald, FB, and SB are important in regulating carbon flux through the CBB cycle and it was suggested these enzymes should be considered in crop improvement strategies [113]. Further analysis of these developmental datasets would likely add to our understanding of how the components of the CBB and  $C_4$  cycles become integrated to produce a full  $C_4$  syndrome.

Information on the mesophyll and BS cell compartmentation of transcripts encoding CBB enzymes has also been provided for diverse  $C_4$  species, including maize, *S. viridis*, *G. gynandra*, and *Panicum virgatum* [108,109,114–116]. This cell-specific information has highlighted some differences and similarities within and between decarboxylation subtypes of  $C_4$  species. A high correlation was reported between transcriptomes from laser-capture microdissected maize mesophyll and BS cells and previous mesophyll and BS cell proteome datasets, with transcripts encoding CBB cycle enzymes enriched in BS cells [109,116]. Consistent with enzyme activity data (Section 3.1) and maize proteome and transcriptome studies [41,42,109,116], transcripts encoding the CBB cycle-associated GAPDH, PGK, and TPI were more abundant in mesophyll cells than in BS cells of the two NAD-ME-type species *G. gynandra* and *P. virgatum*, while mRNAs coding for other CBB cycle enzymes preferentially accumulated in the BS cells [108,114]. Compartmentation of transcripts encoding CBB cycle enzymes between mesophyll and BS cells from *S. viridis* was overall very similar to that of maize [115]; however, two exceptions indicate differences exist in CBB cycle operation and regulation in these two NADP-ME-type species. Transcripts encoding PGK are more abundant in maize mesophyll cells, as is the cognate protein [41,42], whereas in *S. viridis*, BS cells preferentially accumulate mRNA coding for PGK [115]. It was suggested that this difference may be due to posttranscriptional or posttranslational mechanisms controlling PGK expression in the mesophyll of *S. viridis* [115]. Posttranscriptional or translational regulation was suggested to explain the difference in accumulation of transcripts encoding ribose 5-phosphate isomerase (RPI) in the two species. In *S. viridis*, mRNA coding for RPI is enriched in BS cells [115], which agrees with the compartmentation of the maize RPI protein [41,42]; however, in maize, transcripts encoding RPI show preferential accumulation in the mesophyll [109,115].

#### 4.2. Crassulacean acid metabolism

Despite the extensive nature of the published quantitative RNA-Seq transcriptome analysis datasets for a wide diversity of CAM species that represent both obligate/constitutive CAM and facultative/inducible CAM, and that span many independent origins, few insights into the regulation of transcripts encoding the individual enzymes of the CBB cycle have been highlighted to date. In some cases, a decline in the transcript abundance of genes encoding the small subunit of Rubisco (*RBCS*) has been documented (e.g., [91,93]). Furthermore, in *Portulaca oleracea*, a species with  $C_4$  photosynthesis that induces weak CAM (low, but detectable, net atmospheric  $CO_2$  fixation to malate in the dark) in response to drought-stress, both *RBCS* transcripts, and transcripts for most of the enzymes of the photorespiratory pathway, were down-regulated in response to drought and thus coincident with increased CAM [91]. However, detailed insights into the transcript-level regulation of all genes encoding CBB cycle enzymes in CAM species are hiding in plain sight in the published RNA-Seq datasets. Re-analysis of these datasets with a specific focus on the genes encoding CBB cycle enzymes will yield new understanding that can guide more detailed studies into the regulation of associated CBB cycle enzymes and their regulators, and the associated metabolite levels of the CBB cycle intermediates and levels of flux through the pathway during phase III of CAM.

Within our own quantitative RNA-Seq datasets for *K. fedtschenkoi* leaves, comparing the 12-h-light/12-h-dark temporal profile of all transcripts detected from leaf pair 1 (the youngest visible leaves at the shoot apical meristem,  $C_3$ ) and leaf pair 6 (fully expanded leaves that use

full CAM defined by only performing phase I and phase III), a number of intriguing observations can be made regarding the regulation of transcripts encoding proteins associated with the CBB cycle. The most abundant *RBCS* transcript shifts its daily temporal peak by 12 hours from the end of the dark/beginning of the light period in leaf pair 1  $C_3$  to the end of the light/dusk in leaf pair 6 full CAM (Boxall, Dever, Kneřová and Hartwell, unpublished). The transcript levels of *PGK* display a very similar pattern to *RBCS*, both in terms of differential regulation at different diel times between leaf pair 1  $C_3$  and leaf pair 6 full CAM, and the 12-h shift in the timing of the daily peak transcript level; *PGK* transcripts also peak at the end of the light in leaf pair 6 full CAM. In addition, *RCA* displays a dramatic, up to 8-fold induction in leaf pair 6 full CAM relative to leaf pair 1  $C_3$ , with the transcript level rising sharply throughout the dark period, reaching a peak 2 h before dawn. Genes encoding various chloroplast *GAPDH* subunits, two *ALD* transcripts, a ribose 5-phosphate epimerase transcript, *PRK*, *FB*, and transcripts for two *SB* isogenes were all induced in leaf pair 6 CAM to differing extents, and at a range of times across the 24 h cycle. By contrast, transcript levels for *TPI*, one of two isogenes encoding TK, and RPI were all repressed in leaf pair 6 CAM relative to leaf pair 1  $C_3$  (Boxall, Dever, Kneřová and Hartwell, unpublished). Although one of the major changes coincident with leaf development in *K. fedtschenkoi* is the transition from  $C_3$  photosynthesis in leaf pair 1 to full CAM in leaf pair 6, it is important to note that this developmental range also spans from leaf pair 1 being sink leaves to leaf pair 6 being source leaves, and leaf pair 1 are unable to generate clonal plantlets from the notches in their margins when detached from the plant, whereas leaf pair 6 have gained the competence to develop leaf margin adventitious plantlets. Thus, the detected changes in the regulation of transcript levels for genes encoding CBB cycle enzymes between leaf pair 1 and leaf pair 6 may also be associated with these developmental changes in the biology of the leaves either in concert with a role in relation to CAM, or distinct from the development of full CAM in leaf pair 6.

These findings contrast with the current assumption that the CBB cycle is universally down-regulated following CAM development and/or induction due to the increased efficiency of Rubisco following the establishment of the CAM CCM. Furthermore, these results highlight the value of exploring the regulation of CBB cycle gene transcript levels in the transcriptome datasets for the many other diverse CAM species that are already published and awaiting re-analysis. Such studies can help to identify key genes coding for CBB cycle enzymes in CAM species that merit more detailed investigation through the generation of transgenic mutant lines using techniques such as RNAi and gene editing [36,117, 118].

#### 5. Engineering improvement

Overcoming stagnating crop productivity for current and future food, fodder, biofuel, and bioproduct demand is a major goal for plant scientists [119]. Identification of targets and development of strategies to improve  $CO_2$  assimilation in  $C_4$  plants are complicated by the separation of the  $CO_2$  assimilation reactions between two cell types, including flux through the PGA/TP shuttle and differing electron transport capacity between mesophyll and BS cells, as well as the presence of mixed decarboxylation pathways. Nevertheless, several recent reviews have highlighted targets and strategies to improve both  $C_4$  and  $C_3$  crop performance [68,120–127]; however, little or no attention has been directed at enhancing CAM. Below are summarized transgenic experiments in  $C_3$  and  $C_4$  species that have given insights into the regulation of the CBB cycle (Fig. 2) that may lead to crop improvement.



## 5.1. Current knowledge gleaned from transgenic studies in C<sub>3</sub> and C<sub>4</sub> plants

### 5.1.1. Rubisco

As the primary carboxylase in C<sub>3</sub> plants, Rubisco is limiting for CO<sub>2</sub> assimilation and CBB cycle function in these species [128]; however, antisense experiments in *Flaveria bidentis* indicated Rubisco is also a major determinant limiting CO<sub>2</sub> assimilation at saturating light levels in this C<sub>4</sub> eudicot species, with photosynthetic rate decreasing in transformants as Rubisco content and activity decreased (Fig. 2; Table 1) [129,130].

Further evidence of the importance of Rubisco in the photosynthetic flux of C<sub>4</sub> plants has come from the overexpression of genes encoding maize Rubisco SSU, LSU, and Rubisco activation factor 1 (RAF1) in maize (Table 1) [132]. These LSSS-RAF1 transformants showed increases in Rubisco content, CO<sub>2</sub> assimilation, growth, and biomass production relative to wild type (WT) maize. Moreover, while most C<sub>4</sub> plants demonstrate decreases in CO<sub>2</sub> assimilation under chilling conditions [133], the LSSS-RAF1 transformants outperformed WT maize in photosynthetic and growth characteristics as well as recovery following the chilling period [131]. The LSSS-RAF1 transformants also demonstrated better recovery, e.g., higher CO<sub>2</sub> assimilation and biomass, following drought stress than WT maize although no benefits in photosynthesis or growth relative to the control plants were observed during the stress treatment [134]. These results support earlier work that implicated Rubisco content and/or activity as limiting photosynthetic capacity in several C<sub>4</sub> species under chilling stress [133,135–141] and water limitation [142,143].

Either partial or complete replacement of rice Rubisco SSU with sorghum homologues led to a Rubisco with C<sub>4</sub>-like kinetics, including a higher  $k_{cat}$  and  $K_m$  for CO<sub>2</sub> and a reduced specificity for CO<sub>2</sub> (Table 1) [50,52]. No change in CO<sub>2</sub> assimilation at the various CO<sub>2</sub> concentrations tested was apparent in the plants containing a mixture of rice and sorghum SSU isoforms [50] and, except for a reduction in non-productive tillers in the transformants, there were no obvious growth differences relative to WT rice. By contrast, transformants containing only sorghum SSU demonstrated higher CO<sub>2</sub> assimilation rates than WT rice and equal biomass production when CO<sub>2</sub> concentrations were higher than ambient levels [52]. The above results indicate manipulation of Rubisco has the potential to increase CO<sub>2</sub> assimilation, growth, and yield under non-stress and stress conditions in both C<sub>3</sub> and C<sub>4</sub> crops.

### 5.1.2. Sedoheptulose-1,7-bisphosphatase

Knockdown experiments in a number of C<sub>3</sub> plants, including tobacco (*Nicotiana glauca*), have shown that small reductions in SB (Fig. 1)

activity (>9%) result in a >12% reduction in carbon assimilation ( $A_{max}$ ), growth rates, and biomass yields [144–147]. These data demonstrate that SB exerts strong regulation over CBB cycle flux, identifying it as a good target for crop improvement strategies.

Several studies have overexpressed the target enzyme SB, in an effort to increase plant biomass (Fig. 2; Table 2). In the model plant *A. thaliana*, transgenic plants showed a 42% increase in plant biomass and 39–53% increase in seed yield compared to WT [148]. In tobacco, constitutive overexpression of SB resulted in a 40% increase in shoot biomass under controlled greenhouse conditions [149,150]. More recently, the overexpression of SB in wheat was shown to result in an increase in biomass yield and more importantly, a 30–40% increase in seed yield [151] and in tomato, photosynthesis increased by 20–25%, increasing vegetative biomass by 30%, resulting in early onset flowering [152]. These results demonstrate that the manipulation of this CBB cycle enzyme can result in biomass/yield increases in model crops, a cereal crop, and a fruiting crop, suggesting it is a viable route for the manipulation of carbon assimilation in plants (see [122,127] for review). However, a contrasting result was observed when SB was overexpressed in the C<sub>4</sub> plant *Setaria viridis* (Fig. 2; Table 2).

Transgenic plants with 1.5- to 3.2-times increase in SB levels showed no correlation between SB content and saturating rates of CO<sub>2</sub> assimilation [155]. This result suggests that alternate limiting steps may exist in C<sub>4</sub> plants compared to C<sub>3</sub> species. It should be noted that targeting multiple steps in the CBB cycle in C<sub>3</sub> plants can result in a step change in assimilation increase [122,127]. In C<sub>4</sub> plants, it may be necessary to target multiple steps to illicit the same response observed in C<sub>3</sub> plants, while also considering that limiting steps in CBB cycle function in C<sub>4</sub> plants may be different to those in C<sub>3</sub> plants.

## 5.2. Other targets and opportunities to enhance photosynthesis

For C<sub>4</sub> species, non-CBB cycle targets have been suggested or predicted to improve C<sub>4</sub> photosynthesis and increase yield under non-stress and stress conditions, including chloroplast electron transport, transpirational water loss, bundle-sheath leakiness, C<sub>4</sub> cycle enzyme activity, and transport/diffusion of metabolites between mesophyll and BS cells [58,68,120,124,156,157]. While enhancing CAM has not been a research focus, potential non-CBB targets for engineering CAM in C<sub>3</sub> plants to improve photosynthesis have been identified [20,158–163] and include genes encoding proteins involved in the C<sub>3</sub>-to-CAM transition, stomatal regulation, and PWUE. In addition, the genetic variation in wild relatives of modern-day C<sub>4</sub> and CAM crop species is largely an untapped resource and offers potential novel strategies to overcome limitations in productivity [124,158,160,164–166].

**Table 1**

The impact of manipulating ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) expression on CO<sub>2</sub> assimilation, biomass, and yield in C<sub>3</sub> and C<sub>4</sub> plants. Nuclear located transgenes introduced under the control of photosynthetic tissue specific promoters, or a constitutive promoter are specified. Growth conditions are indicated: Controlled Environment \*, Greenhouse \*\*. SSU, small subunit; LSU, large subunit; RAF1, Rubisco activation factor 1.

Plant Type	Plant	Transgene (s) Expressed		Functional Description	Biomass and Yield	Ref
C <sub>4</sub>	<i>Flaveria bidentis</i>	Anti-sense Rubisco SSU	-	-	Constitutive expression. Decrease in CO <sub>2</sub> assimilation.	Decrease in plant growth rates and biomass in plants with < 20% wild type activity. [129, 130] **
	Maize	Maize SSU	maize	maize	Constitutive expression. Significant increase in CO <sub>2</sub> assimilation.	Increase in plant growth and biomass. [131] **
C <sub>3</sub>	Tobacco	<i>F. pringlei</i> or <i>F. floridana</i> or <i>F. bidentis</i> LSU	-	-	Plastome transformation. Decrease in CO <sub>2</sub> assimilation.	Decrease in growth in air. [51] *
	Rice	Sorghum SSU	-	-	Tissue-specific expression. Partial replacement of rice SSU. No change in CO <sub>2</sub> assimilation. C <sub>4</sub> Rubisco kinetic traits.	Decreases in non-productive tillers. [50] *
	Rice	Sorghum SSU	-	-	CRISPR/Cas9. Complete replacement of rice SSU. Increases in CO <sub>2</sub> assimilation at ambient CO <sub>2</sub> . C <sub>4</sub> Rubisco kinetic traits.	Decreases in total dry weight at ambient CO <sub>2</sub> . [52] **

**Table 2**

The impact of the overexpression of sedoheptulose 1,7-bisphosphatase (SB) on plant growth in C<sub>3</sub> and C<sub>4</sub> plants. Transgenes were either under the control of photosynthetic tissue specific promoters or a constitutive promoter. Growth conditions are indicated: Controlled environment \*, Greenhouse \*\*, Field experiments \*\*\*.

Plant Type	Plant	Functional Description	Biomass and Yield	Ref
C <sub>3</sub>	Arabidopsis	Tissue specific expression. 37–85% increase in SB activity, 37% increase in CO <sub>2</sub> assimilation.	42% increase in dry weight and a 53% increase in seed yield.	[148]*
	Tobacco	Constitutive expression. 90–110% average increase in SB activity, Increase in photosynthetic rates. Increases in sucrose and starch content.	30–34% increase in dry weight.	[149, 150]**
	Tomato	Constitutive expression. 55–139% increase in SB activity. An approximately 25% increase in CO <sub>2</sub> assimilation. Increases in sucrose and starch content.	Up to 39% increase in dry weight in best lines.	[153]**
	Wheat	Constitutive expression. Up to 90% increase in SB activities in some lines. Increase in CO <sub>2</sub> assimilation.	Up to 40% increase in grain yield.	[151]**
	Rice	Constitutive expression. Up to 200% increase in SB activities in some lines. Increased CO <sub>2</sub> assimilation rates under elevated temperature.	Higher growth rates under elevated temperature.	[154]**
C <sub>4</sub>	<i>Setaria viridis</i>	Bundle-sheath cell-preferential promoter. Transgenic plants with 1.5- to 3.2-times higher SB content. No correlation between SB content and saturating rates of CO <sub>2</sub> assimilation.	No increase in development or biomass was reported.	[155]*

## 6. Conclusions and future prospects

It has been well over half a century since the reactions making up the CBB cycle and the C<sub>4</sub> and CAM photosynthetic pathways were elucidated [1,11,167]. Nevertheless, there are significant gaps in our knowledge of how the ancestral C<sub>3</sub> biochemistry became integrated and operates in C<sub>4</sub> and CAM syndromes and these limitations restrict our abilities to improve plant photosynthetic performance, biomass, and yield.

Experiments to increase our understanding of CBB cycle operation more fully in CAM plants include instantaneous online discrimination measurements to ascertain the temporal interplay between the two carboxylases across the 24-h light/dark cycle. These studies should include a diverse range of CAM types representing independent CAM origins as well as facultative CAM species under well-watered and drought-stressed conditions. Similar species surveys are also needed to fill the knowledge gap regarding PNUE in CAM species. For C<sub>4</sub> plants, resolution of the presence and operation of mixed decarboxylation pathways and/or a 3-PGA/TP shuttle in individual species will contribute to attempts to enhance C<sub>4</sub> photosynthetic efficiency. The future work needs to include a focus on the integration of the CBB cycle with these pathways in species from diverse C<sub>4</sub> monocot and eudicot lineages. For both C<sub>4</sub> and CAM species, the kinetics and regulation of

CBB cycle enzymes reflect significant gaps in our knowledge.

The abundance and regulation of transcripts encoding CBB cycle enzymes have not been a focus of most transcriptome studies of C<sub>4</sub> and CAM photosynthetic tissues; consequently, the level of information reported in the scientific literature for these transcripts is variable. In addition, metabolite profiles and proteome datasets have been generated for only a very limited number of C<sub>4</sub> and CAM species. These information gaps highlight opportunities to not only re-examine and further mine existing datasets, but to also expand omics technologies to additional CAM types and C<sub>4</sub> decarboxylation subtypes from diverse evolutionary lineages, including more taxa with closely related congeners that use different photosynthetic pathways and wild relatives of crop plants. The resulting knowledge will increase our molecular toolbox, aiding attempts to bioengineer improvements in photosynthetic efficiency and crop yields.

A multi-target approach will likely be needed to enhance photosynthesis in crop plants. Generating bespoke C<sub>4</sub>, CAM, and C<sub>3</sub> crops for growth conditions and environments of the future to meet global food and fodder, fiber, and fuel demands will undoubtedly involve manipulating the expression of a combination of enzymes and pathways using both modern breeding strategies and targeted genetic manipulation.

## Role of funding sources

The funding sources had no role in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the article for publication.

## Declaration of Competing Interest

As corresponding author of the manuscript entitled, “**The Calvin-Benson-Bassham cycle in C<sub>4</sub> and Crassulacean acid metabolism species**”, I submit this declaration of interest on behalf of all the authors: Martha Ludwig, James Hartwell, Christine A Raines, and Andrew J Simkin. For all the authors, there are no declarations of interest.

## Acknowledgements

ML was supported by the Australian Research Council, (DP180102747). JH was supported by the Biotechnology and Biological Sciences Research Council, UK (BBSRC grant no. BB/F009313/1), the US Department of Energy (DOE) Office of Science, Genomic Science Program under Award Number DE-SC0008834, and a Newton Advanced Fellowship funded by the Royal Society, UK (grant no. NA140007 awarded to Prof. Luciano Freschi and JH). The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of DOE. CAR was supported by the European Union’s Horizon 2020 research and innovation programme (No. 862201) project CAPITALISE. AJS was supported by the Innovate UK Growing Kent and Medway Program, UK (Ref 107139).

## References

- [1] T.D. Sharkey, Discovery of the canonical Calvin-Benson cycle, *Photosynth. Res.* 140 (2) (2019) 235–252.
- [2] R.C. Leegood, 2 - Enzymes of the Calvin cycle, in: P.J. Lea (Ed.), *Methods in Plant Biochemistry* vol. 3, Academic Press, 1990, pp. 15–37.
- [3] C.A. Raines, The Calvin cycle revisited, *Photosynth. Res.* 75 (1) (2003) 1–10.
- [4] A.R. Portis Jr., M.A. Parry, Discoveries in Rubisco (ribulose 1,5-bisphosphate carboxylase/oxygenase): a historical perspective, *Photosynth. Res.* 94 (1) (2007) 121–143.
- [5] T.D. Sharkey, Estimating the rate of photorespiration in leaves, *Physiol. Plant.* 73 (1) (1988) 147–152.
- [6] F.A. Busch, Current methods for estimating the rate of photorespiration in leaves, *Plant Biol.* 15 (4) (2013) 648–655.
- [7] C. Peterhansel, V.G. Maurino, Photorespiration redesigned, *Plant Physiol.* 155 (1) (2011) 49–55.
- [8] S.P. Long, X.G. Zhu, S.L. Naidu, D.R. Ort, Can improvement in photosynthesis increase crop yields? *Plant Cell Environ.* 29 (3) (2006) 315–330.

- [9] M.J. West-Eberhard, J.A. Smith, K. Winter, Plant science. Photosynthesis, reorganized, *Science* 332 (6027) (2011) 311–312.
- [10] S.C. Wong, Elevated atmospheric partial pressure of CO<sub>2</sub> and plant growth: I. Interactions of nitrogen nutrition and photosynthetic capacity in C<sub>3</sub> and C<sub>4</sub> plants, *Oecologia* 44 (1) (1979) 68–74.
- [11] R.F. Sage, A portrait of the C<sub>4</sub> photosynthetic family on the 50th anniversary of its discovery: species number, evolutionary lineages, and Hall of Fame, *J. Exp. Bot.* 67 (14) (2016) 4039–4056.
- [12] R.M. Sharpe, S. Offermann, One decade after the discovery of single-cell C<sub>4</sub> species in terrestrial plants: what did we learn about the minimal requirements of C<sub>4</sub> photosynthesis? *Photosynth Res* 119 (1–2) (2014) 169–180.
- [13] S.C. Huber, T.C. Hall, G.E. Edwards, Differential localization of fraction I protein between chloroplast types, *Plant Physiol.* 57 (5) (1976) 730–733.
- [14] M.D. Hatch, C<sub>4</sub> photosynthesis: a unique blend of modified biochemistry, anatomy and ultrastructure, *Biochim. Et. Biophys. Acta (BBA) - Rev. Bioenerg.* 895 (2) (1987) 81–106.
- [15] J.E. Lunn, R.T. Furbank, Localisation of sucrose-phosphate synthase and starch in leaves of C<sub>4</sub> plants, *Planta* 202 (1) (1997) 106–111.
- [16] J.E. Lunn, R.T. Furbank, Sucrose biosynthesis in C<sub>4</sub> plants. *N. Phytol.* 143 (2) (1999) 221–237.
- [17] R.T. Furbank, S. Kelly, Finding the C<sub>4</sub> sweet spot: cellular compartmentation of carbohydrate metabolism in C<sub>4</sub> photosynthesis, *J. Exp. Bot.* 72 (17) (2021) 6018–6026.
- [18] R.T. Furbank, Evolution of the C<sub>4</sub> photosynthetic mechanism: are there really three C<sub>4</sub> acid decarboxylation types? *J. Exp. Bot.* 62 (9) (2011) 3103–3108.
- [19] Y. Wang, A. Bräutigam, A.P.M. Weber, X.-G. Zhu, Three distinct biochemical subtypes of C<sub>4</sub> photosynthesis? A modelling analysis, *J. Exp. Bot.* 65 (13) (2014) 3567–3578.
- [20] A.M. Borland, H. Griffiths, J. Hartwell, J.A. Smith, Exploiting the potential of plants with crassulacean acid metabolism for bioenergy production on marginal lands, *J. Exp. Bot.* 60 (10) (2009) 2879–2896.
- [21] C.B. Osmond, Crassulacean acid metabolism: a curiosity in context, *Annu. Rev. Plant Physiol.* 29 (1) (1978) 379–414.
- [22] K. Winter, M. Garcia, A. Virgo, J.A.C. Smith, Low-level CAM photosynthesis in a succulent-leaved member of the Urticaceae, *Pilea peperomioides*, *Funct. Plant Biol.* 48 (7) (2021) 683–690.
- [23] K. Winter, Ecophysiology of constitutive and facultative CAM photosynthesis, *J. Exp. Bot.* 70 (22) (2019) 6495–6508.
- [24] J. Males, H. Griffiths, Stomatal Biology of CAM Plants, *Plant Physiol.* 174 (2) (2017) 550–560.
- [25] U. Lüttge, Photorespiration in Phase III of Crassulacean Acid Metabolism: Evolutionary and Ecophysiological Implications, in: U.E. Lüttge, W. Beyschlag, B. Büdel, D. Francis (Eds.), *Progress in Botany* 72, Springer Berlin Heidelberg, Berlin, Heidelberg, 2011, pp. 371–384.
- [26] M. Spalding, D. Stumpf, M. Ku, R. Burris, G. Edwards, Crassulacean Acid Metabolism and Diurnal Variations of Internal CO<sub>2</sub> and O<sub>2</sub> Concentrations in *Sedum praealtum* </I> DC, *Funct. Plant Biol.* 6 (4) (1979) 557–567.
- [27] J. Ehleringer, R.W. Pearcy, Variation in quantum yield for CO<sub>2</sub> uptake among C<sub>3</sub> and C<sub>4</sub> plants, *Plant Physiol.* 73 (3) (1983) 555–559.
- [28] J.B. Skillman, Quantum yield variation across the three pathways of photosynthesis: not yet out of the dark, *J. Exp. Bot.* 59 (7) (2008) 1647–1661.
- [29] S.P. Long, Environmental responses, in: R.F. Sage, R. Monson (Eds.), *C<sub>4</sub> Plant Biology*, Academic, San Diego, CA, 1999, pp. 215–250.
- [30] P.N. Pereira, J.C. Cushman, Exploring the relationship between crassulacean acid metabolism (CAM) and mineral nutrition with a special focus on nitrogen, *Int. J. Mol. Sci.* 20 (18) (2019) 4363.
- [31] K. Winter, Crassulacean acid metabolism, in: J. Barber, N.R. Baker (Eds.), *Photosynthetic Mechanisms and the Environment*, Elsevier, Amsterdam, The Netherlands, 1985, pp. 329–387.
- [32] U. Lüttge, Ecophysiology of crassulacean acid metabolism (CAM), *Ann. Bot.* 93 (6) (2004) 629–652.
- [33] O. Ghannoum, J.R. Evans, S. von Caemmerer, Chapter 8 Nitrogen and water use efficiency of C<sub>4</sub> plants, in: A.S. Raghavendra, R.F. Sage (Eds.), *C<sub>4</sub> Photosynthesis and Related CO<sub>2</sub> Concentrating Mechanisms*, Springer Netherlands, Dordrecht, 2011, pp. 129–146.
- [34] S.C. Wong, I.R. Cowan, G.D. Farquhar, Leaf conductance in relation to rate of CO<sub>2</sub> assimilation: I. Influence of nitrogen nutrition, phosphorus nutrition, photon flux density, and ambient partial pressure of CO<sub>2</sub> during Ontogeny, *Plant Physiol.* 78 (4) (1985) 821–825.
- [35] C.P. Osborne, L. Sack, Evolution of C<sub>4</sub> plants: a new hypothesis for an interaction of CO<sub>2</sub> and water relations mediated by plant hydraulics, *Philos. Trans. R. Soc. Lond. B* 367 (1588) (2012) 583–600.
- [36] S.F. Boxall, N. Kadu, L.V. Dever, J. Kneřová, J.L. Waller, P.J.D. Gould, J. Hartwell, Kalanchoë PPC1 Is essential for crassulacean acid metabolism and the regulation of core circadian clock and guard cell signaling genes, *Plant Cell* 32 (4) (2020) 1136–1160.
- [37] C.R. Slack, M.D. Hatch, D.J. Goodchild, Distribution of enzymes in mesophyll and parenchyma-sheath chloroplasts of maize leaves in relation to the C<sub>4</sub>-dicarboxylic acid pathway of photosynthesis, *Biochem J.* 114 (3) (1969) 489–498.
- [38] M.D. Hatch, T. Kagawa, Enzymes and functional capacities of mesophyll chloroplasts from plants with C<sub>4</sub>-pathway photosynthesis, *Arch. Biochem. Biophys.* 159 (2) (1973) 842–853.
- [39] S.B. Ku, G.E. Edwards, Photosynthesis in mesophyll protoplasts and bundle sheath cells of various types of C<sub>4</sub> plants IV. Enzymes of respiratory metabolism and energy utilizing enzymes of photosynthetic pathways, *Z. für Pflanzenphysiol.* 77 (1) (1975) 16–32.
- [40] S. Aubry, N.J. Brown, J.M. Hibberd, The role of proteins in C<sub>3</sub> plants prior to their recruitment into the C<sub>4</sub> pathway, *J. Exp. Bot.* 62 (9) (2011) 3049–3059.
- [41] W. Majeran, Y. Cai, Q. Sun, K.J. van Wijk, Functional differentiation of bundle sheath and mesophyll maize chloroplasts determined by comparative proteomics, *Plant Cell* 17 (11) (2005) 3111–3140.
- [42] G. Friso, W. Majeran, M. Huang, Q. Sun, K.J. van Wijk, Reconstruction of metabolic pathways, protein expression, and homeostasis machineries across maize bundle sheath and mesophyll chloroplasts: Large-scale quantitative proteomics using the first maize genome Assembly, *Plant Physiol.* 152 (3) (2010) 1219–1250.
- [43] T.J. Erb, J. Zarzycki, A short history of RubisCO: the rise and fall (?) of Nature's predominant CO<sub>2</sub> fixing enzyme, *Curr. Opin. Biotechnol.* 49 (2018) 100–107.
- [44] T.D. Sharkey, The discovery of rubisco, *J. Exp. Bot.* 74 (2) (2022) 510–519.
- [45] C. Bathellier, G. Tcherkez, G.H. Lorimer, G.D. Farquhar, Rubisco is not really so bad, *Plant, Cell Environ.* 41 (4) (2018) 705–716.
- [46] A. Bracher, S.M. Whitney, F.U. Hartl, M. Hayer-Hartl, Biogenesis and Metabolic Maintenance of Rubisco, *Annu. Rev. Plant Biol.* 68 (2017) 29–60.
- [47] M.S.B. Ku, M.R. Schmitt, G.E. Edwards, Quantitative determination of RuBP carboxylase-oxygenase protein in leaves of several C<sub>3</sub> and C<sub>4</sub> plants, *J. Exp. Bot.* 30 (1979) 89–98.
- [48] C. Iñiguez, S. Capó-Bauçà, Ü. Niinemets, H. Stoll, P. Aguiló-Nicolau, J. Galmés, Evolutionary trends in RuBisCO kinetics and their co-evolution with CO<sub>2</sub> concentrating mechanisms, *Plant J.: Cell Mol. Biol.* 101 (4) (2020) 897–918.
- [49] W.A. Iqbal, A. Lisitsa, M.V. Kapralov, Predicting plant Rubisco kinetics from RbcL sequence data using machine learning, *J. Exp. Bot.* 74 (2) (2023) 638–650.
- [50] C. Ishikawa, T. Hatanaka, S. Misoo, C. Miyake, H. Fukayama, Functional incorporation of sorghum small subunit increases the catalytic turnover rate of Rubisco in transgenic rice, *Plant Physiol.* 156 (3) (2011) 1603–1611.
- [51] S.M. Whitney, R.E. Sharwood, D. Orr, S.J. White, H. Alonso, J. Galmés, Isoleucine 309 acts as a C<sub>4</sub> catalytic switch that increases ribulose-1,5-bisphosphate carboxylase/oxygenase (rubisco) carboxylation rate in *Flaveria*, *Proc. Natl. Acad. Sci.* 108 (35) (2011) 14688–14693.
- [52] H. Matsumura, K. Shiomi, A. Yamamoto, Y. Taketani, N. Kobayashi, T. Yoshizawa, S.-i Tanaka, H. Yoshikawa, M. Endo, H. Fukayama, Hybrid Rubisco with complete replacement of rice Rubisco small subunits by Sorghum counterparts confers C<sub>4</sub> plant-like high catalytic activity, *Mol. Plant* 13 (11) (2020) 1570–1581.
- [53] H. Usuda, G.E. Edwards, Localization of glycerate kinase and some enzymes for sucrose synthesis in C<sub>3</sub> and C<sub>4</sub> plants, *Plant Physiol.* 65 (5) (1980) 1017–1022.
- [54] K. Meierhoff, P. Westhoff, Differential biogenesis of photosystem II in mesophyll and bundle-sheath cells of monocotyledonous NADP-malic enzyme-type C<sub>4</sub> plants: the non-stoichiometric abundance of the subunits of photosystem II in the bundle-sheath chloroplasts and the translational activity of the plastome-encoded genes, *Planta* 191 (1) (1993) 23–33.
- [55] W. Majeran, G. Friso, L. Ponnala, B. Connolly, M. Huang, E. Reidel, C. Zhang, Y. Asakura, N.H. Bhuiyan, Q. Sun, et al., Structural and metabolic transitions of C<sub>4</sub> leaf development and differentiation defined by microscopy and quantitative proteomics in maize, *Plant Cell* 22 (11) (2010) 3509–3542.
- [56] P. Gao, P. Wang, B. Du, P. Li, B.-H. Kang, Accelerated remodeling of the mesophyll-bundle sheath interface in the maize C<sub>4</sub> cycle mutant leaves, *Sci. Rep.* 12 (1) (2022) 5057.
- [57] E. Omoto, M. Kawasaki, M. Taniguchi, H. Miyake, Salinity induces granal development in bundle sheath chloroplasts of NADP-malic enzyme type C<sub>4</sub> plants, *Plant Prod. Sci.* 12 (2) (2009) 199–207.
- [58] M. Ermakova, C. Bellasio, D. Fitzpatrick, R.T. Furbank, F. Mamedov, S. von Caemmerer, Upregulation of bundle sheath electron transport capacity under limiting light in C<sub>4</sub> *Setaria viridis*, *Plant J.: Cell Mol. Biol.* 106 (5) (2021) 1443–1454.
- [59] F.R. Danila, W.P. Quick, R.G. White, S. von Caemmerer, R.T. Furbank, Response of plasmodesmata formation in leaves of C<sub>4</sub> grasses to growth irradiance, *Plant Cell Environ.* 42 (8) (2019) 2482–2494.
- [60] W.M. Laetsch, I. Price, Development of the dimorphic chloroplasts of sugar cane, *Am. J. Bot.* 56 (1) (1969) 77–87.
- [61] C.R.G. Sales, R.V. Ribeiro, A.H. Hayashi, P.E.R. Marchiori, K.I. Silva, M. O. Martins, J.A.G. Silveira, N.M. Silveira, E.C. Machado, Flexibility of C<sub>4</sub> decarboxylation and photosynthetic plasticity in sugarcane plants under shading, *Environ. Exp. Bot.* 149 (2018) 34–42.
- [62] K.S. Andersen, J.M. Bain, D.G. Bishop, R.M. Smillie, Photosystem II activity in agranal bundle sheath chloroplasts from *Zea mays*, *Plant Physiol.* 49 (4) (1972) 461–466.
- [63] W.J.S. Downton, N.A. Pylotiis, Loss of photosystem II during ontogeny of sorghum bundle sheath chloroplasts, *Can. J. Bot.* 49 (1) (1971) 179–180.
- [64] Edwards G.E., Voznesenskaya E.V.: C<sub>4</sub> photosynthesis: Kranz forms and single-cell C<sub>4</sub> in terrestrial plants. In: *C<sub>4</sub> Photosynthesis and Related CO<sub>2</sub> Concentrating Mechanisms*. Edited by (eds.) IASRaRFS; 2011: 29–61.
- [65] M. Stitt, G. Luca Borghi, S. Arrivault, Targeted metabolite profiling as a top-down approach to uncover interspecies diversity and identify key conserved operational features in the Calvin-Benson cycle, *J. Exp. Bot.* 72 (17) (2021) 5961–5986.
- [66] R.C. Leegood, R.P. Walker, 4 - regulation of the C<sub>4</sub> pathway, in: R.F. Sage, R. K. Monson (Eds.), *C<sub>4</sub> Plant Biology*, Academic Press, San Diego, 1999, pp. 89–131.
- [67] S.C. Huber, G.E. Edwards, C<sub>4</sub> Photosynthesis: Light-dependent CO<sub>2</sub> fixation by mesophyll cells, protoplasts, and protoplast extracts of *Digitaria sanguinalis*, *Plant Physiol.* 55 (5) (1975) 835–844.
- [68] S. von Caemmerer, R.T. Furbank, Strategies for improving C<sub>4</sub> photosynthesis, *Curr. Opin. Plant Biol.* 31 (2016) 125–134.

- [69] M. Stitt, X.G. Zhu, The large pools of metabolites involved in intercellular metabolite shuttles in C<sub>4</sub> photosynthesis provide enormous flexibility and robustness in a fluctuating light environment, *Plant Cell Environ.* 37 (9) (2014) 1985–1988.
- [70] M. Sommer, A. Bräutigam, A.P. Weber, The dicotyledonous NAD malic enzyme C<sub>4</sub> plant *Cleome gynandra* displays age-dependent plasticity of C<sub>4</sub> decarboxylation biochemistry, *Plant Biol.* 14 (4) (2012) 621–629.
- [71] T.R. Pick, A. Bräutigam, U. Schlüter, A.K. Denton, C. Colmsee, U. Scholz, H. Fahnenstich, R. Pieruschka, U. Rascher, U. Sonnewald, et al., Systems analysis of a maize leaf developmental gradient redefines the current C<sub>4</sub> model and provides candidates for regulation, *Plant Cell* 23 (12) (2011) 4208–4220.
- [72] V.K. Gupta, L.E. Anderson, Light modulation of the activity of carbon metabolism enzymes in the crassulacean acid metabolism plant *Kalanchoë*, *Plant Physiol.* 61 (3) (1978) 469–471.
- [73] S.W. Hutcheson, B.B. Buchanan, Enzyme regulation in crassulacean acid metabolism photosynthesis. Studies on the ferredoxin/thioredoxin system of *Kalanchoë daigremontiana*, *Plant Physiol.* 72 (3) (1983) 870–876.
- [74] J.A. Cruz, C. Emery, M. Wüst, D.M. Kramer, B.M. Lange, Metabolite profiling of Calvin cycle intermediates by HPLC-MS using mixed-mode stationary phases, *Plant J.* 55 (6) (2008) 1047–1060.
- [75] K. Maxwell, Resistance is useful: diurnal patterns of photosynthesis in C<sub>3</sub> and crassulacean acid metabolism epiphytic bromeliads, *Funct. Plant Biol.* 29 (6) (2002) 679–687.
- [76] H. Griffiths, B. Helliker, A. Roberts, R.P. Haslam, J. Girnus, W.E. Robe, A. M. Borland, K. Maxwell, Regulation of Rubisco activity in crassulacean acid metabolism plants: better late than never, *Funct. Plant Biol.* 29 (6) (2002) 689–696.
- [77] H. Griffiths, A.B. Cousins, M.R. Badger, S. von Caemmerer, Discrimination in the dark. Resolving the interplay between metabolic and physical constraints to phosphoenolpyruvate carboxylase activity during the crassulacean acid metabolism cycle, *Plant Physiol.* 143 (2) (2007) 1055–1067.
- [78] H. Griffiths, M.S.J. Broadmeadow, A.M. Borland, C.S. Hetherington, Short-term changes in carbon-isotope discrimination identify transitions between C<sub>3</sub> and C<sub>4</sub> carboxylation during Crassulacean acid metabolism, *Planta* 181 (4) (1990) 604–610.
- [79] J.S. Gillon, A.M. Borland, K.G. Harwood, A. Roberts, M.S.J. Broadmeadow, H. Griffiths, Carbon isotope discrimination in terrestrial plants: carboxylations and decarboxylations, in: H. Griffiths (Ed.), *Stable Isotopes: Integration of Biological, Ecological and Geochemical Processes*, BIOS Scientific, Oxford, 1998, pp. 111–131.
- [80] E. van Tongerlo, G. Trouwborst, S.W. Hogewoning, W. van Ieperen, J. A. Dieleman, L.F.M. Marcelis, Crassulacean acid metabolism species differ in the contribution of C<sub>3</sub> and C<sub>4</sub> carboxylation to end of day CO<sub>2</sub> fixation, *Physiol. Plant* 172 (1) (2021) 134–145.
- [81] B.N. Davies, H. Griffiths, Competing carboxylases: circadian and metabolic regulation of Rubisco in C<sub>3</sub> and CAM *Mesembryanthemum crystallinum* L, *Plant, Cell Environ.* 35 (7) (2012) 1211–1220.
- [82] P.S. Schnable, D. Ware, R.S. Fulton, J.C. Stein, F. Wei, S. Pasternak, C. Liang, J. Zhang, L. Fulton, T.A. Graves, et al., The B73 maize genome: complexity, diversity, and dynamics, *Science* 326 (5956) (2009) 1112–1115.
- [83] A.H. Paterson, J.E. Bowers, R. Bruggmann, I. Dubchak, J. Grimwood, H. Gundlach, G. Haberer, U. Hellsten, T. Mitros, A. Poliakov, et al., The Sorghum bicolor genome and the diversification of grasses, *Nature* 457 (7229) (2009) 551–556.
- [84] J. Cai, X. Liu, K. Vanneste, S. Proost, W.-C. Tsai, K.-W. Liu, L.-J. Chen, Y. He, Q. Xu, C. Bian, et al., The genome sequence of the orchid *Phalaenopsis equestris*, *Nat. Genet.* 47 (1) (2015) 65–72.
- [85] L. Yan, X. Wang, H. Liu, Y. Tian, J. Lian, R. Yang, S. Hao, X. Wang, S. Yang, Q. Li, et al., The genome of *Dendrobium officinale* illuminates the biology of the important traditional chinese orchid herb, *Mol. Plant* 8 (6) (2015) 922–934.
- [86] R. Ming, R. VanBuren, C.M. Wai, H. Tang, M.C. Schatz, J.E. Bowers, E. Lyons, M. L. Wang, J. Chen, E. Biggers, et al., The pineapple genome and the evolution of CAM photosynthesis, *Nat. Genet.* 47 (12) (2015) 1435–1442.
- [87] Y. Sun, L. Shang, Q.H. Zhu, L. Fan, L. Guo, Twenty years of plant genome sequencing: achievements and challenges, *Trends Plant Sci.* 27 (4) (2022) 391–401.
- [88] U. Schlüter, A.P.M. Weber, Regulation and Evolution of C<sub>4</sub> Photosynthesis, *Annu. Rev. Plant Biol.* 71 (2020) 183–215.
- [89] K. Heyduk, M. Hwang, V. Albert, K. Silvera, T. Lan, K. Farr, T.-H. Chang, M.-T. Chan, K. Winter, J. Leebens-Mack, Altered gene regulatory networks are associated with the transition from C<sub>3</sub> to crassulacean acid metabolism in *Erycina (Oncidiinae: Orchidaceae)*, *Front. Plant Sci.* (2019) 9.
- [90] K. Heyduk, E.V. McAssey, J. Leebens-Mack, Differential timing of gene expression and recruitment in independent origins of CAM in the Agavoideae (*Asparagaceae*), *N. Phytol.* 235 (5) (2022) 2111–2126.
- [91] R.C. Ferrari, P.P. Bittencourt, M.A. Rodrigues, J.J. Moreno-Villena, F.R.R. Alves, V.D. Gastaldi, S.F. Boxall, L.V. Dever, D. Demarco, S.C.S. Andrade, et al., C<sub>4</sub> and crassulacean acid metabolism within a single leaf: deciphering key components behind a rare photosynthetic adaptation, *N. Phytol.* 225 (4) (2020) 1699–1714.
- [92] R.C. Ferrari, A.B. Kawabata, S.S. Ferreira, J. Hartwell, L. Freschi, A matter of time: regulatory events behind the synchronization of C<sub>4</sub> and crassulacean acid metabolism in *Portulaca oleracea*, *J. Exp. Bot.* 73 (14) (2022) 4867–4885.
- [93] J.C. Cushman, R.L. Tillett, J.A. Wood, J.M. Branco, K.A. Schlauch, Large-scale mRNA expression profiling in the common ice plant, *Mesembryanthemum crystallinum*, performing C<sub>3</sub> photosynthesis and Crassulacean acid metabolism (CAM), *J. Exp. Bot.* 59 (7) (2008) 1875–1894.
- [94] E. Maleckova, D. Brilhaus, T.J. Wrobel, A.P.M. Weber, Transcript and metabolite changes during the early phase of abscisic acid-mediated induction of crassulacean acid metabolism in *Talinum triangulare*, *J. Exp. Bot.* 70 (22) (2019) 6581–6596.
- [95] P.E. Abraham, H. Yin, A.M. Borland, D. Weighill, S.D. Lim, H.C. De Paoli, N. Engle, P.C. Jones, R. Agh, D.J. Weston, et al., Transcript, protein and metabolite temporal dynamics in the CAM plant *Agave*, *Nat. Plants* vol. 2 (2016) 16178.
- [96] Brilhaus D., Bräutigam A., Mettler-Altman T., Winter K., Weber A.P.M.: Reversible burst of transcriptional changes during induction of crassulacean acid metabolism in *Talinum triangulare Plant Physiology*; 2015, 170(1):102–122.
- [97] T. Nelson, The grass leaf developmental gradient as a platform for a systems understanding of the anatomical specialization of C<sub>4</sub> leaves, *J. Exp. Bot.* 62 (9) (2011) 3039–3048.
- [98] C.G. de Oliveira Dal’Molin, C. Orellana, L. Gebbie, J. Steen, M.P. Hodson, P. Chrysanthopoulos, M.R. Plan, R. McQualter, R.W. Palfreyman, L.K. Nielsen, Metabolic Reconstruction of *Setaria italica*: A Systems Biology Approach for Integrating Tissue-Specific Omics and Pathway Analysis of Bioenergy Grasses, *Front. Plant Sci.* 7 (2016) 1138.
- [99] S. Arrivault, T. Alexandre Moraes, T. Obata, D.B. Medeiros, A.R. Fernie, A. Boulouis, M. Ludwig, J.E. Lunn, G.L. Borghi, A. Schlereth, et al., Metabolite profiles reveal interspecific variation in operation of the Calvin-Benson cycle in both C<sub>4</sub> and C<sub>3</sub> plants, *J. Exp. Bot.* 70 (6) (2019) 1843–1858.
- [100] Clapero V., Arrivault S., Stitt M.: Natural variation in metabolism of the calvin-benson cycle *Seminars in Cell and Developmental Biology*; 2023.
- [101] G.L. Borghi, S. Arrivault, M. Günther, D. Barbosa Medeiros, E. Dell’Aversana, G. M. Fusco, P. Carillo, M. Ludwig, A.R. Fernie, J.E. Lunn, et al., Metabolic profiles in C<sub>3</sub>, C<sub>3</sub>-C<sub>4</sub> intermediate, C<sub>4</sub>-like, and C<sub>4</sub> species in the genus *Flaveria*, *J. Exp. Bot.* 73 (5) (2022) 1581–1601.
- [102] R.F. Sage, T.L. Sage, F. Kocacinar, Photorespiration and the evolution of C<sub>4</sub> photosynthesis, *Annu. Rev. Plant Biol.* 63 (2012) 19–47.
- [103] U. Schlüter, A. Bräutigam, U. Gowik, M. Melzer, P.-A. Christin, S. Kurz, T. Mettler-Altman, A.P. Weber, Photosynthesis in C<sub>3</sub>-C<sub>4</sub> intermediate *Moricandia* species, *J. Exp. Bot.* 68 (2) (2017) 191–206.
- [104] U. Gowik, A. Bräutigam, K.L. Weber, A.P. Weber, P. Westhoff, Evolution of C<sub>4</sub> photosynthesis in the genus *Flaveria*: how many and which genes does it take to make C<sub>4</sub>? *Plant Cell* 23 (6) (2011) 2087–2105.
- [105] A. Bräutigam, S. Schliesky, C. Külahoglu, C.P. Osborne, A.P. Weber, Towards an integrative model of C<sub>4</sub> photosynthetic subtypes: insights from comparative transcriptome analysis of NAD-ME, NADP-ME, and PEP-CK C<sub>4</sub> species, *J. Exp. Bot.* 65 (13) (2014) 3579–3593.
- [106] A. Bräutigam, K. Kajala, J. Wullenweber, M. Sommer, D. Gagneul, K.L. Weber, K. M. Carr, U. Gowik, J. Mass, M.J. Lercher, et al., An mRNA blueprint for C<sub>4</sub> photosynthesis derived from comparative transcriptomics of closely related C<sub>3</sub> and C<sub>4</sub> species, *Plant Physiol.* 155 (1) (2011) 142–156.
- [107] R.F. Sage, P.-A. Christin, E.J. Edwards, The C<sub>4</sub> plant lineages of planet Earth, *J. Exp. Bot.* 62 (9) (2011) 3155–3169.
- [108] S. Aubry, S. Kelly, B.M.C. Kümpers, R.D. Smith-Unna, J.M. Hibberd, Deep evolutionary comparison of gene expression identifies parallel recruitment of trans-factors in two independent origins of C<sub>4</sub> photosynthesis, *PLOS Genet.* 10 (6) (2014), e1004365.
- [109] P. Li, L. Ponnala, N. Gandotra, L. Wang, Y. Si, S.L. Tausta, T.H. Kebrom, N. Provart, R. Patel, C.R. Myers, et al., The developmental dynamics of the maize leaf transcriptome, *Nat. Genet.* 42 (12) (2010) 1060–1067.
- [110] C. Külahoglu, A.K. Denton, M. Sommer, J. Maß, S. Schliesky, T.J. Wrobel, B. Berckmans, E. Gongora-Castillo, C.R. Buell, R. Simon, et al., Comparative transcriptome atlases reveal altered gene expression modules between two Cleomaceae C<sub>3</sub> and C<sub>4</sub> plant species, *Plant Cell* 26 (8) (2014) 3243–3260.
- [111] L. Mattiello, D.M. Riano-Pachón, M.C. Martins, L.P. da Cruz, D. Bassi, P. E. Marchiori, R.V. Ribeiro, M.T. Labate, C.A. Labate, M. Menossi, Physiological and transcriptional analyses of developmental stages along sugarcane leaf, *BMC Plant Biol.* 15 (2015) 300.
- [112] S. Prochetto, A.J. Studer, R. Reinheimer, De novo transcriptome assemblies of C<sub>3</sub> and C<sub>4</sub> non-model grass species reveal key differences in leaf development, *BMC Genom.* 24 (1) (2023) 64.
- [113] Z. Ding, S. Weissmann, M. Wang, B. Du, L. Huang, L. Wang, X. Tu, S. Zhong, C. Myers, T.P. Brutnell, et al., Identification of photosynthesis-associated C<sub>4</sub> candidate genes through comparative leaf gradient transcriptome in multiple lineages of C<sub>3</sub> and C<sub>4</sub> species, *PLoS One* 10 (10) (2015), e0140629.
- [114] X. Rao, N. Lu, G. Li, J. Nakashima, Y. Tang, R.A. Dixon, Comparative cell-specific transcriptomics reveals differentiation of C<sub>4</sub> photosynthesis pathways in switchgrass and other C<sub>4</sub> lineages, *J. Exp. Bot.* 67 (6) (2016) 1649–1662.
- [115] C.R. John, R.D. Smith-Unna, H. Woodfield, S. Covshoff, J.M. Hibberd, Evolutionary convergence of cell-specific gene expression in independent lineages of C<sub>4</sub> grasses, *Plant Physiol.* 165 (1) (2014) 62–75.
- [116] S.L. Tausta, P. Li, Y. Si, N. Gandotra, P. Liu, Q. Sun, T.P. Brutnell, T. Nelson, Developmental dynamics of Kranz cell transcriptional specificity in maize leaf reveals early onset of C<sub>4</sub>-related processes, *J. Exp. Bot.* 65 (13) (2014) 3543–3555.
- [117] L.V. Dever, S.F. Boxall, J. Kneřová, J. Hartwell, Transgenic perturbation of the decarboxylation phase of Crassulacean acid metabolism alters physiology and metabolism but has only a small effect on growth, *Plant Physiol.* 167 (1) (2015) 44–59.
- [118] S.F. Boxall, L.V. Dever, J. Kneřová, P.D. Gould, J. Hartwell, Phosphorylation of phosphoenolpyruvate carboxylase is essential for maximal and sustained dark

- CO<sub>2</sub> fixation and core circadian clock operation in the obligate crassulacean acid metabolism species *Kalanchoë fedtschenkoi*, *Plant Cell* 29 (10) (2017) 2519–2536.
- [119] D.K. Ray, N.D. Mueller, P.C. West, J.A. Foley, Yield trends are insufficient to double global crop production by 2050, *PLoS One* 8 (6) (2013).
- [120] C.R.G. Sales, Y. Wang, J.B. Evers, J. Kromdijk, Improving C<sub>4</sub> photosynthesis to increase productivity under optimal and suboptimal conditions, *J. Exp. Bot.* 72 (17) (2021) 5942–5960.
- [121] J. Bailey-Serres, J.E. Parker, E.A. Ainsworth, G.E.D. Oldroyd, J.I. Schroeder, Genetic strategies for improving crop yields, *Nature* 575 (7781) (2019) 109–118.
- [122] A.J. Simkin, P.E. Lopez-Calcano, C.A. Raines, Feeding the world: improving photosynthetic efficiency for sustainable crop production, *J. Exp. Bot.* 70 (4) (2019) 1119–1140.
- [123] P.E. Verslues, J. Bailey-Serres, C. Brodersen, T.N. Buckley, L. Conti, A. Christmann, J.R. Dinneny, E. Grill, S. Hayes, R.W. Heckman, et al., Burning questions for a warming and changing world: 15 unknowns in plant abiotic stress, *Plant Cell* 35 (1) (2023) 67–108.
- [124] N.A. Eckardt, E.A. Ainsworth, R.N. Bahuguna, M.R. Broadley, W. Busch, N. C. Carpita, G. Castrillo, J. Chory, L.R. DeHaan, C.M. Duarte, et al., Climate change challenges, plant science solutions, *Plant Cell* 35 (1) (2023) 24–66.
- [125] C.A. Raines, A.P. Cavanagh, A.J. Simkin, Chapter 9. Improving carbon fixation, in: A. Ruban, E. Murchie, C. Foyer (Eds.), *Photosynthesis in Action*, 1 ed., Academic Press, 2022.
- [126] C.A. Raines, Improving plant productivity by re-tuning the regeneration of RuBP in the Calvin–Benson–Bassham cycle, *N. Phytol.* 236 (2) (2022) 350–356.
- [127] A.J. Simkin, Genetic engineering for global food security: photosynthesis and biofortification, *Plants* 8 (12) (2019) 586.
- [128] T.D. Sharkey, D.A. Walker, C.B. Osmond, Evaluating the role of Rubisco regulation in photosynthesis of C<sub>3</sub> plants, *Philos. Trans. R. Soc. Lond. B, Biol. Sci.* 323 (1216) (1989) 435–448.
- [129] R.T. Furbank, J.A. Chitty, S. Von Caemmerer, C. Jenkins, Antisense RNA inhibition of RbcS gene expression reduces rubisco level and photosynthesis in the C<sub>4</sub> Plant *Flaveria bidentis*, *Plant Physiol.* 111 (3) (1996) 725–734.
- [130] S. Von Caemmerer, A. Millgate, G.D. Farquhar, R.T. Furbank, Reduction of ribulose-1,5-bisphosphate carboxylase/oxygenase by antisense RNA in the C<sub>4</sub> Plant *Flaveria bidentis* leads to reduced assimilation rates and increased carbon isotope discrimination, *Plant Physiol.* 113 (2) (1997) 469–477.
- [131] C.E. Saless-Smith, R.E. Sharwood, F.A. Busch, D.B. Stern, Increased Rubisco content in maize mitigates chilling stress and speeds recovery, *Plant Biotechnol. J.* 18 (6) (2020) 1409–1420.
- [132] C.E. Saless-Smith, R.E. Sharwood, F.A. Busch, J. Kromdijk, V. Bardal, D.B. Stern, Overexpression of Rubisco subunits with RAF1 increases Rubisco content in maize, *Nat. Plants* 4 (10) (2018) 802–810.
- [133] S.P. Long, A.K. Spence, Toward cool C<sub>4</sub> crops, *Annu. Rev. Plant Biol.* 64 (1) (2013) 701–722.
- [134] L. Doron, L. Xu, S. Rachmilevitch, D.B. Stern, Transgenic overexpression of rubisco subunits and the assembly factor RAF1 are beneficial to recovery from drought stress in maize, *Environ. Exp. Bot.* 177 (2020), 104126.
- [135] A.H. Kingston-Smith, J. Harbinson, J. Williams, C.H. Foyer, Effect of chilling on carbon assimilation, enzyme activation, and photosynthetic electron transport in the absence of photoinhibition in Maize leaves, *Plant Physiol.* 114 (3) (1997) 1039–1046.
- [136] J. Pittermann, R.F. Sage, Photosynthetic performance at low temperature of *Bouteloua gracilis* Lag., a high-altitude C<sub>4</sub> grass from the Rocky Mountains, USA, *Plant, Cell Environ.* 23 (8) (2000) 811–823.
- [137] J. Pittermann, R.F. Sage, The response of the high altitude C<sub>4</sub> grass *Muhlenbergia montana* (Nutt.) A.S. Hitchc. to long- and short-term chilling, *J. Exp. Bot.* 52 (357) (2001) 829–838.
- [138] D.S. Kubien, S. von Caemmerer, R.T. Furbank, R.F. Sage, C<sub>4</sub> photosynthesis at low temperature. A study using transgenic plants with reduced amounts of Rubisco, *Plant Physiol.* 132 (3) (2003) 1577–1585.
- [139] S.L. Naidu, S.P. Long, Potential mechanisms of low-temperature tolerance of C<sub>4</sub> photosynthesis in *Miscanthus x giganteus*: an in vivo analysis, *Planta* 220 (1) (2004) 145–155.
- [140] D. Wang, S.L. Naidu, A.R. Portis Jr., S.P. Moose, S.P. Long, Can the cold tolerance of C<sub>4</sub> photosynthesis in *Miscanthus x giganteus* relative to *Zea mays* be explained by differences in activities and thermal properties of Rubisco? *J. Exp. Bot.* 59 (7) (2008) 1779–1787.
- [141] D. Wang, A.R. Portis Jr., S.P. Moose, S.P. Long, Cool C<sub>4</sub> photosynthesis: pyruvate Pi dikinase expression and activity corresponds to the exceptional cold tolerance of carbon assimilation in *Miscanthus x giganteus*, *Plant Physiol.* 148 (1) (2008) 557–567.
- [142] T.W. Becker, H.P. Fock, Effects of water stress on the gas exchange, the activities of some enzymes of carbon and nitrogen metabolism, and on the pool sizes of some organic acids in maize leaves, *Photosynth. Res.* 8 (2) (1986) 175–181.
- [143] A. Lal, G. Edwards, Analysis of inhibition of photosynthesis under water stress in the C<sub>4</sub> species *Amaranthus cruentus* and *Zea mays*: Electron transport, CO<sub>2</sub> fixation and carboxylation capacity, *Funct. Plant Biol.* 23 (4) (1996) 403–412.
- [144] E.P. Harrison, N.M. Willingham, J.C. Lloyd, C.A. Raines, Reduced sedoheptulose-1,7-bisphosphatase levels in transgenic tobacco lead to decreased photosynthetic capacity and altered carbohydrate accumulation, *Planta* 204 (1) (1998) 27–36.
- [145] E.P. Harrison, H. Olcer, J.C. Lloyd, S.P. Long, C.A. Raines, Small decreases in SBPase cause a linear decline in the apparent RuBP regeneration rate, but do not affect Rubisco carboxylation capacity, *J. Exp. Bot.* 52 (362) (2001) 1779–1784.
- [146] H. Olcer, J.C. Lloyd, C.A. Raines, Photosynthetic capacity is differentially affected by reductions in sedoheptulose-1,7-bisphosphatase activity during leaf development in transgenic tobacco plants, *Plant Physiol.* 125 (2) (2001) 982–989.
- [147] T. Lawson, B. Bryant, S. Lefebvre, J.C. Lloyd, C.A. Raines, Decreased SBPase activity alters growth and development in transgenic tobacco plants, *Plant Cell Environ.* 29 (1) (2006) 48–58.
- [148] A.J. Simkin, P.E. Lopez-Calcano, P.A. Davey, L.R. Headland, T. Lawson, S. Timm, H. Bauwe, C.A. Raines, Simultaneous stimulation of sedoheptulose 1,7-bisphosphatase, fructose 1,6-bisphosphate aldolase and the photorespiratory glycine decarboxylase-H protein increases CO<sub>2</sub> assimilation, vegetative biomass and seed yield in Arabidopsis, *Plant Biotechnol. J.* 15 (7) (2017) 805–816.
- [149] A.J. Simkin, L. McAusland, L.R. Headland, T. Lawson, C.A. Raines, Multigene manipulation of photosynthetic carbon assimilation increases CO<sub>2</sub> fixation and biomass yield in tobacco, *J. Exp. Bot.* 66 (13) (2015) 4075–4090.
- [150] S. Lefebvre, T. Lawson, M. Fryer, O.V. Zakhleniuk, J.C. Lloyd, C.A. Raines, Increased sedoheptulose-1,7-bisphosphatase activity in transgenic tobacco plants stimulates photosynthesis and growth from an early stage in development, *Plant Physiol.* 138 (1) (2005) 451–460.
- [151] S.M. Driever, A.J. Simkin, S. Alotaibi, S.J. Fisk, P.J. Madgwick, C.A. Sparks, H. D. Jones, T. Lawson, M.A.J. Parry, C.A. Raines, Increased SBPase activity improves photosynthesis and grain yield in wheat grown in greenhouse conditions, *Philos. Trans. R. Soc. B* 372 (2017) 1730.
- [152] F. Ding, M. Wang, S. Zhang, X. Ai, Changes in SBPase activity influence photosynthetic capacity, growth, and tolerance to chilling stress in transgenic tomato plants, *Sci. Rep.* 6 (2016) 32741.
- [153] F. Ding, M. Wang, S. Zhang, X. Ai, Changes in SBPase activity influence photosynthetic capacity, growth, and tolerance to chilling stress in transgenic tomato plants, *Sci. Rep.* 6 (2016) 32741.
- [154] L.L. Feng, Y.J. Han, G. Liu, B.G. An, J. Yang, G.H. Yang, Y.S. Li, Y.G. Zhu, Overexpression of sedoheptulose-1,7-bisphosphatase enhances photosynthesis and growth under salt stress in transgenic rice plants, *Funct. Plant Biol.* 34 (9) (2007) 822–834.
- [155] Ermakova M., Lopez-Calcano P.E., Furbank R.T., Raines C.A., von Caemmerer S.: Increased sedoheptulose-1,7-bisphosphatase content in *Setaria viridis* does not affect C<sub>4</sub> photosynthesis. *Plant Physiology*; 2022.
- [156] Y. Wang, S.S. Stutz, C.J. Bernacchi, R.A. Boyd, D.R. Ort, S.P. Long, Increased bundle-sheath leakiness of CO<sub>2</sub> during photosynthetic induction shows a lack of coordination between the C<sub>4</sub> and C<sub>3</sub> cycles, *N. Phytol.* 236 (5) (2022) 1661–1675.
- [157] M. Ermakova, P.E. Lopez-Calcano, C.A. Raines, R.T. Furbank, S. von Caemmerer, Overexpression of the Rieske FeS protein of the Cytochrome *b<sub>6</sub>f* complex increases C<sub>4</sub> photosynthesis in *Setaria viridis*, *Commun. Biol.* 2 (1) (2019) 314.
- [158] S.C. Davis, R. Ming, D.S. LeBauer, S.P. Long, Toward systems-level analysis of agricultural production from crassulacean acid metabolism (CAM): scaling from cell to commercial production, *N. Phytol.* 208 (1) (2015) 66–72.
- [159] A.M. Borland, S.D. Wullschlegel, D.J. Weston, J. Hartwell, G.A. Tuskan, X. Yang, J.C. Cushman, Climate-resilient agroforestry: physiological responses to climate change and engineering of crassulacean acid metabolism (CAM) as a mitigation strategy, *Plant Cell Environ.* 38 (9) (2015) 1833–1849.
- [160] A.M. Borland, J. Hartwell, D.J. Weston, K.A. Schlauch, T.J. Tschaplinski, G. A. Tuskan, X. Yang, J.C. Cushman, Engineering crassulacean acid metabolism to improve water-use efficiency, *Trends Plant Sci.* 19 (5) (2014) 327–338.
- [161] A.M. Borland, V.A. Barrera Zambrano, J. Ceusters, K. Shorrocks, The photosynthetic plasticity of crassulacean acid metabolism: an evolutionary innovation for sustainable productivity in a changing world, *N. Phytol.* 191 (3) (2011) 619–633.
- [162] S.D. Lim, S. Lee, W.-G. Choi, W.C. Yim, J.C. Cushman, Laying the foundation for crassulacean acid metabolism (CAM) biodesign: expression of the C<sub>4</sub> metabolism cycle genes of cam in Arabidopsis, *Front. Plant Sci.* (2019) 10.
- [163] X. Yang, D. Liu, T.J. Tschaplinski, G.A. Tuskan, Comparative genomics can provide new insights into the evolutionary mechanisms and gene function in CAM plants, *J. Exp. Bot.* 70 (22) (2019) 6539–6547.
- [164] J. Kromdijk, A.J. McCormick, Genetic variation in photosynthesis: many variants make light work, *J. Exp. Bot.* 73 (10) (2022) 3053–3056.
- [165] A. Bohra, B. Kilian, S. Sivasankar, M. Caccamo, C. Mba, S.R. McCouch, R. K. Varshney, Reap the crop wild relatives for breeding future crops, *Trends Biotechnol.* 40 (4) (2022) 412–431.
- [166] S.C. Davis, J. Simpson, Kd.C. Gil-Vega, N.A. Niechayev, Ev Tongerlo, N. H. Castano, L.V. Dever, A. Búrquez, Undervalued potential of crassulacean acid metabolism for current and future agricultural production, *J. Exp. Bot.* 70 (22) (2019) 6521–6537.
- [167] K.R. Hultine, J.C. Cushman, D.G. Williams, New perspectives on crassulacean acid metabolism biology, *J. Exp. Bot.* 70 (22) (2019) 6489–6493.
- [168] W. Yamori, K. Hikosaka, D.A. Way, Temperature response of photosynthesis in C<sub>3</sub>, C<sub>4</sub>, and CAM plants: temperature acclimation and temperature adaptation, *Photosynth. Res.* 119 (1–2) (2014) 101–117.
- [169] E.A. Ainsworth, Photosynthesis, in: J.L. Hatfield, M.V.K. Sivakumar, J.H. Prueger (Eds.), *Agroclimatology: Link. Agric. Clim.* vol. 60 (2020) 129–151.