

# **ORCA – Online Research @ Cardiff**

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:https://orca.cardiff.ac.uk/id/eprint/161913/

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Evans, Sam, Farnell, Damian J. J., Carson-Stevens, Andy and Kemp, Alison 2023. Survey of practices for documenting evidence of bruises from physical abuse during child protection proceedings. BMJ Paediatrics Open 7 (1), e002047. 10.1136/bmjpo-2023-002047

Publishers page: http://dx.doi.org/10.1136/bmjpo-2023-002047

## Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See http://orca.cf.ac.uk/policies.html for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# BMJ Paediatrics Open

# Survey of practices for documenting evidence of bruises from physical abuse during child protection proceedings

Sam Evans <sup>(D)</sup>, <sup>1</sup> Damian J J Farnell, <sup>1</sup> Andy Carson-Stevens, <sup>2</sup> Alison Kemp <sup>(D)</sup> <sup>3</sup>

### ABSTRACT

**To cite:** Evans S, Farnell DJJ, Carson-Stevens A, *et al.* Survey of practices for documenting evidence of bruises from physical abuse during child protection proceedings. *BMJ Paediatrics Open* 2023;**7**:e002047. doi:10.1136/ bmjpo-2023-002047

 Additional supplemental material is published online only. To view, please visit the journal online (http://dx.doi.org/ 10.1136/bmjpo-2023-002047).

Received 16 May 2023 Accepted 22 June 2023

## Check for updates

© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.


<sup>1</sup>School of Dentistry, Cardiff University, Cardiff, UK <sup>2</sup>Division of Population Medicine, School of Medicine, Cardiff University, Cardiff, UK <sup>3</sup>Department of Child Health, School of Medicine, Cardiff University, Cardiff, UK

Correspondence to Dr Sam Evans; evansst@cardiff. Accurate recording of forensically important information on bruises is vital in child protection proceedings (Royal College of Paediatrics and Child Health (RCPCH)). An online survey was distributed to the RCPCH child protection committees to assess compliance with guidance. 56 individuals were contacted by email, 47 (84%) completed the survey. Results showed that the paediatricians always or usually record size (n=41; 87%), site (n=45; 96%), shape (n=32; 68%) and colour (n=36; 77%); n=10; 22% of the paediatricians stated that they 'always' used a ranking system for likelihood of abuse; n=12; 35% of those surveyed 'sometimes' estimated the size of the bruise. Results showed that paediatric bruise reporting is inconsistent and incomplete for some fields compared with national guidance.

The robust and accurate recording of forensically important information on injuries such as bruises is vital in child protection proceedings. This is highlighted in the Child Protection Companion,<sup>1</sup> designed and developed by the Royal College of Paediatrics and Child Health (RCPCH), which specifies best practice in paediatric bruise recording in the UK.

An online survey was created and distributed between November 2018 and March 2019 to assess an overview of the practices of paediatricians in the UK. Ten questions (online supplemental appendix 1) were created to determine which information doctors carrying out child protection medicals record following history and examination, what images they take and what post hoc image analysis they undertake on injuries that may have been sustained from physical abuse. Members from the RCPCH child protection committees provide a fair representation of clinicians involved in child protection in the UK. Fifty-six individuals were contacted by email and 47 (83.9%) of them completed the survey.


The majority of those surveyed stated that they always or usually record size (n=41; 87% (46 responders)), site (n=45; 96%), shape (n=32; 68%) and colour (n=36; 77%), in



**Figure 1** Results for the question 'When taking history and visual analysis of a cutaneous injury of a child with suspected physical abuse do you record the following information?' Results are shown as a stacked bar chart.

accordance with best practice outlined in the Child Protection Companion<sup>1</sup> (figure 1). The higher percentage for colour compared with shape may indicate that bruise colour has some relevance to paediatricians in their daily practice contrary to the evidence presented in prior research.<sup>23</sup> Other indicators of abuse such as proposed implement (n=18; 39% (46 responders)), possible mechanism (n=22; 48% (46 responders)) and alleged perpetrator (n=22; 47%) were 'always' recorded by less than half of the paediatricians.

Survey data demonstrated that a quarter (n=10; 22% (45 responders)) of those surveyed 'always' used a ranking system<sup>4</sup> when offering an opinion on the likelihood of



**Figure 2** Results from question 'When taking such measurements what method do you employ?' Results are shown as a stacked bar chart.

ac.uk

abuse. A third (n=14; 30%) of respondents measured the size of all injuries at every examination, including those deemed to be accidental; n=6; 17% (35 responders) of respondents stated they always use a forensic standard (ABFO No.2) scale (figure 2) when measuring bruises. Over a third (n=12; 35% (34 responders)) of respondents 'sometimes' estimated the size of the bruise. One respondent stated that they 'sometimes' use software to measure a bruise from a digital photograph. Results showed that n=36 (77%) respondents would use user-friendly software to digitally analyse bruises; n=41; 89% (46 responders) of respondents confirmed they would use a reporting service (from a medical photography department) to digitally measure a bruise, if such a service was available.

Our results show that paediatric bruise reporting is inconsistent and incomplete for some fields with respect to recommended guidelines in the Child Protection Companion.<sup>1</sup> These results also suggest that an enhanced medical photography service would be acceptable to the vast majority of respondents. We acknowledge the hiatus between data collection and reporting these findings, although no recent publications suggest that an improvement in practice has occurred. Our findings indicate that child protection teams should critically review their existing procedures for reliably recording such evidence; future studies of ours will introduce a standard proforma to achieve this aim.

**Contributors** All four authors have provided substantial contribution to the work, including the design, acquisition, analysis and interpretation of data for the work. All four authors assisted in drafting the work, approval of the final version and are accountable for all aspects of the work.

Funding This study was funded by Cardiff University.

#### Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval REC reference: 14/WA/0070; IRAS project ID: 136073.

Provenance and peer review Not commissioned; internally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

### **ORCID** iDs

Sam Evans http://orcid.org/0000-0002-2657-7436 Alison Kemp http://orcid.org/0000-0002-1359-7948

### REFERENCES

- RCPCH. 2020 child protection evidence reviews. 2020. Available: https://www.rcpch.ac.uk/key-topics/child-protection/evidencereviews [Accessed 9 Oct 2020].
- 2 Maguire S, Mann M. Systematic reviews of bruising in relation to child abuse-what have we learnt: an overview of review updates. *Evid Based Child Health* 2013;8:255–63.
- 3 Langlois NEI. The science behind the quest to determine the age of bruises – a review of the English language literature. *Forensic Sci Med Pathol* 2007;3:241–51.
- 4 Lindberg DM, Lindsell CJ, Shapiro RA. Variability in expert assessments of child physical abuse likelihood. *Pediatrics* 2008;121:e945–53.