

Chapman University

Chapman University Digital Commons

Physical Therapy Faculty Articles and Research

Physical Therapy

6-1-2021

Using Biofeedback to Reduce Spatiotemporal Asymmetry Impairs Dynamic Balance in People Post-Stroke

Sungwoo Park
University of Southern California

Chang Liu
University of Southern California

Natalia Sánchez
Chapman University, sanchezaldana@chapman.edu

Julie K. Tilson
University of Southern California

Sara J. Mulroy
Rancho Los Amigos National Rehabilitation Center

See next page for additional authors

Follow this and additional works at: https://digitalcommons.chapman.edu/pt_articles



Part of the [Neurosciences Commons](#), and the [Physical Therapy Commons](#)

Recommended Citation

Park S, Liu C, Sánchez N, Tilson JK, Mulroy SJ, Finley JM. Using Biofeedback to Reduce Step Length Asymmetry Impairs Dynamic Balance in People Poststroke. *Neurorehabilitation and Neural Repair*. 2021;35(8):738-749. <https://doi.org/10.1177/15459683211019346>

This Article is brought to you for free and open access by the Physical Therapy at Chapman University Digital Commons. It has been accepted for inclusion in Physical Therapy Faculty Articles and Research by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Using Biofeedback to Reduce Spatiotemporal Asymmetry Impairs Dynamic Balance in People Post-Stroke

Comments

This article was originally published in *Neurorehabilitation and Neural Repair*, volume 35, issue 8, in 2021.
<https://doi.org/10.1177/1545968321101934>

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).


Copyright


The authors

Authors

Sungwoo Park, Chang Liu, Natalia Sánchez, Julie K. Tilson, Sara J. Mulroy, and James M. Finley

Using Biofeedback to Reduce Step Length Asymmetry Impairs Dynamic Balance in People Poststroke

Neurorehabilitation and
Neural Repair
2021, Vol. 35(8) 738–749
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/15459683211019346
journals.sagepub.com/home/nnr


Sungwoo Park, PhD¹, Chang Liu, PhD¹ , Natalia Sánchez, PhD¹ ,
Julie K. Tilson, MS¹, Sara J. Mulroy, PhD², and James M. Finley, PhD¹ 

Abstract

Background. People poststroke often walk with a spatiotemporally asymmetric gait, due in part to sensorimotor impairments in the paretic lower extremity. Although reducing asymmetry is a common objective of rehabilitation, the effects of improving symmetry on balance are yet to be determined. **Objective.** We established the concurrent validity of whole-body angular momentum as a measure of balance, and we determined if reducing step length asymmetry would improve balance by decreasing whole-body angular momentum. **Methods.** We performed clinical balance assessments and measured whole-body angular momentum during walking using a full-body marker set in a sample of 36 people with chronic stroke. We then used a biofeedback-based approach to modify step length asymmetry in a subset of 15 of these individuals who had marked asymmetry and we measured the resulting changes in whole-body angular momentum. **Results.** When participants walked without biofeedback, whole-body angular momentum in the sagittal and frontal plane was negatively correlated with scores on the Berg Balance Scale and Functional Gait Assessment supporting the validity of whole-body angular momentum as an objective measure of dynamic balance. We also observed that when participants walked more symmetrically, their whole-body angular momentum in the sagittal plane increased rather than decreased. **Conclusions.** Voluntary reductions of step length asymmetry in people poststroke resulted in reduced measures of dynamic balance. This is consistent with the idea that after stroke, individuals might have an implicit preference not to deviate from their natural asymmetry while walking because it could compromise their balance. Clinical Trials Number: NCT03916562.

Keywords

stroke, gait, asymmetry, balance, biofeedback, validation, angular momentum

Introduction

Neuromotor impairments associated with stroke often result in asymmetric walking patterns and reducing step length asymmetry is a common objective of rehabilitation.^{1–3} However, the effects of reducing asymmetry on the energy cost of walking poststroke are heterogeneous^{4,5} and depend on individual differences in the level of impairment.⁶ These findings demonstrate that reducing step length asymmetry does not always lead to reductions in energetic cost. One alternative advantage of reducing step length asymmetry is that it may improve balance and reduce fall risk, which is critically important poststroke.^{7,8} One recent study found that people with lower scores on the Berg Balance Scale (BBS) and greater step widths had more asymmetric step lengths and swing times during walking.⁹ However, the BBS and measures of step width provide a limited and indirect assessment of dynamic balance during walking. Therefore, it is unclear if step length asymmetry is associated with

objective measures of dynamic balance during walking or if reducing asymmetry would improve dynamic balance in either the sagittal or frontal planes.

One way to characterize dynamic balance during walking is by measuring the rotational dynamics of the body using measures of whole-body angular momentum.^{10–14} Whole-body angular momentum captures the combined effects of the rotational dynamics of the body and the relative rotations of the limbs about an axis of rotation that is commonly considered to project through the body's center of mass (CoM). People without disabilities regulate their

¹University of Southern California, Los Angeles, CA, USA

²Rancho Los Amigos National Rehabilitation Center, Downey, CA, USA

Corresponding Author:

James M. Finley, Division of Biokinesiology and Physical Therapy,
University of Southern California, 1540 East Alcazar Street, CHP 155,
Los Angeles, CA 90033, USA.
Email: jmfiney@usc.edu

balance, in part, by maintaining the peak-to-peak range of angular momentum during walking within a small range.^{10,11} In addition, angular momentum in the frontal plane has been validated as a measure of mediolateral balance in people poststroke as higher momentum is associated with lower scores on clinical balance assessments.^{13,14}

People poststroke have a higher peak-to-peak range of whole-body angular momentum in both the frontal plane¹³ and the sagittal plane¹² compared with age- and speed-matched controls. Increased frontal plane momentum may result from compensatory strategies such as lateral trunk lean, hip hiking, and circumduction.^{13,15,16} In the sagittal plane, regulation of angular momentum is achieved by the cancellation of limb angular momenta between sides.¹¹ Cancellation of angular momenta in the sagittal plane occurs when contralateral limb segments move in anti-phase. This corresponds to a kinematic pattern where the moment of peak flexion for segments on one side of the body happens at the same time as the moment of peak extension for contralateral segments and vice-versa. However, if the contralateral limb segments do not move in anti-phase, the degree of segmental angular momenta cancellation decreases, and whole-body angular momentum increases.¹⁷ Angular momentum in the frontal plane is primarily regulated by anti-phase movement of trunk and swing leg during mediolateral weight shift.¹¹ Therefore, one might expect to observe greater angular momentum during walking in stroke survivors in part because their gait asymmetries may reduce the sagittal plane momentum cancellation between the left and right limbs. Although asymmetries may increase angular momentum in the sagittal plane, it is yet to be determined if restoring symmetry could reduce angular momentum.

The strategy that people poststroke use to reduce step length asymmetry may vary widely due to heterogeneity in the level of motor impairment and in the direction and magnitude of step length asymmetry. Asymmetries in step length and step timing can result from difficulty advancing the paretic leg,^{18,19} reduced propulsion with the paretic leg,^{20,21} or both. Depending on the source of impairment, people poststroke may walk with asymmetries characterized by longer step lengths with either the paretic or nonparetic leg.^{6,22,23} People poststroke retain the capacity to reduce this step length asymmetry, but this capacity depends on the magnitude and direction of asymmetry.⁶ Therefore, the strategies used to reduce step length asymmetry are not consistent across the poststroke population. For example, individuals who have longer step lengths on the paretic side may need to advance the nonparetic limb and increase paretic stance time²⁴ to reduce step length asymmetry. In contrast, individuals with longer nonparetic steps may rely on strategies such as increasing mediolateral trunk lean,²⁵ hip hiking, or circumduction^{15,16} to advance the paretic limb to compensate for weakness of the paretic hip and knee

flexors^{23,26-28} inadequate ankle plantar flexor power generation,²⁹ or spasticity in the quadriceps or hamstrings.³⁰ What remains to be seen is if reducing step length asymmetry alters dynamic balance and whether those effects depend on the direction of each individual's asymmetry.

The primary goal of this work is to determine whether reducing step length asymmetry improves dynamic balance by decreasing whole-body angular momentum. Before addressing this goal, we first established the concurrent validity of whole-body angular momentum as a measure of balance for people poststroke. If angular momentum is a valid measure of balance during walking, the baseline angular momentum in both the sagittal and frontal planes would be inversely correlated with clinical assessments of balance and balance confidence such that a higher angular momentum would be associated with lower balance measures. We also tested the hypothesis that participants would reduce angular momentum as step length asymmetry decreases. This would be consistent with the notion that reductions in asymmetry would improve dynamic balance. We expected that participants who walked with longer nonparetic step lengths would show smaller reductions in angular momentum compared with participants who walked with longer paretic step lengths given that they would rely more on compensatory strategies to advance the paretic limb when reducing asymmetry. In the frontal plane, we hypothesized that participants would increase mediolateral peak angular momentum as step length asymmetry decreases as participants might rely on compensatory strategies such as circumduction to increase step length.

Methods

Participants

A total of 36 individuals poststroke (92 ± 84.5 months poststroke) (Table 1) participated in this study. We computed a necessary sample size of 20 from a set of pilot data where we observed a mean difference in the peak angular momentum in the sagittal plane of 0.004 during asymmetric versus symmetric walking with a standard deviation of 0.006. Inclusion criteria for this study were (1) a history of unilateral stroke more than 6 months before the study, (2) paresis confined to one side, (3) ability to provide informed consent and communicate with the investigators, and (4) ability to walk more than 5 minutes without the assistance of another individual or walking aids (eg, a cane or walker). The use of an ankle-foot orthosis was permitted during the experiment. All experimental procedures were approved by the University of Southern California Institutional Review Board, and each participant provided written, informed consent before the experiment began. All aspects of the study conformed to the principles described in the Declaration of Helsinki.

Table 1. Participants' Demographics.

Subject	Sex	Age, y	Months poststroke	Paretic side	ABC scale	Fall efficacy	Berg Balance	FGA	FM-LE	Overground speed, m/s	Direction SLA	Fall history	Treadmill speed, m/s	Baseline step length asymmetry
1	F	66	394	R	90	20	53	19	22	0.83	LNP	No	0.4	0.02
2	M	60	127	L	87.5	18	45	14	15	0.94	LP	Yes	0.55	-0.12
3	F	57	114	R	77	22	50	20	30	0.66	LP	No	0.4	-0.21
4	M	50	86	R	75	26	53	24	29	0.915	LNP	No	0.62	0.00
5	M	75	173	L	75.7	41	54	23	28	0.63	LNP	Yes	0.65	0.02
6	F	59	74	R	66	24	49	17	26	0.9	LNP	Yes	0.55	0.01
7	M	78	103	R	89	17	53	23	34	1.3	LNP	Yes	0.8	0.01
8	M	63	189	R	83	23	49	16	29	0.79	LP	No	0.38	-0.05
9	M	35	65	R	80	27	55	29	30	0.97	LNP	Yes	0.9	0.04
10	M	63	80	R	66	23	47	29	30	1.01	LP	No	0.65	-0.08
11	M	71	123	L	56	36	40	11	29	0.44	LNP	No	0.3	0.38
12	M	73	55	L	73	28	54	20	28	0.9	LP	No	0.6	-0.07
13	M	63	127	L	65	26	44	15	20	0.53	LNP	Yes	0.45	0.01
14	M	54	16	L	53	42	48	22	24	0.7	LP	Yes	0.5	-0.18
15	M	60	16	R	64	37	53	22	24	0.95	LP	No	0.5	-0.10
16	M	66	88	R	62	42	50	25	28	0.93	LNP	Yes	0.6	0.10
17	F	62	63	L	95	16	55	28	31	1.15	LNP	No	0.95	0.03
18	M	49	89	R	100	16	53	26	25	0.92	LNP	No	0.85	0.05
19	F	62	24	L	82	61	53	16	28	0.61	LP	Yes	0.35	-0.19
20	M	68	24	R	54	32	27	6	27	0.22	LNP	No	0.25	0.45
21	F	64	56	R	83.75	23	54	25	28	0.91	LNP	Yes	0.55	0.02
22	M	77	67	L	73	22	54	28	28	1.09	LNP	No	0.7	0.03
23	F	32	136	R	80.5	22	55	28	31	0.96	LP	No	0.6	-0.06
24	M	55	28	R	86	53	55	29	33	1.24	LP	No	0.75	-0.07
25	M	42	25	R	52.5	38	53	23	25	0.91	LP	Yes	0.55	-0.01
26	F	50	58	R	55.6	64	47	21	27	0.84	LP	Yes	0.6	-0.07
27	F	62	270	R	78	24	52	18	20	0.58	LNP	Yes	0.22	0.02
28	M	66	31	L	90	18	41	13	16	0.28	LNP	No	0.3	0.05
29	M	57	55	L	75	30	54	25	26	1.06	LP	Yes	0.72	-0.01
30	M	62	14	R	79	28	49	14	18	0.55	LNP	Yes	0.35	0.08
31	F	76	67	L	82.5	35	49	24	33	1.2	LNP	No	0.76	0.01
32	F	52	95	L	93	20	54	26	31	0.98	LP	No	0.65	-0.09
33	F	49	10	R	68.7	30	55	18	18	0.64	LNP	No	0.28	0.06
34	F	46	19	L	73.8	33	55	26	30	0.9	LP	Yes	0.75	-0.01
35	M	52	43	R	91	19	54	24	20	1.05	LNP	No	0.85	0.06
36	M	63	96	L	98	17	51	19	22	0.71	LP	Yes	0.45	-0.15
13 F		58 ± 11	92 ± 84.5	15 L	77 ± 13.2	29 ± 11.5	50 ± 5.5	21 ± 5.6	26 ± 4.8	0.82 ± 0.26	16 LP	17 yes	0.55 ± 0.20	

Abbreviations: F, female; M, male; L, left; R, right; ABC scale, Activities-Specific Balance Confidence Scale; FGA, Functional Gait Assessment; FM-LE, lower extremity portion of the Fugl-Meyer; SLA, step length asymmetry; LNP, longer nonparetic step length; LP, longer paretic step length.




Figure 1. Experimental protocol. We provided online visual feedback to show the position of the ankle markers during swing phase as round points and a vertical bar on each side, which represented the target step length. Black horizontal lines near the top of each bar in the Baseline asymmetry trial represent the target range that was equal to the standard deviation of the step length measured during baseline. Scores were provided on the top left or right corner of the display to encourage participants to achieve the desired step lengths.

Experimental Protocol

All participants visited the lab once and performed a 10-m walk test before the experiment to determine their self-selected walking speed. We performed the following assessments: (1) the Berg Balance Scale (BBS)^{31,32} to assess static balance, (2) the Functional Gait Assessment (FGA) to assess dynamic balance,^{32,33} (3) the Activity-Specific Balance Confidence (ABC) test to evaluate balance self-efficacy,³⁴ (4) the Fall Efficacy Scale to assess fear of falling,³⁵ and (5) the lower extremity motor domain of the Fugl-Meyer (FM) assessment to evaluate level of motor impairment.^{36,37}

After the clinical assessments, we determined the participants' self-selected walking speed on the treadmill. We started the treadmill at 70% of a participant's speed measured during the 10-m walk test, and progressively increased or decreased the speed in increments of 0.05 m/s until the participant felt that the speed was comfortable. The experiment began with three minutes of walking at their self-selected treadmill speed to determine the participant's natural step length asymmetry (*Baseline*) (Figure 1). Participants then walked with visual feedback of actual and desired step lengths for 3 different trials.^{6,38} Visual feedback was displayed on a screen in front of the participant and showed the position of reflective markers placed on the ankle during the swing phase and a vertical bar on each side, which represented the target step length.

The order of the three feedback trials, baseline asymmetry (*Baseline FB*), symmetry (*Sym*), and exaggerated asymmetry (*ExggAsym*) (Figure 1), was randomized. Each trial lasted three minutes. During *Baseline FB*, participants

walked with visual feedback of their natural step length asymmetry as this allowed us to assess the effects of walking with visual feedback on their angular momentum. During *Sym*, participants were instructed to walk such that the ankle marker reached the target step length at each foot strike, and we set the target length of the shorter step equal to the length of the longer step. During *ExggAsym*, we lengthened the target length of the longer step by adding the original difference in step lengths so that their step length asymmetry was doubled. We provided participants with scores to motivate them to achieve the target asymmetry level during each condition. The scores were calculated at every step based on the difference between the actual and target step lengths as described previously.⁶ If the difference between the actual and target step length was less than 10 mm, the participant would receive a score of 10. The scores were shown adjacent to the top of the bars at every foot-strike. All participants wore a safety harness, and no handrail support was provided during the experiment. Participants also received rest breaks of approximately 5 minutes between each trial to minimize fatigue.

Data Acquisition

Segmental kinematics were recorded using a full-body marker set placing retroreflective markers on bony landmarks.^{39,40} In addition, sets of cluster markers were placed over the upper arms, lower arms, thighs, shanks, and heels. Marker kinematics were recorded using a 10-camera Qualisys Oqus motion capture system (Qualisys AB). We first collected 5 seconds of a static trial and removed all markers on bony landmarks during the walking trials.

Data Processing

Markers were labeled using Qualisys Track Manager and then exported to Visual 3D (C-Motion) to construct a full-body model. Three-dimensional marker positions were low-pass filtered with a cutoff frequency of 6 Hz. We then created a full-body model in Visual 3D, which included 12 segments: head, trunk/pelvis, left and right upper arms, forearms, thighs, shanks, feet. Since participants wore a safety harness over their pelvis, we removed the reflective markers from the pelvis. As this prevented us from tracking the pelvis, we assumed that the pelvis and trunk were rigidly connected in our model. We validated this 12-segment model by comparing the peak-to-peak angular momentum to a 13-segment model in 5 participants without disabilities. The root-mean-square error for the peak positive and negative whole-body angular momentum during a gait cycle in the 12-segment model compared with the 13-segment model was $2.1\% \pm 1.5\%$ and $0.95\% \pm 0.70\%$ in the sagittal plane, respectively. Thus, the 12-segment model used in this study introduced relatively small errors in the whole-body angular momentum compared with 13-segment model.

Step length asymmetry (SLA) was computed as the normalized difference in step lengths based on Equation 1.

$$SLA = \frac{SL_{nonparetic} - SL_{paretic}}{SL_{nonparetic} + SL_{paretic}} \quad (1)$$

Here, $SL_{nonparetic}$ is the fore-aft distance between the heel markers at nonparetic limb foot-strike and $SL_{paretic}$ is the fore-aft distance between heel markers at paretic limb foot-strike. A positive SLA indicates that the participant takes longer nonparetic steps, while a negative SLA indicates a longer paretic step.

Marker data were also used to compute whole-body angular momentum about the CoM in the sagittal and frontal plane as measures of dynamic balance. Whole-body angular momentum (L) is the sum of the angular momenta of individual limb segments with respect to the whole-body CoM¹¹ (Equation 2).

$$\bar{L} = \frac{\sum_{i=1}^{limbs} [\bar{r}_i \times m_i \bar{v}_i + I_i \bar{\omega}_i]}{MHV} \quad (2)$$

Whole-body angular momentum was normalized by body mass (M), the CoM height (H), and self-selected walking speed (V).¹² The angular momentum of the individual limb segments in the sagittal and frontal plane was calculated as the sum of the angular momentum of each segment about a mediolateral and fore-aft axis, respectively, projecting through the body's CoM (\bar{r}_i = position of the segmental CoM, m_i = segment mass, \bar{v}_i = linear velocity of the segment relative to body's CoM) and the angular momentum of each segment about its own CoM (I_i = segmental moment of inertia, $\bar{\omega}_i$ = segmental angular velocity).

Data Analysis

We calculated whole-body angular momentum (L) during each stride in the sagittal and frontal planes. In the sagittal plane, negative values of angular momentum represented forward rotation, while positive values represented backward rotation. In the frontal plane, we specified rotation toward the paretic side as positive and rotation toward the nonparetic side as negative. A stride was defined as starting with paretic foot-strike and ending with the subsequent paretic foot-strike. Since angular momentum captures the contribution of inertia and velocity, greater angular momentum in any direction would require more effort to restore the body toward its equilibrium position. Thus, we calculated peak negative (L_{Sag}^{min} , L_{Fr}^{min}) and positive (L_{Sag}^{max} , L_{Fr}^{max}) angular momentum during each stride in the sagittal and frontal planes as our measures of dynamic balance.

Statistical Analysis

All statistical analyses were performed in MATLAB version R2018a (MathWorks). We first tested for correlations between *Baseline* peak angular momentum and each clinical balance assessment (BBS, ABC, and FGA) with the whole dataset to determine the validity of angular momentum as a measure of dynamic balance. We then used linear regression models to determine if *Baseline* angular momentum in the sagittal and frontal plane was associated with *Baseline* step length asymmetry and the direction of asymmetry.

We used mixed-effects regression models to determine if changes in step length asymmetry influenced measures of angular momentum in the sagittal or frontal plane. We used a step length asymmetry of 0.06 as the minimum asymmetry necessary to be included in this analysis based on previous studies,^{26,41} and we included 15 participants who had a step length asymmetry greater than the cutoff in the regression model. We restricted our analysis to this subset because we were only interested in how changes in asymmetry influenced angular momentum. The independent variables in this model included (1) changes in step length asymmetry magnitude ($\Delta|SLA|$) from *Baseline*, (2) whether participants took longer paretic or longer nonparetic steps ($SLA_{direction}$), and (3) level of impairment measured by overground walking speed (OGW).⁴² We also included a random intercept for each participant to account for differences in angular momentum across participants. We used the mean values of all variables for each participant during each trial in the regression models. We fit a range of models with main effects and interactions between the independent variables. The most complex potential model included $\Delta|SLA|$, $SLA_{direction}$, and OGW as main effects and interactions between $\Delta|SLA|$ and $SLA_{direction}$ and between $\Delta|SLA|$ and OGW (Equation 3).




Figure 2. Angular momentum across the gait cycle. (A) Angular momentum in the sagittal plane. Participants rotated forward most around nonparetic foot-strike (40%-60% of a gait cycle) and rotated backward most around nonparetic mid-swing (75%-85% of a gait cycle). The line represents the average across participants and the shaded area represents the standard deviation. The gait cycle starts with foot-strike on the paretic side. (B) Angular momentum in the frontal plane. Participants rotated toward the nonparetic side near nonparetic foot-strike (40%-60% of a gait cycle) and rotated toward the paretic side most near paretic foot-strike (0%-15% of a gait cycle).

$$\begin{aligned} \Delta L \sim & \Delta|SLA| + SLA_{direction} \\ & + OGW + \Delta|SLA| \times SLA_{direction} \\ & + \Delta|SLA| \times OGW + (1|Subj) \end{aligned} \quad (3)$$

The interaction between $\Delta|SLA|$ and $SLA_{direction}$ captures how differences in SLA direction influence the association between changes in SLA magnitude and angular momentum. The interaction between $\Delta|SLA|$ and OGW represents how different levels of impairment influence the association between changes in SLA magnitude and angular momentum. We selected the model with the lowest Akaike information criterion (AIC)⁴³ as the best-fit model,^{44,45} and we conducted post hoc comparisons using the Bonferroni correction for multiple comparisons. Significance was set at the $P < .05$ level for all analyses.

Results

Timing of Peak Angular Momentum in the Sagittal and Frontal Planes

The magnitude and direction of angular momentum changed systematically during each stride cycle. Whole-body angular momentum in the sagittal plane was most negative during the transition from swing to stance, which corresponded to the moment of peak forward

momentum (Figure 2A). Participants then increased angular momentum until it became positive during the swing phase and then fell forward again at foot-strike. The peak forward angular momentum in the sagittal plane (L_{Sag}^{min}) occurred between ~40% and 60% of the gait cycle, indicating that participants fell forward most rapidly immediately before non-paretic foot-strike. Furthermore, the 2 peaks of backward angular momentum (L_{Sag}^{max}) occurred within 25% and 35% and 75% and 85% of the gait cycle, indicating that the largest backward body rotation occurred during mid-swing of both the paretic and non-paretic sides. In the frontal plane, positive angular momentum peaked at paretic foot-strike (L_{Fr}^{min}) and negative angular momentum peaked at non-paretic foot-strike (L_{Fr}^{max}) (Figure 2B).

Associations Between Baseline Angular Momentum and Clinical Assessments of Balance

We first assessed the concurrent validity of measures of whole-body angular momentum by determining if *Baseline* angular momentum was associated with clinical balance assessments. We found significant correlations between measures of peak angular momentum and participants' scores on clinical assessments of balance. For the sagittal plane, L_{Sag}^{min} was positively correlated with




Figure 3. Associations between the peak forward and backward angular momentum in the sagittal (L_{Sag}^{min} and L_{Sag}^{max}) and frontal plane (L_{Fr}^{min}) and clinical balance assessments during Baseline. L_{Sag}^{min} was positively associated with (A) the Functional Gait Assessment (FGA) and L_{Sag}^{max} was negatively associated with (B) Berg Balance Scale (BBS) and (C) FGA. L_{Fr}^{min} was positively associated with (D) FGA.

FGA ($r = 0.53$, $P < .001$) and L_{Sag}^{max} was negatively correlated with BBS ($r = -0.43$, $P = .009$) and FGA ($r = -0.68$, $P < .001$) (Figure 3A-C). This indicates that participants who scored better on clinical assessments of balance had less forward and backward angular momentum in the sagittal plane. In the frontal plane, L_{Fr}^{min} was positively correlated with FGA ($r = 0.40$, $P < .001$, Figure 3D). This indicates that individuals who performed better on clinical assessments of dynamic balance during gait tended to rotate more slowly toward the nonparetic side while walking. We did not observe associations between ABC and L_{Sag}^{min} ($r = 0.004$, $P = .98$), L_{Sag}^{max} ($r = -0.23$, $P = .17$), L_{Fr}^{min} ($r = 0.12$, $P = .48$), or L_{Fr}^{max} ($r = 0.01$, $P = .95$). Overall, these results support the use of peak angular momentum as a measure of dynamic balance in people poststroke.

Baseline Backward Angular Momentum Was Associated With Step Length Asymmetry Magnitude

There was a significant effect of step length asymmetry magnitude on backward angular momentum (L_{Sag}^{max}) during Baseline ($F_{1,32} = 6.641$, $P = .015$) (Figure 4), but there was no effect of $SLA_{direction}$ ($F_{1,32} = 1.748$, $P = .195$) or interaction between $|SLA|$ and $SLA_{direction}$ ($F_{1,32} = 1.522$, $P = .226$). This association between L_{Sag}^{max} and $|SLA|$ remained significant after removing 2 participants who had asymmetry magnitude greater than 0.3 ($P = .04$). In contrast, forward angular momentum (L_{Sag}^{min}) during Baseline was not associated with $|SLA|$ ($F_{1,32} = 2.663$, $P = .112$) or $SLA_{direction}$ ($F_{1,32} = 1.307$, $P = .262$), nor was there an interaction between $|SLA|$ and $SLA_{direction}$ ($F_{1,32} = 0.946$, $P = .338$).




Figure 4. Association between step length asymmetry and baseline peak angular momentum in the sagittal plane. Participant's natural step length asymmetry (SLA) magnitude was positively associated with L_{Sag}^{max} magnitude as measured during the Baseline trial. This indicates that larger SLA magnitude is associated with larger peak backward angular momentum in the sagittal plane.

Baseline angular momentum in the frontal plane was not associated with asymmetry magnitude or the direction of asymmetry. Angular momentum toward the nonparetic side (L_{Fr}^{min}) during *Baseline* was not associated with $|SLA|$ ($F_{1,32} = 0.441$, $P = .511$) or $SLA_{direction}$ ($F_{1,32} = 0.131$, $P = .720$), nor was there an interaction between $|SLA|$ and $SLA_{direction}$ ($F_{1,32} = 0.003$, $P = .955$). Similarly, angular momentum toward the nonparetic side (L_{Fr}^{max}) during *Baseline* was not associated with $|SLA|$ ($F_{1,32} = 0.010$, $P = .921$) or $SLA_{direction}$ ($F_{1,32} = 0.043$, $P = .836$), nor was there an interaction between $|SLA|$ and $SLA_{direction}$ ($F_{1,32} = 0.019$, $P = .89$).

Participants Modified Step Length Asymmetry Using Visual Feedback

Participants successfully modified SLA using visual feedback. Performance of one representative participant with longer nonparetic steps demonstrates that during *Baseline FB*, SLA was maintained at the same level as *Baseline* since the target asymmetry during *Baseline FB* was the mean of their natural asymmetry (Figure 5A). The participant reduced SLA during *Sym* when walking with feedback to take steps of equal length. Last, SLA increased during *ExggAsym* compared with *Sym* since the participant walked with feedback of exaggerated asymmetry. Similarly, a representative participant with longer paretic steps maintained a consistent level of asymmetry during *Baseline*, *Baseline FB*, but modified their asymmetry in the *Sym* and *ExggAsym* conditions (Figure 5B). At the group level, participants reduced SLA magnitude during *Sym* (0.08 ± 0.06) compared with *Baseline FB* ($P = .004$), and increased SLA during *ExggAsym* (0.18 ± 0.09) relative to *Baseline FB* ($P = .02$) (Figure 5C).

Peak Angular Momentum in the Sagittal Plane Varied With Changes in SLA Magnitude

Reductions in SLA were associated with increases in forward angular momentum in the sagittal plane. The model with the best fit (lowest AIC) for predicting changes in L_{Sag}^{min} magnitude only included SLA magnitude as a significant predictor. Changes in L_{Sag}^{min} magnitude were negatively correlated with $\Delta|SLA|$ ($\beta = -0.040$, $P < .001$), indicating that reductions in asymmetry were associated with an increase in the magnitude of forward angular momentum (Figure 5D). However, no measures of asymmetry or impairment were associated with changes in the magnitude of the peak backward angular momentum in the sagittal plane and angular momentum in the frontal plane.

Discussion

We sought to determine how manipulation of SLA magnitude and the direction of asymmetry influence changes in angular momentum during walking in people post-stroke. We used online visual feedback to reduce or exaggerate step length asymmetry during walking. We first validated the use of whole-body angular momentum as a measure of dynamic balance by demonstrating that measures of angular momentum were correlated with scores on clinical balance assessments. Then, we demonstrated that as participants reduced step length asymmetry during the feedback trials, they increased the magnitude of the peak forward angular momentum in the sagittal plane. We also found that the peak backward angular momentum was greater in individuals who were more impaired and took longer nonparetic step lengths.

Associations Between Clinical Balance Assessments and Baseline Angular Momentum

Participants' peak forward angular momentum in the sagittal plane at *Baseline* was associated with clinical assessments of balance. We expected that participants with smaller peaks in angular momentum would score higher on clinical assessments as this would be consistent with the idea that people who are at a lower risk of falls would have less body rotation during walking. Consistent with this prediction, we found that participants with less angular momentum in the sagittal plane generally had higher scores on BBS and FGA. In the frontal plane, we found that FGA was positively correlated with the peak negative angular momentum indicating that people who had less mediolateral angular momentum at nonparetic foot-strike tended to score better on a clinical assessment of their balance control while walking. This extends previous observations demonstrating that people with a lower rate of change of angular momentum in the frontal plane also scored better




Figure 5. Variation in step length asymmetry (SLA) across trials. (A) A representative participant with longer nonparetic step lengths had positive SLA during Baseline, Baseline FB, and ExggAsym. During Sym, the participant reduced SLA. (B) A representative participant with longer paretic step lengths had negative SLA during Baseline, Baseline FB, and ExggAsym. During Sym, the participant reduced SLA. (C) SLA magnitude for all participants during Baseline, Baseline FB, Sym, and ExggAsym. $*P < .05$, $**P < .01$, and $***P < .001$. (D) Changes in the magnitude of L_{Sag}^{min} in the sagittal plane were negatively correlated with changes in SLA magnitude relative to Baseline indicating that reductions in asymmetry led to increases in angular momentum. $*P < .05$.

on clinical assessments of balance.¹³ Since exaggerated mediolateral sway in people poststroke often results from compensatory movements such as hip hiking and circumduction,^{15,16,25} these compensatory strategies may be reflected by measures of frontal plane angular momentum. Overall, our findings suggest that whole-body angular momentum in the sagittal and frontal planes can be used as a valid measure of balance control during walking in people poststroke.

The degree of association between BBS and FGA and angular momentum was fair and we did not find a significant association between balance confidence and angular momentum. This should be expected since there are other factors that influence these clinical assessments that are not captured by angular momentum. For example, the BBS evaluates an individual's ability to transition between chairs, retrieve an object from the floor, and perform alternating steps on a stool. We would not expect that the ability

to perform these tasks would be captured by measures of angular momentum during normal walking. Similarly, we did not find significant associations between balance confidence measured by ABC and whole-body angular momentum. This is likely because angular momentum captures people's actual balance capability rather than their perceived ability. Previous work has established that there was discordance between the actual balance capacity and the perceived balance in people poststroke who showed low BBS and high ABC and vice versa.⁴⁶

Effects of Manipulating Step Length Asymmetry on Angular Momentum

Contrary to our hypothesis, the peak forward angular momentum in the sagittal plane increased when people walked more symmetrically. We expected that angular momentum would reduce as SLA decreased since walking with more symmetric step lengths would result in more symmetrical, antiphase limb rotation and better cancellation of the contralateral limb angular momenta. However, we found that participants increased their forward angular momentum as they reduced SLA, indicating that improving asymmetry did not reduce whole-body rotational behavior during walking. Although changes in asymmetry led to changes in angular momentum in the sagittal plane, peak angular momentum in the frontal plane did not change when walking more symmetrically. This suggests that changes in step length asymmetry in the sagittal plane do not systematically influence frontal plane momentum. However, it is possible that changes in asymmetry could have caused changes in both frontal and sagittal plane balance that we could not detect. For example, our study was not designed to capture changes in the reactive control of balance³⁸ and this, among other balance-related measures, could be sensitive to changes in asymmetry. Since manipulating asymmetry in our study was not a type of perturbation requiring reactive balance control, the changes in angular momentum in the frontal plane may have not been significant.

Eight of 15 participants increased forward angular momentum near the time of foot-strike on the side where they increased step lengths. Since we manipulated asymmetry by increasing the shorter step length, this may have been associated with an increased excursion of the CoM relative to the trailing leg and higher CoM velocity when lengthening the short step. This increase in velocity may have contributed to increases in momentum that we observed in these individuals. Seven participants increased forward angular momentum near the time of foot-strike on the side contralateral to shorter step length. These individuals may have relied on increasing propulsion before lift-off of the limb on the side where they attempted to increase step lengths. This attempt to generate more swing leg

momentum may have caused increased forward angular momentum near the time of foot-strike on the side contralateral to the shorter step length.

Another potential reason why participants did not reduce angular momentum is that reductions in SLA may not have produced meaningful changes in momentum cancellation between limbs. A recent study found that people poststroke continue to walk with asymmetric interlimb coordination even when reducing step length asymmetry.⁵ This suggests that their legs may not move in perfect antiphase relative to the CoM as they reduce SLA. Overall, our results suggest that balance-related responses to reductions in step length asymmetry in people poststroke may vary based on the strategy they use to change their step lengths.

The observed increases in angular momentum during more symmetric walking may reflect impairments in dynamic balance. In our study, the increase in the peak forward angular momentum during foot-strike was about 0.008 (~16%) as SLA decreased by 0.2 using the estimated coefficient ($\beta = -0.04$) in the linear model relating $\Delta|\text{SLA}|$ and $\Delta|L_{\text{Sag}}^{\text{min}}|$. The size of the changes in angular momentum when walking more symmetrically was smaller than the angular momentum changes when exposed to a perturbation large enough to cause a fall during walking.⁴⁷ When participants without disabilities experienced fore-aft perturbations characterized by abrupt increases in treadmill speed of approximately 0.6 m/s, they increased peak angular momentum in the sagittal plane by ~0.03 (~50%). Thus, the changes in momentum we observed in the current study were approximately equal to the changes that one might expect in individuals without disabilities responding to an unexpected 0.2 m/s increase in treadmill speed. Although the change in angular momentum that we observed is unlikely to be as large as one might expect in a near fall, such changes could be perceived as sizable for people poststroke depending on their level of sensorimotor function.

However, it remains to be seen how longer-term practice or interventions leading to reductions in SLA affects the regulation of momentum during walking. Longer-term improvements in symmetry could be achieved through methods such as repeated practice on a split-belt treadmill,² using auditory feedback of stance time asymmetry,⁴⁸ or gait training with functional electrical stimulation.⁴⁹ Although participants in these studies retained improvements in symmetry after practice, it remains to be seen if these improvements in symmetry impact dynamic balance. Overall, our findings may inform future research efforts that seek to understand how novel interventions affect quantitative measures of balance control during walking poststroke.

Conclusion

We found that whole-body angular momentum in the sagittal and frontal plane during walking is a valid measure of

dynamic balance in people poststroke. Although we expected that improving symmetry would improve balance poststroke, we found that the reductions in asymmetry impair dynamic balance by increasing whole-body angular momentum. This may imply, more generally, that people poststroke have an implicit preference not to deviate from their natural asymmetry because it may perturb their balance. However, additional investigation is necessary to determine the relative contributions of other factors such as energetic optimization, discomfort, and aesthetics in the selection of self-selected walking patterns in people poststroke. Future studies should determine if the findings observed here generalize to overground walking where participants' walking speed is less constrained. This is critical from a clinical perspective as existing clinical approaches to increase symmetry may rely on providing instructive guidance during overground walking. An additional clinical implication of our results is that if clinicians choose to focus on guiding their patients to improve symmetry, they may also need to consider how to reduce heightened destabilization resulting from these improvements or increase their patients' capacity to withstand the instability.

Declaration of Conflicting Interests


The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Eunice Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of Health under Award Number R01HD091184 and SC CTSI (NIH/NCATS) through Grant UL1TR001855.

ORCID iDs

Chang Liu  <https://orcid.org/0000-0002-0381-4136>

Natalia Sánchez  <https://orcid.org/0000-0002-6467-1781>

James M. Finley  <https://orcid.org/0000-0003-2679-2221>

References

1. Patterson KK, Mansfield A, Biasin L, Brunton K, Inness EL, McIlroy WE. Longitudinal changes in poststroke spatiotemporal gait asymmetry over inpatient rehabilitation. *Neurorehabil Neural Repair*. 2015;29:153-162. doi:10.1177/1545968314533614
2. Reisman DS, McLean H, Keller J, Danks KA, Bastian AJ. Repeated split-belt treadmill training improves poststroke step length asymmetry. *Neurorehabil Neural Repair*. 2013;27:460-468. doi:10.1177/1545968312474118
3. Silver KH, Macko RF, Forrester LW, Goldberg AP, Smith GV. Effects of aerobic treadmill training on gait velocity, cadence, and gait symmetry in chronic hemiparetic stroke: a preliminary report. *Neurorehabil Neural Repair*. 2000;14:65-71. doi:10.1177/154596830001400108
4. Nguyen TM, Jackson RW, Aucie Y, Kam D de, Collins SH, Torres-Oviedo G. Self-selected step length asymmetry is not explained by energy cost minimization in individuals with chronic stroke. Accessed April 3, 2020. <https://www.medrxiv.org/content/10.1101/19013854v1>
5. Padmanabhan P, Rao KS, Gulhar S, Cherry-Allen KM, Leech KA, Roemmich RT. Persons post-stroke restore step length symmetry by walking asymmetrically. Accessed May 11, 2021. <https://www.biorxiv.org/content/10.1101/799775v1>
6. Sánchez N, Finley JM. Individual differences in locomotor function predict the capacity to reduce asymmetry and modify the energetic cost of walking poststroke. *Neurorehabil Neural Repair*. 2018;32:701-713. doi:10.1177/1545968318787913
7. Harris JE, Eng JJ, Marigold DS, Tokuno CD, Louis CL. Relationship of balance and mobility to fall incidence in people with chronic stroke. *Phys Ther*. 2005;85:150-158. doi:10.1093/ptj/85.2.150
8. Weerdesteyn V, de Niet M, van Duijnhoven HJR, Geurts ACH. Falls in individuals with stroke. *J Rehabil Res Dev*. 2008;45:1195-1213.
9. Lewek MD, Bradley CE, Wutzke CJ, Zinder SM. The relationship between spatiotemporal gait asymmetry and balance in individuals with chronic stroke. *J Appl Biomech*. 2014;30:31-36. doi:10.1123/jab.2012-0208
10. Elftman H. The function of the arms in walking. *Hum Biol*. 1939;11:529.
11. Herr H, Popovic M. Angular momentum in human walking. *J Exp Biol*. 2008;211(pt 4):467-481. doi:10.1242/jeb.008573
12. Honda K, Sekiguchi Y, Muraki T, Izumi SI. The differences in sagittal plane whole-body angular momentum during gait between patients with hemiparesis and healthy people. *J Biomech*. 2019;86:204-209. doi:10.1016/j.jbiomech.2019.02.012
13. Nott CR, Neptune RR, Kautz SA. Relationships between frontal-plane angular momentum and clinical balance measures during post-stroke hemiparetic walking. *Gait Posture*. 2014;39:129-134. doi:10.1016/j.gaitpost.2013.06.008
14. Vistamehr A, Kautz SA, Bowden MG, Neptune RR. Correlations between measures of dynamic balance in individuals with post-stroke hemiparesis. *J Biomech*. 2016;49:396-400. doi:10.1016/j.jbiomech.2015.12.047
15. Cruz TH, Lewek MD, Dhaher YY. Biomechanical impairments and gait adaptations post-stroke: multi-factorial associations. *J Biomech*. 2009;42:1673-1677. doi:10.1016/j.jbiomech.2009.04.015
16. Stanhope VA, Knarr BA, Reisman DS, Higginson JS. Frontal plane compensatory strategies associated with self-selected walking speed in individuals post-stroke. *Clin Biomech (Bristol, Avon)*. 2014;29:518-522. doi:10.1016/j.clinbiomech.2014.03.013
17. Popovic M, Hofmann A, Herr H. Angular momentum regulation during human walking: biomechanics and control. In: *Proceedings of the IEEE International Conference on Robotics and Automation, 2004 ICRA '04*. Vol 3. IEEE; 2004:2405-2411. doi:10.1109/ROBOT.2004.1307421
18. Patten C, Lexell J, Brown HE. Weakness and strength training in persons with poststroke hemiplegia: rationale, method, and efficacy. *J Rehabil Res Dev*. 2004;41(3A):293-312. doi:10.1682/jrjd.2004.03.0293

19. Rybar MM, Walker ER, Kuhnen HR, Ouellette DR, Hunter SK, Hyngstrom AS. The stroke-related effects of hip flexion fatigue on over ground walking. *Gait Posture*. 2014;39:1103-1108. doi:10.1016/j.gaitpost.2014.01.012
20. Allen JL, Kautz SA, Neptune RR. Forward propulsion asymmetry is indicative of changes in plantarflexor coordination during walking in individuals with post-stroke hemiparesis. *Clin Biomech (Bristol Avon)*. 2014;29:780-786. doi:10.1016/j.clinbiomech.2014.06.001
21. Lauzière S, Miéville C, Betschart M, Aissaoui R, Nadeau S. Plantarflexor weakness is a determinant of kinetic asymmetry during gait in post-stroke individuals walking with high levels of effort. *Clin Biomech (Bristol Avon)*. 2015;30:946-952. doi:10.1016/j.clinbiomech.2015.07.004
22. Finley JM, Long A, Bastian AJ, Torres-Oviedo G. Spatial and temporal control contribute to step length asymmetry during split-belt adaptation and hemiparetic gait. *Neurorehabil Neural Repair*. 2015;29:786-795. doi:10.1177/1545968314567149
23. Roerdink M, Beek PJ. Understanding inconsistent step-length asymmetries across hemiplegic stroke patients: impairments and compensatory gait. *Neurorehabil Neural Repair*. 2011;25:253-258. doi:10.1177/1545968310380687
24. Varraine E, Bonnard M, Pailhous J. Intentional on-line adaptation of stride length in human walking. *Exp Brain Res*. 2000;130:248-257. doi:10.1007/s002219900234
25. Titus AW, Hillier S, Louw QA, Inglis-Jassiem G. An analysis of trunk kinematics and gait parameters in people with stroke. *Afr J Disabil*. 2018;7:310. doi:10.4102/ajod.v7i0.310
26. Balasubramanian CK, Bowden MG, Neptune RR, Kautz SA. Relationship between step Length asymmetry and walking performance in subjects with chronic hemiparesis. *Arch Phys Med Rehabil*. 2007;88:43-49. doi:10.1016/j.apmr.2006.10.004
27. Hsu AL, Tang PF, Jan MH. Analysis of impairments influencing gait velocity and asymmetry of hemiplegic patients after mild to moderate stroke. *Arch Phys Med Rehabil*. 2003;84:1185-1193. doi:10.1016/S0003-9993(03)00030-3
28. Olney SJ, Richards C. Hemiparetic gait following stroke. Part I: characteristics. *Gait Posture*. 1996;4:136-148. doi:10.1016/0966-6362(96)01063-6
29. Merlo A, Campanini I. Impact of instrumental analysis of stiff knee gait on treatment appropriateness and associated costs in stroke patients. *Gait Posture*. 2019;72:195-201. doi:10.1016/j.gaitpost.2019.06.009
30. Kerrigan DC, Gronley J, Perry J. Stiff-legged gait in spastic paresis. A study of quadriceps and hamstrings muscle activity. *Am J Phys Med Rehabil*. 1991;70:294-300.
31. Blum L, Korner-Bitensky N. Usefulness of the Berg Balance Scale in stroke rehabilitation: a systematic review. *Phys Ther*. 2008;88:559-566. doi:10.2522/ptj.20070205
32. Langley F, Mackintosh S. Functional balance assessment of older community dwelling adults: a systematic review of the literature. *Internet J Allied Health Sci Pract*. 2007;5(4), 13. Accessed May 11, 2021. <https://nsuworks.nova.edu/ijahsp/vol5/iss4/13>
33. Leddy AL, Crowner BE, Earhart GM. Functional gait assessment and balance evaluation system test: Reliability, validity, sensitivity, and specificity for identifying individuals with Parkinson disease who fall. *Phys Ther*. 2011;91:102-113. doi:10.2522/ptj.20100113
34. Powell LE, Myers AM. The Activities-specific Balance Confidence (ABC) Scale. *J Gerontol A Biol Sci Med Sci*. 1995;50A:M28-M34. doi:10.1093/gerona/50a.1.m28
35. Dewana N, MacDermid JC. Fall Efficacy Scale—International (FES-I). Accessed April 3, 2021. <https://www.sciencedirect.com/science/article/pii/S1836955314000265?via%3Dihub>
36. Fugl-Meyer AR, Jääskö L, Leyman I, Olsson S, Steglind S. The post-stroke hemiplegic patient. 1. a method for evaluation of physical performance. *Scand J Rehabil Med*. 1975;7:13-31.
37. Sullivan KJ, Tilson JK, Cen SY, et al. Fugl-Meyer Assessment of sensorimotor function after stroke. *Stroke*. 2011;42:427-432. doi:10.1161/STROKEAHA.110.592766
38. Liu C, Macedo LD, Finley JM. Conservation of reactive stabilization strategies in the presence of step length asymmetries during walking. *Front Hum Neurosci*. 2018;12:251. doi:10.3389/fnhum.2018.00251
39. Havens KL, Mukherjee T, Finley JM. Analysis of biases in dynamic margins of stability introduced by the use of simplified center of mass estimates during walking and turning. *Gait Posture*. 2018;59:162-167. doi:10.1016/j.gaitpost.2017.10.002
40. Song J, Sigward S, Fisher B, Salem GJ. Altered dynamic postural control during step turning in persons with early-stage Parkinson's disease. *Parkinsons Dis*. 2012;2012:386962. doi:10.1155/2012/386962
41. Allen JL, Kautz SA, Neptune RR. Step length asymmetry is representative of compensatory mechanisms used in post-stroke hemiparetic walking. *Gait Posture*. 2011;33:538-543. doi:10.1016/j.gaitpost.2011.01.004
42. Mulroy S, Gronley J, Weiss W, Newsam C, Perry J. Use of cluster analysis for gait pattern classification of patients in the early and late recovery phases following stroke. *Gait Posture*. 2003;18:114-125. doi:10.1016/s0966-6362(02)00165-0
43. Akaike H. Likelihood of a model and information criteria. *J Econom*. 1981;16:3-14. doi:10.1016/0304-4076(81)90071-3
44. Hocking RR, Leslie RN. Selection of the best subset in regression analysis. *Technometrics*. 1967;9(4):531-540. doi:10.2307/1266192
45. James G, Witten D, Hastie T, Tibshirani R. *An Introduction to Statistical Learning: With Applications in R*. Springer-Verlag; 2013. Accessed September 9, 2019. <https://www.springer.com/gp/book/9781461471370>
46. Liphart J, Gallichio J, Tilson JK, Pei Q, Wu SS, Duncan PW. Concordance and discordance between measured and perceived balance and the effect on gait speed and falls following stroke. *Clin Rehabil*. 2016;30:294-302. doi:10.1177/0269215515578294
47. Martelli D, Monaco V, Luciani LB, Micera S. Angular momentum during unexpected multidirectional perturbations delivered while walking. *IEEE Trans Biomed Eng*. 2013;60:1785-1795. doi:10.1109/TBME.2013.2241434
48. Krishnan V, Khoo I, Marayong P, DeMars K, Cormack J. Gait training in chronic stroke using walk-even feedback device: a pilot study. *Neurosci J*. 2016;2016:6808319. doi:10.1155/2016/6808319
49. Awad LN, Palmer JA, Pohlig RT, Binder-Macleod SA, Reisman DS. Walking speed and step length asymmetry modify the energy cost of walking after stroke. *Neurorehabil Neural Repair*. 2015;29:416-423. doi:10.1177/1545968314552528