



UvA-DARE (Digital Academic Repository)

MAXI J1659-152 fading in optical

Russell, D.M.; Lewis, F.; Schreuder, L.; Yang, Y.J.; Wijnands, R.; Tripp, A.

Publication date

2011

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Russell, D. M., Lewis, F., Schreuder, L., Yang, Y. J., Wijnands, R., & Tripp, A. (2011). MAXI J1659-152 fading in optical. *The astronomer's telegram*, 3517. <https://www.astronomerstelegam.org/?read=3517>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

ATel On

Patreon
Mastodon
Twitter

The Astronomer's Telegram

Post | Search | Policies
Credential | Feeds | Email

24 Jul 2023; 13:15 UT

This space for free for your conference.



Thanks to Patrons, The Astronomer's Telegram is free to read, free to publish and always will be. Thank you.

[[Previous](#) | [Next](#) | [ADS](#)]

MAXI J1659-152 fading in optical

ATel #3517; *D. M. Russell (University of Amsterdam), F. Lewis, (Univ. of Glamorgan, Faulkes Telescope Project, Open Univ.), L. Schreuder, Y. J. Yang, R. Wijnands (University of Amsterdam), A. Tripp (Faulkes Telescope Project)*

on 28 Jul 2011; 15:37 UT

Credential Certification: David M. Russell (D.M.Russell@uva.nl)

Subjects: Optical, X-ray, Binary, Black Hole, Transient

Referred to by ATel #: [3524](#)

[Tweet](#)

It was recently reported that the black hole candidate X-ray binary MAXI J1659-152 has faded to an X-ray flux (0.5-10 keV) of $< 3.0e-13$ ergs cm⁻² s⁻¹ (ATel #[3506](#)). This implies the source may be in quiescence after an outburst that lasted almost one year (GCN #[11296](#), ATel #[2873](#)) including a late re-brightening in the last three months (ATels #[3298](#), #[3339](#), #[3358](#), #[3379](#)). Here we report on recent optical observations with the two 2-m Faulkes Telescopes North and South (see ATel #[2884](#)).

MAXI J1659-152 has faded in optical but is still detected in V, R and i'-bands as of 2011-07-27 (MJD 55769.4). The magnitudes on this date were V = 20.8 +/- 0.1, R ~ 20.1. Flux calibration in V-band was achieved using field star magnitudes measured from Swift UVOT images. In R-band, the magnitudes of field stars in the GSC 2.2 digital sky survey were used to calibrate the data, and there may be systematic errors of several tenths of a magnitude. Flux calibration in i'-band is not available.

Soon after the outburst began in September 2010, the optical flux peaked at V ~ 16.5 mag (ATel #[2884](#); [Kennea et al. 2011, ApJ, 736, 22](#)). In 2011 our Faulkes Telescope monitoring shows that by 2011-05-21 (MJD 55702.8) it had faded by only 2.5 mag to V = 19.0 +/- 0.1. During the next 15 days the source remained at a fairly steady flux level; V ~ 18.9 - 19.0. This was during the period of X-ray re-brightenings. At a magnitude of V =

Related

- [4044](#) MAXI J1305-704: Swift/XRT detection of dips, possible 1.5 hour period
- [3524](#) Lulin Optical Observations of MAXI J1659-152
- [3517](#) MAXI J1659-152 fading in optical
- [3506](#) MAXI J1659-152 has likely returned back into quiescence
- [3379](#) Further re-brightening of the black hole candidate MAXI J1659-152
- [3358](#) X-ray and radio observations of the re-brightening event in MAXI J1659-152
- [3339](#) Re-brightening of the black hole candidate MAXI J1659-152
- [3298](#) Continued Swift monitoring of the black hole candidate MAXI J1659-152
- [3251](#) RXTE detection of Rapidly Varying Transient Swift J164449.3+573451
- [3250](#) Swift J164449.3+573451/GRB 110328A: Continued Swift Monitoring
- [3249](#) Swift XRT/UVOT monitoring of MAXI J1659-152 during its low luminosity phase
- [3244](#) MAXI/GSC observation of Swift J164449.3+573451
- [3242](#) Swift J164449.3+573451 (AKA GRB110328A): A new candidate SFXT?
- [3201](#) Swift XRT/UVOT follow-up of the Black Hole Candidate MAXI J1659-152 during a low luminosity state
- [2999](#) Soft-to-Hard transition in MAXI J1659-152
- [2976](#) Optical emission of the black hole X-ray transient MAXI J1659-152 during quiescence
- [2951](#) RXTE shows a transition to the high-soft state in MAXI J1659-152
- [2927](#) Transition to the soft-intermediate state of MAXI J1659-152
- [2926](#) RXTE dips yield better orbital period determination for MAXI J1659-152
- [2918](#) Sudden radio flux decline in MAXI J1659-152
- [2912](#) MAXI J1659-152: the shortest period black-hole binary?

20.8 +/- 0.1 on 2011-07-27, MAXI J1659-152 is now ~4.3 mag fainter than its brightest outburst magnitude. This is a fade of a factor of 50 in optical flux; for comparison the X-ray flux of $< 3.0e-13$ ergs cm⁻² s⁻¹ (ATel #3506) is $> 10^5$ times fainter than the peak 0.5-10 keV X-ray flux (Kennea et al. 2011). The optical colour has also changed; V - i' is 0.5 mag redder on 2011-07-27 compared to 2011-05-21. A link to the light curves is provided below.

The upper limit on the quiescent magnitude of MAXI J1659-152 is V > 21 mag from its non-detection in the USNO B1 catalogue (Kennea et al. 2011). The source is now almost as faint as this upper limit; we aim to continue monitoring the optical counterpart to test whether the source continues to fade or remains at this magnitude. A wide range of quiescent magnitudes for MAXI J1659-152 have been predicted; from r ~ 22.4 (ATel #2976) to V ~ 28 (Kuulkers et al. 2011, arXiv:1102.2102).

The Faulkes Telescope observations are part of an on-going monitoring campaign of ~ 30 low-mass X-ray binaries (Lewis et al. 2008, arXiv:0712.2751). The Faulkes Telescope South is maintained and operated by Las Cumbres Observatory Global Telescope Network. FL acknowledges support from the Dill Faulkes Educational Trust.

MAXI J1659-152 optical light curves and finding charts

| | |
|------|--|
| 2906 | EVN e-VLBI detections of MAXI J1659-152 |
| 2900 | REM optical/NIR observations of MAXI J1659-152 |
| 2890 | INTEGRAL shows MAXI J1659-152 further declines in hard X-rays |
| 2888 | INTEGRAL TOO observations of MAXI J1659-152 |
| 2887 | XMM-Newton observations of MAXI J1659-152 |
| 2884 | Optical variability in MAXI J1659-152 |
| 2881 | MAXI J1659-152 is a BH candidate |
| 2880 | AGILE upper limits above 100 MeV regarding the recent transient MAXI J1659-152 |
| 2877 | MAXI J1659-152: Swift localization and monitoring |
| 2875 | INTEGRAL detection of the new MAXI transient MAXI J1659-152 |
| 2874 | WSRT Radio and Polarization Detection of GRB 100925A / MAXI J1659-152 |
| 2873 | MAXI/GSC detection of a new hard X-ray transient source MAXI J1659-152 |
| 2734 | Radio and X-ray monitoring of Cygnus X-1 during the recent state change |

[[Telegram Index](#)]

R. E. Rutledge, Editor-in-Chief

rrutledge@astronomerstelegam.org

Derek Fox, Editor

dfox@astronomerstelegam.org