



GLOBAL JOURNAL OF HUMAN-SOCIAL SCIENCE: A
ARTS & HUMANITIES - PSYCHOLOGY
Volume 15 Issue 1 Version 1.0 Year 2015
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-460X & Print ISSN: 0975-587X

Relations of Anthropometric and Conative (Normal and Pathological) Dimensions of Volleyball Junior Players

By Miroljub Ivanović, Srdjan Milosavljević & Uglješa Ivanović

Business Informatics College, Serbia

Abstract- The main objective of this transversal study was to examine the relation of morphological and conative characteristics (normal and pathologic) factors in volleyball junior players. The data were collected on the sample of examinees (N = 126), aged 16 to 17 years from five volleyball clubs. The applied measuring instruments consisted of 23 anthropometric measures, 16 normal and 12 pathological conative variables. Descriptive statistics, Pearson's correlation coefficient and canonical correlation analysis were applied for data processing. The results showed that among the set of anthropometric characteristics and set of normal conative dimensions, with 51% of explained variance, there is a statistically significant canonical correlation of high level ($R_c = .77$, $p < .05$), whereas among the set of normal and pathological conative dimensions, with the interpreted variability of 59%, two significant correlations of average intensity ($R_c = .73$, $p < .01$, and $R_c = .70$, $p < .05$) were defined.

Keywords: *anthropometric characteristics, conative variables, volleyball players.*

GJHSS-A Classification : *FOR Code: 110699*



Strictly as per the compliance and regulations of:



Relations of Anthropometric and Conative (Normal and Pathological) Dimensions of Volleyball Junior Players

Miroljub Ivanović ^α, Srdjan Milosavljević ^σ & Uglješa Ivanović ^ρ

Abstract- The main objective of this transversal study was to examine the relation of morphological and conative characteristics (normal and pathologic) factors in volleyball junior players. The data were collected on the sample of examinees (N = 126), aged 16 to 17 years from five volleyball clubs. The applied measuring instruments consisted of 23 anthropometric measures, 16 normal and 12 pathological conative variables. Descriptive statistics, Pearson's correlation coefficient and canonical correlation analysis were applied for data processing. The results showed that among the set of anthropometric characteristics and set of normal conative dimensions, with 51% of explained variance, there is a statistically significant canonical correlation of high level ($R_c = .77$, $p < .05$), whereas among the set of normal and pathological conative dimensions, with the interpreted variability of 59%, two significant correlations of average intensity ($R_c = .73$, $p < .01$, and $R_c = .70$, $p < .05$) were defined. Following relevant canonical factors were isolated: I) morphological ectomesomorphic factors (longitudinal dimensionality of the skeleton and transversal dimensionality of the skeleton), II) normal conative factors (extraversion-introversion and anxiety, intelligence, self-awareness, courage, a sharp temper and protension; introspection and insight-honesty) and III) pathological conative factors (performance of systems for regulating and monitoring the biological functions and automatic functions and editing of responses of defense and attack). The structure of the relations between anthropometric characteristics and conative (normal and pathological) latent dimensions was presented for a probability level of 0.05.

Keywords: anthropometric characteristics, conative variables, volleyball players.

I. INTRODUCTION

In the last decade of the XXI century, relations between morphological characteristics and conative (normal and pathologic) factors of athletes arouse more and more attention in anthropological studies, as it is reflected in surveys conducted by: (Blaževik, 2006; Djurković, 2009; Djurkovic, 2009; Sing & Rathore, 2013; Sheppard *et al.*, 2008; Toskić, Okičić & Stankovic, 2012). Morphological anthropometry - somatometry - provides information about the processes of growth and

Author α: Preschool Teacher Training and Business Informatics College – Sirmium, Sremska Mitrovica, Serbia.
e-mail: miroljub.ivanovic@gmail.com

Author σ: Sports and Health College, Belgrade, Serbia.

Author ρ: Faculty of Sports Management, Belgrade, Alpha University, Belgrade, Serbia.

ontogenetic development in volleyball population, and specialization, i.e. defining characteristic team positions with specific tasks, which is proven by the research conducted by Malacko & Stankovic (2009).

Significant number of authors investigated anthropometric characteristics of volleyball players Dopsaj *et al.*, 2010; Djurkovic, Marelic & Rešetar 2011; Ivanovic, Ivanovic, 2007; Ivanovic & Ivanovic, 2012; Small *et al.*, 2010; Marelič, Matkovic & Antekolović, 2010; Martin Matillas *et al.*, 2014; Rodríguez-Ruiz *et al.*, 2012; Popovic *et al.*, 2010; Cabral *et al.*, 2011). Applying methods of multivariate analysis, the researchers isolated latent dimensions in the field of anthropometry and determined their interdependence with other anthropological skills and characteristics of volleyball players. It has been determined that in this sports game, tall athletes, the ones with a greater arm and leg length have a certain advantage when compared to short athletes, or those with short extremities. Furthermore, athletes with greater amounts of fatty tissue are inferior in comparison to the ones with the same weight but unburdened by the unwanted fatty tissue.

Taking into account the influence of genetic and exogenous factors on anthropometric measures of volleyball players, a significant number of authors emphasizes their importance: Bandyopadhyay, 2007; Vrbik, Čížmek & Gruić, 2011; Marelic, Djurković & Rešetar, 2007; Popovic-Ilic *et al.*, 2010; Trajkovic *et al.*, 2011; Pori & Sibiu, 2009. Considering their latent morphological dimension, the authors emphasize that the anthropometric characteristics are responsible for the longitudinal dimensionality of the skeleton (the growth of bones in terms of length), transversal dimensionality of the skeleton (the growth of bones in terms of width), the circular dimensionality and body weight (in terms of volume and total body mass) and subcutaneous fatty tissue.

The influence of psychological factors on sports results represents the a precondition for efficient accomplishment of the motor task, according to the survey done by (Duncan, Woodfield, & Nakeeb, 2006; Jurko, Nestic & Stojanovic, 2013; Ivanovic, Ivanovic, 2011; Ivanovic Ivanovic, 2012a; Ivanovic & Ivanovic, 2012b, Ivanovic, Ivanovic, 2012v; Ivanovic & Ivanovic, 2013; Ivanovic, Ivanovic & Samardzic, 2012; Stojanović, Milenkoski & Stojanovic, 2011; Tubić, 2010). Conative

(motivational) component of attitudes refers to an individual's activities directed towards a specific object. Its effect can be positive or negative and harmonized with the knowledge and emotions related to that object. If the knowledge and emotions are favorable, people will try to bring positive actions in relation to the object, for example conative component can be directed from a tendency to help, support and protect the object to the tendency to destroy the object, as it is shown by research done by Blaževik, (2009), Kalach & Gontarev (2013), Muric et al., (2012), Sindik i Nazor (2011), Sindik et al., (2011), Stankovic & Popovic (2011), and Toskic et al., (2012). Conative (normal and pathologic) factors are structured by personality traits and responsible for its modes of behavior. Normal conative factor is characterized by moderate intensity acting neutral on person's adjustment disorders. On the other hand, pathological conative factor reduces the level of adjustment of the individual and causes personality disorder, states Stankovic et al., (2011) in his research.

A large number of studies have been based on identifying relations between morphological and conative characteristics of athletes, as it is shown by research conducted by: Karalejić et al., (2010), Joško i Joško (2011), Stankovic et al., (2011), Stankovic et al., (2013) and Stojanovic et al., (2011). In these studies it is emphasized that the action and achievement of effective results during the match demand specific needs integrity of afore mentioned anthropological qualities. In his study, Pajević (2003) suggests that normal conative factors cause different forms of human behavior, and integrate ego and improve effective communication with the environment. Popovic & Simonovic (2008) find that normal conative characteristics are mutually related in the processes of excitation and prevention of potential of adjustment. On the other hand, inherited pathological conative characteristics cause disorders in unifying the personality, or disturb the balance between reducing and increasing of adaptive powers, according to a survey conducted by Horga, Momirović & Jankovic (1983).

The results of previous studies that have observed the relationship of cognitive and anthropometric variables in adolescence are not consistent, many aspects still do not have final explanation, and this transversal study investigates their linear relations. Further empirical research on the interaction of these anthropological characteristics in junior volleyball players population are needed to fully shed light on the mechanism of their correlates.

The aim of this research was to evaluate whether there were any statistically significant relations between the system of morphological characteristics, normal conative characteristics and pathological conative characteristics in the population of junior volleyball players. In order to achieve this goal, the authors of this paper assumed that the systems of

anthropometric characteristics and normal and pathological conative dimensions are statistically significantly correlated.

II. RESEARCH METHOD AND PROCEDURES

a) Examinees

The research involved 126 junior volleyball players, aged 16 to 17 years from five volleyball teams: *super league* "Spartak" (Ljig); *I League* "Železničar" (Lajkovac); *II League* "Loznica" (Loznica), "Bravo" (Valjevo) and "Ub" (UB). The average age of a suitable sample of examinees was 16.68 years (SD = 7.24). All volleyball players had at least two years of systematic and organized training and competition.

b) The sample of morphological variables

In order to evaluate latent morphological characteristics, the following variables were used: Body height (MS), Leg length (DNO), Hand length (DSA), Foot length (DST), Arm length (DRU), Biacromial range (BIA), - *longitudinal dimensionality of the skeleton*; Bicristal range (BIK), Hand width (SSA), Wrist diameter (DIR), Elbow diameter (DIL), Knee diameter (DIK), Foot width (SST) - *transversal dimensionality of the skeleton*; Body mass (MAS), Upper arm volume (ONA), Lower arm volume (OPO), Upper leg volume (ONT), Lower leg volume (OPT), Average thorax volume (OGK) - *circular dimensionality and body mass*; Upper arm skinfold (KNA), Back skinfold (KLE), Armpit skinfold (KPA), Abdominal skinfold (KTR), Lower leg skinfold (KPT) - *subcutaneous fatty tissue*.

Anthropometric measurements were performed according to the instructions and regulations of the International Biological Program, which was formed by (Norton & Olds, 2004). Applied standard measuring instruments (anthropometer according to Martin's scale for measuring height, medical decimal scale with sliding weights to measure body weight, Holtain flexible measuring tape for body volume, Holtain Bicondylar Vernier caliper to measure the diameters of the bones and caliper by John Bull for measuring skin folds), are calibrated before measuring. All morphological characteristics were measured again, with the exception of skin folds, which were presented as the mean value after the three consecutive measurements. The study was conducted in groups during regular trainings in May 2014.

Prior to the implementation of anthropometric measurements, the examinees were explained the aim of the research, they were asked to participate and explained that they could give up whenever they want. On average, anthropometric measurements lasted for 45 min. Examinees were first informed about the research and protection of the anonymity of the data, after which they signed an agreement on participation in the study.

c) *The sample of conative (normal and pathological variables)*

In order to evaluate the normal conative characteristics, the 16PF battery was used (Cattell, Eber, & Tatsuoka, 1970) with first-order factors: Outgoing – reserved (A), High intelligence – low intelligence (B), Higher ego strength – lower ego strength (C), Dominance – submissiveness (E), Surgency – desurgency (F), Stronger superego – weaker superego (G), Bold (parmia) – shy (threctia) (H), Tender-minded (premsia) – tough-minded (I), Protension (L), Autia (imaginative) – practical (M), Shrewdness – artlessness (N), Apprehensive (O), Radicalism – conservatism (Q1), Self-sufficiency – group dependence (Q2), High self-concept – low self-concept (Q3), Tense – relaxed (Q4).

Pathological conative characteristics were operationalized through the score for selected scales on a test of pathological conative characteristics (Momirović, 1971), C.I.-N4 - *the efficiency of the regulatory system and control of organic functions* (H1); Cardiovascular conversion (K10), Gastrointestinal conversion (G11), Inhibitory conversion (I7),

Hypochondria (H13), *the effectiveness of the regulatory system and the control of defensive reactions* (ALPHA), Anxiety (A1), Obsession (O3), Hypersensitivity (S5), Phobias (F2), *the efficiency of the regulatory system and the control of attack reactions* (SIGMA), Impulsiveness (N14), Aggression (T15) *and the effectiveness of the system for the coordination of regulatory functions* (DELTA), Paranoia (P18), Depression (D6).

III. RESULTS

a) *Descriptive data and correlations between examined variables*

Results of morphological variables examined through descriptive statistical parameters are shown in Table 1a, 1b. Obtained values skewness and kurtosis indicate that there is no significant dispersion of the distribution from normal Gaussian distribution (Tenjović, 2002). This indicates that the measuring instruments were well suited for the measurement of anthropometric variables in volleyball junior players population, so that further statistical analysis can be performed.

Table 1 : Basic descriptive-statistical parameters of anthropometric variables

Variables	M	SD	Min	Max	Sk	Ku
Body height (cm)	1820.08	61.785	1631	1963	-.257	.295
Leg length (cm)	1033.06	47.929	942	1182	.263	.301
Habd length (cm)	196.34	11.231	171	239	.572	1.287
Foot length (cm)	280,35	13,005	261	319	.799	.558
Arm length (cm)	783.06	50.040	689	879	-.207	-.219
Biacromial range (cm)	409.86	29.871	311	539	.390	4.661
Foot width (cm)	92.19	7.916	68	111	-.395	.592
Knee diameter (cm)	92.56	5.878	75	106	.182	-.401
Elbow diameter (cm)	75.06	6.606	61	89	.230	-.492
Wrist diameter (cm)	59.04	2.927	52	69	.193	.149
Body mass (kg)	78.543	111.065	549	1079	.393	-.312
Bicristal range (cm)	300.03	26.008	244	389	.728	2.212
Nad width (cm)	86.00	5.437	74	99	.088	-.353

Upper arm volume (cm)	284.06	24.975	229	371	.714	.967
Lower arm volume (cm)	270.04	20.98	219	321	.306	-.071
Upper leg volume (cm)	562.03	44.923	479	681	.648	-.015
Lower leg volume (cm)	391.87	49.016	319	599	1.918	7.012
Average thorax volume (cm)	957.01	70.012	839	1111	.252	-.694
Upper arm skinfold (mm)	93.98	35.894	35	199	1.017	.735
Back skinfold (mm)	105.01	34.018	65	215	1.753	3.023
Armpit skinfold (mm)	75.39	45	163	24.004	1.565	2.891
Abdominal skinfold (mm)	117.03	63.013	49	431	2.356	9.437
Lower leg skinfold (mm)	103.05	48.692	34	257	.797	.492

Legend: AS – arithmetic mean; SD – standard deviation; Min – minimum; Max – maximum; Sk – skewness, Ku – Kurtosis

Table 2 shows descriptive statistical indicators (arithmetic means and standard deviations) for individual pathological conative characteristics of junior volleyball players. Calculated average and variable values are consistent with those obtained in the earlier study, as shown by the study done by Stankovic et al (Stankovic *et al.*, 2013).

Table 2: Basic descriptive statistical parameters of cognitive (pathological) variables

Variables	M	SD
Cardiovascular conversion	13.01	8.613
Gastrointestinal conversion	.85	1.330
Inhibitory conversion	1.29	1.302
Hypochondria	.23	.796
Anxiety	.19	.682
Obsession	.30	.896
Hypersensitivity	.39	.899
Phobias	1.24	1.495
Impulsiveness	.60	.890
Aggression	.49	.839
Paranoia	1.51	1.493
Depression	1.17	.702

Table 3 shows basic descriptive statistical parameters of a set of variables for assessing normal conative characteristics of volleyball players aged from

16 to 17, where the calculated values represent measures of central tendency.

Table 3: Basic descriptive statistical parameters of normal conative variables (raw results)

Variables	M	SD
Affectia – sizia (A)	13.05	9.114
High intelligence – low intelligence (B)	6.96	1.985
Stronger ego – weaker ego (C)	15.92	3.786
Dominance – submissiveness (E)	14.784	3.992
Surgency – desurgency (F)	14.888	3.651
Stronger superego – weaker superego (G)	10.03	2.063
Boldness – timidity (H)	16.06	4.115
Tender-minded – premeditation (I)		
Protension (L)	9.876	3.006
Autia – imaginative	9.993	2.787
Shrewdness – artlessness (M)		
Shrewdness – artlessness (N)	11.04	2.443
Apprehensive (O)	9.982	2.899
Traditionalism – liberalism (Q ₁)	8.98	3.007
Self-sufficiency – group adherence (Q ₂)	9.99	2.826
High self-control – self conflict (Q ₃)	13.01	2.900
Tense – relaxed (Q ₄)	10.12	4.05

b) Canonical correlation analysis of examined variables

Based on the results Bartlett Chi-square test ($\chi^2 = 439.05$ a statistically significant function, with a canonical correlation of high intensity ($R_c = .77$, $p < .05$), 59% and the proportion of the mutual variance that explains maximum possible level of linear combinations of single isolated pair of canonical factors is presented in Table 4. The calculated values, with the error probability of 5% suggest that analyzed multivariate system of conative (normal) variables depends on the system of original anthropometric characteristics.

Table 4: Statistical significance of calculated pair of canonical factors

Canonical function	R_c	Rc^2	χ^2	p
1	.77	.59	439.05	.05

Legend: canonical correlation coefficient (R_c), canonical coefficient of determination (R), the value of Bartlett's χ^2 -test, level of significance, i.e. the proportion of error in reasoning (p)

To test the nature of the relations between original normal and pathological variables we applied model of canonical correlation analysis. In Table 5, the left set of variables represents variables in normal conative domain, and the right set of variables in pathological conative domain. Statistically significant coefficients of correlation between the tested variables

had values of -0.19 (negative correlation between normal conative factor affectia-sizia and pathological conative characteristics of depression) to 0.42 (positive correlation of normal conative factor tense-relaxed and pathological conative characteristics of inhibitory conversion).

Based on the analysis of the cross-correlation matrix (Table 6) we can note statistically significant correlations between normal and pathological conative characteristics for the following variables: affectia - sizia and depression ($p = -.19$), dominance - submissiveness and cardiovascular conversion ($r = .28$), dominance - submissiveness and gastrointestinal conversion ($r = .30$), dominance - submissiveness and aggression ($r = -.33$), surgency - desurgency and depression ($r = .40$),); stronger superego - weaker superego and depression ($r = .29$),); stronger superego - weaker superego and hypersensitivity ($r = .30$), boldness-shy and obsession ($r = .28$), protension and cardiovascular conversion ($r = .31$), shrewdness - artlessness and inhibitory conversion ($r = .32$), apprehensive and aggression ($r = -.33$), traditionalism-limeralism and hypochondria ($r = -.31$), self-sufficiency - group-adherence and obsession ($r = -.39$) high self-control-self-conflict and gastrointestinal conversion ($r = -.36$) and tense - relaxed and gastrointestinal conversion ($r = -.42$). Therefore, obtained results (with the probability level of $p < .05$) show low correlations between normal and pathological characteristics in conative space.

Table 5: Crosscorrelations between normal and pathological variables

	KK ₁₀	GG	I7	HH	AA ₁	OQ ₃	SS ₅	FF ₂	NN	TT ₁₅	PP ₁₈	DD ₆
		11		13					14			
A	.10	-.12	-.20	.03	-.01	-.07	-.08	-.11	.06	-.09	-.04	-.19
BB	.05	.02	.09	.05	-.10	.15	-.06	-.04	-.07	.09	.21	.01
CC	.09	.05	.02	.08	.04	.06	-.08	.05	.03	.02	.06	.05
EE	.28	.30	.32	.09	.40	-.06	.12	.15	.01	-.01	.05	.03
FF	-.09	.07	.16	-.05	-.12	.15	.06	-.09	.08	.14	-.02	.29
GG	.05	-.09	.07	.03	.08	.06	.30	.03	.06	.09	.07	.10
HH	.09	.13	.16	.03	.16	.28	-.08	-.04	-.02	.09	.00	.06
II	.11	-.05	-.10	-.16	-.14	.01	.00	-.08	.12	-.16	-.06	.13
LL	.31	.08	.15	.14	.16	.16	.05	.14	-.10	-.06	-.08	-.06
MM	.06	.03	-.06	-.07	.04	.16	-.00	.06	.07	.05	.03	.10
NN	.10	.16	.32	.14	.15	.06	.11	.14	.12	-.05	.01	.12
OO	-.03	.10	-.06	-.03	.02	-.08	.05	-.06	-.12	-.33	.04	.06
QQ ₁	-.08	-.12	-.05	-.31	.12	.16	.03	-.09	.16	-.14	.12	-.06
QQ ₂	-.08	-.05	-.07	-.00	.12	-.39	.07	.04	.15	-.06	.15	-.03
QQ1	-.03	.36	.16	.14	.03	-.02	.15	.07	.14	.13	.09	.05
3												
QQ ₄	.05	.10	.42	.13	.09	-.03	.15	.15	.14	.03	.05	-.04

Note: statistically significant correlations with p-value < .05 are presented in bold

Linear correlations obtained in this research are similar to those obtained in the research done by Stankovic et al., (2013).

By solving the characteristic equations of the cross-correlation matrix (Table 6), as the roots of this equation, two canonical factors were isolated. While studying the relations between the systems of normal conative variables and systems of pathological conative variables relatively high canonical correlations were determined for both isolated canonical factors ($R_c = .74$ and $R_c = .68$), and are statistically significant at the $p < .05$ level. The squares of the canonical correlation (R_c^2), which explain the common variance of the variables of the two groups of the overall variability of the analyzed systems of variables have a value of $R_c^2 = .51$, и $R_c^2 = .45$. Having considered significant intensity of canonical correlation and percent of mutual variance, it can be concluded, with the probability level

higher than 5%, that pathological conative characteristics of examiners were manifested more than normal conative characteristics.

Table 6 : Statistical significance of calculated pairs of canonical factors

Canonical function	R_c	R_c^2	χ^2	p
1	.74	.51	256.12	.05
2	.68	.45	201.26	.05

Legend: canonical correlation coefficient (R_c), canonical coefficient of determination (R_c^2), the value of Bartlett's χ^2 -test, level of significance, i.e. the proportion of error in reasoning (p)

The results presented in (Table 7) show mutual relations between individual morphological and normal conative variables and the first isolated canonical function.

Table 7 : Structure of canonical function of anthropometric and normal conative variables

Morphological variables	Canonical function 1	Normal conative factors	Canonical function 2
Body height	-.10	Affectia-cizia	-.38
Leg length	-.15	High intelligence-low intelligence	-.39
Hand length	.11	Higher ego-weaker ego	.09
Foot length	-.61	Dominance-submissiveness	-.18
Arm length	-.40	Surgency-desurgency	.05
Biacromial range	-.12	Higher super ego-weaker super ego	-.04
Bicristal range	.10	Bold (parmia)-shy	-.46
Hand width	.37	Tender-minded (premsia)-tough-minded	-.35
Wrist diameter	.33	Protension	-.48
Elbow diameter	.09	Autia (imaginative) -practical	.03
Knee diameter	.32	Shrewdness (phitmia) -artlessness	-.21
Foot width	.03	Apprehensive	-.01
Body mass	-.01	Traditionalism -liberalism	.11
Upper arm volume	.07	Self-sufficiency-group dependence	-.30
Lower arm volume	.13	High self-control-self conflict	-.04
Upper leg volume	.11	Tense-relaxed	-.37
Lower leg volume	.30		
Average thorax volume	.02		
Morphological variables	Canonical function 1	Normal conative factors	Canonical function 2
Upper arm skinfold	-.06		
Back skinfold	-.00		
Armpit skinfold	.08		
Abdominal skinfold	-.09		
Lower leg skinfold	-.04		

Coefficients within the set of anthropometric variables that show correlation between manifested variable with a canonical function, range from 0,30 to -0,61. *Foot length* (-.61) and *arm length* (-.40), show dominant negative correlations of low and moderate intensity with the first canonical factor, whereas *hand width* (.37), *wrist diameter* (.33) and *knee diameter* (.32), have positive correlation, whereas *lower leg volume* (.30)

is placed at the sixth position. Other anthropometric variables are not statistically significant for this canonical dimension. Having considered significant results, as well as the classification of morphological types in the research done by Misigoj-Durakovic (2008), this latent dimension can be hypothetically defined as a bipolar **ectomesomorphic morphological factor (longitudinal**

dimensionality of the skeleton and transverse dimensionality of the skeleton).

The second canonical function is presented in the same matrix of canonical structure. It involves seven normal conative variables. They range from 0,30 to 0,46 and show negative mutual linear correlation between original variable and isolated canonical function. "Most responsible" variables with negative and low correlation with this canonical function are: *protensia* (-.48) and *boldness (parmia)- shy* (-.46), *higher intelligence-lower intelligence* (-.39), *affectia-sizia* (-.38), *tense-relaxed* (-.37), whereas the results *tender-minded (premsia)-tough-minded* (-.35) and *self-sufficiency-group dependence* (-.30). are at the last position. The set of the second pair of normal conative factors cannot be precisely explained because of the combinations of complex and latent 'psychological nature'. However, obtained results show that it can be theoretically interpreted as a **factor of extrovert-introvert and anxiety**.

Obtained results presented in Table 8 show statistically significant and low linear correlation between analyzed normal and pathological variables of the first and second canonical function.

Coefficients that show mutual linear relations between normal conative variables and the first

canonical function range from 0,32 до -0,46. On the basis of the obtained results contained within the matrix of the canonical structure of normal and pathological conative variables, we can note statistically significant correlations between the applied variables of the first and second canonical factor. The first canonical factor in the space of normal conative variables was defined with the variables of *high intelligence - low intelligence* (-.46), *higher ego - lower ego* (-.43), *bold (parmia) - shy* (.41), *tender-minded (premsia) - tough-minded* (.37), whereas, *protension* is at the final position (.30). so that it could be defined as the canonical factor which characterizes volleyball players in relation to **intelligence, ego, boldness, tough-mindedness and protension**.

The second canonical function is defined by four normal conative variables with positive correlation coefficients ranging from 0,32 до 0,46. Most dominant variables are: *tense-relaxed* (.61), *self-sufficiency - group adherence* (.47) *high self-control - self-conflict* (-.46), whereas the result *shrewdness - artlessness* (.32) has the minimum influence on this function. Therefore, latent structure of junior volleyball players can be theoretically explained as a **factor of introspection and shrewdness - artlessness**.

Table 8 : The canonical structure of normal and pathological conative variables

	Normal conative factors		Pathological conative factor	
	<i>Fc₁</i>	<i>Fc₂</i>	<i>Fc₁</i>	<i>Fc₂</i>
Affectia-sizia	-.011	.06	Cardiovascular conversion	.43 -.05
High Intelligence-low intelligence	.32	-.23	Gastrointestinal conversion	.39 .23
Higher ego-weaker ego	.43	-.12	Inhibitory conversion	.35 .15
Dominance-submissiveness	.19	.10	Hypochondria	.38 .60
Surgency-desurgency	.03	-.07	Anxiety	.21 .18
Higher super ego-weaker super ego	-.06	-.03	Obsession	.14 .12
Bold (parmia)-shy	.41	-.20	Hypersensitivity	-.01 .14
Tender-minded (premsia)-tough-minded	.37	-.16	Phobias	.35 .59
Protension	.35	.18	Impulsiveness	-.28 .53
Autia (imaginative)-practical	.06	.17	Agression	.24 .12
Shrewdness (phitmia)-artlessness	-.05	.32	Paranoia	.33 -.01
Apprehensive	.30	-.15	Depression	-.05 .48
Traditionalism-liberalism	-.16	.06		
Self-sufficiency-group dependence	-.46	.47		
High self-control-self conflict	.20	.36		
Tense-relaxed	.20	.61		

The structure of the second canonical function is presented in the same matrix. It involves positive signs of four pathological conative variables. Their

coefficients range from 0,36 to 0,62. Most significant variables for this canonical function are: *obsession* (.62), *phobia* (.59), and *impulsiveness* (.53),, whereas

depression has the lowest contribution (.48). Therefore, latent structure of junior volleyball players can be theoretically labeled as a **factor of the regulation of reactions such as defense and attack**.

The results of this transversal study are similar to those obtained by the research done by Stankoci et al. (2013).

IV. DISCUSSIONS

Investigation and identification of potential mechanisms for the realization of morphological, normal and pathological conative dimensions in volleyball population is of great theoretical and practical importance. Success in performing sports activities depends on the personality dimensions governed by the modalities of its behavior. Researches done by the authors: Blažević, (2009), Kalach & Gontarev (2013), Murić *et al* (2012), Popović & Simonović (2008), Sindik *et al.*, (2011), Sindi i Nazor (2011), Stanković *et al.*, (2011), Stanković *et al.*, (2013), Stanković, Malacko & Doder (2011), Stankovića & Popovića (2011), Toskić *et al.*, (2012), have shown that conative characteristics directly and indirectly limit the effectiveness of different sports activities because of their impact on other anthropological characteristics. Also, the same researchers believe that in some sports activities identical conative characteristics represent limiting factor, while in others they have stimulating effect, and that there are no two persons with identical conative structures (Jurko *et al.*, 2013; Ivanovic & Ivanovic, 2011; Ivanovic & Ivanovic, 2012a; Ivanovic & Ivanovic, 2012b; Ivanovic & Ivanovic, 2012v; Ivanović & Ivanović, 2013, Ivanovic, Samardjic & Ivanovic, 2012, Stojanovic *et al.*, 2011). Therefore, the identification of conative factors represents an important assumption for efficient implementation of activities in volleyball.

For an explanation of canonically statistically significant relations between morphological and conative factors standard rule that a linear increase in the value of the resulting vector of variables of the first canonical factor from the first analyzed area corresponds to a proportional linear increase of the value of the resultant vector of the variables of canonical factors from another analyzed area and vice versa (Fajgelj, 2003) should be taken into account.

Based on the results contained in the matrix of the canonical structure of anthropometric and normal conative variables we can note a statistically significant correlation between the anthropometric variables and the first canonical factor. The relations between the first canonical factor from the system of anthropometric variables, interpreted as the bipolar canonical factor of the ectomesomorphic morphological type and the first canonical factor from the system of normal conative factors, interpreted as the canonical factor of extraversion – introversion and anxiety indicates that

volleyball players of the ectomesomorphic type are characterized at one end by longitudinal measures and on the other by transversal dimensionality of the volume of the femoral region with an increase in the value of the conative variables of extraversion and decreased anxiety.

On the basis of the results obtained and contained within the matrix of the canonical structure of normal and pathological conative variables, we can note statistically significant correlations between the measured variables in the case of the first and second canonical factor. The first canonical factor in the space of normal conative variables can be defined as the canonical factor which characterizes volleyball players in relation to intelligence, ego, boldness, tough mindedness and protension. The second one can be interpreted as a factor of introspection and tender-mindedness – tough-mindedness.

The first canonical factor in the space of pathological conative variables can be defined as a canonical factor of effectiveness of the system for the regulation and control of organ function and regulatory function, while the second canonical factor can be interpreted as a canonical factor of the regulator of the reactions of defense and attack. The structure of the correlations indicate that among the volleyball players with more pronounced warm types of behavior, who are more polite, considerate of others, there was an increase in the scores for depression, as with those who indicated more pronounced liveliness and impulsiveness and expressiveness. Volleyball players who usually manifest dominant, competitive, assertive or even aggressive behavior face a greater danger of developing one of the conversions (cardiovascular, gastrointestinal or inhibitory) and are more prone to anxiety. Those who are more moral individuals and who have more respect for the rules are more prone to hypersensitivity, and those who are shyless and adventurous score higher for obsession. Those who are suspicious and skeptical indicate a higher inclination towards cardiovascular conversion. The more agile, inquisitive, and discrete participants indicated a greater tendency for inhibitory conversion, anxiety, phobia and impulsive reactions. Those who were more worried, more insecure, often blaming themselves for everything were less prone to aggression. Those who were more open to change, more liberal, more analytical and critical volleyball players scored lower values for hypochondria, obsession and paranoia. The participants who were more self-satisfied and more self-confident were more prone to obsession, and those with a tendency for perfection, compulsion and self-discipline, who showed a tendency towards more dominant behavior with a lot of self-pity were at a higher risk of gastrointestinal and inhibitory conversion and were more impulsive. Tense, energetic, impatient volleyball players more frequently

develop gastrointestinal conversion, hypochondria and phobias.

Although the canonical results of this study section are expected and correspond with previous research studies, it is important to note several methodological shortcomings of this study. The biggest limitation is the relatively small size of the sample of examinees limited by age (16 and 17 year old), composed exclusively of male volleyball players. This structure of the sample greatly reduces the generalization of the results so that the obtained results are to be taken with some reserve. Realistic data could be obtained by experimental research in laboratory and real life conditions.

Finally, we should add that future longitudinal research should include other morphological and conative variables, an equal number of males and females in the population of volleyball players, without age limitation, which would apply the same or similar methodology, in order to get a greater probability of more reliable statistical conclusions about the complex relations of the examined phenomena.

V. CONCLUSIONS AND RECOMMENDATIONS

The results of an empirical research emphasize some important aspects of linear correlations between morphological, normal and pathological conative characteristics. Several conclusions are made according to these findings and their interpretation.

According to the results of applied canonical correlation analysis, this transversal study isolated following statistically significant factors: a) *ectomesomorfic morphological factor* (longitudinal dimensionality of the skeleton and transversal dimensionality of the skeleton), b) *normal conative factors* (extraversion-introversion and anxiety, intelligence, self-awareness, boldness, sharp temper and protensia; introspection and insight-honesty) and c) *pathological conative factors* (performance of the system for regulation and supervision of physical functions and automatic functions and editing of responses of defense and attack).

The results obtained in this study speak of the fact that volleyball players of the ectomesomorphic type are characterized at the one end, by longitudinal measures and at the other the dimensionality and volume of the femoral region if they are extrovert and non-anxious. In the case of volleyball players of this rank of competition, who show pronounced signs of warm behavior, who are more polite, considerate of others, there is an increase in the scores for depression, as in the case of those who indicate pronounced liveliness and impulsiveness and expressiveness. Volleyball players who frequently manifest dominant, competitive, assertive or even aggressive behavior are more prone to anxiety. Those who have higher morals

and more respect for the rules are more prone to hypersensitivity, and those who are shyless and adventurous score higher values on the score for obsession. Tense, energetic, impatient volleyball players more frequently develop gastrointestinal conversion, hypochondria and phobia.

Thus, the initial hypothesis of this transversal study on significantly linked set of anthropometric characteristics and a set of normal and pathological conative dimensions, is absolutely proven. The findings are partially consistent with previous studies of relations between morphological and conative (normal and pathologic) factors in our country and abroad.

Based on the results of relations, it is concluded that despite certain methodological limitations (section study design, relatively small and suitable sample, age limitation, mostly male examinees), our research can contribute to the development of predictive model of morphological, normal and pathological conative dimensions in volleyball junior population. Finally, additional longitudinal studies may provide more accurate results, which might significant at the theoretical level - in terms of further development of theoretical and methodological approach to these issues, and in the empirical sense, since statistically significant linear correlations between examined constructs could be noticed if greater population of both genders would be included.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Bandyopadhyay, A. (2007). Anthropometry and body composition in soccer and volleyball players in West Bengal, India. *Journal of Physiological Anthropology*, 26 (4), 501–505.
2. Blažević, S. (2006). *Relaciji pomeću specifičnite motorički sposobnosti i morfološke karakteristiki, bazičnite motorički sposobnosti, kognitivnite sposobnosti i konativnite karakteristiki kaj vrvnite bokseri*. Doktorska disertacija. Skopje: Fakultet za fizička kultura.
3. Blažević, S. (2009). Evaluation of boxer's conative status contribution to success of specific motor structures. *Sport Science*, 2 (2), 96–101.
4. Vrbik, I., Čižmek, A., i Gručić I. (2011). Morfološke karakteristike između igračkih pozicija kod vrhunskih rukometaša. *Hrvatski športskomedicinski vjesnik*, 26 (2), 94–99.
5. Dopsaj, M., Nešić G. & Čopić, N. (2010). The multicentroid position of the anthropomorphological profile of female volleyball players at different competitive levels. *Facta Universitatis – series: Physical Education and Sport*, 8 (1), 47–57.
6. Đurković T., Marelc, N., i Rešetar, T., (2011). Latent structure of anthropometric dimensions in elite croatian volleyball players. U D. Milanović i G.

- Sporiš (Ed.), *Intergrative power of kinesiology* (pp.500–504). Zagreb: Kineziološki fakultet.
7. Ivanović, M. (2007). *Odbojka na časovima fizičkog vaspitanja*. Valjevo: Valjevo-print.
 8. Ivanović, M., i Ivanović, U. (2011). Relations of psychological needs, life goals and happiness orientation of junior volleyball player's subjective benefit. *Rezime radova: III međunarodni naučni kongres „Antropološki aspekti sporta i fizičkog vaspitanja i rekreacije* (str. 26–27). Banja Luka: Fakultet fizičkog vaspitanja i sporta.
 9. Ivanović, M., i Ivanović, U. (2012a). Anthropometric structure – predictor of junior volleyball players' explosive strength. *Rezime radova: LI. kongres antropološkog društva Srbije* (str. 65). Niš: Antropološko društvo Srbije.
 10. Ivanović, M., i Ivanović, U. (2012b). Relations of junior volleyball player's value orientation and behavior model of social clashes. *Sportski logos*, 10 (18), 10–18.
 11. Ivanović, M., Samardžić, B., i Ivanović, U. (2012v). Aggression – criteria of junior volleyball players' attachment and self-esteem. *Zbornik radova: VIII međunarodna naučna konferencija „Menadžment u sportu“* (str. 144–155). Beograd: Alfa Univerzitet, Fakultet za menadžment u sportu.
 12. Ivanović, M., i Ivanović, U. (2013). Individual and family dimensions as emotional determinants of volleyball male and female cadets regulation. *Sportske nauke i zdravlje*, 3(1), 25–37.
 13. Jakovljević, S., Karalejić, M., i Lazarević, LJ. (2010). The latent structure of conative dimensions of elite senior and junior basketball players. *Facta Universitatis Physical Education and Sport*, 8 (1), 21–30.
 14. Joško, S., i Joško, V. (2011). Razlike ukupne situacione efikasnosti vrhunskog košarkaša u odnosu na konativne karakteristike. *Facta universitas-series: Physical Education and Sport*, 9 (1), 99–112.
 15. Jurko, D., Nešić, G., i Stojanović, T. (2013). Does precompetitive anxiety affect perceived competitive efficacy of volleyball players? *Facta Universitatis Series: Physical Education and Sport*, 11(1), 57–64.
 16. Kalach, R., & Gontarev, S. (2013). Motor skill, cognitive and conative variables on the performance of motor capacity by throwing off 100 combined punches at a heavy bag among boxers. *Research in Kinesiology Federation of the Sports Pedagogues of the Republic of Macedonia*, 41(2), 169–172.
 17. Karalejić, M., Jakovljević, S., i Lazarević, LJ. (2010). Latente structure of conative dimensions of elite senior and junior basketball players. *Facta Universitatis Series: Physical Education and Sport*, 8 (1), 21– 30.
 18. Cattell, R. B., Eber, H. W., & Tatsuoka, M. (1970). *Handbook for the Sixteen Personality Factor Questionnaire*. Champaign, ILL: Institute for Personality and Ability Testing.
 19. Malacko, J., i Stanković V. (2009). The latent structure of morphological characteristics of top handball, basketball and football players. *Sport Science*, 1 (2), 111–116
 20. Mala, L., Maly, T., Záhalka, F., & Bunc, V. (2010). The profile and comparison of body composition of elitefemale volleyball players. *International Journal of Fundamental and Applied Kinesiology*, 42 (1), 90–97.
 21. Marelić, N., Matković, B. R. i Antekolović, Lj. (2010). Antropološke karakteristike igračnika i treniranih odbojkaša. *Hrvatski Športskomedicinski Vjesnik*, 25 (1), 23–27.
 22. Martín-Matillas, M., Valadés, D., Hernández-Hernández, E., Olea-Serrano, F., Sjöström, M., Delgado-Fernández, M., & Ortega, F. B. (2014). Anthropometric, body composition and somatotype characteristics of elite female volleyball players from the highest Spanish league. *J Sports Sci.*, 32 (2), 137–48.
 23. Mišigoj-Duraković, M. (2008). *Kinantropologija-biološki aspekti tjelesnog vježbanja*. Zagreb: Kineziološki fakultet.
 24. Momirović, K. (1971). *Struktura i mjerenje patoloških konativnih faktora*. Zagreb, HR: Republički zavod za zapošljavanje.
 25. Nejić, D. Stanković, R. & Joksimović, A. (2009). Razlike u prostoru morfoloških karakteristika kod odbojkaša i fudbalera. *Glasnik Antropološkog društva Srbije*, 44, 191–199.
 26. Popović-Ilić T., Stanković V., Vitošević B., & Ilić S. (2010). Struktura morfoloških dimenzija vrhunskih sportista. *Praxis medica*, 38 (3-4), 63–70.
 27. Rodríguez-Ruiz, D., Rodríguez-Matoso, D., Quiroga, M. E., Sarmiento, S., Da Silva-Grigoletto, M. E., García-Manso, J. M. (2012). Study of extensor and flexor musculature in the knees of male and female volleyball players. *European Journal of Sport Science*, 12 (5), 399–407.
 28. Sing, A. B., & Rathore, V. S. (2013). Kinematic Factors of Off-Speed and Power Spike Techniques in Volleyball. *Journal of Education and Practice*, 4 (7), 112–118.
 29. Sindik, J. i Nazor, D. (2011). Differences in Conative Characteristics and Perceived Group Cohesion of the Basketball Players Playing in Different Positions in the Team, *Collegium Antropologicum*, 35(3), 895–904.
 30. Sindik, J., Nazor, D., i Vukosav, J. (2011). Correlation between the conative characteristics at top senior basketball players, *Sport Science* 4(1), 78–83.
 31. Stanković, V., & Malacko, J. (2008). Relations between systems of motor, cognitive and conative variables

32. of top-class handball players. *Kinesiologia Slovenica*, 14 (3), 33–43.
33. Stanković, V., Malacko, J., Bojić, I., i Ilić, J. (2013). Relacije morfoloških varijabli i normalnih i patoloških konativnih karakteristika rukometaša. *SportLogia*, 9(2), 89–99.
34. Stanković, V., Malacko, J., & Doder, D. (2011). Comparative analysis of factor structures of latent variables of pathological connate characteristics in case of handball players belonging to different competition ranks. *Acta Kinesiologica*, 5(2), 58–62.
35. Stanković, V., i Popović, D. (2011). Canonical relations between the system of motor and conative variables of female adolescents. *Proceedings*, 3, 29–37.
36. Stanković, V., i Popović, D. (2012). The relations between the motor and cognitive abilities of adolescent females. *Facta universitatis, Series: Physical Education and Sport*, 10(3), 211 – 219.
37. Stojanović, T., Milenkoski, J., i Stojanović, N. (2011). Konativne karakteristike odbojkaša različitog nivoa kvaliteta. *Sport i zdravlje*, 1, 67–71.
38. Sheppard, J. M., Cronin, J. B., Gabbett, T. J., McGuigan, M. R., Etxebarria, N., & Newton, R. U. (2008). Relative importance of strength, power and anthropometric measures to jump performance of elite volleyball players. *Journal of Strength and Conditioning Research*, 22(3), 758–765.
39. Toskić, D., Tomislav Okičić, T., i Stanković, V. (2012). Canonic relations of conative characteristics and success in water polo. *Acta Medica Medianae*, 51 (2), 15–18.
40. Trajković, N., Milanović, Z., Sporiš, G., & Radisavljević, M. (2011). Pozicijske razlike u tjelesnoj kompoziciji i izvodenju skokova mladih elitnih odbojkaša. *Acta Kinesiologica* 5(1), 62–66.
41. Tubić, T. (2010). *Psihologija i sport*. Novi Sad: Fakultet sporta i fizičkog vaspitanja.
42. Fajgelj, S. (2003). *Psihometrija – Metod i teorija psihološkog merenja*. Beograd: Centar za primenjenu psihologiju.
43. Cabral de Araújo Tónico, B. G., Cabral de Araújo Tónico, S., de Miranda, Dantas & Reis (2011). Discriminant effect of morphology and range of attack on the performance level of volleyball players. *Revista Brasileira de Cineantropometria & Desempenho Humano*, 13 (3), 223–229.
44. Šibila, M., & Pori, P. (2009). Position-related differences in selected morphological body characteristics of top-level handball players. *Collegium antropologicum*, 33(4), 1079–1086.
45. Warner, R. M. (2012). *Applied statistics: From bivariate through multivariate techniques* (2nd ed.). Thousand Oaks, CA: Sage Publications.