

University of Groningen

The time evolution of gaps in tidal streams in axisymmetric potentials

Koppelman, Helmer H.; Helmi, Amina

Published in:
 Proceedings of the International Astronomical Union

DOI:
[10.1017/S1743921317006135](https://doi.org/10.1017/S1743921317006135)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Koppelman, H. H., & Helmi, A. (2018). The time evolution of gaps in tidal streams in axisymmetric potentials. In A. Recio-Blanco, P. de Laverny, A. G. A. Brown, & T. Prusti (Eds.), *Proceedings of the International Astronomical Union: Astrometry and Astrophysics in the Gaia sky* (Vol. 12, pp. 229-230). Cambridge University Press. <https://doi.org/10.1017/S1743921317006135>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The time evolution of gaps in tidal streams in axisymmetric potentials

Helmer H. Koppelman and Amina Helmi

Kapteyn Instituut, University of Groningen,
P.O. Box 800, 9700 AV, Groningen, the Netherlands
email: h.h.koppelman@rug.nl

Abstract. Our goal is to understand the evolution and properties of gaps produced by dark matter subhalos in stellar tidal streams. Here we explore how gaps grow in spherical potentials in comparison to axisymmetric potentials. We develop a model that uses the divergence of two orbits, one on each side of the gap, to describe the size of the gap and how this varies with time and depends on the characteristics of the encounter with the dark subhalo. To this end we use a formalism based on action-angle variables.

Keywords. Galaxy: halo, Galaxy: structure, Galaxy: kinematics and dynamics

1. Introduction & Motivation

Λ CDM simulations typically predict thousands of satellites orbiting the Milky Way to be completely dark matter dominated (Springel *et al.* 2008). Possibly interesting objects to probe this population of (otherwise invisible) dark satellites are tidal streams. We thus aim to infer the properties of the dark matter satellites from the properties of the gaps formed as a dark subhalo comes sufficiently close to a tidal stream. Modeling the evolution of a gap as the divergence of two orbits has shown good agreement with N-body experiments carried out in a spherical host potential (Helmi & Koppelman, 2016). We present here the results obtained for a more realistic Stäckel Galactic potential, now including a disk and a halo component.

2. Model

Building on Helmi & Koppelman (2016) we model the size of the gap as the separation of two orbits A & B which correspond to particles in the stream that have been perturbed the most because of the interaction with the subhalo. To obtain the initial separation of these two orbits we use the impulse approximation (Erkal & Belokurov, 2015). By assuming a Plummer profile for the dark matter we can obtain analytic expressions for their separations in phase-space, ΔX_0 and ΔV_0 . This initial separation can be transformed to a separation in action-angle space with a matrix: $M_0 = \partial(\Theta, J)/\partial(X, V)$, which is the Jacobian of the coordinate transformation. In action-angle space, time evolution is simple: $\Delta\Theta(t) = \Delta\Theta_0 + \Omega(J)t$, where $\Omega(J)$ are the frequencies of motion, and $J(t) = J_0$. We encode this evolution in a matrix $\Omega' = \begin{bmatrix} I_3 & \partial\Omega/\partial\mathbf{J}t \\ [0.3em]0 & I_3 \end{bmatrix}$. At each timestep the separation vector in action-angle space is transformed back locally to Cartesian coordinates with matrix M_t^{-1} . For a schematic overview of the model see Fig.1a.




Figure 1: Panel (a): A schematic overview of the model. First we calculate the initial size of the gap from the impulse approximation, then we evolve this separation in time using actions and angles. Panel (b): Gaps growing in tidal streams. The colored lines show the predicted sizes of the gaps from our model, the solid black lines show the size measured in N-body experiments. The fitted dashed lines show the linear rate of growth of the size of the gap with time for these experiments run in a realistic (axisymmetric) Galactic potential.

3. Results & Conclusions

Fig. 1b shows the evolution in time of the four different gaps produced by encounters with different mass subhalos. The solid colored lines show the prediction for the size of the gap from our model, the solid black lines are the measured size of the gaps in the corresponding N-body experiments. The fitted dashed lines show that a linear fit to the growth of the gap size in time works well on average. The oscillations seen depend on the orbital phase of the gap in the stream.

Our model thus shows very nice agreement with gap sizes measured in the N-body experiments. The rate of growth is dependent on the properties of the dark matter subhalo, e.g. its mass and scale radius, on the geometry of the impact, and on the age of the stream. The latter two have not been varied for the results shown in Fig. 1b. The evolution in time of the size of a gap is thus linear, also in the case of an axisymmetric potential.

References

- Helmi, A. & Koppelman, H. H. 2016, *ApJ* (Letters), 828, L10
 Helmi, A. & White, S. D. M. 1999, *MNRAS*, 307, 3
 Erkal, D. & Belokurov, V. 2015, *MNRAS*, 450, 1
 Springel, V., Wang, J., Vogelsberger, M., *et al.* 2008, *MNRAS*, 391, 1685