



**UNIVERSIDAD PERUANA DE CIENCIAS APLICADAS**

**FACULTAD DE CIENCIAS DE LA SALUD**

**PROGRAMA ACADÉMICO DE MEDICINA**

**Associated factors, inequalities, and spatial distribution of the use of  
modern contraceptive methods among women of reproductive age in  
Peru: a population-based cross-sectional study**

**TESIS**

Para optar el título profesional de Médico Cirujano

**AUTORES**

Ana Lucia Diaz Alvites (ORCID 0000-0002-2020-5623)

Gonzalo Yrala Castillo (ORCID 0000-0003-4067-3712)

**ASESOR**

Munayco Escate, Cesar Vladimir (ORCID 0000-0001-7872-8913)

**Lima, 28 de diciembre del 2022**

## *DEDICATORIA*

*A nuestros padres, por su apoyo incondicional durante la carrera, y a nuestros maestros, por los conocimientos adquiridos a través de todos estos años.*

## AGRADECIMIENTOS

Agradecemos al Dr. Ali Al-Kassab Córdova y al Dr. Cesar Vladimir Munayco Escate, por aconsejarnos durante todo el proceso de esta investigación y por alentarnos a hacer de este proyecto, un excelente trabajo. También agradecer a la Universidad Peruana de Ciencias Aplicadas, por proveernos las mejores facilidades y recursos que nos permitieron desempeñarnos como buenos estudiantes e investigadores.

## CONTENIDO

- I. RESUMEN
- II. INTRODUCCIÓN
- III. MÉTODOS
- IV. RESULTADOS
- V. DISCUSIÓN
- VI. CONCLUSIONES
- VII. REFERENCIAS

## RESUMEN

**Antecedentes:** El uso de métodos anticonceptivos en el Perú se ha incrementado notablemente en las últimas décadas. Sin embargo, a pesar de los esfuerzos en lograr accesibilidad a métodos de planificación familiar, la utilización de métodos anticonceptivos modernos en el Perú se mantiene por debajo del promedio sudamericano. Este estudio tuvo como objetivo dilucidar los factores asociados al uso de anticonceptivos modernos, así como la presencia de desigualdades y la distribución espacial en mujeres peruanas de 15 a 49 años en 2019.

**Métodos:** Se realizó un análisis de datos secundarios utilizando información de la Encuesta Demográfica y de Salud del Perú 2019. Realizamos estadística descriptiva, análisis bivariado y regresión múltiple de Poisson. Las desigualdades se estimaron mediante curvas de concentración e índice de concentración normalizado de Erreygers. El análisis espacial incluyó un mapa coroplético, índice de Moran global, interpolación Kriging y estadística Getis-Ord-Gi\*.

**Resultados:** La prevalencia del uso de anticonceptivos modernos fue de 39,3% entre las mujeres peruanas en edad reproductiva. El uso de anticonceptivos modernos se asoció directamente con la juventud (RP 1,39), a mujeres que tuvieron su primera relación sexual antes de los 18 años (RP 1,41) y al estar casadas pero no conviviendo (RP 1,87). Además, hablar quechua o aimara (RP 0,87) y no tener hijos (RP 0,59) se asoció inversamente con la utilización de anticonceptivos modernos. Encontramos la presencia de desigualdades en el uso de métodos anticonceptivos (distribución pro-ricos), aunque la magnitud fue baja. El análisis espacial reveló la presencia de un patrón de distribución agrupado (Índice de Moran = 0,009); sin embargo, hubo heterogeneidad interdepartamental e intradepartamental en la prevalencia prevista del uso de anticonceptivos modernos. Además, se encontraron importantes puntos calientes y fríos en Perú.

**Conclusiones:** El uso de anticonceptivos modernos fue del 39,3% entre las mujeres peruanas en edad reproductiva. Esto se asoció con menor edad de la mujer, menor edad de la primera relación sexual, estar casada o conviviente, entre otros. No se encontró una desigualdad sustancial en el uso de anticonceptivos modernos. La prevalencia fue heterogénea a nivel intra e interdepartamental. Los departamentos ubicados en el sur, sureste y noreste tuvieron la prevalencia más baja. Por lo tanto, las barreras no financieras deben abordarse a través de esfuerzos multi e intersectoriales y continuar brindando anticonceptivos modernos de manera universal.

**Palabras clave:** Planificación Familiar; métodos anticonceptivos; Desigualdades en Salud; Análisis espacial; Encuestas Demográficas y de Salud; Perú

## ABSTRACT

**Background:** The use of contraceptive methods in Peru has remarkably increased in recent decades. Nevertheless, despite the completeness and accessibility of family planning methods, modern contraceptive methods utilization in Peru remains below the South American average. Thus, this study aimed to elucidate the factors associated with modern contraceptive use, as well as the presence of inequalities and the spatial distribution in Peruvian women aged 15–49 years in 2019.

**Methods:** A secondary data analysis was conducted using information from the 2019 Peruvian Demographic and Health Survey. We performed descriptive statistics, bivariate analysis, and Poisson multiple regression. Inequalities were estimated through concentration curves and Erreygers' normalized concentration index. Spatial analysis included choropleth map, Global Moran's I, Kriging interpolation and Getis-Ord-Gi\* statistic.

**Results:** The prevalence of modern contraceptive use was 39.3% among Peruvian women of reproductive age. Modern contraceptive use was directly associated with youth (aPR 1.39), women having their first sexual intercourse before the age of 18 (aPR 1.41), and being married but not together (aPR 1.87). In addition, speaking Quechua or Aymara (aPR 0.87) and having no children (aPR 0.59) were inversely associated with utilization of modern contraceptives. We found the presence of inequalities in the use of contraceptive methods (pro-rich distribution), although the magnitude was low. Spatial analysis unveiled the presence of a clustered distribution pattern (Moran's Index = 0,009); however, there was inter-departmental and intra-departmental heterogeneity in the predicted prevalence of the use of modern contraceptives. In addition, significant hot and cold spots were found in Peru.

**Conclusions:** The use of modern contraceptives was 39.3% among Peruvian women of reproductive age. This was associated with younger women's age, younger age at first sexual intercourse, being married or cohabitant, among others. No substantial inequality was found in modern contraceptive use. The prevalence was heterogeneous at the intra- and inter-departmental level. Those departments located in the south, south-east, and north-east had the lowest prevalence. Therefore, nonfinancial barriers must be tackled through multi- and cross-sectoral efforts and continue to universally provide modern contraceptives.

**Keywords:** Family Planning; Contraceptive Methods; Healthcare Inequalities; Spatial Analysis; Demographic and Health Surveys; Peru



# N°3176\_Associated factors, inequalities, and spatial distribution of the use of modern contraceptive methods among women of reproductive age in Peru: a population-based cross-sectional study

## INFORME DE ORIGINALIDAD

11%

INDICE DE SIMILITUD

10%

FUENTES DE INTERNET

10%

PUBLICACIONES

3%

TRABAJOS DEL ESTUDIANTE

## FUENTES PRIMARIAS

1	<a href="https://bmjopen.bmj.com">bmjopen.bmj.com</a> Fuente de Internet	3%
2	<a href="http://www.rrh.org.au">www.rrh.org.au</a> Fuente de Internet	2%
3	Ali Al-kassab-Córdova, Claudia Silva-Perez, Carolina Mendez-Guerra, Lucero Sangster-Carrasco et al. "Inequalities in infant vaccination coverage during the COVID-19 pandemic: a population-based study in Peru", <i>Vaccine</i> , 2022 Publicación	1%
4	Submitted to Washington University in St. Louis Trabajo del estudiante	1%
5	<a href="http://www.track20.org">www.track20.org</a> Fuente de Internet	1%

[pubmed.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov)

6	Fuente de Internet	1%
7	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Fuente de Internet	1%
8	<a href="http://repositorio.urp.edu.pe">repositorio.urp.edu.pe</a> Fuente de Internet	<1%
9	<a href="http://www.researchgate.net">www.researchgate.net</a> Fuente de Internet	<1%
10	Ayal Debie, Ayenew Molla Lakew, Koku Sisay Tamirat, Getasew Amare, Getayeneh Antehunegn Tesema. "Complete vaccination service utilization inequalities among children aged 12–23 months in Ethiopia: a multivariate decomposition analyses", International Journal for Equity in Health, 2020 Publicación	<1%
11	Cabieses, Baltica, Richard Cookson, Manuel Espinoza, Gillian Santorelli, and Iris Delgado. "Did Socioeconomic Inequality in Self-Reported Health in Chile Fall after the Equity-Based Healthcare Reform of 2005? A Concentration Index Decomposition Analysis", PLoS ONE, 2015. Publicación	<1%
12	David R. Soriano-Moreno, Anderson N. Soriano-Moreno, Angela Mejia-Bustamante, Cristhian A. Guerrero-Ramirez et al. "Factors	<1%

associated with highly effective contraceptive use among reproductive-age women in Peru: Evidence from a nationwide survey", European Journal of Obstetrics & Gynecology and Reproductive Biology, 2020

Publicación

13

[live.guttmacher.org](https://live.guttmacher.org)

Fuente de Internet

<1 %

14

Submitted to International Institute of Social Studies - Erasmus University Rotterdam

Trabajo del estudiante

<1 %

15

Laísa Rodrigues Moreira, Cauane Blumenberg, Beatriz Elena Caicedo Velasquez, Fernanda Ewerling et al. "The role of gender inequality and health expenditure on the coverage of demand for family planning satisfied by modern contraceptives: a multilevel analysis of cross-sectional studies in 14 LAC countries", The Lancet Regional Health - Americas, 2023

Publicación

<1 %

16

[fileservice.eea.comacloud.net](https://fileservice.eea.comacloud.net)

Fuente de Internet

<1 %

Excluir citas

Apagado

Exclude assignment

Activo

Excluir bibliografía

Activo

Exclude template

Excluir coincidencias < 20 words

## **BACKGROUND**

This article was originally published in Díaz-Alvites, A.L., Yrala-Castillo, G., Al-kassab-Córdova, A. et al. Associated factors, inequalities, and spatial distribution of the use of modern contraceptive methods among women of reproductive age in Peru: a population-based cross-sectional study. *BMC Public Health* 22, 2267 (2022). <https://doi.org/10.1186/s12889-022-14629-0>

Family planning (FP) is the health strategy that empowers individuals to decide whether to have children, how many they want, and the spacing between pregnancies (1). FP is possible through the use of contraceptive methods (CM). It encompasses a set of interventions that take place prior to the prescription and provision of CM, such as information dissemination, education, and counseling (2). In this way, individuals or couples can decide freely and responsibly about their reproduction.

A CM is any method, medication or device used to prevent pregnancy (3). Primarily, there are two types of major CM, viz., the traditional contraceptive methods (TCM) and modern contraceptive methods (MCM). TCMs use different signals from a woman's body to identify the times when she is most fertile so that couples may avoid sexual intercourse (4). TCM includes the rhythm method (periodic abstinence), withdrawal (coitus interruptus), fertility awareness-based methods, the lactational amenorrhea method and folk methods (5). A MCM defines as a product or medical procedure that interferes with reproduction from acts of sexual intercourse. MCM includes barrier methods, hormonal contraceptives and intrauterine device (6). Unfortunately, there is no consensus on the definition of MCM, thus the measurement of MCM differs between studies (6). For example, the United Nations Population Fund and the Guttmacher Institute (7) name

lactational amenorrhea as a traditional method, while the World Health Organization (WHO) and researchers with the Demographic and Health Surveys label it as a modern method (8). MCM were designed to permit complete sexual freedom so that couples may have sexual intercourse at any time. The job of the health provider is to inform and counsel individuals or couples about the CM options that they have, taking into account individual factors such as sociodemographic, religious, cultural and medical determinants (9). Nonetheless, the decision to use CM is at the discretion of the individual or couple.

Various studies have highlighted the association between certain sociodemographic factors and the use of CM, such as the woman's age, educational level, employment status, and socioeconomic status (10-15). Soriano-Moreno DR, et al. conducted a study in Peruvian women and they found that having one or more children and having health insured children were associated with the use of highly effective contraceptive methods (HECM) (16).

Globally, almost one in two women of reproductive age used a form of CM in 2019, a slight increase compared to previous decades. However, there remains a significant unmet demand and regional gaps persist (17). In Peru, the use of MCM has remarkably increased in recent decades (18,19). FP programs seek to reach the entire population under approaches of interculturality, comprehensiveness, gender equity and social inclusion. Nevertheless, despite the completeness and accessibility of FP methods due to the Ministry of Health (MINSA, from Spanish acronym) policies (2), MCM utilization in Peru remains below the South American average (68%) and the use of TCM remains high (20,21). In addition, the majority of Peruvian women reported having more children than

desired (22).

Access to CM is a human right (23,24). The United Nations aims to eliminate all the unmet demands for FP by 2030. Improvements in reproductive health, including voluntary FP, can bolster economies, contribute to sustainable development, and reduce pregnancy-related costs (25). To do so, it is essential to implement targeted strategies to reduce geographic and socioeconomic gaps in access. In Peru, there has been improvements in access and coverage of health services. However, significant inequalities remain that require resolution, especially in the most vulnerable populations (26). By elucidating the determinants of MCM use and its geographic pattern, policymakers would redirect their policies. Otherwise, the fertility rate may increase, which entails risks to the health of children and their mothers, undermines investment in human capital, dampens economic growth, and aggravates environmental threats (27). Therefore, we conducted this study to elucidate factors associated with the use of MCM, in addition to the magnitude of inequality and the spatial distribution among Peruvian women of reproductive age.

## **METHODS**

### ***Study design and data sources***

We conducted a secondary data analysis using information from the 2019 Peruvian Demographic and Health Survey (DHS). The DHS is annually conducted by the National Institute of Statistics and Informatics (INEI, from the Spanish acronym) of Peru. It has national, departmental, and area of residence representativeness. Administratively, the Peruvian territory is divided into 24 departments and one constitutional province, which are subsequently subdivided into provinces and districts. The survey design was

probabilistic, two-stage, balanced, stratified, independent, and self-weighted. The 2019 survey included 36,745 households, from which 35,522 individuals were interviewed (28). Although the Peruvian DHS collects information at the household level, it is mapped at the sampling cluster level.

### ***Selection criteria***

The DHS includes women aged 12–49 years. However, our study included only women of reproductive age. According to the World Health Organization (WHO), women of reproductive age are those aged between 15–49 years (29). Participants with incomplete data for the variables of interest were excluded.

### ***Outcome definition***

The outcome variable (use of MCM) was defined according to the WHO definition (30) and based on the reviewed literature (6,11,43,44) which considered MCM as an outcome variable. This allowed us to standardize that definition and thus achieving greater comparability to the study. It was categorized into MCM utilization and MCM non-utilization. MCM utilization included oral contraceptive pills, intrauterine device, injectables, female and male condoms, female and male sterilization, implants, lactational amenorrhea method, vaginal barrier methods, and emergency contraception pills. MCM non-utilization included traditional and folkloric methods, such as abstinence, periodic abstinence, and withdrawal, and no method use.

### ***Independent variables***

We included 11 independent categorical variables related to social determinants of inequality, which were selected based on an extensive literature review and the

availability of variables obtained in the DHS database. Age was divided into three groups: 15–19, 20–34, and 35–49 years. Natural region was categorized into coast, highlands, rainforest, and Metropolitan Lima. In addition, we included other sociodemographic variables such as residence area, marital status, education, employment status, language, wealth index, age at first sexual intercourse, number of children alive, and family members. DHS uses the variable "family members" to refer to the number of people currently living in the household. The cut-off point of 4 family members was regarding previous studies, this also let us achieve greater comparability to the study.

### ***Socioeconomic status***

The wealth index was used as a proxy variable to socioeconomic status. The DHS does not directly measure living standard (for instance income); it is a measurement of relative socioeconomic position of a household and is based on household data of ownership and housing characteristics. Subsequently, it is calculated through principal component analysis (31,32).

### ***Statistical analysis***

The DHS databases were downloaded from the "Microdatos" webpage of the INEI (33). Descriptive, bivariate, multiple regression, and inequality analyses were conducted using STATA version 16.0 (Stata Corporation, College Station, TX, US). Estimates were made by considering the complex design of the survey (strata, weights, and primary sampling units) through the svy module. P-values of  $< 0.05$  were considered to be significant, and confidence intervals were computed to 95% (95% CI).



Descriptive analysis was performed to obtain absolute and relative frequencies. The prevalence of MCM use was estimated at national and departmental levels. Bivariate analysis was performed to evaluate the prevalence of MCM use among independent variables for which fixed chi-square test was used. Chi-square test was fixed with Rao Scott fix. Prevalence ratios were estimated to evaluate the magnitude of association between independent variables and MCM use, crude (cPR), and adjusted (aPR). Consequently, the generalized linear model (glm) Poisson family log link function was used. We computed this model as it assumes adequate probability distribution, there is no numerical instability, variances are smaller, and PR is more interpretable than other measures of association (34,35).

### ***Concentration curve and index***

The lorenz and conindex commands were used for the analysis of inequalities (36,37). The magnitude of wealth inequality in use of MCM was estimated through concentration curves (CC) and concentration index (CI). CC represents the distribution of health among the cumulative proportions of a specific population classified according to their socioeconomic level: from the poorest to the richest. This curve has the distribution of women surveyed ordered from the lower to higher socioeconomic level on its X-axis and the health variable (in our case, the use of MCM) on its Y-axis. If the proportion of health was equally distributed among the population based on their income, a 45° diagonal would be generated, and the CI would equal zero. A deviation of the curve to either side indicates the existence of inequality. The separation of the curve from the diagonal generates an area under the curve (AUC), which will be the value of the CI. A positive value of the CI (curve below the diagonal) implies that inequality in access to health is more concentrated among the rich groups, and a negative CI value (curve above the

diagonal) implies greater inequality among poor groups (38). CI values close to zero represent the existence of very little inequality, whereas CI values close to +/- 1 indicate the existence of greater inequality (39). The greater the AUC (represented by CI), the greater the inequality.

Considering that MCM use is a binary variable, Erreygers' normalized concentration index (ECI) was used in our study instead of CI. This is mathematically depicted below.

$$ECI(h) = \frac{1}{n} \sum_{i=1}^n 4h_i(2R_i - 1)$$

Where: n represents the sample size,  $h_i$  is the binary outcome of interest for person  $i$  (with limit values of 0 and 1), and  $R_i$  is the individuals rank by wealth index. Weighted ECI standardizes the uncorrected index by adjusting the CI to allow for the bounded nature of the variable under study. Therefore, certain axiomatic properties for an inequality index (transfer, level independence, cardinal invariance, and mirror) are satisfied [37,40–42].

### ***Spatial analysis***

All spatial analyses were performed in ArcGIS version 10.8 (ESRI, Redlands, CA, US). A choropleth was plotted to represent the regional prevalence of MCM utilization. To evaluate the spatial autocorrelation of the outcome variable, Global Moran's I was calculated. It ranges from -1 to 1. A positive value implies a clustered pattern, a negative value implies a dispersed pattern, and zero implies a random pattern. In addition, we conducted ordinary Kriging interpolation analysis to predict the prevalence of MCM utilization in unsampled locations. Furthermore, we evaluated the presence of hot and cold spots through Getis-Ord- $G_i^*$  statistic.

## RESULTS

### *Characteristics of the study population*

A total of 33,311 women aged 15–49 years were included in the analysis. Their mean age was 31.29 years (SD: 9.99). Almost half of the participants were aged between 20–34 years (43.9%). The majority of participants were from Metropolitan Lima (43.2%) and lived in urban areas (82.6%), and over half were currently married or living with their partner (55.3%). The majority of participants had reached the secondary level education (44.9%), was employed (71.5%), spoke Spanish (94.5%) and had a middle wealth index (21.7%). The majority of women had experienced their first sexual intercourse before the age of 18 years (40.5%), had 1–2 living children (42.5%), and had  $\leq 4$  members in their family (54.4%) (**Table 1**).

**Table 1.** Descriptive and bivariate analysis of MCM use among Peruvian women of reproductive age

Population characteristics	Variable categories	Frequency n (%)	Modern contraceptive methods utilization		p-value*
			Yes (%)	No (%)	
Age	15–19	4 668 (15.7)	11.8	88.2	<0.001
	20–34	16 994 (43.9)	45.3	54.7	
	35–49	11 627 (40.4)	43.5	56.5	
Natural region	Coast	11 216 (27.9)	37.7	62.3	<0.001
	Highlands	10 005 (20.3)	35.6	64.4	
	Rainforest	6 370 (8.6)	44.7	55.3	
	Metropolitan Lima	5 698 (43.2)	41.0	59.0	
Residence area	Urban	23 859 (82.6)	39.6	60.4	0.1049
	Rural	9 430 (17.4)	38.0	62.0	
Marital status	Never married	7 418 (31.4)	15.0	85.0	<0.001
	Married/Cohabitant	21 907 (55.3)	55.6	44.4	
	Married but not together	3 964 (13.4)	29.4	70.6	
Education	No education/Primary	6 651 (15.9)	40.5	59.5	0.0043
	Secondary	15 671 (44.9)	37.5	62.5	
	Higher	10 967 (39.2)	41.0	59.0	


Employment status	Unemployed	10 416 (28.5)	36.7	63.3	0.0013
	Employed	22 873 (71.5)	40.4	59.6	
Language	Quechua/Aymara	2 649 (4.7)	31.6	68.4	0.0003
	Spanish	30 161 (94.5)	39.7	60.3	
	Other native and/or foreign languages	479 (0.8)	36.7	63.3	
Wealth index	Poorest	8 671 (16.5)	35.5	64.5	0.0104
	Poor	8 618 (20.2)	41.1	58.9	
	Middle	6 815 (21.7)	38.9	61.1	
	Rich	5 290 (21.1)	39.6	60.4	
	Richest	3 895 (20.5)	40.9	59.1	
Age at first sexual intercourse	Never had sexual intercourse	3 620 (15.5)	0.1	99.9	<0.001
	<18	15 919 (40.5)	50.4	49.6	
	18-24	12 249 (38.4)	44.6	55.4	
	≥25	1 501 (5.6)	31.9	68.1	
Number of children alive	Has no children	6 991 (33.9)	16.7	83.3	<0.001
	1-2	16 382 (42.5)	49.3	50.7	
	3-4	7 586 (18.6)	56.4	43.6	
	≥5	2 330 (5.0)	44.9	55.1	
Family members	≤4	15 581 (54.4)	38.3	61.7	0.0246
	>4	17 708 (45.7)	40.6	59.4	

\*Pearson's chi-squared test

In 2019, the prevalence of MCM use was 39.3% among Peruvian women of reproductive age. The most used MCM were injectables (32.3%), male condoms (27.5%), female sterilization (15.4%), and oral contraceptive pills (12.9%) (**Figure 1**).

**Figure 1.** Proportion of MCM use among Peruvian women of reproductive age

IDU: Intrauterine device. LAM: Lactation amenorrhea method. ECP: Emergency contraceptive pill.



### *Bivariate analysis*

**Table 1** shows the prevalence of MCM use according to each independent variable. MCM were used by 11.8% of women aged 15–19 years. In all the natural regions of Peru, the prevalence of MCM use was low (37.7% on the coast, 35.6% in the highlands, 44.7% in the rainforest, and 41% in Metropolitan Lima). Regarding marital status, the majority of women who used MCM were married or lived with their partner (55.6%). Regarding education, women with a higher degree of education primarily used MCM (41%). In addition, MCM utilization was higher among employed women (40.4%). Furthermore, those speaking Spanish used more MCM (39.7%), than those speaking Quechua/Aymara (31.6%). In addition, women who had a poor wealth index primarily used these methods (41.1%), and MCM was higher among those who had their first sexual intercourse before the age of 18 years (50.4%) and among women who had 3–4 living children (56.4%). Finally, women who had over four members in their family also had higher MCM use (40.6%). Aside from residence area, all of these variables showed statistically significant differences.

### *Multiple regression analysis*

We performed a multiple regression analysis to identify independent predictors of MCM use among Peruvian women of reproductive age. Being young (aged 15–19 years) was associated to a 39% greater chance (aPR 1.39; 95% CI: 1.20–1.59) of MCM utilization than older women (aged 35–49 years). Similarly, married women, or those who lived with their partner had 87% more likelihood of using MCM than women who were married but not together (aPR 1.87; 95% CI: 1.69–2.06). Having higher education was associated with 23% higher probability (aPR 1.23; 95% CI: 1.13–1.34) of using MCM, compared to having no education or primary. Other sociodemographic variables associated with higher MCM use were richest wealth index (aPR 1.33; 95% CI: 1.19–1.48) and having had their first sexual intercourse at < 18 years of age (aPR 1.41; 95% CI: 1.22–1.62). However, certain variables demonstrated a protective effect concerning the use of MCM. Women living in the highlands were 13% less likely (aPR 0.87; 95% CI: 0.82–0.93) to use MCM than those living in Metropolitan Lima. Being employed was associated with 0.93 times less likelihood (aPR 0.93; 95% CI: 0.89–0.98) of using MCM than being unemployed. Similarly, Quechua or Aymara speakers were 13% less likely (aPR 0.87; 95% CI: 0.79–0.95) to use MCM compared to those who speak Spanish. In addition, having 1–2 children alive was associated with 0.88 times less likelihood (aPR: 0.88; 95% CI: 0.79–0.98) of using MCM compared with having over 4 children (**Table 2**).

**Table 2.** Multiple regression analysis of MCM use among Peruvian women of reproductive age

Population characteristics	Crude analysis		Adjusted analysis	
	PR <sup>a</sup>	95% CI	PR <sup>b</sup>	95% CI
Age				
15 -19	0.27	0.24-0.32 <sup>§</sup>	1.39	1.20-1.59 <sup>§</sup>
20-34	1.04	0.99-1.10	1.30	1.23-1.37 <sup>§</sup>

35-49	Ref	Ref	Ref	Ref
Natural region				
Coast	0.92	0.86-0.98*	0.91	0.86-0.96 <sup>†</sup>
Highlands	0.87	0.81-0.93 <sup>§</sup>	0.87	0.82-0.93 <sup>§</sup>
Rainforest	1.09	1.02-1.17*	0.96	0.90-1.03
Metropolitan Lima	Ref	Ref	Ref	Ref
Residence area				
Urban	1.04	0.99-1.10	0.96	0.91-1.02
Rural	Ref	Ref	Ref	Ref
Marital status				
Never married	0.51	0.44-0.58 <sup>§</sup>	1.15	0.99-1.33
Married/Cohabitant	1.89	1.71-2.09 <sup>§</sup>	1.87	1.69-2.06 <sup>§</sup>
Married but not together	Ref	Ref	Ref	Ref
Education				
No education/Primary	Ref	Ref	Ref	Ref
Secondary	0.93	0.87-0.99*	1.12	1.05-1.19 <sup>†</sup>
Higher	1.01	0.94-1.09	1.23	1.13-1.34 <sup>§</sup>
Employment status				
Unemployed	Ref	Ref	Ref	Ref
Employed	1.10	1.04-1.17 <sup>§</sup>	0.93	0.89-0.98 <sup>†</sup>
Language				
Quechua/Aymara	0.80	0.72-0.88 <sup>§</sup>	0.87	0.79-0.95 <sup>†</sup>
Spanish	Ref	Ref	Ref	Ref
Other native and/or foreign languages	0.92	0.70-1.22	0.83	0.62-1.11
Wealth index				
Poorest	Ref	Ref	Ref	Ref
Poor	1.16	1.09-1.23 <sup>§</sup>	1.15	1.08-1.23 <sup>§</sup>
Middle	1.10	1.02-1.18 <sup>†</sup>	1.17	1.08-1.28 <sup>§</sup>
Rich	1.12	1.03-1.21 <sup>†</sup>	1.22	1.11-1.34 <sup>§</sup>
Richest	1.15	1.06-1.26 <sup>§</sup>	1.33	1.19-1.48 <sup>§</sup>
Age at first sexual intercourse				
Never had sexual intercourse	0.00	0.00-0.01 <sup>§</sup>	0.00	0.00-0.01 <sup>§</sup>
<18	1.58	1.36-1.83 <sup>§</sup>	1.41	1.22-1.62 <sup>§</sup>
18-24	1.40	1.20-1.62 <sup>§</sup>	1.29	1.12-1.49 <sup>§</sup>
≥25	Ref	Ref	Ref	Ref
Number of children alive				
Has no children	0.37	0.32-0.43 <sup>§</sup>	0.59	0.50-0.70 <sup>§</sup>
1-2	1.10	1.00-1.20*	0.88	0.79-0.98*
3-4	1.26	1.15-1.38 <sup>§</sup>	1.09	0.99-1.20


Family members	$\geq 5$	Ref	Ref	Ref	Ref
	$\leq 4$	Ref	Ref	Ref	Ref
	$> 4$	1.06	1.01-1.12*	1.01	0.96-1.07

<sup>a</sup>PR: crude prevalence ratio. <sup>b</sup>PR: adjusted prevalence ratio. \*p-value <0.05, †p-value<0.01, §p-value<0.001

### *Inequalities analysis*


The prevalence of MCM use indicated a pro-rich distribution among Peruvian women, albeit low in magnitude (ECI = 0.026). Moreover, the inequalities in MCM use were higher in rural areas (ECI = 0.079) than in urban areas (ECI = 0.015). Similarly, the concentration curve indicated that the distribution of MCM use was concentrated in rich households (**Figure 2 and Figure 3**).

**Figure 2.** Concentration curves of wealth-related inequalities for MCM use among Peruvian women of reproductive age



**Figure 3.** Concentration curves of wealth-related inequalities for MCM use among Peruvian women of reproductive age stratified by residence area






### *Spatial analysis*

The spatial distribution of MCM use in Peruvian women had a clustered pattern (Moran's Index=0,009, p-value<0.001) (**Figure 4**).

**Figure 4.** Spatial autocorrelation (Global Moran's I) of MCM use among Peruvian women of reproductive age

Given the z-score of 41.6693389136, there is a less than 1% likelihood that this clustered pattern could be the results of random chance.




The choropleth map represents the prevalence of MCM utilization at the departmental level. The departments with the highest prevalence of MCM use were Tumbes (50.7%), San Martín (48.4%), and Ucayali (46%). Puno (25.8%), Huancavelica (28.7%), and Tacna (33.1%) reported the lowest use of MCM (**Figure 5a**).

Kriging interpolation analysis depicts the predicted prevalence of MCM use. The prevalence increases from red (low prevalence) to green-colored (high prevalence) areas. Those departments located in the south, south-east, and north-east had the lowest predicted prevalence of use of MCM (**Figure 5b**).

Hot spot (Getis-Ord-Gi\*) analysis shows red and blue points, which represent a more intense clustering of high and low proportion of MCM use, respectively. A high proportion of MCM use was found in Tumbes, Lima, Ucayali, Amazonas, Moquegua, and San Martin regions. Meanwhile, a low proportion of use of MCM was found in Puno, Huancavelica, Cajamarca, Lambayeque, Tacna, and La Libertad (**Figure 5c**).

**Figure 5.** Spatial analysis of MCM use among Peruvian women of reproductive age

**5a.** Departmental prevalence of MCM. **5b.** Kriging interpolation. **5c.** Hot spot analysis of MCM



## DISCUSSION

### *Main findings*

Several sociodemographic factors were associated with MCM utilization, although the prevalence was low and with high variability between departments among Peruvian women of reproductive age. The strongest association was found with women's age, age at first sexual intercourse, marital status, and language, even after adjusting for multiple potential cofounders. Other associated variables were natural region, education, wealth index, employment status, and number of living children. Moreover, the most used MCM were injections, male condoms, and female sterilization. Regarding the CI, our study revealed the presence of inequalities in the use of MCM (pro-rich distribution), although

the magnitude was low. However, spatial analysis unveiled the presence of a clustered distribution pattern (albeit low in magnitude), but there was inter-departmental and intra-departmental heterogeneity in the predicted prevalence of the MCM use. In addition, we found significant hot and cold spots of MCM utilization across Peru.

### ***Comparison with previous studies***

Approximately two out of five Peruvian women of reproductive age use MCM. Although the prevalence has increased, it is below the South American average (68.2%) and that of other Latin American countries, such as El Salvador (66.8%), the Dominican Republic (67.1%), Nicaragua (68.8%), and Costa Rica (73.9%) (11,43,44). However, this may be due to the large indigenous population in Peru. A study reported that the use of TCM was higher in countries with larger indigenous populations while the use of MCM was lower (45), and Peru has the highest proportion of TCM use in Latin America (11). Other explanatory factors include limited access to FP due to geographic and language differences, or a different stage of the demographic transition.

Soriano-Moreno DR, et al. investigated the factors associated with the use of HECM among Peruvian women of reproductive age (16). They reported a prevalence of 29.9% concerning the use of HECM, whereas we reported a prevalence of MCM utilization of 39.3%, as our outcome definition included more FP methods. Similar associations were found in both studies. However, our study included a spatial and inequality analysis, and they did not include language, age at first sexual intercourse, and the number of family members as exposure variables. In addition, our outcome variable is broader as it encompasses more CM (beyond those that are highly effective). Finally, the database used in our study is more updated: at the end of 2017, the FP Technical Standard was

implemented in Peru, which could have significantly impacted the use of MCM (2).

### ***Factors associated with MCM use***

Centralization has been problematic in Peru for decades. Women living in the coast and highlands were less likely to use MCM than women living in Metropolitan Lima (the capital of Peru). The Peruvian health system is fragmented and segmented, with large gaps (46,47). For instance, Lima has the largest number of health facilities and physicians per inhabitant countrywide (48). Centralization fuels health inequalities, therefore, the deconcentration of health resources is necessary.

Younger women were more likely to use MCM, which may be due to the recent implementation of FP policies that include health guidance and counseling towards adolescents nationwide (49,50). However, a number of previous studies have established that older women are typically associated with greater use of MCM (51). Therefore, data regarding the prevalence of MCM utilization between age groups are controversial (15,52–55).

Education is essential to increase the use of MCM. Those women with a higher degree were more prone to use MCM, which aligns with several previous studies (10,15,56,57). Indeed, those with lower educational level are the most affected by existing inequalities in Peru (26).

Having their first sexual intercourse at below 18 years of age was associated with greater use of MCM. Similar results were found in Ethiopia (58). This could be because younger women are more sexually active than older women and are economically dependent (in

most cases). Economic independence is important because having children implies a significant financial burden. Interestingly, sex education (if provided before first experience of sexual intercourse), which is included in the Peruvian school's curriculum, protects youth from having sex at an early age (59).

Both marital status and number of children were associated with the use of MCM. Women who were married or lived with their partner had more chances to use MCM. This aligns with the results from a study from Uganda, which found that married adolescents were more likely to use MCM than unmarried adolescents. It would have been interesting to analyze the level of education of the partner and its influence on MCM use in our study, as undertaken by the study in Uganda (60). Nevertheless, this variable was excluded from our study as there were missing data in the Peruvian DHS.

Quechua and Aymara speakers were less likely to use MCM (compared with Spanish speakers). The majority of these speakers belong to indigenous ethnical populations, which maintain ancestral behaviors on specific territories (61). Quechua and Aymara are the primary Peruvian native languages; however, there are others such as Ashaninka, Awajun and Shipibo, etc. Furthermore, we believe that indigenous language speakers were experiencing a prior stage of the demographic transition; however, we did not find studies supporting this hypothesis. Likewise, speaking Quechua or Aymara was associated with a higher prevalence of mistreatment in health services (62). In addition, due to their customs they are prone to use TCM, which are ineffective (63). Although Spanish is the most spoken language in Peru, MINSA must guarantee access to FP information for native speakers. Its approach proposes interculturality, integrity, and social inclusion (2); however, this may not be enough.

### *Inequalities analysis*

The higher the wealth index, the higher the prevalence of MCM utilization. However, the inequality analysis, at a nationwide level, indicated the presence of inequalities (but in small magnitude). However, when this analysis was stratified by area of residence, we found that rural areas had a higher magnitude of inequalities, even more than at the national level. This may be because the public health care system is oversaturated and faces expenditure shortages, subsequently, there is a high out-of-pocket spending on FP, particularly in rural areas (64), where health resources are scarce. Moreover, the majority of people living in rural areas are indigenous, and they typically use TCM (28). Governmental social programs, such as the FP program, are of paramount importance for reducing inequality gaps of MCM utilization. Overall, the low magnitude of inequalities is the result of MINSA's continued efforts to universally provide MCM.

### *Spatial analysis*

Common sociodemographic factors may underlie the observed spatial patterns in the regions with the lowest MCM use as the spatial distribution was clustered. Huancavelica, Cajamarca, and Puno are among the 10 poorest departments in our country: Huancavelica is the poorest (65). In addition, Cajamarca and Puno are the least urbanized departments (66). Women from rural areas are less educated and the majority of their health facilities are remote and poorly equipped (67,68). Furthermore, the time to health-care facilities was estimated to be 5.3 times longer in rural settings than in urban settings (69). Differences in education are also indicated by the use of MCM. In fact, school attendance in Huancavelica, La Libertad, and Cajamarca is low (70). There was also intra-departmental heterogeneity in the use of MCM. These geographic disparities may also be attributed to multiculturalism.

### ***Implication for policy and research***

Several strategies must be implemented to improve Peruvian women's access and use of MCM. Although decentralization is challenging, it can be achieved through efficient allocation of health resources. Health facilities and providers should be placed in rural areas, especially in native communities. Besides, the multicultural approach should be continued, and joint work between social actors and health workers should be boosted.

Unfortunately, most MCM are obtained from private health care providers (64,71). Therefore, social health insurance must enlarge its coverage, which must be accompanied by the inclusion of comprehensive FP strategies. In addition, this should encompass the expansion of MCM options and the improvement of sex education programs and counseling services among sexually active people, particularly targeted at all young people. Sexual education policies and programs should be developed on the basis of evidence-based thesis based on modern adolescent development theories and ecological models (72). Furthermore, it is essential to ensure sufficient well-trained health providers nationwide.

The reduction of nonfinancial barriers is crucial. This could be remedied by establishing multi- and cross-sectoral efforts, such as the implementation of health centers in remote locations, improving of highways and roads, and improving FP services.

### ***Strengths and limitations***

This study has several limitations. First, the DHS did not specify whether all women included in its survey were sexually active at the time of the interview. As the Peruvian DHS collects information from women 12 years and over, we included only those women



who were of reproductive age. Second, we used a secondary database and thus had no data quality control. However, DHS interviewers received training courses and employed rigorous procedures for data quality control. Third, as it is a secondary database, there were interesting variables regarding the partner or the family that were excluded in the DHS. Fourth, due to the cross-sectional design, causality cannot be determined. Fifth, although the Peruvian DHS lacks a direct measure of socioeconomic status, we used an asset-based wealth index as a proxy variable, which is suitable for inequality studies in the absence of a direct measure (73). However, to the best of our knowledge, this is the first study assessing inequalities and spatial distribution in the use of MCM among Peruvian women. Our results are derived from a large sample size, which implies a great statistical power and representativeness at the national level. In addition, the standardized definition of our outcome allows us to compare our results with other studies. Besides, we calculated ECIs, which satisfy some shortcomings of the traditional CI (40).

## **CONCLUSION**

The use of MCM was 39.3% among Peruvian women of reproductive age. The use of MCM was directly associated with younger women's age, younger age at first sexual intercourse, and being married or cohabitant, among other factors. However, it was inversely associated among those speaking Quechua or Aymara. No substantial inequality was found in MCM utilization at national level; however, it was higher in rural areas. The prevalence of use of MCM was heterogeneous at the intra- and inter-departmental level. Those departments located in the south, south-east, and north-east had the lowest prevalence of MCM utilization. Therefore, it is paramount to tackle nonfinancial barriers through multi- and cross-sectoral efforts and continue to universally provide MCM.

## LIST OF ABBREVIATIONS

FP: Family planning

CM: Contraceptive methods

TCM: Traditional contraceptive methods

MCM: Modern contraceptive methods

HECM: Highly effective contraceptive methods

MINSA: Ministry of Health

DHS: Demographic and Health Survey

INEI: National Institute of Statistics and Informatics

WHO: World Health Organization

CC: Concentration curve

CI: Concentration index

AUC: Area under the curve

ECI: Erreygers' normalized concentration index

95% CI: 95% Confidence interval

## REFERENCES

1. Shaw D. The ABC's of family planning [Internet]. The Partnership for Maternal, Newborn & Child Health. [citado 21 de junio 2022]. Disponible en: [https://www.who.int/pmnch/media/news/2010/20100322\\_d\\_shaw\\_oped/en/](https://www.who.int/pmnch/media/news/2010/20100322_d_shaw_oped/en/)
2. Plataforma digital única del Estado Peruano [Internet]. Lima: Biblioteca Central del Ministerio de Salud; 2017 [citado 22 de junio 2022]. Disponible en: <https://www.gob.pe/institucion/minsa/informes-publicaciones/280860-norma-tecnica-de-salud-de-planificacion-familiar>
3. Oficina para la Salud de la Mujer (OASH) [Internet]. EE.UU: Oficina para la Salud de la Mujer en el Departamento de Salud y Servicios Humanos de EE. UU; 2017 [citado 22 de junio 2022]. Disponible en:

<https://espanol.womenshealth.gov/a-z-topics/birth-control-methods>

4. Marie Stopes Timor-Leste. Traditional family planning [Internet]. Marie Stopes Timor-Leste [citado 21 de junio 2022]. Disponible en: <https://www.mariestopes.tl/services/traditional-family-planning/>
5. Almalik M, Mosleh S, Almasarweh I. Are users of modern and traditional contraceptive methods in Jordan different? East Mediterr Heal J. 2018;24(4):377–84.
6. Hubacher D, Trussell J. A Definition of Modern Contraceptive Methods. Contraception. 2015;92:420–1.
7. Guttmacher Institute [Internet]. Guttmacher Institute, United Nations Population Fund (UNFPA); 2014 [citado 24 de diciembre 2022]. Disponible en: <https://www.guttmacher.org/report/adding-it-costs-and-benefits-investing-sexual-and-reproductive-health-2014>
8. World Health Organization (WHO) [Internet]. WHO; 2020 [citado 24 de diciembre 2022]. Disponible en: <https://www.who.int/news-room/fact-sheets/detail/family-planning-contraception>
9. Mayo Clinic [Internet]. EE.UU: Mayo Clinic; 2022 [citado 24 de diciembre 2022]. Disponible en: <https://www.mayoclinic.org/healthy-lifestyle/birth-control/in-depth/birth-control-options/art-20045571>
10. Çalikoğlu EO, Yerli EB, Kavuncuoğlu D, Yılmaz S, Koşan Z, Aras A. Use of family planning methods and influencing factors among women in Erzurum. Medical Science Monitor. 2018;24:5027–34.
11. Ponce de Leon RG, Ewerling F, Serruya SJ, Silveira MF, Sanhueza A, Moazzam A, et al. Contraceptive use in Latin America and the Caribbean with a focus on long-acting reversible contraceptives: prevalence and inequalities in 23 countries. Lancet Glob Health. 2019;7:e227-235.

12. Li Z, Patton G, Sabet F, Zhou Z, Subramanian S v., Lu C. Contraceptive Use in Adolescent Girls and Adult Women in Low- and Middle-Income Countries. *JAMA Netw Open*. 2020;3:e1921437.
13. Asif MF, Pervaiz Z. Socio-demographic determinants of unmet need for family planning among married women in Pakistan. *BMC Public Health*. 2019;19:1–8.
14. Munakampe MN, Zulu JM, Michelo C. Contraception and abortion knowledge, attitudes and practices among adolescents from low and middle-income countries: A systematic review. *BMC Health Serv Res*. 2018;18:1–13.
15. Al Kindi RM, Al Sumri HH. Prevalence and sociodemographic determinants of contraceptive use among women in oman. *Eastern Mediterranean Health Journal*. 2019;25:495–502.
16. Soriano-Moreno DR, Soriano-Moreno AN, Mejia-Bustamante A, Guerrero-Ramirez CA, Toro-Huamanchumo CJ. Factors associated with highly effective contraceptive use among reproductive-age women in Peru: Evidence from a nationwide survey. *European Journal of Obstetrics and Gynecology and Reproductive Biology*. 2020;245:114–20.
17. United Nations Department of Economic and Social Affairs. World Fertility and Family Planning 2020: Highlights [Internet]. New York: United Nations Publication; 2020 [citado 24 de diciembre 2022]. Disponible en: <https://www.unlibrary.org/content/books/9789210043694>
18. Instituto Nacional de Estadística e Informática (INEI) [Internet]. Perú: INEI; 2016 [citado 24 de diciembre 2022]. Disponible en: [https://www.inei.gob.pe/media/MenuRecursivo/publicaciones\\_digitales/Est/Lib1356/](https://www.inei.gob.pe/media/MenuRecursivo/publicaciones_digitales/Est/Lib1356/)
19. United States Agency for International Development (USAID) [Internet]. EE.UU:

- USAID; 2016 [citado 24 de diciembre 2022]. Disponible en: <https://www.usaid.gov/sites/default/files/documents/1864/peru-508.pdf>
20. Department of Economic and Social Affairs, Population Division [Internet]. New York: United Nations; 2015 [citado 24 de diciembre 2022]. Disponible en: [https://www.un.org/development/desa/pd/sites/www.un.org.development.desa.pd/files/undesapd\\_report\\_2015\\_trends\\_contraceptive\\_use.pdf](https://www.un.org/development/desa/pd/sites/www.un.org.development.desa.pd/files/undesapd_report_2015_trends_contraceptive_use.pdf)
21. Health Policy Plus (HP+) [Internet]. EE.UU: HP+; 2016 [citado 24 de diciembre 2022]. Disponible en: [http://www.healthpolicyplus.com/ns/pubs/2068-2101\\_HPSFIBriefPeruEnglish.pdf](http://www.healthpolicyplus.com/ns/pubs/2068-2101_HPSFIBriefPeruEnglish.pdf)
22. Mendez-Guerra C, Al-Kassab-Córdova A, Cornejo-Venegas G, Quevedo-Ramirez A, Alva K, Napanga-Saldaña E. Factors associated with unwanted additional fertility: secondary analysis of a national survey in Peru. *Rev Peru Med Exp Salud Publica*. 2020;37:645–53.
23. International Federation of Gynecology and Obstetrics (FIGO). FIGO; 10 de julio 2018 [citado 21 de diciembre 2022]. Disponible en: <https://www.who.int/news/family-planning-human-right>
24. United Nations Population Fund (UNFPA). UNFPA; 2021 [citado 21 de diciembre 2022]. Disponible en: <https://www.unfpa.org/family-planning>
25. United Nations Population Fund (UNFPA). Turkey: UNFPA; 11 de julio 2017 [citado 21 de junio 2022]. Disponible en: <https://turkey.unfpa.org/en/news/world-population-day-2017>
26. Organización Panamericana de la Salud (OPS). Equidad en la atención en salud en Perú 2005 – 2015 [Internet]. Lima: OPS; 2017 [citado 23 de junio 2022]. Disponible en: <https://iris.paho.org/handle/10665.2/34357>
27. The World Bank. Determinants and Consequences of High Fertility: A Synopsis

- of the Evidence [Internet]. Washington, DC: The World Bank; 2010 [citado 25 de octubre 2022]. Disponible en: <https://openknowledge.worldbank.org/handle/10986/27497>
28. Instituto Nacional de Estadística e Informática (INEI) [Internet]. Perú: INEI; 2020 [citado 23 de enero 2022]. Disponible en: [https://www.inei.gob.pe/media/MenuRecursivo/publicaciones\\_digitales/Est/Endes2019/](https://www.inei.gob.pe/media/MenuRecursivo/publicaciones_digitales/Est/Endes2019/)
29. World Health Organization (WHO). WHO; 2022 [citado 24 de mayo 2022]. Disponible en: [https://www.who.int/data/gho/indicator-metadata-registry/imr-details/women-of-reproductive-age-\(15-49-years\)-population-\(thousands\)](https://www.who.int/data/gho/indicator-metadata-registry/imr-details/women-of-reproductive-age-(15-49-years)-population-(thousands))
30. World Health Organization (WHO). WHO; 2022 [citado 22 de enero 2022]. Disponible en: <https://www.who.int/data/gho/indicator-metadata-registry/imr-details/3334>
31. The Demographic and Health Surveys Program. Riccardo Gangale USAID; 2016 [citado 24 de mayo 2022]. Disponible en: <https://dhsprogram.com/topics/wealth-index/Wealth-Index-Construction.cfm>
32. Rutstein SO, Johnson K. DHS Comparative Reports No. 6: The DHS Wealth Index [Internet]. Maryland, USA: ORC Macro; 2004 [citado 22 de enero 2022]. Disponible en: <https://www.dhsprogram.com/publications/publication-cr6-comparative-reports.cfm>
33. Instituto Nacional de Estadística e Informática (INEI) [Internet]. Perú: INEI; [citado 22 de enero 2022]. Disponible en: <https://proyectos.inei.gob.pe/microdatos/>
34. Lee J, Tan CS, Chia KS. A practical guide for multivariate analysis of dichotomous outcomes. Ann Acad Med Singap. 2009;38:714–9.

35. Bastos LS, Oliveira R de VC de, Velasque L de S. Obtaining adjusted prevalence ratios from logistic regression models in cross-sectional studies. *Cad Saude Publica*. 2015;31:487–95.
36. Jann B. Estimating Lorenz and concentration curves. *Stata Journal*. 2016;16:837–66.
37. O'Donnell O, O'Neill S, Van Ourti T, Walsh B. coindex: Estimation of concentration indices. *Stata Journal*. 2016;16:112–38.
38. Ahmed S, Tariqujjaman M, Rahman MA, Hasan MZ, Hasan MM. Inequalities in the prevalence of undiagnosed hypertension among Bangladeshi adults: Evidence from a nationwide survey. *Int J Equity Health*. 2019;18:1–12.
39. Schneider MC, Castillo-Salgado C, Bacallao J, Loyola E, Mujica OJ, Vidaurre M, et al. Métodos de medición de las desigualdades de salud. *Revista Panamericana de Salud Publica/Pan American Journal of Public Health*. 2002;12:398–415.
40. Erreygers G. Correcting the Concentration Index. *J Health Econ*. 2009;28:504–15.
41. Erreygers G, Van Ourti T. Measuring socioeconomic inequality in health, health care and health financing by means of rank-dependent indices: A recipe for good practice. *J Health Econ*. 2011;30:685–94.
42. Kjellsson G, Gerdtham UG. On correcting the concentration index for binary variables. *J Health Econ*. 2013;32:659–70.
43. Fagan T, Dutta A, Rosen J, Olivetti A, Klein K. Family planning in the context of Latin America's universal health coverage Agenda. *Glob Health Sci Pract*. 2017;5:382–98.
44. United Nations (UN). *Trends in Contraceptive Use Worldwide 2015*. New York; 2016.

45. Fagan T, Dutta A, Rosen J, Olivetti A, Klein K. Family planning in the context of Latin America's universal health coverage Agenda. *Glob Health Sci Pract.* 2017;5:382–98.
46. Alcalde-rabanal JE, Lazo-gonzález O, Nigenda G, Je A, Nigenda G. Sistema de salud de Perú. *Salud Publica Mex.* 2011;53:243–54.
47. Sánchez-Moreno F. El sistema nacional de salud en el Perú. *Rev Peru Med Exp Salud Publica.* 2014;31:747–53.
48. Instituto Nacional de Estadística e Informática (INEI) [Internet]. Perú: INEI; 2020 [citado 26 de enero 2022]. Disponible en: <https://www.inei.gob.pe/estadisticas/indice-tematico/health/>
49. Gobierno del Perú [Internet]. Perú; 2021 [citado 8 de febrero 2022]. Disponible en: <https://www.gob.pe/15121-anticoncepcion-en-la-adolescencia>
50. Ministerio de Salud del Perú, Dirección General de Intervenciones Estratégicas en Salud Pública. Norma técnica de salud para la atención integral de salud de adolescentes 2019 [Internet]. Perú: MINSA; 25 de octubre 2019 [citado 8 de febrero 2022]. Disponible en: <http://bvs.minsa.gob.pe/local/MINSA/4942.pdf>
51. Al Kindi RM, al Sumri HH. Prevalence and sociodemographic determinants of contraceptive use among women in oman. *Eastern Mediterranean Health Journal.* 2019;25:495–502.
52. Lun CN, Aung T, Mya KS. Utilization of modern contraceptive methods and its determinants among youth in Myanmar: Analysis of Myanmar Demographic and Health Survey (2015-2016). *PLoS One* [Internet]. 2021 [citado 21 de enero 2022];16(10):19. Disponible en: <http://dx.doi.org/10.1371/journal.pone.0258142>
53. Nkunzimana E, Sufiyan Babale M, Ndoreraho A, Nyandwi J. Uptake of Modern Contraceptive Methods among Burundian Women and Associated Factors:



- Analysis of Demographic and Health Survey Data, Burundi 2016–2017. *East African Health Research Journal*. 2021;5:75–81.
54. Adokiya MN, Boah M, Adampah T. Women's autonomy and modern contraceptive use in Ghana: a secondary analysis of data from the 2014 Ghana Demographic and Health Survey. *European Journal of Contraception and Reproductive Health Care*. 2021;26:383–9.
55. Bakibinga P, Matanda D, Kisia L, Mutombo N. Factors associated with use of injectables, long-acting and permanent contraceptive methods (iLAPMs) among married women in Zambia: Analysis of demographic and health surveys, 1992-2014. *Reprod Health*. 2019;16:1–12.
56. Saurina C, Vall-Llosera L, Saez M. Factors determining family planning in Catalonia. Sources of inequity. *Int J Equity Health*. 2012;11:1–8.
57. Ponce de Leon RG, Ewerling F, Serruya SJ, Silveira MF, Sanhueza A, Moazzam A, et al. Contraceptive use in Latin America and the Caribbean with a focus on long-acting reversible contraceptives: prevalence and inequalities in 23 countries. *Lancet Glob Health*. 2019;7:e227-235.
58. Gebre MN, Edossa ZK. Modern contraceptive utilization and associated factors among reproductive-age women in Ethiopia: Evidence from 2016 Ethiopia demographic and health survey. *BMC Womens Health*. 2020;20:1–14.
59. Mueller TE, Gavin LE, Kulkarni A. The Association Between Sex Education and Youth's Engagement in Sexual Intercourse, Age at First Intercourse, and Birth Control Use at First Sex. *Journal of Adolescent Health*. 2008;42:89–96.
60. Sserwanja Q, Musaba MW, Mukunya D. Prevalence and factors associated with modern contraceptives utilization among female adolescents in Uganda. *BMC Womens Health*. 2020;21:1–7.

61. Montenegro RA, Stephens C. Indigenous health in Latin America and the Caribbean. *Lancet*. 2006;367:1859–69.
62. Herrera-Añazco P, Benites-Zapata VA, Hernández A. Association between the Non-use of Health Services and Maltreatment Based on Ethnicity in Peru. *J Health Care Poor Underserved*. 2022;33:234–52.
63. Hassoun D. Methodes de contraception naturelle et methodes barrieres. *RPC contraception CNGOF. Gynecol Obstet Fertil Senol*. 2018;46(12):873–82.
64. Flachsbarth I, Schotte S, Lay J, Garrido A. Rural structural change, poverty and income distribution: evidence from Peru. *J Econ Inequal*. 2018;16:631–53.
65. Instituto Peruano de Economía [Internet]. Perú: INEI; 2021 [citado 22 de mayo 2022]. Disponible en: <https://www.ipe.org.pe/portal/pobreza-2020-el-peru-retrocede-10-anos/>
66. Instituto Nacional de Estadística e Informática (INEI) [Internet]. Perú: INEI; 2018 [citado 22 de mayo 2022]. Disponible en: [https://www.inei.gob.pe/media/MenuRecursivo/publicaciones\\_digitales/Est/Lib1539/](https://www.inei.gob.pe/media/MenuRecursivo/publicaciones_digitales/Est/Lib1539/)
67. Kandala NB, Lukumu FK, Mantempa JN, Kandala JD, Chirwa T. Disparities in modern contraception use among women in the democratic Republic of Congo: A cross-sectional spatial analysis of provincial variations based on household survey data. *J Biosoc Sci*. 2015;47:345–62.
68. Mathe JK, Kasonia KK, Maliro AK. Barriers to Adoption of Family Planning among Women in Eastern Democratic Republic of Congo. *Afr. J. Reprod. Health*. 2011 March;15(1):69-77.
69. Carrasco-Escobar G, Manrique E, Tello-Lizarraga K, Miranda JJ. Travel Time to Health Facilities as a Marker of Geographical Accessibility Across

- Heterogeneous Land Coverage in Peru. *Front Public Health*. 2020;8:1–10.
70. Instituto Nacional de Estadística e Informática (INEI). Perú: Indicadores de Educación por Departamentos, 2009-2019 [Internet]. Perú: INEI; 2020 [citado 22 de mayo 2022]. Disponible en: [https://www.inei.gob.pe/media/MenuRecursivo/publicaciones\\_digitales/Est/Lib1751/libro.pdf](https://www.inei.gob.pe/media/MenuRecursivo/publicaciones_digitales/Est/Lib1751/libro.pdf)
71. Class D, Cavagnero E, AS Rajkumar, Ferl K. Peru - Health financing profile (English) [Internet]. Washington, D.C.: The World Bank Group; 2015 [citado 22 de mayo 2022]. Disponible en: <http://documents.worldbank.org/curated/en/242851468099278957/Peru-Health-financing-profile>
72. Leung H, Shek DTL, Leung E, Shek EYW. Development of Contextually-relevant Sexuality Education: Lessons from a Comprehensive Review of Adolescent Sexuality Education Across Cultures. *Int J Environ Res Public Health*. 2019;16.
73. McKenzie DJ. Measuring inequality with asset indicators. *J Popul Econ*. 2005;18:229–60.