

DNA Barcoding for Selected Mangrove-Based Estuary Fishes from Way Kambas National Park, Lampung Province, Indonesia

Yanti Ariyanti*, Ika Agus Rini, Indah Oktaviani, Sovia Santi Leksikowati

Department of Biology, Institut Teknologi Sumatera, Lampung 35365, Indonesia

Article history:

Submission November 2020

Revised January 2021

Accepted March 2021

*Corresponding author:

E-mail: yanti.ariyanti@bi.itera.ac.id

ABSTRACT

Over the past decade, DNA barcoding has provided new insight into fish ecology and biosystematics and led to new species' discovery. DNA barcoding is a method for the recognition and identification of species using short, standardised DNA fragments. The correct taxonomic identification of species is critical for the assessment and monitoring of biodiversity. This study applied DNA barcoding techniques to identify selected fish species from a mangrove-based estuary in Way Kambas National Park, Lampung Province, Indonesia. The gene encoding cytochrome c oxidase subunit I (COI) was amplified and bi-directionally sequenced from 22 specimens. The resulting 680 base pairs (bp) sequence was used to identify species, obtain phylogenetic information, and analyse genetic distances. A neighbour-joining tree was constructed based on the mitochondrial COI gene using the Kimura two-parameter model. This study also exhibits conservation status for those identified species. Our findings will facilitate future studies of fish species diversity in mangrove estuary-based ecosystems and provide preliminary data in policymaking in conservation areas such as National Park.

Keywords: Biodiversity, COI, Estuarine, Genetic source, Mangrove

Introduction

DNA barcoding remain a practical tool for species identification due to its rapid, accuracy, cost-effective features, and functionality [1, 2]. DNA barcodes using a standardized DNA region that allows objective and universal comparable results, which can quickly be repeated even by non-taxonomist specialists [3]. Furthermore, this method can analyze inadequate, fragmented specimens and at different life stages [4–6]. The mitochondrial DNA region at the 5' ends of cytochrome oxidase c subunit I (COI) was used as a molecular marker to delineate species; hence, this region is commonly used as a “barcode” [7]. The diversity in the amino acid sequence [about 648 base pairs (bp)] of the COI gene was sufficient to reliably place species into higher taxonomic levels [1]. Over the past decade, DNA barcoding has been used by diverse taxa, including birds [8], fishes [9–11], spiders [12], invertebrates [13], and mammals [14–16]. DNA barcoding has also been

used to evaluate food quality and authenticity [17], resolve taxonomic uncertainty [18], and monitoring fish biodiversity [19, 20]. Moreover, this method could provide broader insight into the ecology and biosystematics, for example, by revealing cryptic organisms and discovering new species [21–23].

Way Kambas National Park (WKNP) is located in eastern Lampung province, Sumatera, Indonesia. It covers about 1,300 km² of swamp forest and lowland rain forest. Mangrove ecosystem sites can be found in the eastern region of WKNP. Mangrove-based estuarine ecosystems have numerous of finfish species. Around 267 species of fish from 81 families have been reported in two major rivers in India [24]. An estuarine habitat is a partially enclosed, coastal water body where freshwater from rivers and streams mixes with saltwater from the ocean. River mouths, lagoons, and bays often constitute estuarine habitats. The

How to cite:

Ariyanti Y, Rini IA, Oktaviani I, Leksikowati SS (2021) DNA Barcoding for Selected Mangrove-Based Estuary Fishes from Way Kambas National Park, Lampung Province, Indonesia. *Journal of Tropical Life Science* 11 (2): 151 – 160. doi: [10.11594/jtls.11.02.04](http://dx.doi.org/10.11594/jtls.11.02.04).




Figure 1. Map of sample collection site in Way Kambas National Park.

mixing of seawater and freshwater provides high levels of nutrients both in the water column and sediment, making estuaries the most productive natural habitat [25]. Estuaries provide habitat for life-cycle completion by various species, including mollusks, crustaceans, and fishes. These unique ecosystems are essential for maintaining biodiversity. The accurate taxonomic identification of species is critical for the assessment and monitoring of biodiversity.

The eastern side of Way Kambas National Park is directly facing the Java Sea, and this area can be accessed freely by many fishermen. Most of them are part of the local community who still often catch fish in the national park estuaries. However, they do not know the conservation status of the various types of fish caught. This study provides information on the conservation status and species of fish originating from the national park area. This information can be used as material for consideration in the making of conservation policies by related agencies. DNA barcode technique was applied to identify and confirm selected fish species in the mangrove-based estuaries in WKNP.

Material and Methods

Study area

Fish were captured with permission from the WKNP Agency (National Park Entrance Permit, sanction order no. SI.990/BTNWK-1/2018, dated 28 July 2018). Ethics approval for this study was unrequired because no endangered or protected

fish species were involved. Fish samples were collected from three sampling stations at the following estuarine river mangrove bases: Kuala Penet ($5^{\circ}15'18.01''\text{S}$ $105^{\circ}51'38.09''\text{E}$), Kuala Kambas ($5^{\circ} 3'49.43''\text{S}$ $105^{\circ}51'25.07''\text{E}$), and Wako ($4^{\circ}50'17.79''\text{S}$ $105^{\circ}51'45.00''\text{E}$) (Figure 1). The samples were photographed immediately after collection. Approximately 3–5 g of dorsal muscle tissue or fin were excised and immersed in 96% ethanol for genomic analysis. Voucher specimens were stored in 70% alcohol and morphologically identified based on the FAO Species Identification Guide for Fishery Purposes [26], Kottelat *et al.* [27] and <http://www.fishbase.org> [28].

DNA isolation and sequencing

Total genomic DNA was extracted from stored epaxial muscle tissue or fins using a GeneAid GT300 Genomic DNA Mini Kit (Tissue) following the manufacturer's protocols. The DNA concentration was estimated using a NanoDrop™ 2000/c spectrophotometer (Thermo Fisher). The target region of COI was amplified by PCR using the primers FishF1 (5'-TCAACCAACCACAAAGACATTGGCAC-3') and FishR1 (5'-TAGACTTCTGGGTGGCCAAGAATCA-3') [29]. PCR was performed in a total volume of 25 μL consisting of 1 μL of DNA template, 12.5 μL of GoTaq® Green Master Mix (Promega), 1 μL of each primer, and distilled water. The amplification conditions were as follows: initial denaturation for 2 min at 95°C followed by 35 cycles of 0.5 min at 94°C , 0.5 min at 54°C , and 1 min at 72°C , with a

Table 1. Samples analysed in this study and the GenBank accession numbers of the COI mtDNA sequences

Sample/Genus/Species	No. of specimens (n)	Source	Collection sites	GenBank Accession number	
				References	In this study
<i>Lutjanus argentimaculatus</i>	2	Wild	Wako		MN243478, MN243479
	3	GenBank		MG923374.1	
				EU502685.1	
				KC970482.1	
<i>Tetraodon nigroviridis</i>	1	Wild	Kuala Kambas		MN243476
	3	GenBank		DQ019313.1	
				KC959930.1	
				JQ681838.1	
<i>Photopectoralis bindus</i>	1	Wild	Kuala Kambas		MN243471
	1	Wild	Wako		MN243480
	2	GenBank		KJ013055.1	
MG677547.1					
<i>Scatophagus argus</i>	1	Wild	Kuala Kambas		MN243469
	3	GenBank		KY634864.1	
				MG923404.1	
				KY634866.1	
<i>Ambassis</i> sp.	1	Wild	Kuala Kambas		MN243472
	3	GenBank		KU692232.1	
				KU692234.1	
				KU692233.1	
<i>Stigmatogobius sadanundio</i>	1	Wild	Wako		MN243486
	2	GenBank		MG495948.1	
				MF594606.1	
<i>Planiliza parmata</i>	1	Wild	Kuala Kambas		MN243475
	2	GenBank		KX977548.1	
				KX977546.1	
				HQ564490.1	
<i>Planiliza subviridis</i>	1	GenBank			
<i>Mugil cephalus</i>	1	Wild	Wako		MN243481
	1	GenBank			
<i>Osteochilus hasseltii</i>	1	GenBank			
<i>Osteochilus vittatus</i>	1	Wild	Wako		MN243482
<i>Osteochilus</i> sp.	1	GenBank			
<i>Parachela oxygastroides</i>	1	Wild	Wako		MN243487
	1	GenBank			
<i>Parachela hypophthalmus</i>	2	GenBank			
	2	Wild	Wako		MN243484, MN243488
<i>Hemibagrus nemurus</i>	3	GenBank		KM213068.1	
				KM213067.1	
					KJ573466.1
<i>Hemibagrus capitulum</i>	1	GenBank			
	4	GenBank			
<i>Mystus cavasius</i>				KU870465.1	
				JX983383.1	
				JX983379.1	
<i>Mystus wolffii</i>	1	Wild	Kuala Kambas		MN243470
	1	Wild	Wako		MN243483
<i>Mystus bleekeri</i>	1	GenBank			
<i>Mystus singaringan</i>	1	GenBank			
	1	Wild	Kuala Kambas		MN243474
<i>Eleutheronema tetradactylum</i>	1	Wild	Wako		MN243477
	3	GenBank		MG923347.1	
				MG923350.1	
				MG923349.1	

Continue...

Table 1. Samples analysed in this study and the GenBank accession numbers of the COI mtDNA sequences

Sample/Genus/Species	No. of specimens (n)	Source	Collection sites	GenBank Accession number	
				References	In this study
<i>Hexanematichthys saqor</i>	1	Wild	Kuala Penet		MN243467
	1	GenBank		JX198212.1	
<i>Netuma cf. thalassina</i>	1	GenBank		HQ564482.1	
<i>Arius</i>	1	GenBank		KX211965.1	
<i>Arius microcephalus</i>	1	GenBank		MK604248.	
	2	Wild	Kuala Kambas		MN243468, MN243473
<i>Arius maculatus</i>	1	Wild	Wako		MN243485
	2	GenBank		KY849505.1 KY849504.1	

final hold for 10 min at 72°C. The products were electrophoresed in 1% agarose gels and stained with ethidium bromide; DNA bands were visualised under an ultraviolet transilluminator. Intense bands were sent for sequencing on the BigDye® Terminator v. 3.1 platforms provided by 1st BASE Laboratories (Singapore).

Data analysis

The 22 bi-directional sequences were initially checked by eye using the sequence editor BioEdit v. 7.0.9.0 [30]. Next, the sequences were aligned using ClustalW in MEGA X [31]. The final alignment comprised 680 bp. The sequences were submitted to GenBank. A similarity search of the generated sequences was performed using the Basic Local Alignment Search Tool (BLAST; <http://blast.ncbi.nlm.nih.gov/>) and the Barcode of Life Data System (BOLD) (<http://www.barcodinglife.org/index.php/IDS-OpenIdEngine>) [32]. The generated sequences and 44 published sequences of related species were aligned as a dataset (Table 1); equal sequence lengths were used to prevent incongruent outcomes between genetic distances and the neighbour-joining (NJ) tree. The genetic distances within and between species were calculated using the Kimura two-parameter model [33]. A phylogenetic tree was constructed using the NJ method [34] and the maximum parsimony (MP) algorithm. Homologous COI sequences available in GenBank were included in the phylogenetic analysis. The clade confidence in the tree was tested by 1,000-replicate bootstrapping to determine support values for the clade nodes.

Result and Discussion

A total of 68 mitochondrial COI barcode sequences were obtained from 11 families, 15 genera, and 27 species. After editing, the consensus

length of all barcode sequences was > 500 bp. Stop codons, insertions, and deletions were undetected in any of the sequences.

The analysis results of the nucleotide pair frequency revealed that 339 of 593 (57.17%) sites were conserved, 254 of 593 (42.83%) sites were variable, 252 of 593 (42.49%) sites were parsimony informative, and 2 of 593 (0.33%) were singleton sites. The transitional substitutions rates are shown in bold and those of transversional substitutions in italics (Table 2).

The transitional substitutions rates ranged from 10.41 to 31.3, while those of transversional substitutions ranged from 2.24 to 3.66. The transition/transversion rate ratios were $k1 = 10.022$ (purines) and $k2 = 2.862$ (pyrimidines). The overall transition/transversion bias for the dataset was $R = 2.765$. The nucleotide frequencies were 24.65% (A), 28.93% (T/U), 28.72% (C), and 17.70% (G). A base-composition analysis showed that the average T content was highest, and the average G content was lowest. The AT content (53.58%) was higher than the GC content (46.42%), similar to the results for Australian [35], Canadian [36], Cuban [37], and Taiwan Strait fish species [38].

According to Ward *et al.* [35], the result was relevant in mostly marine fishes, in which the content of AT is higher than that of GC. The mean genetic divergence of the dataset was 22%. The intraspecific distance ranged from 0% (mostly species in the studied dataset) to 0.9%; the greatest distance (0.9%) was in *Scatophagus argus*. The interspecific distance ranged from 7.4% to 30.3%. The smallest interspecific distance was between *Netuma cf. thalassina* (HQ564482.1) and *Arius arius* (KX211965.1), and the largest was between *Eleutheronema tetradactylum* (MN243474) and *Stigmatogobius sadanundio* (MF594606.1).

The mean within-family distance ranged from

Table 2. Maximum composite likelihood estimates of the nucleotide substitution pattern

	A	T	C	G
A	-	3.66	3.64	22.47
T	3.12	-	10.41	2.24
C	3.12	10.49	-	2.24
G	31.3	3.66	3.64	-

Notes:

Rates of transitional substitution are in bold and those of transversal substitution are in italics.

0.20% to 11.62%, and the mean between-family distance from 12.10% to 28.80%. Species identification by DNA barcoding relies on both intraspecific and interspecific divergence. According to Meier *et al.* [39], the barcode gap can be calculated as the smallest versus the largest intraspecific distance. The within-species genetic distance was < 2%, and that between species was > 5%. The 22 sequences were matched with homologs by BLAST searching using a species identity of > 97%. Phylogenetic and molecular evolutionary analyses were conducted using MEGA X [31]. The NJ tree (Figure 2) shows distinct clustering of all studied species. The unrooted NJ tree comprised three clusters.

Most specimens from the same species were grouped into one cluster. The first cluster consisted of seven families: Lutjanidae, Tetraodontidae, Leiognathidae, Scatopagidae, Ambassidae, Gobiidae, and Mugilidae. All members of the first cluster were typically found in marine to brackish water. The second cluster consisted of Cyprinidae; this cluster was placed with confidence between the other two clusters. The third cluster consisted of three families: Bagridae, Polynemidae, and Ariidae. The Cyprinidae are a family of freshwater fishes known as cyprinids and commonly called carp. Cyprinidae is the largest and most diverse fish family and the most abundant vertebrate family (~3,000 species) [27, 40].

Osteochilus and *Parachela* are cyprinid fish genera mainly found in Southeast Asia. Both inhabit freshwater habitats, including rivers [41]. Two families, Bagridae and Ariidae, are catfishes. Ariids are found in shallow temperate and tropical seas around the coastlines of North and South America, Africa, Asia, and Australia. Ariid catfish species mainly live in freshwater and brackish water. Bagridae (naked catfish) are freshwater cat

fish native to Africa and Asia (from Japan to Borneo).

Hexanemathychthys sagor (MN243467) was genetically close (0.002) to the other Ariid catfish species, *H. sagor* (JX198212.1). The two *Mystus* sp. sequences (MN243470 and MN243483) had an identity of ~88%, meaning that species are missing from our reference library. Those sequences matched (99%) *Mystus wolffii* according to a sequence alignment using the BOLD ID database (<http://www.barcodinglife.org/index.php/IDS-OpenIdEngine>). Figure 3 shows the phylogenetic tree constructed from the BOLD ID database sequences; the *Mystus* sp. was confidently grouped with *M. wolffii*. Two *M. wolffii* sequences were added to GenBank. The sequences added for these species could previously only be found in the BOLD database as private records.

The eastern coast of Sumatera has a vast area of mangrove swamp. This study was conducted in a mangrove ecosystem within the protected area of WKNP. Data on species diversity, especially fish, originating from the mangrove ecosystem in this area, is still lacking. Barcode data for most of the studied species are available in GenBank. However, the library still lacks suitable COI homologous sequences for some species.

Based on BOLD ID results, three sequences (*Osteochilus vittatus*, *M. wolffii*, and *Ambassis* sp.) had 100% similarity scores. However, one *Ambassis* sp. was unnamed to the species level due to a lack of appropriate specimens and a reference sequence in GenBank or BOLD ID. There were eight species with the Least Concern status according to the IUCN Red List—*Lutjanus argentimaculatus*, *S. argus*, *Ambassis* sp., *Mugil cephalus*, *O. vittatus*, *Parachela oxygastroides*, *Hemibagrus nemurus*, and *M. wolffii*. *Photopectoralis bindus* and *Arius maculatus* were categorised as Data-Deficient species due to insufficient information for a proper assessment, while the remaining species were categorised as Not Evaluated [42–50] (Table 3).

Intriguingly, one species of the family Polynemidae, *E. tetradactylum*, was classified as endangered under criterion A (EN A4d). Based on the previous assessment (2014), this species has likely declined by ~50% in the Persian Gulf. The population of this species is likely to decline by 50–87.5% over a three-generation period [51]. *Eleutheronema tetradactylum* possesses several diagnostic traits, including four pectoral filaments;




Figure 2. NJ tree constructed based on COI sequences using Kimura two-parameter distances. Scale bar, 0.020 substitutions per nucleotide position.




Figure 3. NJ tree constructed based on the BOLD ID data. Red, position of *Mystus wolffii* within the BOLD phylogenetic tree

Table 3. Details of the studied fish species from the mangrove ecosystem in WKNP and their IUCN conservation status.

Order	Family	Genus/Species	IUCN status
Perciformes	Lutjanidae	<i>Lutjanus argentimaculatus</i>	LC
Tetraodontiformes	Tetraodontidae	<i>Tetraodon nigroviridis</i>	NE
Perciformes	Leiognathidae	<i>Photopectoralis bindus</i>	DD
Perciformes	Scatophagidae	<i>Scatophagus argus</i>	LC
Perciformes	Ambassidae	<i>Ambassis</i> sp.	LC
Perciformes	Gobiidae	<i>Stigmatogobius sadanundio</i>	NE
Mugiliformes	Mugilidae	<i>Planiliza parmata</i>	NE
		<i>Mugil cephalus</i>	LC
Cypriniformes	Cyprinidae	<i>Osteochilus vittatus</i>	LC
		<i>Parachela oxygastroides</i>	LC
Siluriformes	Bagridae	<i>Hemibagrus nemurus</i>	LC
		<i>Mystus wolffii</i>	LC
Perciformes	Polynemidae	<i>Eleutheronema tetradactylum</i>	E A4d*
Siluriformes	Ariidae	<i>Hexanematichthys sagor</i>	NE
		<i>Arius maculatus</i>	DD

16–18 (mode 17, rarely 15 or 19) pectoral fins rays; 14 (rarely 13 or 15) second dorsal fin soft rays; a vomer with deciduous tooth plates on both sides, except in juveniles; and pectoral fin membranes that are vivid yellow in life, except in large specimens. This species is widely distributed in the Indo-West Pacific and is extant in Bahrain, Iran, Iraq, Kuwait, Qatar, Saudi Arabia, and the United Arab Emirates [52]. However, there is no policy on the susceptibility of these species in the Indo-West Pacific; thus, they should focus on conservation efforts, particularly in Indonesia.

The total number of barcoded Chordata from Indonesia was 20,217, of which 15,631 were Actinopterygii. Based on the Fish Barcode of Life Initiative (FISH-BOL), there are about 12,140 of fish species from Southeast Asia have been barcoded in 2019. The Scientific Committee on Antarctic Research–Marine Biodiversity Information Network in 2012 reported that < 20% of fish species in Southeast Asia have a barcode [53]. The development of comprehensive DNA barcode reference libraries, especially in Indonesian freshwater fishes, was initiated by Hubert *et al.* [9] several years ago.

DNA barcode techniques have been widely used to reveal the diversity of fish in Indonesia, for example, in the endangered species of sharks [54], the substantial economic value of reef fish [55], and fish originating from the peat swamp environment of New Guinea island, Indonesia [56]. Wibowo *et al.* [56] found something unusual, about 68% of the fish larvae sequences could not

determine into species level due to the lack of a suitable COI sequence in the reference dataset. The vast region and various habitats are become challenging to reveal the diversity of fish in Indonesia. In summary, the present study contributes to form a complete DNA barcode library, especially for teleost fish originating from the mangrove ecosystem.

Conclusion

The DNA Barcoding technique enabled to discriminate of selected fish from WKNP into species level. Mitochondrial COI barcode for 22 mangrove-based estuarine fish species from WKNP has been submitted to GenBank. These findings will facilitate future studies on the diversity of fish species in mangrove estuary-based ecosystems and provide valuable preliminary data in policy-making in conservation areas such as National Parks.

Acknowledgement

The authors are thankful to the Directorate General of Natural Resources and Ecosystem Conservation of the Ministry of Environment and Forestry Republic of Indonesia, especially to the Head Officer of Way Kambas National Park, for providing the necessary permission for sampling license. We also express our thankfulness to all staff officers of Way Kambas National Park, particularly Mr. Lukman, for assistance during field sampling. This work was supported by the Institut Teknologi Sumatera (ITERA) through ITERA

Smart Research Grant [Grant number 134f/IT9.C1/PP/2018].

References

1. Hebert PDN, Ratnasingham S, DeWaard JR (2003) Barcoding animal life: Cytochrome c oxidase subunit 1 divergences among closely related species. *Proc R Soc B Biol Sci.* doi: 10.1098/rsbl.2003.0025
2. Pereira LHG, Hanner R, Foresti F, Oliveira C (2013) Can DNA barcoding accurately discriminate megadiverse Neotropical freshwater fish fauna? *BMC Genet.* doi: 10.1186/1471-2156-14-20
3. Hollingsworth PM (2007) DNA barcoding: potential users. *Genomics, Soc Policy.* doi: 10.1186/1746-5354-3-2-44
4. Wibowo A, Panggabaian AS, Zamroni A et al. (2018) Using DNA barcode to improve the identification of marine fish larvae, case study coastal water near Jakarta and Banda Sea, Indonesia. *Indones Fish Res J.* doi: 10.15578/ifrj.24.1.2018.37-44
5. Sakaguchi SO, Shimamura S, Shimizu Y et al. (2017) Comparison of morphological and DNA-based techniques for stomach content analyses in juvenile chum salmon *Oncorhynchus keta*: a case study on diet richness of juvenile fishes. *Fish Sci.* doi: 10.1007/s12562-016-1040-6
6. Collin R, Venera-Pontón DE, Driskell AC et al. (2020) DNA barcoding of echinopluteus larvae uncovers cryptic diversity in neotropical echinoids. *Invertebr Biol.* doi: 10.1111/ivb.12292
7. Roe AD, Sperling FAH (2007) Patterns of evolution of mitochondrial cytochrome c oxidase I and II DNA and implications for DNA barcoding. *Mol Phylogenet Evol.* doi: 10.1016/j.ympev.2006.12.005
8. Hebert PDN, Stoeckle MY, Zemplak TS, Francis CM (2004) Identification of birds through DNA barcodes. *PLoS Biol.* doi: 10.1371/journal.pbio.0020312
9. Hubert N, Kadarusman, Wibowo A et al. (2016) DNA Barcoding Indonesian freshwater fishes: challenges and prospects. *DNA Barcodes.* doi: 10.1515/dna-2015-0018
10. Zhang J Bin, Hanner R (2011) DNA barcoding is a useful tool for the identification of marine fishes from Japan. *Biochem Syst Ecol.* doi: 10.1016/j.bse.2010.12.017
11. Sachithanandam V, Mohan PM, Muruganandam N et al. (2012) DNA barcoding, phylogenetic study of *Epinephelus* spp. from Andaman coastal region, India. *Indian J. Mar. Sci.*
12. Barrett RDH, Hebert PDN (2005) Identifying spiders through DNA barcodes. *Can J Zool.* doi: 10.1139/z05-024
13. Heimeier D, Lavery S, Sewell MA (2010) Using DNA barcoding and phylogenetics to identify Antarctic invertebrate larvae: Lessons from a large scale study. *Mar Genomics.* doi: 10.1016/j.margen.2010.09.004
14. Borisenko A V., Lim BK, Ivanova N V. et al. (2008) DNA barcoding in surveys of small mammal communities: A field study in Suriname. *Mol Ecol Resour.* doi: 10.1111/j.1471-8286.2007.01998.x
15. Clare EL, Lim BK, Engstrom MD et al. (2007) DNA barcoding of Neotropical bats: Species identification and discovery within Guyana: Barcoding. *Mol Ecol Notes.* doi: 10.1111/j.1471-8286.2006.01657.x
16. Clare EL, Lim BK, Fenton MB, Hebert PDN (2011) Neotropical bats: Estimating species diversity with DNA barcodes. *PLoS One.* doi: 10.1371/journal.pone.0022648
17. Wong EHK, Hanner RH (2008) DNA barcoding detects market substitution in North American seafood. *Food Res Int.* doi: 10.1016/j.foodres.2008.07.005
18. Mabrugaña E, de Astarloa JMD, Hanner R et al. (2011) DNA barcoding identifies argentine fishes from marine and brackish waters. *PLoS One.* doi: 10.1371/journal.pone.0028655
19. Rosas U, Menendez F, Cornejo R et al. (2018) Fish DNA barcoding around large marine infrastructure for improved biodiversity assessment and monitoring. *Mitochondrial DNA Part A DNA Mapping, Seq Anal.* doi: 10.1080/24701394.2018.1431225
20. Kundu S, Rath S, Laishram K et al. (2019) DNA barcoding identified selected ornamental fishes in Murti river of East India. *Mitochondrial DNA Part B Resour.* doi: 10.1080/23802359.2018.1561220
21. Janzen DH, Hajibabaei M, Burns JM et al. (2005) Wedding biodiversity inventory of a large and complex Lepidoptera fauna with DNA barcoding. *Philos Trans R Soc B Biol Sci.* doi: 10.1098/rstb.2005.1715
22. Wang T, Zhang Y ping, Yang Z yu et al. (2020) DNA barcoding reveals cryptic diversity in the underestimated genus *Triplophysa* (Cypriniformes: Cobitidae, Nemacheilinae) from the northeastern Qinghai-Tibet Plateau. *BMC Evol Biol.* doi: 10.1186/s12862-020-01718-0
23. Léger T, Kehlmaier C, Vairappan CS, Nuss M (2020) Twenty-six new species of hoploscopa (Lepidoptera, crambidae) from South-East Asia revealed by morphology and DNA barcoding. *Zookeys.* doi: 10.3897/zookeys.907.36563
24. Mandal B, Mukherjee A, Banerjee S (2013) A review on the ichthyofaunal diversity in mangrove based estuary of Sundarbans. *Rev Fish Biol Fish.* doi: 10.1007/s11160-012-9300-8
25. Day JW, Crump BC, Kemp WM, Yáñez-Arancibia A (2012) *Estuarine Ecology.* *Estuar Ecol.* doi: 10.1002/9781118412787
26. Carpenter KE, Niem VH (1999) The living marine resources of the Western Central Pacific. Volume 4. Bony fishes part 2 (Mugilidae to Carangidae). *FAO Species Identif. Guid. Fish. Purp.*
27. Kottelat M, Whitten AJ, Kartikasari SN, Wirjoatmodjo S (1993) *Freshwater fishes of western Indonesia and Sulawesi.* Hong Kong, Periplus Editions.
28. Froese R, Pauly D (2018) *www.fishbase.org.* (2018) Accessed:
29. Ward RD, Hanner R, Hebert PDN (2009) The campaign to DNA barcode all fishes, FISH-BOL. *J Fish Biol.* doi: 10.1111/j.1095-8649.2008.02080.x
30. Hall TA (1999) BIOEDIT: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/ NT. *Nucleic Acids Symp. Ser.*
31. Kumar S, Stecher G, Li M et al. (2018) MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol.* doi: 10.1093/molbev/msy096
32. RATNASINGHAM S, HEBERT PDN (2007) BARCODING: bold: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Mol Ecol Notes.* doi: 10.1111/j.1471-8286.2007.01678.x

33. Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol.* doi: 10.1007/BF01731581
34. Tamura K, Nei M, Kumar S (2004) Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc Natl Acad Sci U S A.* doi: 10.1073/pnas.0404206101
35. Ward RD, Zemlak TS, Innes BH et al. (2005) DNA barcoding Australia's fish species. *Philos Trans R Soc B Biol Sci.* doi: 10.1098/rstb.2005.1716
36. Hubert N, Hanner R, Holm E et al. (2008) Identifying Canadian freshwater fishes through DNA barcodes. *PLoS One.* doi: 10.1371/journal.pone.0002490
37. Lara A, Ponce de León JL, Rodríguez R et al. (2010) DNA barcoding of Cuban freshwater fishes: Evidence for cryptic species and taxonomic conflicts. *Mol Ecol Resour.* doi: 10.1111/j.1755-0998.2009.02785.x
38. Bingpeng X, Heshan L, Zhilan Z et al. (2018) Dna barcoding for identification of fish species in the taiwan strait. *PLoS One.* doi: 10.1371/journal.pone.0198109
39. Meier R, Zhang G, Ali F (2008) The use of mean instead of smallest interspecific distances exaggerates the size of the "barcoding gap" and leads to misidentification. *Syst Biol.* doi: 10.1080/10635150802406343
40. He S, Mayden RL, Wang X et al. (2008) Molecular phylogenetics of the family Cyprinidae (Actinopterygii: Cypriniformes) as evidenced by sequence variation in the first intron of S7 ribosomal protein-coding gene: Further evidence from a nuclear gene of the systematic chaos in the family. *Mol Phylogenet Evol.* doi: 10.1016/j.ympev.2007.06.001
41. Hui TH, Kottelat M (2009) The fishes of the Batang Hari drainage, Sumatra, with description of six new species. *Ichthyol. Explor. Freshwaters*
42. Russell B, Carpenter, KE, Smith-Vaniz, WF, Lawrence A SJ (2016) *Lutjanus argentimaculatus*. The IUCN Red List of Threatened Species 2016: e.T61250A3101831. <http://dx.doi.org/10.2305/IUCN.UK.20163.RLTS.T61250A3101831.en>. Accessed: August 2019.
43. Vidthayanon C (2012) *Osteochilus vittatus*. The IUCN Red List of Threatened Species 2012:e.T180750A1658850. <http://dx.doi.org/10.2305/IUCN.UK.20121.RLTS.T180750A165880.en>. Accessed: August 2019.
44. Jenkins A, Kullander FF TH (2009) *Parachela oxygastroides*. The IUCN Red List of Threatened Species 2009:e.T169547A6645702. <http://dx.doi.org/10.2305/IUCN.UK.2009-2.RLTS.T169547A6645702.en>. Accessed: August 2019.
45. Jenkins A, Kullander FF TH (2009) *Mystus wolffii*. The IUCN Red List of Threatened Species 2009:e.T169551A6646308. <http://dx.doi.org/10.2305/IUCN.UK.20092.RLTS.T169551A6646308.en>. Accessed: August 2019.
46. Collen B, Richman N, Beresford A, Chenery A RM (Sampled RLICT (2010) *Scatophagus argus*. The IUCN Red List of Threatened Species 2010:e.T155268A4761779. <http://dx.doi.org/10.2305/IUCN.UK.20104.RLTS.T155268A4761779.en>. Accessed: August 2019.
47. N D (2012) *Ambassis nalua*. The IUCN Red List of Threatened Species 2012:e.T172359A1340093. Accessed: August 2019.
48. HH N (2012) *Hemibagrus nemurus*. The IUCN Red List of Threatened Species 2012:e.T180954A1681839. <http://dx.doi.org/10.2305/IUCN.UK.20121.RLTS.T180954A1681839.en>. Accessed: August 2019.
49. Kaymaram F, Hartmann S, Al-Husaini M, Alam S AM (2015) *Photopectoralis bindus*. The IUCN Red List of Threatened Species 2015:e.T50903077A57142170. Accessed: August 2019.
50. Kaymaram F, Bishop J, Al-Husaini M, Almkhtar M AS *Arius maculatus*. The IUCN Red List of Threatened Species 2015: e.T196796A56996099. Accessed: August 2019.
51. Motomura H, Iwatsuki Y, Kimura S, Yoshino T (2002) Revision of the Indo-West Pacific polynemid fish genus *Eleutheronema* (Teleostei: Perciformes). *Ichthyol Res.* doi: 10.1007/s102280200005
52. Motomura H, Matsuura K, Bishop J KF (2015) *Eleutheronema tetradactylum*. The IUCN Red List of Threatened Species 2015:e.T46087646A57168342. Accessed: August 2019.
53. Bhattacharya M, Sharma AR, Patra BC et al. (2016) DNA barcoding to fishes: Current status and future directions. *Mitochondrial DNA.* doi: 10.3109/19401736.2015.1046175
54. Sembiring A, Pertiwi NPD, Mahardini A et al. (2015) DNA barcoding reveals targeted fisheries for endangered sharks in Indonesia. *Fish Res.* doi: 10.1016/j.fishres.2014.11.003
55. Fadli N, Mohd Nor SA, Othman AS et al. (2020) DNA barcoding of commercially important reef fishes in Weh Island, Aceh, Indonesia. *PeerJ.* doi: 10.7717/peerj.9641
56. Wibowo A, Wahlberg N, Vasemägi A (2017) DNA barcoding of fish larvae reveals uncharacterised biodiversity in tropical peat swamps of New Guinea, Indonesia. *Mar Freshw Res.* doi: 10.1071/MF16078.