



# Correction to “Association between urinary chloride excretion and progression of coronary artery calcification in patients with non-dialysis chronic kidney disease: results from KNOW-CKD study”

Sang Heon Suh<sup>1</sup>, Tae Ryom Oh<sup>1</sup>, Hong Sang Choi<sup>1</sup>, Chang Seong Kim<sup>1</sup>, Eun Hui Bae<sup>1</sup>, Seong Kwon Ma<sup>1</sup>, Kook-Hwan Oh<sup>2</sup>, Tae-Hyun Yoo<sup>3</sup>, Dong-Wan Chae<sup>4</sup>, Soo Wan Kim<sup>1</sup>; the Korean Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD) Investigators

<sup>1</sup>Department of Internal Medicine, Chonnam National University Hospital, Chonnam National University Medical School, Gwangju, Republic of Korea

<sup>2</sup>Department of Internal Medicine, Seoul National University Hospital, Seoul, Republic of Korea

<sup>3</sup>Department of Internal Medicine and Institute of Kidney Disease Research, Yonsei University College of Medicine, Seoul, Republic of Korea

<sup>4</sup>Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam, Republic of Korea

**Correction to:** Suh SH, Oh TR, Choi HS, et al. Association between urinary chloride excretion and progression of coronary artery calcification in patients with nondialysis chronic kidney disease: results from the KNOW-CKD study. *Kidney Res Clin Pract* 2023;42:251-261. DOI: 10.23876/j.krcp.22.072

The original version of this article contained an error in the Funding section.

“This work was supported by the Research Program funded by the Korea Centers for Disease Control and Prevention (2011E3300300, 2012E3301100, 2013E3301600, 2013E3301601, 2013E3301602, 2016E3300200, 2016E3300201, 2016E3300202, and 2019E320100), by the National Research Foundation of Korea (NRF) funded by the Korean government (MSIT) (NRF-2019R1A2C2086276), and a grant (BCRI22042 and BCRI22079) from Chonnam National University Hospital Biomedical Research Institute.”

The corrected version of the article’s Funding section is as follows:

“This work was supported by the Research Program funded by the **Korea Disease Control and Prevention Agency (2011E3300300, 2012E3301100, 2013E3301600, 2013E3301601, 2013E3301602, 2016E3300200, 2016E3300201, 2016E3300202, 2019E320100, 2019E320101, 2019E320102, 2022-11-007)**, by the National Research Foundation of Korea (NRF) funded by the Korea government (MSIT) (NRF-2019R1A2C2086276), and a grant (BCRI22042 and BCRI22079) of Chonnam National University Hospital Biomedical Research Institute.”

The original version has been rectified. The authors would like to apologize for any inconvenience caused.

**Correspondence:** Soo Wan Kim

Department of Internal Medicine, Chonnam National University Hospital, Chonnam National University Medical School, 42 Jebong-ro, Dong-gu, Gwangju 61469, Republic of Korea. E-mail: skimw@chonnam.ac.kr  
ORCID: <https://orcid.org/0000-0002-3540-9004>

Copyright © 2023 by The Korean Society of Nephrology

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial and No Derivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits unrestricted non-commercial use, distribution of the material without any modifications, and reproduction in any medium, provided the original works properly cited.