



Western Washington University
Western CEDAR

Salish Sea Ecosystem Conference

2022 Salish Sea Ecosystem Conference
(Online)

Apr 27th, 4:00 PM - 4:30 PM

PREVALENCE, DISTRIBUTION, AND CONTROL OF SHELL-BORING POLYCHAETES ON OYSTER FARMS FROM CALIFORNIA TO ALASKA

Dr. Julieta Martinelli
University of Washington

Follow this and additional works at: <https://cedar.wwu.edu/ssec>

Martinelli, Dr. Julieta, "PREVALENCE, DISTRIBUTION, AND CONTROL OF SHELL-BORING POLYCHAETES ON OYSTER FARMS FROM CALIFORNIA TO ALASKA" (2022). *Salish Sea Ecosystem Conference*. 28. <https://cedar.wwu.edu/ssec/2022ssec/allsessions/28>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.


PREVALENCE, DISTRIBUTION, AND CONTROL OF SHELL-BORING POLYCHAETES ON OYSTER FARMS FROM CALIFORNIA TO ALASKA

Julieta Martinelli, Considine M, Tarpey C, King TL, Hauser L, Padilla-Gamiño JL, Rumrill S & CL Wood

 @julishells /julimar@uw.edu

RESEARCH QUESTIONS

- 1) What is the prevalence and distribution of shell-boring worms in the US west coast?
- 2) Which treatments can oyster growers use to control them?
- 3) Can environmental variables help predict cold and hotspots of infestation?



PREVALENCE & DISTRIBUTION

We found burrows and blisters on oyster shells from all states (CA to WA & AK). Prevalence was highly variable across farm, growth method (on/off bottom), season and state. We confirmed the presence of *P. websteri* and *P. hoplura* among other shell-boring polychaetes.





Fig 2. (A) Haplotype network of species found in WA, and (B) Pooled prevalence of shell-boring worms in the US west coast.

CONTROL TREATMENTS

We co-developed 3 treatments with oyster farmers to apply on farms: (i) cold-air drying for 3 days, (ii) outdoor drying for 2 days, (iii) and freshwater dip + 2 days of drying. Drying was the most effective one but led to 4% oyster mortality in the Fall.



ENVIRONMENTAL VARIABLES

We are developing models to test whether salinity, water temperature, and pH, among others, are significant explanatory factors to predict the success of shell-boring polychaetes in different places.

Fig 1. (A) Shell-boring parasite, *Polydora websteri*, and (B) the mud-filled blisters it creates on oyster shells.