

Article

Explicit Properties of Apostol-Type Frobenius–Euler Polynomials Involving q -Trigonometric Functions with Applications in Computer Modeling

Yongsheng Rao ¹, Waseem Ahmad Khan ^{2,*}, Serkan Araci ^{3,*} and Cheon Seoung Ryoo ⁴

¹ Institute of Computing Science and Technology, Guangzhou University, Guangzhou 510006, China; rysheng@gzhu.edu.cn

² Department of Mathematics and Natural Sciences, Prince Mohammad Bin Fahd University, P.O. Box 1664, Al Khobar 31952, Saudi Arabia

³ Department of Basic Sciences, Faculty of Engineering, Hasan Kalyoncu University, TR-27010 Gaziantep, Turkey

⁴ Department of Mathematics, Hannam University, Daejeon 34430, Republic of Korea; ryoo@hnu.kr

* Correspondence: wkhan1@pmu.edu.sa (W.A.K.); serkan.araci@hku.edu.tr (S.A.)

Abstract: In this article, we define q -cosine and q -sine Apostol-type Frobenius–Euler polynomials and derive interesting relations. We also obtain new properties by making use of power series expansions of q -trigonometric functions, properties of q -exponential functions, and q -analogues of the binomial theorem. By using the Mathematica program, the computational formulae and graphical representation for the aforementioned polynomials are obtained. By making use of a partial derivative operator, we derived some interesting finite combinatorial sums. Finally, we detail some special cases for these results.

Keywords: q -trigonometric functions; q -exponential functions; Frobenius–Euler polynomials; Apostol Frobenius–Euler polynomials; generating functions; combinatorial sums

MSC: 11B68; 11B73; 05A15; 05A19



Citation: Rao, Y.; Khan, W.A.; Araci, S.; Ryoo, C.S. Explicit Properties of Apostol-Type Frobenius–Euler Polynomials Involving q -Trigonometric Functions with Applications in Computer Modeling. *Mathematics* **2023**, *11*, 2386. <https://doi.org/10.3390/math11102386>

Academic Editor: Sitnik Sergey

Received: 4 May 2023

Revised: 17 May 2023

Accepted: 18 May 2023

Published: 20 May 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Recently, many authors have considered and applied the generating functions techniques to new families of special polynomials, including two parametric kinds of polynomials, such as Bernoulli, Euler, Genocchi, etc. (see [1–10]). They have firstly derived the basic identities of these polynomials. Additionally, they have established more identities and relations among trigonometric functions, using two parametric kinds of polynomials by using generating functions. By applying the partial derivative operator to these generating functions, derivative formulae, and finite combinatorial sums involving the special polynomials and numbers are obtained. We would like to note that these special polynomials facilitate the derivation of various helpful properties in a fairly straightforward way and lead to introducing new families of special polynomials. The Apostol-type polynomials appear in combinatorial mathematics and play an important role in theory, generalization, applications and modeling; thus, many number theorists and combinatorics experts have extensively investigated their properties and obtained a series of interesting results (see [5,8,9,11–13]). Inspired by the above polynomials, in this study, we are in a position to state the parametric kinds of Apostol-type Frobenius–type Euler polynomials by introducing the two specific q -analogues of exponential generating functions. Additionally, we prove many formulas and relations for these polynomials, including some implicit summation formulas, differentiation rules and correlations with the earlier polynomials by utilizing some series manipulation methods. Additionally, as an application, we show the

zero values of q -Apostol-type Frobenius-type Euler polynomials using tables and draw some graphical representations.

We begin by stating the following definitions and notations of q -calculus reviewed here, which are taken from (see [14]):

A q -analogue of the shifted factorial $(a)_n$ is given by

$$(a; q)_0 = 1, (a; q)_n = \prod_{m=0}^{n-1} (1 - q^m a), n \in \mathbb{N}.$$

A q -analogue of a complex number a and of the factorial function are given by

$$[a]_q = \frac{1 - q^a}{1 - q}, (q \in \mathbb{C} \setminus \{1\}; a \in \mathbb{C}),$$

$$[n]_q! = \prod_{m=1}^n [m]_q = [1]_q [2]_q \cdots [n]_q = \frac{(q; q)_n}{(1 - q)^n}, q \neq 1; n \in \mathbb{N},$$

$$[0]_q! = 1, q \in \mathbb{C}; 0 < |q| < 1.$$

The Gauss q -binomial coefficient $\binom{n}{k}_q$ is given by

$$\binom{n}{k}_q = \frac{[n]_q!}{[k]_q! [n - k]_q!} = \frac{(q; q)_n}{(q; q)_k (q; q)_{n - k}}, k = 0, 1, \dots, n.$$

The q -analogue of the function $(x + y)_q^n$ is given by

$$(x + y)_q^n = \sum_{k=0}^n \binom{n}{k}_q q^{k(k-1)/2} x^{n-k} y^k, n \in \mathbb{N}_0. \tag{1}$$

The q -analogues of exponential functions are given by

$$e_q(x) = \sum_{n=0}^{\infty} \frac{x^n}{[n]_q!} = \frac{1}{((1 - q)x; q)_{\infty}}, 0 < |q| < 1; |x| < |1 - q|^{-1}, \tag{2}$$

$$E_q(x) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{[n]_q!} x^n = (- (1 - q)x; q)_{\infty}, 0 < |q| < 1; x \in \mathbb{C}. \tag{3}$$

These two functions are related by the equation (see [14])

$$e_q(x) E_q(-x) = 1.$$

A q -derivative operator of a function is defined by

$$D_{q,z} f(z) := D_q f(z) = \frac{f(qz) - f(z)}{qz - z}, 0 < |q| < 1,$$

and $D_q f(0) = f'(0)$ provided that f is differentiable at $x = 0$.

A q -derivative fulfills the following product and quotient rules

$$D_{q,z} (f(z)g(z)) = f(z)D_{q,z}g(z) + g(qz)D_{q,z}f(z), \tag{4}$$

$$D_{q,z} \left(\frac{f(z)}{g(z)} \right) = \frac{g(qz)D_{q,z}f(z) - f(qz)D_{q,z}g(z)}{g(z)g(qz)}. \tag{5}$$

The Apostol-type q -Bernoulli polynomials $\mathbb{B}_{n,q}^{(\alpha)}(x; \lambda)$ of order α , the Apostol-type q -Euler polynomials $\mathbb{E}_{n,q}^{(\alpha)}(x; \lambda)$ of order α and the Apostol-type q -Genocchi polynomials $\mathbb{G}_{n,q}^{(\alpha)}(x; \lambda)$ of order α are defined by (see [15,16]):

$$\left(\frac{t}{\lambda e_q(t) - 1}\right)^\alpha e^{xt} = \sum_{n=0}^\infty \mathbb{B}_{n,q}^{(\alpha)}(x; \lambda) \frac{t^n}{[n]_q!}, \tag{6}$$

$$\left(\frac{2}{\lambda e_q(t) + 1}\right)^\alpha e^{xt} = \sum_{n=0}^\infty \mathbb{E}_{n,q}^{(\alpha)}(x; \lambda) \frac{t^n}{[n]_q!}, \tag{7}$$

$$\left(\frac{2t}{\lambda e_q(t) + 1}\right)^\alpha e^{xt} = \sum_{n=0}^\infty \mathbb{G}_{n,q}^{(\alpha)}(x; \lambda) \frac{t^n}{[n]_q!}, \tag{8}$$

respectively.

Clearly, we can obtain

$$\mathbb{B}_{n,q}^{(\alpha)}(\lambda) = \mathbb{B}_{n,q}^{(\alpha)}(0; \lambda), \mathbb{E}_{n,q}^{(\alpha)}(\lambda) = \mathbb{E}_{n,q}^{(\alpha)}(0; \lambda), \mathbb{G}_{n,q}^{(\alpha)}(\lambda) = \mathbb{G}_{n,q}^{(\alpha)}(0; \lambda),$$

and

$$\mathbb{B}_{n,q}^{(1)}(x; \lambda) = \mathbb{B}_{n,q}(x; \lambda), \mathbb{E}_{n,q}^{(1)}(x; \lambda) = \mathbb{E}_{n,q}(x; \lambda), \mathbb{G}_{n,q}^{(1)}(x; \lambda) = \mathbb{G}_{n,q}(x; \lambda).$$

Let $u \in \mathbb{C}$ with $u \neq 1$ and $\xi \in \mathbb{R}$. The Apostol-type q -Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha)}(x, y; u; \lambda)$ of order $\alpha \in \mathbb{C}$ are defined by (see [11,12]):

$$\left(\frac{1-u}{\lambda e_q(t) - u}\right)^\alpha e_q(xt)E_q(yt) = \sum_{n=0}^\infty \mathbb{H}_{n,q}^{(\alpha)}(x, y; u; \lambda) \frac{t^n}{[n]_q!}, \quad |z| < \left|\ln\left(\frac{\lambda}{u}\right)\right|. \tag{9}$$

It is obvious that

$$\mathbb{H}_{n,q}^{(\alpha)}(u; \lambda) = \mathbb{H}_{n,q}^{(\alpha)}(0, 0; u; \lambda) \quad \lim_{\lambda \rightarrow 1^-} \mathbb{H}_{n,q}^{(\alpha)}(x, y; u; \lambda) = \mathbb{H}_n^{(\alpha)}(x + y; u; \lambda).$$

Kang et al. [2,4] introduced the q -Bernoulli and q -Euler polynomials defined by

$$\frac{z}{e_q(z) - 1} e_q(\xi z) \text{COS}_q(\eta z) = \sum_{j=0}^\infty \frac{\mathbb{B}_{j,q}((\xi + i\eta)_q) + \mathbb{B}_{j,q}((\xi - i\eta)_q)}{2} \frac{z^j}{[j]_q!} = \sum_{j=0}^\infty \mathbb{B}_{j,q}^{(C)}(\xi, \eta) \frac{z^j}{[j]_q!}, \tag{10}$$

$$\frac{z}{e_q(z) - 1} e_q(\xi z) \text{SIN}_q(\eta z) = \sum_{j=0}^\infty \frac{\mathbb{B}_{j,q}((\xi + i\eta)_q) - \mathbb{B}_{j,q}((\xi - i\eta)_q)}{2i} \frac{z^j}{[j]_q!} = \sum_{j=0}^\infty \mathbb{B}_{j,q}^{(S)}(\xi, \eta) \frac{z^j}{[j]_q!}, \tag{11}$$

and

$$\frac{2}{e_q(z) + 1} e_q(\xi z) \text{COS}_q(\eta z) = \sum_{j=0}^\infty \frac{\mathbb{E}_{j,q}((\xi + i\eta)_q) + \mathbb{E}_{j,q}((\xi - i\eta)_q)}{2} \frac{z^j}{[j]_q!} = \sum_{j=0}^\infty \mathbb{E}_{j,q}^{(C)}(\xi, \eta) \frac{z^j}{[j]_q!}, \tag{12}$$

$$\frac{2}{e_q(z) + 1} e_q(\xi z) \text{SIN}_q(\eta z) = \sum_{j=0}^\infty \frac{\mathbb{E}_{j,q}((\xi + i\eta)_q) - \mathbb{E}_{j,q}((\xi - i\eta)_q)}{2i} \frac{z^j}{[j]_q!} = \sum_{j=0}^\infty \mathbb{E}_{j,q}^{(S)}(\xi, \eta) \frac{z^j}{[j]_q!}, \tag{13}$$

respectively.

Additionally, they have proved that (see [2,4]):

$$e_q(\xi z) \text{COS}_q(\eta z) = \sum_{r=0}^\infty C_{r,q}(\xi, \eta) \frac{z^r}{[r]_q!}, \tag{14}$$

and

$$e_q(\xi z)SIN_q(\eta z) = \sum_{r=0}^{\infty} S_{r,q}(\xi, \eta) \frac{z^r}{[r]_q!}, \tag{15}$$

where

$$C_{r,q}(\xi, \eta) = \sum_{j=0}^{\lfloor \frac{r}{2} \rfloor} \binom{r}{2j}_q (-1)^j q^{2j-1} \xi^{r-2j} \eta^{2j}, \tag{16}$$

and

$$S_{r,q}(\xi, \eta) = \sum_{j=0}^{\lfloor \frac{r-1}{2} \rfloor} \binom{r}{2j+1}_q (-1)^j q^{(2j+1)j} \xi^{r-2j-1} \eta^{2j+1}. \tag{17}$$

2. q -Apostol-Type Frobenius–Euler Polynomials of Complex Variable

In this section, we consider the q -Cosine and q -Sine Apostol-type Frobenius–Euler polynomials of a complex variable and deduce some identities of these polynomials. First, we present the following definition.

$$\left(\frac{1-u}{\lambda e_q(t)-u} \right)^\alpha e_q(xt)E_q(iy) = \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(\alpha)}((\xi + iy)_q; u; \lambda) \frac{t^n}{[n]_q!}. \tag{18}$$

It is well-known from ([4] Definition 5) that

$$e_q(xt)E_q(iy) = e_q(xt)(COS_q(yt) + iSIN_q(yt)). \tag{19}$$

Thus, by (18) and (19), we have

$$\begin{aligned} \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(\alpha)}((x + iy)_q; u; \lambda) \frac{t^n}{n!} &= \left(\frac{1-u}{\lambda e_q(t)-u} \right)^\alpha e_q(xt)E_q(iy) \\ &= \left(\frac{1-u}{\lambda e_q(t)-u} \right)^\alpha e_q(xt)(COS_q(yt) + iSIN_q(yt)), \end{aligned} \tag{20}$$

and

$$\begin{aligned} \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(\alpha)}((x - iy)_q; u; \lambda) \frac{t^n}{[n]_q!} &= \left(\frac{1-u}{\lambda e_q(t)-u} \right)^\alpha e_q(xt)E_q(-iy) \\ &= \left(\frac{1-u}{\lambda e_q(t)-u} \right)^\alpha e_q(xt)(COS_q(yt) - iSIN_q(yt)). \end{aligned} \tag{21}$$

From (20) and (21), we get

$$\left(\frac{1-u}{\lambda e_q(t)-u} \right)^\alpha e_q(xt)COS_q(yt) = \sum_{n=0}^{\infty} \left(\frac{\mathbb{H}_{n,q}^{(\alpha)}((x + iy)_q; u; \lambda) + \mathbb{H}_{n,q}^{(\alpha)}((x - iy)_q; u; \lambda)}{2} \right) \frac{t^n}{[n]_q!}, \tag{22}$$

and

$$\left(\frac{1-u}{\lambda e_q(t)-u} \right)^\alpha e_q(xt)SIN_q(yt) = \sum_{n=0}^{\infty} \left(\frac{\mathbb{H}_{n,q}^{(\alpha)}((x + iy)_q; u; \lambda) - \mathbb{H}_{n,q}^{(\alpha)}((x - iy)_q; u; \lambda)}{2} \right) \frac{t^n}{[n]_q!}. \tag{23}$$

Definition 1. Let $j \geq 0$. We define two parametric kinds of q -Cosine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda)$ and q -Sine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda)$, for a non negative integer n , by

$$\left(\frac{1-u}{\lambda e_q(t)-u} \right)^\alpha e_q(xt)COS_q(yt) = \sum_{n=0}^{\infty} \mathbb{H}_{j,q}^{(\alpha,c)}(x, y; u; \lambda) \frac{t^n}{[n]_q!}, \tag{24}$$

and

$$\left(\frac{1-u}{\lambda e_q(t)-u}\right)^\alpha e_q(xt)SIN_q(yt) = \sum_{n=0}^\infty \mathbb{H}_{n,q}^{(\alpha,s)}(x,y;u;\lambda) \frac{t^n}{[n]_q!}, \tag{25}$$

respectively.

Note that $\mathbb{H}_{n,q}^{(\alpha,c)}(0,0;u;\lambda) = \mathbb{H}_{n,q}(u;\lambda)$, $\mathbb{H}_{n,q}^{(\alpha,s)}(0,0;u;\lambda) = 0, (n \geq 0)$.
 From (22)–(25), we have

$$\mathbb{H}_{n,q}^{(\alpha,c)}(x,y;u;\lambda) = \frac{\mathbb{H}_{n,q}((x+iy)_q;u;\lambda) + \mathbb{H}_{n,q}((x-iy)_q;u;\lambda)}{2}, \tag{26}$$

$$\mathbb{H}_{n,q}^{(\alpha,s)}(x,y;u;\lambda) = \frac{\mathbb{H}_{n,q}((x+iy)_q;u;\lambda) - \mathbb{H}_{n,q}((x-iy)_q;u;\lambda)}{2i}. \tag{27}$$

Remark 1. For $x = 0$ in (24) and (25), we obtain

$$\left(\frac{1-u}{\lambda e^t-u}\right)^\alpha \text{COS}_q(yt) = \sum_{j=0}^\infty \mathbb{H}_{n,q}^{(\alpha,c)}(y;u;\lambda) \frac{t^n}{[n]_q!}, \tag{28}$$

and

$$\left(\frac{1-u}{\lambda e^t-u}\right)^\alpha \text{SIN}_q(yt) = \sum_{n=0}^\infty \mathbb{H}_{n,q}^{(\alpha,s)}(y;u;\lambda) \frac{t^n}{[n]_q!}, \tag{29}$$

respectively.

It is clear that

$$\mathbb{H}_{n,q}^{(\alpha,c)}(0;u;\lambda) = \mathbb{H}_{n,q}(u;\lambda), \quad \mathbb{H}_{n,q}^{(\alpha,s)}(0;u;\lambda) = 0, (n \geq 0).$$

Now, we provide some basic properties of these polynomials.

Theorem 1. Let $n \geq 0$. Then,

$$\mathbb{H}_{n,q}^{(\alpha,c)}(y;u;\lambda) = \sum_{v=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n+v}{2v}_q (-1)^v q^{(2v-1)v} y^{2v} \mathbb{H}_{n-2v,q}(u;\lambda), \tag{30}$$

and

$$\mathbb{H}_{j,q}^{(\alpha,s)}(u;\lambda) = \sum_{v=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n+v}{2v+1}_q (-1)^v q^{(2v+1)v} y^{2v+1} \mathbb{H}_{n-2v-1,q}(u;\lambda). \tag{31}$$

Proof. By (28) and (29), we can derive the following equations

$$\begin{aligned} \sum_{n=0}^\infty \mathbb{H}_{n,q}^{(\alpha,c)}(y;u;\lambda) \frac{t^n}{[n]_q!} &= \left(\frac{1-u}{\lambda e_q(t)-u}\right)^\alpha \text{COS}_q(yt) \\ &= \sum_{n=0}^\infty \mathbb{H}_{n,q}^{(\alpha)}(u;\lambda) \frac{t^n}{[n]_q!} \sum_{v=0}^\infty (-1)^v q^{(2v-1)v} y^{2v} \frac{t^v}{[2v]_q!}. \\ &= \sum_{n=0}^\infty \left(\sum_{v=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n+v}{2v}_q (-1)^v q^{(2v-1)v} y^{2v} \mathbb{H}_{n-2v,q}(u;\lambda) \right) \frac{t^n}{[n]_q!}, \end{aligned} \tag{32}$$

and

$$\sum_{n=0}^\infty \mathbb{H}_{n,q}^{(\alpha,s)}(y;u;\lambda) \frac{t^n}{n!} = \left(\frac{1-u}{\lambda e_q(t)-u}\right)^\alpha \text{SIN}_q(yt)$$

$$= \sum_{n=0}^{\infty} \left(\sum_{v=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{2v+1}_q (-1)^v q^{(2v+1)v} y^{2v+1} \mathbb{H}_{j-2v-1,q}(u; \lambda) \right) \frac{t^n}{[n]_q!}. \tag{33}$$

Therefore, with (32) and (33), we get (30) and (31). \square

Theorem 2. Let $n \geq 0$. Then,

$$\begin{aligned} \mathbb{H}_{n,q}^{(\alpha)}((x + iy)_q; u; \lambda) &= \sum_{k=0}^n \binom{n}{k}_q (x + iy)_q^k \mathbb{H}_{n-k,q}^{(\alpha)}(u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q (iy)^k \mathbb{H}_{j-k,q}^{(\alpha)}(x; u; \lambda), \end{aligned} \tag{34}$$

and

$$\begin{aligned} \mathbb{H}_{n,q}^{(\alpha)}((x - iy)_q; u; \lambda) &= \sum_{k=0}^n \binom{n}{k}_q (x - iy)_q^k \mathbb{H}_{n-k,q}^{(\alpha)}(u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q (-1)^k (iy)^k \mathbb{H}_{n-k,q}^{(\alpha)}(x; u; \lambda). \end{aligned} \tag{35}$$

Proof. By using (20) and (21), we obtain (34) and (35). So, we omit the proof. \square

Theorem 3. Let $n \geq 0$. Then,

$$\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) = \sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{k,q}^{(\alpha)}(u; \lambda) C_{n-k,q}(x, y), \tag{36}$$

and

$$\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) = \sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{k,q}^{(\alpha)}(u; \lambda) S_{n-k,q}(x, y). \tag{37}$$

Proof. Consider

$$\left(\sum_{n=0}^{\infty} a_n \frac{t^n}{n!} \right) \left(\sum_{k=0}^{\infty} b_k \frac{t^k}{k!} \right) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^j a_{n-k} b_k \right) \frac{t^n}{n!}.$$

Now,

$$\begin{aligned} \sum_{n=0}^{\infty} \mathbb{H}_{j,q}^{(\alpha,c)}(x, y; u; \lambda) \frac{t^n}{[n]_q!} &= \left(\frac{1-u}{\lambda e_q(t) - u} \right)^\alpha e_q(xt) \text{COS}_q(yt) \\ &= \left(\sum_{k=0}^{\infty} \mathbb{H}_{k,q}^{(\alpha)}(u; \lambda) \frac{t^k}{k!} \right) \left(\sum_{n=0}^{\infty} C_{n,q}(x, y) \frac{t^n}{[n]_q!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{k,q}^{(\alpha)}(u; \lambda) C_{n-k,q}(x, y) \right) \frac{t^n}{[n]_q!}, \end{aligned}$$

which proves (36). The proof of (37) is similar. \square

By using Definition 1, we can easily obtain the following Theorems. So, we omit the proofs.

Theorem 4. Let $j \geq 0$. Then,

$$\mathbb{H}_{j,q}^{(\alpha,c)}(x + s, y; u; \lambda) = \sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{k,q}^{(\alpha,c)}(x, y; u; \lambda) r^{n-k}, \tag{38}$$

and

$$\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) = \sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{k,q}^{(\alpha,s)}(x, y; u; \lambda) r^{n-k}. \tag{39}$$

Theorem 5. Let $j \geq 1$. Then,

$$\frac{\partial}{\partial x} \mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) = [n]_q \mathbb{H}_{n-1,q}^{(\alpha,c)}(x, y; u; \lambda), \tag{40}$$

$$\frac{\partial}{\partial y} \mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) = -[n]_q \mathbb{H}_{n-1,q}^{(\alpha,s)}(x, qy; u; \lambda), \tag{41}$$

and

$$\frac{\partial}{\partial x} \mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) = [n]_q \mathbb{H}_{n-1,q}^{(\alpha,s)}(x, y; u; \lambda), \tag{42}$$

$$\frac{\partial}{\partial y} \mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) = [n]_q \mathbb{H}_{n-1,q}^{(\alpha,c)}(x, qy; u; \lambda). \tag{43}$$

Theorem 6. Let n be a nonnegative integer, the following formulas hold true.

$$\begin{aligned} &\lambda \mathbb{H}_{n,q}^{(\alpha,c)}(1, y; u; \lambda) - u \mathbb{H}_{n,q}^{(\alpha,c)}(0, y; u; \lambda) \\ &= (1 - u) \mathbb{H}_{n,q}^{(\alpha-1,c)}(0, y; u; \lambda), \end{aligned} \tag{44}$$

$$\begin{aligned} &\lambda \mathbb{H}_{n,q}^{(\alpha,s)}(1, y; u; \lambda) - u \mathbb{H}_{n,q}^{(\alpha,s)}(0, y; u; \lambda) \\ &= (1 - u) \mathbb{H}_{n,q}^{(\alpha-1,s)}(0, y; u; \lambda), \end{aligned} \tag{45}$$

Theorem 7. The following relations hold true.

$$\mathbb{H}_{n,q}^{(\alpha+\beta,c)}(x, y; u; \lambda) = \sum_{m=0}^n \binom{n}{m}_q \mathbb{H}_{n-m,q}^{(\alpha,c)}(x, y; u; \lambda) \mathbb{H}_{m,q}^{(\beta)}(u; \lambda), \tag{46}$$

$$\mathbb{H}_{n,q}^{(\alpha+\beta,s)}(x, y; u; \lambda) = \sum_{m=0}^n \binom{n}{m}_q \mathbb{H}_{n-m,q}^{(\alpha,s)}(x, y; u; \lambda) \mathbb{H}_{m,q}^{(\beta)}(u; \lambda), \tag{47}$$

$$\mathbb{H}_{n,q}^{(\alpha-\beta,c)}(x, y; u; \lambda) = \sum_{m=0}^n \binom{n}{m}_q \mathbb{H}_{n-m,q}^{(\alpha,c)}(x, y; u; \lambda) \mathbb{H}_{m,q}^{(-\beta)}(u; \lambda), \tag{48}$$

$$\mathbb{H}_{n,q}^{(\alpha-\beta,s)}(x, y; u; \lambda) = \sum_{m=0}^n \binom{n}{m}_q \mathbb{H}_{n-m,q}^{(\alpha,s)}(x, y; u; \lambda) \mathbb{H}_{m,q}^{(-\beta)}(u; \lambda), \tag{49}$$

Theorem 8. Let x, y , and r be any real numbers. Then, we have

(i)

$$\begin{aligned} &\mathbb{H}_{n,q}^{(\alpha,c)}((x+r)_q, y; u; \lambda) + \mathbb{H}_{n,q}^{(\alpha,s)}((x-r)_q, y; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{l}_q q^{\binom{n-l}{2}} r^{n-l} \left(\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) + (-1)^{n-k} \mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) \right), \end{aligned} \tag{50}$$

(ii)

$$\begin{aligned} &\mathbb{H}_{n,q}^{(\alpha,s)}((x+r)_q, y; u; \lambda) + \mathbb{H}_{n,q}^{(\alpha,c)}((x-r)_q, y; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{l}_q q^{\binom{n-l}{2}} r^{n-l} \left(\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) + (-1)^{n-k} \mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) \right). \end{aligned} \tag{51}$$

Corollary 1. Let $j \geq 0$. Then,

$$\begin{aligned} & \mathbb{H}_{n,q}^{(c)}((x+r)_q, y; u; \lambda) + \mathbb{H}_{n,q}^{(c)}((x-r)_q, y; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q q^{\binom{k}{2}} r^k \left(\mathbb{H}_{n-k,q}^{(c)}(x, y; u; \lambda) + (-1)^k \mathbb{H}_{n-k,q}^{(c)}(x, y; u; \lambda) \right), \end{aligned} \tag{52}$$

and

$$\begin{aligned} & \mathbb{H}_{n,q}^{(s)}((x+r)_q, y; u; \lambda) + \mathbb{H}_{n,q}^{(s)}((x-r)_q, y; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q q^{\binom{k}{2}} r^k \left(\mathbb{H}_{n-k,q}^{(s)}(x, y; u; \lambda) + (-1)^k \mathbb{H}_{n-k,q}^{(s)}(x, y; u; \lambda) \right). \end{aligned} \tag{53}$$

Corollary 2. For $r = 1$ in Theorem 8, we obtain

$$\begin{aligned} & \mathbb{H}_{n,q}^{(c)}((x+1)_q, y; u; \lambda) + \mathbb{H}_{n,q}^{(s)}((x-1)_q, y; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q q^{\binom{k}{2}} r^k \left(\mathbb{H}_{n-k,q}^{(c)}(x, y; u; \lambda) + (-1)^k \mathbb{H}_{n-k,q}^{(s)}(x, y; u; \lambda) \right), \end{aligned} \tag{54}$$

and

$$\begin{aligned} & \mathbb{H}_{n,q}^{(s)}((x+1)_q, y; u; \lambda) + \mathbb{H}_{n,q}^{(c)}((x-1)_q, y; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q q^{\binom{n-k}{2}} r^{n-k} \left(\mathbb{H}_{k,q}^{(s)}(x, y; u; \lambda) + (-1)^{n-k} \mathbb{H}_{k,q}^{(c)}(x, y; u; \lambda) \right). \end{aligned} \tag{55}$$

3. Summation Formulas for q -Cosine and q -Sine Apostol-Type Frobenius–Euler Polynomials

In this section, we derive some correlations for the q -cosine and q -sine Apostol-type Frobenius–Euler polynomials of order α associated with the q -Bernoulli, Euler, and Genocchi polynomials and the q -Stirling numbers of the second kind. We first provide the following theorems.

Theorem 9. The following results hold true:

$$\begin{aligned} & (2u-1) \sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{k,q}(x, 0; u; \lambda) \mathbb{H}_{n-k,q}^{(c)}(0, y; 1-u; \lambda) \\ &= u \mathbb{H}_{n,q}^{(c)}(x, y; u; \lambda) - (1-u) \mathbb{H}_{n,q}^{(c)}(x, y; 1-u; \lambda), \end{aligned} \tag{56}$$

and

$$\begin{aligned} & (2u-1) \sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{k,q}(x, 0; u; \lambda) \mathbb{H}_{n-k,q}^{(s)}(0, y; 1-u; \lambda) \\ &= u \mathbb{H}_{n,q}^{(s)}(x, y; u; \lambda) - (1-u) \mathbb{H}_{n,q}^{(s)}(x, y; 1-u; \lambda). \end{aligned} \tag{57}$$

Proof. We set

$$\frac{(2u-1)}{(\lambda e_q(t) - u)(\lambda e_q(t) - (1-u))} = \frac{1}{\lambda e_q(t) - u} - \frac{1}{\lambda e_q(t) - (1-u)}.$$

From the above equation, we see that

$$\begin{aligned} & (2u-1) \frac{(1-u)e_q(xt)(1-(1-u))\text{COS}_q(yt)}{(\lambda e_q(t) - u)(\lambda e_q(t) - (1-u))} \\ &= \frac{(1-u)e_q(xt)u\text{COS}_q(yt)}{\lambda e_q(t) - u} - \frac{(1-u)e_q(xt)\text{COS}_q(yt)(1-(1-u))}{\lambda e_q(t) - (1-u)}, \end{aligned}$$

which when using Equations (9) and (24) on both sides, we can obtain

$$(2u - 1) \left(\sum_{k=0}^{\infty} \mathbb{H}_{k,q}(x, 0; u; \lambda) \frac{t^k}{[k]_q!} \right) \left(\sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(c)}(0, y; 1 - u; \lambda) \frac{t^n}{[n]_q!} \right) \\ = u \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(c)}(x, y; u; \lambda) \frac{t^n}{[n]_q!} - (1 - u) \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(c)}(x, y; 1 - u; \lambda) \frac{t^n}{[n]_q!}.$$

By applying the Cauchy product rule in the above equation and then equating the coefficients of like powers of t on both sides of the resultant equation, assertion (56) follows. Similarly, we obtain (57). \square

Theorem 10. *The following relations hold true:*

$$u \mathbb{H}_{n,q}^{(c)}(x, y; u; \lambda) = \sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{n,q}^{(c)}(x, y; u; \lambda) - (1 - u) C_{n,q}(x, y), \tag{58}$$

and

$$u \mathbb{H}_{n,q}^{(s)}(x, y; u; \lambda) = \sum_{k=0}^n \binom{n}{k}_q \mathbb{H}_{n,q}^{(s)}(x, y; u; \lambda) - (1 - u) S_{n,q}(x, y). \tag{59}$$

Proof. Consider the following identity

$$\frac{u}{\lambda(\lambda e_q(t) - u)e_q(t)} = \frac{1}{(\lambda e_q(t) - u)} - \frac{1}{\lambda e_q(t)}.$$

Evaluating the following fraction using the above identity, we find

$$\frac{u(1 - u)e_q(xt) \text{COS}_q(yt)}{\lambda(\lambda e_q(t) - u)e_q(t)} = \frac{(1 - u)e_q(xt) \text{COS}_q(yt)}{\lambda e_q(t) - u} - \frac{(1 - u)e_q(xt) \text{COS}_q(yt)}{\lambda e_q(t)} \\ u \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(c)}(x, y; u; \lambda) \frac{t^n}{[n]_q!} = \lambda \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(c)}(x, y; u; \lambda) \frac{t^n}{[n]_q!} \sum_{k=0}^{\infty} \frac{t^k}{[k]_q!} - (1 - u) \sum_{n=0}^{\infty} C_{n,q}(x, y) \frac{t^n}{[n]_q!}.$$

By applying the Cauchy product rule in the above equation and then equating the coefficients of like powers of t on both sides of the resultant equation, assertion (58) follows. Similarly, we obtain (59). \square

The following Theorems can be easily derived by making use of the definitions of used polynomials and series manipulations. So, we omit the proofs.

Theorem 11. *The following relation holds true:*

$$\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) \\ = \frac{1}{1 - u} \sum_{k=0}^n \binom{n}{k}_q \left[\lambda \mathbb{H}_{n-k,q}(u; \lambda) \mathbb{H}_{k,q}^{(\alpha,c)}(x, y; u; \lambda) \right. \\ \left. - u \mathbb{H}_{n-k,q}(u; \lambda) \mathbb{H}_{k,q}^{(\alpha,c)}(x, y; u; \lambda) \right]. \tag{60}$$

and

$$\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) \\ = \frac{1}{1 - u} \sum_{k=0}^n \binom{n}{k}_q \left[\lambda \mathbb{H}_{n-k,q}(u; \lambda) \mathbb{H}_{k,q}^{(\alpha,s)}(x, y; u; \lambda) \right. \\ \left. - u \mathbb{H}_{n-k,q}(u; \lambda) \mathbb{H}_{k,q}^{(\alpha,s)}(x, y; u; \lambda) \right]. \tag{61}$$

Theorem 12. *The following relations hold true:*

$$\begin{aligned} \mathbb{H}_{n,q}^{(\alpha,c)}(x,y;u;\lambda) &= \sum_{k=0}^{n+1} \binom{n+1}{k}_q \left(\lambda \sum_{r=0}^k \binom{k}{r}_q \mathbb{B}_{k-r,q}(x;\lambda) - \mathbb{B}_{k,q}(x;\lambda) \right) \\ &\quad \times \mathbb{H}_{n-k+1,q}^{(\alpha,c)}(0,y;u;\lambda). \end{aligned} \tag{62}$$

and

$$\begin{aligned} \mathbb{H}_{n,q}^{(\alpha,s)}(x,y;u;\lambda) &= \sum_{k=0}^{n+1} \binom{n+1}{k}_q \left(\lambda \sum_{r=0}^k \binom{k}{r}_q \mathbb{B}_{k-r,q}(x;\lambda) - \mathbb{B}_{k,q}(x;\lambda) \right) \\ &\quad \times \mathbb{H}_{n-k+1,q}^{(\alpha,s)}(0,y;u;\lambda). \end{aligned} \tag{63}$$

Theorem 13. *The following relations hold true:*

$$\begin{aligned} \mathbb{H}_{n,q}^{(\alpha,c)}(x,y;u;\lambda) &= \frac{1}{2} \sum_{k=0}^n \binom{n}{k}_q \left(\lambda \sum_{r=0}^k \binom{k}{r}_q \mathbb{E}_{k-r,q}(\lambda) + \mathbb{E}_{k,q}(\lambda) \right) \\ &\quad \times \mathbb{H}_{n-k,q}^{(\alpha,c)}(x,y;u;\lambda). \end{aligned} \tag{64}$$

and

$$\begin{aligned} \mathbb{H}_{n,q}^{(\alpha,s)}(x,y;u;\lambda) &= \frac{1}{2} \sum_{k=0}^n \binom{n}{k}_q \left(\lambda \sum_{r=0}^k \binom{k}{r}_q \mathbb{E}_{k-r,q}(\lambda) + \mathbb{E}_{k,q}(\lambda) \right) \\ &\quad \times \mathbb{H}_{n-k,q}^{(\alpha,s)}(x,y;u;\lambda). \end{aligned} \tag{65}$$

Theorem 14. *The following relations hold true:*

$$\begin{aligned} \mathbb{H}_{n,q}^{(\alpha,c)}(x,y;u;\lambda) &= \frac{1}{2} \sum_{k=0}^{n+1} \binom{n+1}{k}_q \left(\lambda \sum_{r=0}^k \binom{k}{r}_q \mathbb{G}_{k-r,q}(\lambda) + \mathbb{G}_{k,q}(\lambda) \right) \\ &\quad \times \mathbb{H}_{n-k+1,q}^{(\alpha,c)}(x,y;u;\lambda). \end{aligned} \tag{66}$$

and

$$\begin{aligned} \mathbb{H}_{n,q}^{(\alpha,s)}(x,y;u;\lambda) &= \frac{1}{2} \sum_{k=0}^{n+1} \binom{n+1}{k}_q \left(\lambda \sum_{r=0}^k \binom{k}{r}_q \mathbb{G}_{k-r,q}(\lambda) + \mathbb{G}_{k,q}(\lambda) \right) \\ &\quad \times \mathbb{H}_{n-k+1,q}^{(\alpha,s)}(x,y;u;\lambda). \end{aligned} \tag{67}$$

Theorem 15. *Let α and γ be nonnegative integers. The following relations hold true:*

$$\left(\frac{1-u}{u} \right)^\alpha C_{n,q}(x,y) = \alpha! \sum_{l=0}^n \binom{n}{l}_q \mathbb{H}_{n-l,q}^{(\alpha,c)}(x,y;u;\lambda) S\left(l, \alpha, \frac{\lambda}{u} : q\right), \tag{68}$$

$$\left(\frac{1-u}{u} \right)^\alpha S_{n,q}(x,y) = \alpha! \sum_{l=0}^n \binom{n}{l}_q \mathbb{H}_{n-l,q}^{(\alpha,s)}(x,y;u;\lambda) S\left(l, \alpha, \frac{\lambda}{u} : q\right), \tag{69}$$

$$\begin{aligned} &\mathbb{H}_{n,q}^{(\alpha-\gamma,c)}(x,y;u;\lambda) \\ &= \gamma! \left(\frac{u}{1-u} \right)^\gamma \alpha! \sum_{l=0}^n \binom{n}{l}_q \mathbb{H}_{n-l,q}^{(\alpha,c)}(x,y;u;\lambda) S\left(l, \alpha, \frac{\lambda}{u} : q\right), \end{aligned} \tag{70}$$

and

$$\mathbb{H}_{n,q}^{(\alpha-\gamma,s)}(x,y;u;\lambda)$$

$$= \gamma! \left(\frac{u}{1-u}\right)^\gamma \alpha! \sum_{l=0}^n \binom{n}{l}_q \mathbb{H}_{n-l,q}^{(\alpha,s)}(x, y; u; \lambda) S\left(l, \alpha, \frac{\lambda}{u} : q\right). \tag{71}$$

Theorem 16. *The following relations hold true:*

$$\begin{aligned} & \mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) \\ &= \sum_{j=0}^n \sum_{l=j}^n \binom{\alpha+j-1}{j}_q j! \binom{n}{l}_q (1-u)^{-j} S(l, j; \lambda : q) (\lambda-1)^{l-j} C_{n-l,q}(x, y). \end{aligned} \tag{72}$$

and

$$\begin{aligned} & \mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) \\ &= \sum_{j=0}^n \sum_{l=j}^n \binom{\alpha+j-1}{j}_q j! \binom{n}{l}_q (1-u)^{-j} S(l, j; \lambda : q) (\lambda-1)^{l-j} S_{n-l,q}(x, y). \end{aligned} \tag{73}$$

Theorem 17. *The following relationships hold true:*

$$\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) = \sum_{k=0}^n (\alpha)_k (u-1)^{-k} S_{n,q}^{(k,c)}(x, y; \lambda) \tag{74}$$

and

$$\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) = \sum_{k=0}^n (\alpha)_k (u-1)^{-k} S_{n,q}^{(k,s)}(x, y; \lambda), \tag{75}$$

where

$$e_q(xt) \text{COS}_q(yt) \frac{(\lambda e_q(t) - 1)^k}{[k]_q!} = \sum_{n=k}^\infty S_{n,q}^{(k,c)}(x, y; \lambda) \frac{t^n}{[n]_q!}$$

and

$$e_q(xt) \text{SIN}_q(yt) \frac{(\lambda e_q(t) - 1)^k}{[k]_q!} = \sum_{n=k}^\infty S_{n,q}^{(k,s)}(x, y; \lambda) \frac{t^n}{[n]_q!}.$$

4. Symmetry Identities for q -Cosine and q -Sine Apostol-Type Frobenius–Euler Polynomials

In this section, we describe the general symmetry identities for the q -cosine and q -sine Apostol-type Frobenius–Euler polynomials and generalized Apostol-type Frobenius–Euler polynomials by applying the generating functions (9), (24) and (25). We begin with the following theorem.

Theorem 18. *Let $a, b, > 0$ with $a \neq b$ and $j \geq 0$. Then,*

(i)

$$\begin{aligned} & \sum_{k=0}^n \binom{n}{k}_q b^k a^{n-k} \mathbb{H}_{n-k,q}^{(\alpha,c)}(bx, by; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q a^k b^{n-k} \mathbb{H}_{n-k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,c)}(bx, by; u; \lambda), \end{aligned} \tag{76}$$

(ii)

$$\begin{aligned} & \sum_{k=0}^n \binom{n}{k}_q b^k a^{n-k} \mathbb{H}_{n-k,q}^{(\alpha,s)}(bx, by; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,s)}(ax, ay; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q a^k b^{n-k} \mathbb{H}_{n-k,q}^{(\alpha,s)}(ax, ay; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,s)}(bx, by; u; \lambda). \end{aligned} \tag{77}$$

Proof. Let

$$A(z) = \left(\frac{(1-u)^2 (e_q(abxt) \text{COS}_q(abyt))^2}{(\lambda e_q(az) - u)(\lambda e_q(bz) - u)} \right)^{2\alpha}. \tag{78}$$

Then, the expression for $A(z)$ is symmetric in a and b , and we obtain

$$\begin{aligned} A(z) &= \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(\alpha,c)}(bx, by; u; \lambda) \frac{(at)^n}{[n]_q!} \sum_{k=0}^{\infty} H_{k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \frac{(bt)^k}{[k]_q!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{j}{k}_q \right) b^k a^{j-k} \mathbb{H}_{n-k,q}^{(\alpha,c)}(bx, by; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \frac{t^n}{[n]_q!}. \end{aligned}$$

Similarly, we can show that

$$\begin{aligned} A(z) &= \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(\alpha,c)}(ax, ay; u; \lambda) \frac{(bt)^n}{[n]_q!} \sum_{k=0}^{\infty} \mathbb{H}_{k,q}^{(\alpha,c)}(bx, by; u; \lambda) \frac{(at)^k}{k!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_q \right) a^k b^{n-k} \mathbb{H}_{n-k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,c)}(bx, by; u; \lambda) \frac{t^n}{[n]_q!}. \end{aligned}$$

On comparing the coefficients of t^n on the right hand sides of the last two equations, we arrive at the desired result (76). Similarly, we obtain (77). \square

Remark 2. For $\alpha = 1$ in Theorem 18, the result reduces to

(i)

$$\begin{aligned} &\sum_{k=0}^n \binom{n}{k}_q b^k a^{n-k} \mathbb{H}_{n-k,q}^{(c)}(bx, by; u; \lambda) \mathbb{H}_{k,q}^{(c)}(ax, ay; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q a^k b^{n-k} \mathbb{H}_{n-k,q}^{(c)}(ax, ay; u; \lambda) \mathbb{H}_{k,q}^{(c)}(bx, by; u; \lambda), \end{aligned} \tag{79}$$

(ii)

$$\begin{aligned} &\sum_{k=0}^n \binom{n}{k}_q b^k a^{n-k} \mathbb{H}_{n-k,q}^{(s)}(bx, by; u; \lambda) \mathbb{H}_{k,q}^{(s)}(ax, ay; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q a^k b^{n-k} \mathbb{H}_{n-k,q}^{(s)}(ax, ay; u; \lambda) \mathbb{H}_{k,q}^{(s)}(bx, by; u; \lambda). \end{aligned} \tag{80}$$

Remark 3. Assume $q \rightarrow 1$ in Theorem 18, the result reduces to

(i)

$$\begin{aligned} &\sum_{k=0}^n \binom{n}{k} b^k a^{n-k} \mathbb{H}_{n-k}^{(\alpha,c)}(bx, by; u; \lambda) \mathbb{H}_k^{(\alpha,c)}(ax, ay; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k} a^k b^{n-k} \mathbb{H}_{n-k}^{(\alpha,c)}(ax, ay; u; \lambda) \mathbb{H}_k^{(\alpha,c)}(bx, by; u; \lambda), \end{aligned} \tag{81}$$

(ii)

$$\begin{aligned} &\sum_{k=0}^n \binom{n}{k} b^k a^{n-k} \mathbb{H}_{n-k}^{(\alpha,s)}(bx, by; u; \lambda) \mathbb{H}_k^{(\alpha,s)}(ax, ay; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k} a^k b^{n-k} \mathbb{H}_{n-k}^{(\alpha,s)}(ax, ay; u; \lambda) \mathbb{H}_k^{(\alpha,s)}(bx, by; u; \lambda). \end{aligned} \tag{82}$$

Theorem 19. Let $a, b, > 0$ with $a \neq b$ and $n \geq 0$. Then,

$$\begin{aligned}
 & (i) \\
 & \sum_{k=0}^n \binom{n}{k} \sum_{q=0}^{a-1} \sum_{j=0}^{b-1} (-\lambda)^{i+j} a^{n-k} b^k \mathbb{H}_{n-k,q}^{(\alpha,c)} \left(bx + \frac{b}{a}i + j, by; u; \lambda \right) \mathbb{H}_{k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \\
 & = \sum_{k=0}^n \binom{n}{k} \sum_{q=0}^{b-1} \sum_{j=0}^{a-1} (-\lambda)^{i+j} b^{n-k} a^k \mathbb{H}_{n-k,q}^{(\alpha,c)} \left(ax + \frac{a}{b}i + j, ay; u; \lambda \right) \mathbb{H}_{k,q}^{(\alpha,c)}(bx, by; u; \lambda). \tag{83}
 \end{aligned}$$

$$\begin{aligned}
 & (ii) \\
 & \sum_{k=0}^n \binom{n}{k} \sum_{q=0}^{a-1} \sum_{j=0}^{b-1} (-\lambda)^{i+j} a^{n-k} b^k \mathbb{H}_{n-k,q}^{(\alpha,s)} \left(bx + \frac{b}{a}i + j, by; u; \lambda \right) \mathbb{H}_{k,q}^{(\alpha,s)}(ax, ay; u; \lambda) \\
 & = \sum_{k=0}^n \binom{n}{k} \sum_{q=0}^{b-1} \sum_{j=0}^{a-1} (-\lambda)^{i+j} b^{n-k} a^k \mathbb{H}_{n-k,q}^{(\alpha,s)} \left(ax + \frac{a}{b}i + j, ay; u; \lambda \right) \mathbb{H}_{k,q}^{(\alpha,s)}(bx, by; u; \lambda). \tag{84}
 \end{aligned}$$

Proof. Consider the identity

$$B(t) = \left(\frac{(1-u)^2}{(\lambda e_q(at) - u)(\lambda e_q(bt) - u)} \right)^{2\alpha} \frac{1 + \lambda(-1)^{a+1} e^{abz}}{(\lambda e^{at} + 1)(\lambda e^{bz} + 1)} (e_q(abxt) \text{COS}_q(abyt))^2. \tag{85}$$

$$\begin{aligned}
 B(t) & = \left(\frac{1-u}{\lambda e_q(az) - u} \right)^\alpha e_q(abxt) \text{COS}_q(abyt) \left(\frac{1 - \lambda(-e_q(-bt))^a}{\lambda e_q(bt) + 1} \right) \left(\frac{1-u}{\lambda e_q(bz) - u} \right)^\alpha \\
 & \quad \times \left(\frac{1 - \lambda(-e_q(-az))^b}{\lambda e_q(az) + 1} \right) e_q(abxt) \text{COS}_q(abyt) \\
 & = \left(\frac{1-u}{\lambda e_q(az) - u} \right)^\alpha e_q(abxt) \text{COS}_q(abyt) \sum_{i=0}^{a-1} (-\lambda)^i e_q(bzi) \left(\frac{1-u}{\lambda e_q(bt) - u} \right)^\alpha e_q(abxt) \text{COS}_q(abyt) \sum_{j=0}^{b-1} (-\lambda)^j e_q(azj) \\
 & = \left(\frac{1-u}{\lambda e_q(az) - u} \right)^\alpha \text{Cos}_q(abyt) \sum_{i=0}^{a-1} \sum_{j=0}^{b-1} (-\lambda)^{i+j} e_q \left(\left(bx + \frac{b}{a}i + j \right) at \right) \sum_{k=0}^\infty \mathbb{H}_{k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \frac{(bt)^k}{[k]_q!} \\
 & = \sum_{n=0}^\infty \sum_{i=0}^{a-1} \sum_{j=0}^{b-1} (-\lambda)^{i+j} \mathbb{H}_{n,q}^{(\alpha,c)} \left(bx + \frac{b}{a}i + j, by; u; \lambda \right) \frac{(at)^n}{[n]_q!} \sum_{k=0}^\infty \mathbb{H}_{k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \frac{(bt)^k}{[k]_q!} \\
 & = \sum_{n=0}^\infty \sum_{k=0}^n \binom{n}{k} \sum_{q=0}^{a-1} \sum_{j=0}^{b-1} (-\lambda)^{i+j} a^{s-k} b^k \mathbb{H}_{n-k,q}^{(\alpha,c)} \left(byx + \frac{b}{a}i + j, b\eta; u; \lambda \right) \\
 & \quad \times \mathbb{H}_{k,q}^{(\alpha,c)}(ax, ay; u; \lambda) \frac{t^n}{[n]_q!}. \tag{86}
 \end{aligned}$$

On the other hand, we obtain

$$\begin{aligned}
 B(t) & = \sum_{n=0}^\infty \sum_{k=0}^n \binom{n}{k} \sum_{q=0}^{b-1} \sum_{j=0}^{a-1} (-\lambda)^{i+j} b^{n-k} a^k \mathbb{H}_{n-k,q}^{(\alpha,c)} \left(ax + \frac{a}{b}i + j, ay; u; \lambda \right) \\
 & \quad \times \mathbb{H}_{k,q}^{(\alpha,c)}(bx, by; u; \lambda) \frac{t^n}{[n]_q!}. \tag{87}
 \end{aligned}$$

By using (86) and (87), we arrive at the desired result (82). Similarly, we obtain (83). \square

Theorem 20. Let $a, b, > 0$ with $a \neq b$ and $j \geq 0$. Then,

$$\begin{aligned} & \sum_{k=0}^n \binom{n}{k}_q b^k a^{n-k} \mathbb{H}_{n-k,q}^{(\alpha,c)}(bx, by; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,s)}(ax, ay; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k}_q a^k b^{n-k} \mathbb{H}_{n-k,q}^{(\alpha,s)}(ax, ay; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,c)}(bx, by; u; \lambda). \end{aligned} \tag{88}$$

Proof. Suppose that

$$A(t) = \left(\frac{(1-u)^2 (e_q(abt))^2 \text{COS}_q(abyt) \text{SIN}_q(abyt)}{(\lambda e^{az} - u)(\lambda e^{bz} - u)} \right)^{2\alpha}. \tag{89}$$

Then, the expression for $A(t)$ is symmetric in a and b , and we obtain

$$\begin{aligned} C(t) &= \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(\alpha,c)}(bx, by; u; \lambda) \frac{(at)^n}{[n]_q!} \sum_{k=0}^{\infty} \mathbb{H}_{k,q}^{(\alpha,s)}(ax, ay; u; \lambda) \frac{(bt)^k}{[k]_q!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_q b^k a^{j-k} \mathbb{H}_{n-k,q}^{(\alpha,c)}(bx, by; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,s)}(ax, ay; u; \lambda) \right) \frac{t^n}{[n]_q!}. \end{aligned}$$

Similarly, we can show that

$$\begin{aligned} A(z) &= \sum_{n=0}^{\infty} \mathbb{H}_{n,q}^{(\alpha,s)}(ax, ay; u; \lambda) \frac{(bt)^n}{n!} \sum_{k=0}^{\infty} \mathbb{H}_{k,q}^{(\alpha,c)}(bx, by; u; \lambda) \frac{(at)^k}{k!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_q a^k b^{n-k} \mathbb{H}_{n-k,q}^{(\alpha,s)}(ax, ay; u; \lambda) \mathbb{H}_{k,q}^{(\alpha,c)}(bx, by; u; \lambda) \right) \frac{t^n}{n!}. \end{aligned}$$

On comparing the coefficients of t^n on the right hand sides of the last two equations, we arrive at the desired result (88). \square

Remark 4. Assume that $q \rightarrow 1$ in Theorem 18, for which the result reduces to

$$\begin{aligned} & \sum_{k=0}^n \binom{n}{k} b^k a^{n-k} \mathbb{H}_{n-k}^{(\alpha,c)}(bx, by; u; \lambda) \mathbb{H}_k^{(\alpha,s)}(ax, ay; u; \lambda) \\ &= \sum_{k=0}^n \binom{n}{k} a^k b^{n-k} \mathbb{H}_{n-k}^{(\alpha,s)}(ax, ay; u; \lambda) \mathbb{H}_k^{(\alpha,c)}(bx, by; u; \lambda). \end{aligned} \tag{90}$$

5. Symmetric Structure of Approximate Roots for q -Cosine Apostol-Type Frobenius–Euler Polynomials and Their Application

In this section, certain zeros of the q -Cosine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda)$ and graphical representations are shown.

A few of them are as follows:

$$\begin{aligned} \mathbb{H}_{0,q}^{(\alpha,c)}(x,y;u;\lambda) &= \left(\frac{-1+u}{u-\lambda}\right)^\alpha, \\ \mathbb{H}_{1,q}^{(\alpha,c)}(x,y;u;\lambda) &= -\frac{ux\left(\frac{-1+u}{u-\lambda}\right)^\alpha}{-u+\lambda} + \frac{x\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{-u+\lambda} - \frac{\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{-u+\lambda}, \\ \mathbb{H}_{2,q}^{(\alpha,c)}(x,y;u;\lambda) &= \frac{x^2\left(\frac{-1+u}{u-\lambda}\right)^\alpha}{(1-q)(1+q)} - \frac{q^2x^2\left(\frac{-1+u}{u-\lambda}\right)^\alpha}{(1-q)(1+q)} - \frac{y^2\left(\frac{-1+u}{u-\lambda}\right)^\alpha}{(1-q)(1+q)} + \frac{q^2y^2\left(\frac{-1+u}{u-\lambda}\right)^\alpha}{(1-q)(1+q)} \\ &+ \frac{u\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{(1-q)(1+q)(-u+\lambda)^2} - \frac{q^2u\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{(1-q)(1+q)(-u+\lambda)^2} - \frac{\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)(1+q)(-u+\lambda)^2} \\ &+ \frac{q\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)(1+q)(-u+\lambda)^2} + \frac{q^2\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)(1+q)(-u+\lambda)^2} - \frac{q^3\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)(1+q)(-u+\lambda)^2} \\ &+ \frac{\alpha^2\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)(1+q)(-u+\lambda)^2} + \frac{q\alpha^2\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)(1+q)(-u+\lambda)^2} - \frac{q^2\alpha^2\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)(1+q)(-u+\lambda)^2} \\ &- \frac{q^3\alpha^2\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)(1+q)(-u+\lambda)^2} - \frac{x\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{(1-q)(-u+\lambda)} + \frac{q^2x\alpha\left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{(1-q)(-u+\lambda)}. \end{aligned}$$

We investigate the zeros of the q -Cosine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,c)}(x,y;u;\lambda)$ by using a computer. We plot the zeros of the q -Cosine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,c)}(x,y;u;\lambda) = 0$ for $n = 20$ (Figure 1).




Figure 1. Zeros of $\mathbb{H}_{n,q}^{(\alpha,c)}(x,y;u;\lambda)$.

In Figure 1 (top-left), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{1}{10}, y = 3$. In Figure 1 (top-right), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{3}{10}, y = 3$. In Figure 1 (bottom-left), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{5}{10}, y = 3$. In Figure 1 (bottom-right), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{7}{10}, y = 3$.

Stacks of zeros of the q -Cosine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) = 0$ for $1 \leq n \leq 20$, forming a 3D structure, are presented (Figure 2).




Figure 2. Zeros of $\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda)$.

In Figure 2 (top-left), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{1}{10}, y = 3$. In Figure 2 (top-right), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{3}{10}, y = 3$. In Figure 2 (bottom-left), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{5}{10}, y = 3$. In Figure 2 (bottom-right), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{7}{10}, y = 3$.

Next, we calculated an approximate solution satisfying the q -Cosine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,c)}(x, y; u; \lambda) = 0$ for $q = \frac{1}{10}$. The results are provided in Table 1.

Table 1. Approximate solutions of $\mathbb{H}_{n,q}^{(4,c)}(x, 3; 3; 2) = 0$.

Degree n	x
1	−8.0000
2	−4.4000 − 4.8621 i, −4.4000 + 4.8621 i
3	−6.5429, −1.1686 − 5.5743 i, −1.1686 + 5.5743 i
4	−5.1586 − 2.9339 i, −5.1586 + 2.9339 i, 0.7146 − 5.2176 i, 0.7146 + 5.2176 i
5	−5.8344, −3.3214 − 4.2756 i, −3.3214 + 4.2756 i, 1.7942 − 4.6626 i, 1.7942 + 4.6626 i
6	−5.1069 − 2.0266 i, −5.1069 + 2.0266 i, −1.7739 − 4.7670 i, −1.7739 + 4.7670 i, 2.4363 − 4.1334 i, 2.4363 + 4.1334 i
7	−5.4063, −3.9859 − 3.2720 i, −3.9859 + 3.2720 i, −0.5893 − 4.8378 i, −0.5893 + 4.8378 i, 2.8338 − 3.6774 i, 2.8338 + 3.6774 i
8	−4.9566 − 1.5254 i, −4.9566 + 1.5254 i, −2.8747 − 3.9790 i, −2.8747 + 3.9790 i, 0.2982 − 4.7116 i, 0.2982 + 4.7116 i, 3.0887 − 3.2950 i, 3.0887 + 3.2950 i
9	−5.1150, −4.2106 − 2.6028 i, −4.2106 + 2.6028 i, −1.8968 − 4.3377 i, −1.8968 + 4.3377 i, 0.9638 − 4.5004 i, 0.9638 + 4.5004 i, 3.2566 − 2.9754 i, 3.2566 + 2.9754 i

6. Symmetric Structure of Approximate Roots for q-Sine Apostol-Type Frobenius–Euler Polynomials and Their Application

In this section, certain zeros of the q-Sine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda)$ and beautiful graphical representations are shown.

A few of them are as follows:

$$\begin{aligned} \mathbb{H}_{0,q}^{(\alpha,s)}(x, y; u; \lambda) &= 0, \\ \mathbb{H}_{1,q}^{(\alpha,s)}(x, y; u; \lambda) &= y \left(\frac{-1+u}{u-\lambda} \right)^\alpha, \\ \mathbb{H}_{2,q}^{(\alpha,s)}(x, y; u; \lambda) &= -\frac{uxy \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)(-u+\lambda)} + \frac{q^2uxy \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)(-u+\lambda)} + \frac{xy \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda}{(1-q)(-u+\lambda)} \\ &\quad - \frac{q^2xy \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda}{(1-q)(-u+\lambda)} - \frac{y\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda}{(1-q)(-u+\lambda)} + \frac{q^2y\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda}{(1-q)(-u+\lambda)}, \\ \mathbb{H}_{3,q}^{(\alpha,s)}(x, y; u; \lambda) &= \frac{x^2y \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)^2(1+q)} - \frac{q^2x^2y \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)^2(1+q)} - \frac{q^3x^2y \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)^2(1+q)} \\ &\quad + \frac{q^5x^2y \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)^2(1+q)} - \frac{q^3y^3 \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)^2(1+q)(1+q+q^2)} + \frac{q^5y^3 \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)^2(1+q)(1+q+q^2)} \\ &\quad + \frac{q^6y^3 \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)^2(1+q)(1+q+q^2)} - \frac{q^8y^3 \left(\frac{-1+u}{u-\lambda} \right)^\alpha}{(1-q)^2(1+q)(1+q+q^2)} + \frac{uy\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda}{(1-q)^2(1+q)(-u+\lambda)^2} \\ &\quad - \frac{q^2uy\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda}{(1-q)^2(1+q)(-u+\lambda)^2} - \frac{q^3uy\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda}{(1-q)^2(1+q)(-u+\lambda)^2} + \frac{q^5uy\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda}{(1-q)^2(1+q)(-u+\lambda)^2} \\ &\quad - \frac{y\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} + \frac{qy\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} + \frac{q^2y\alpha \left(\frac{-1+u}{u-\lambda} \right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} \end{aligned}$$

$$\begin{aligned}
 & - \frac{q^4 y \alpha \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} - \frac{q^5 y \alpha \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} + \frac{q^6 y \alpha \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} \\
 & + \frac{y \alpha^2 \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} + \frac{q y \alpha^2 \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} - \frac{q^2 y \alpha^2 \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} \\
 & - \frac{q^3 y \alpha^2 \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{(1-q)^2(1+q)(-u+\lambda)^2} - \frac{q^4 y \alpha^2 \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} + \frac{q^5 y \alpha^2 \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} \\
 & + \frac{q^6 y \alpha^2 \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda^2}{2(1-q)^2(1+q)(-u+\lambda)^2} - \frac{x y \alpha \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{(1-q)^2(-u+\lambda)} + \frac{q^2 x y \alpha \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{(1-q)^2(-u+\lambda)} \\
 & + \frac{q^3 x y \alpha \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{(1-q)^2(-u+\lambda)} - \frac{q^5 x y \alpha \left(\frac{-1+u}{u-\lambda}\right)^\alpha \lambda}{(1-q)^2(-u+\lambda)}
 \end{aligned}$$

In Figure 3 (top-left), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{2}{10}, y = 3$. In Figure 3 (top-right), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{4}{10}, y = 3$. In Figure 3 (bottom-left), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{6}{10}, y = 3$. In Figure 3 (bottom-right), we choose $\alpha = 4, \lambda = 2, u = 3$, and $q = \frac{8}{10}, y = 3$.




Figure 3. Zeros of $\mathbb{H}_{n,q}^{(\alpha,s)}(x,y;u;\lambda)$.

Stacks of zeros of the q -Sine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) = 0$ for $2 \leq n \leq 20$, forming a 3D structure, are presented (Figure 4).




Figure 4. Zeros of $\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda)$.

In Figure 4 (top-left), we plot stacks of zeros of $\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) = 0$ for $2 \leq n \leq 30$, $q = \frac{8}{10}$, $\alpha = 4$, $\lambda = 2$, $u = 3$, $y = 3$. In Figure 4 (top-right), we draw x and y axes but no z axis in three dimensions. In Figure 4 (bottom-left), we draw y and z axes but no x axis in three dimensions. In Figure 4 (bottom-right), we draw x and z axes but no y axis in three dimensions.

Next, we calculated an approximate solution satisfying the q -Sine Apostol-type Frobenius–Euler polynomials $\mathbb{H}_{n,q}^{(\alpha,s)}(x, y; u; \lambda) = 0$ for $q = \frac{8}{10}$. The results are given in Table 2.

Table 2. Approximate solutions of $\mathbb{H}_{n,q}^{(4,s)}(x, 3; 3; 2) = 0$.

Degree n	x
2	−8.0000
3	−7.2000 − 5.1256 i, −7.2000 + 5.1256 i
4	−9.0707, −5.2247 − 8.2772 i, −5.2247 + 8.2772 i
5	−8.7853 − 3.8486 i, −8.7853 + 3.8486 i, −3.0227 − 10.2577 i, −3.0227 + 10.2577 i
6	−9.8793, −7.6190 − 6.7293 i, −7.6190 + 6.7293 i, −0.8878 − 11.4618 i, −0.8878 + 11.4618 i
7	−9.7452 − 3.1858 i, −9.7452 + 3.1858 i, −6.0868 − 8.8578 i, −6.0868 + 8.8578 i, 1.0748 − 12.1326 i, 1.0748 + 12.1326 i
8	−10.501, −8.9527 − 5.7661 i, −8.9527 + 5.7661 i, −4.4346 − 10.3981 i, −4.4346 + 10.3981 i, 2.8319 − 12.4327 i, 2.8319 + 12.4327 i
9	−10.4252 − 2.7531 i, −10.4252 + 2.7531 i, −7.8126 − 7.8189 i, −7.8126 + 7.8189 i, −2.7879 − 11.4801 i, −2.7879 + 11.4801 i, 4.3811 − 12.4748 i, 4.3811 + 12.4748 i
10	−10.985, −9.8411 − 5.0793 i, −9.8411 + 5.0793 i, −6.5042 − 9.4234 i, −6.5042 + 9.4234 i, −1.2119 − 12.2069 i, −1.2119 + 12.2069 i, 5.7340 − 12.3389 i, 5.7340 + 12.3389 i
11	−10.9347 − 2.4373 i, −10.9347 + 2.4373 i, −8.9518 − 7.0112 i, −8.9518 + 7.0112 i, −5.1343 − 10.6536 i, −5.1343 + 10.6536 i, 0.2605 − 12.6598 i, 0.2605 + 12.6598 i, 6.9078 − 12.0821 i, 6.9078 + 12.0821 i

7. Conclusions

By making use of q -numbers and q -concepts, Jang et al. [2,4] defined q -Bernoulli polynomials and numbers, q -Genocchi polynomials and numbers and q -Euler polynomials and numbers and provided some new and interesting identities and formulae. With this viewpoint, several authors have introduced q -analogues of special numbers and polynomials and have investigated their properties. In this paper, by making use of the q -cosine polynomials and q -sine polynomials, we have considered a new class of q -analogues of Apostol-type Frobenius–Euler polynomials and have obtained new properties and identities. In addition, we have analysed the behaviour of q -integral and q -derivative representations. Additionally, we have checked the roots and graphical representations of these polynomials by making use of Mathematica software. This approach led us to consider different methods, and special cases of used variables of newly defined polynomial in the paper. In this viewpoint, we will try to continue working on newly considered polynomials in this line.

Author Contributions: All authors contributed equally to the manuscript and written, read, and approved the final manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the National Natural Science Foundation of China (No. 62172116) and the Basic Research Programs of Guizhou Province (No. QianKeHe ZK[2023]279).

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to thank the reviewers who have improved the presentation of the paper substantially.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Alam, N.; Khan, W.A.; Ryoo, C.S. A note on Bell-based Apostol-type Frobenius-Euler polynomials of complex variable with its certain applications. *Mathematics* **2022**, *10*, 2109. [[CrossRef](#)]
2. Kang, J.Y.; Ryoo, C.S. Various structures of the roots and explicit properties of q -cosine Bernoulli polynomials and q -sine Bernoulli polynomials. *Mathematics* **2020**, *8*, 463. [[CrossRef](#)]
3. Muhiuddin, G.; Khan, W.A.; Al-Kadi, D. Construction on the degenerate poly-Frobenius-Euler polynomials of complex variable. *J. Funct. Spaces* **2021**, *2021*, 3115424. [[CrossRef](#)]
4. Ryoo, C.S.; Kang, J.Y. Explicit properties of q -Cosine and q -Sine Euler polynomials containing symmetric structures. *Symmetry* **2020**, *12*, 1247. [[CrossRef](#)]
5. Kim, D.S.; Kim, T.; Lee, H. A Note on Degenerate Euler and Bernoulli Polynomials of Complex Variable. *Symmetry* **2019**, *11*, 1168. [[CrossRef](#)]
6. Kim, T.; Ryoo, C.S. Some Identities for Euler and Bernoulli Polynomials and Their Zeros. *Axioms* **2018**, *7*, 56. 10.3390/axioms7030056. [[CrossRef](#)]
7. Masjed-Jamei, M.; Beyki, M.R.; Koepf, W. A New Type of Euler Polynomials and Numbers. *Mediterr. J. Math.* **2018**, *15*, 138. [[CrossRef](#)]
8. Srivastava, H.M.; Masjed-Jamei, M.; Beyki, M.R. A Parametric Type of the Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi Polynomials. *Appl. Math. Inf. Sci.* **2018**, *12*, 907–916. [[CrossRef](#)]
9. Arjika, S. On q^2 -Trigonometric functions and their q^2 -Fourier transform. *J. Math. Syst. Sci.* **2019**, *9*, 130–135. [[CrossRef](#)]
10. Koekoek, R.; Lesky, P.A.; Swarttouw, R.F. *Hypergeometric Orthogonal Polynomials and Their q -Analogues*; Springer: Berlin, Germany, 2010.
11. Kurt, B. A note on the Apostol type q -Frobenius-Euler polynomials and generalizations of the Srivastava-Pinter addition theorems. *Filomat* **2016**, *30*, 65–72. [[CrossRef](#)]
12. Kang, J.Y.; Khan, W.A. A new class of q -Hermite based Apostol type Frobenius Genocchi polynomials. *Commun. Korean Math. Soc.* **2020**, *35*, 759–771.
13. Kim, T.; Kim, D.S.; Jang, L.C.; Kim, H.-Y. On type 2 degenerate Bernoulli and Euler polynomials of complex variable. *Adv. Differ. Equ.* **2019**, *2019*, 490. [[CrossRef](#)]
14. Kac, V.; Cheung, P. *Quantum Calculus*; Springer: New York, NY, USA, 2001.
15. Mahmudov, N.I. q -analogues of the Bernoulli and Genocchi polynomials and the Srivastava-Pinter addition theorems. *Discrete Dyn. Nat. Soc.* **2012**, *2012*, 169348. [[CrossRef](#)]
16. Mahmudov, N.I. On a class of q -Bernoulli and q -Euler polynomials. *Adv. Differ. Equ.* **2013**, *2013*, 108. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.