



van Tuijl, L., Basten, M., Pan, K-Y., Vermeulen, R., Boyd, A. W., & Teyhan, A. (2023). Depression, anxiety, and the risk of cancer: An individual participant data meta-analysis. *Cancer*.
<https://doi.org/10.1002/cncr.34853>

Publisher's PDF, also known as Version of record

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1002/cncr.34853](https://doi.org/10.1002/cncr.34853)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://doi.org/10.1002/cncr.34853>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research


General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Use Signatera™ ctDNA dynamics to inform earlier treatment decisions in metastatic melanoma patients

Early on-treatment ctDNA dynamics were predictive of PFS in metastatic melanoma patients receiving 1st line ICI treatment¹


At week 6, Signatera™ identified that patients with increasing ctDNA had a 18x higher risk of progression than ctDNA-negative patients



- > Patients with any increase in ctDNA levels from baseline by week 6 of 1st Line ICI treatment (monotherapy and combination ICIs) had a significantly shorter PFS (HR: 18; p=0.013).
- > Signatera™ was able to help distinguish between true vs pseudo-progression

Should treatment be changed or escalated?

Early rise in ctDNA can help inform treatment escalation or change



Covered by Medicare for immunotherapy treatment response monitoring across all stages for solid tumors



PFS = Progression-free survival

References:

- Eroglu Z, Krinshpun S, Kalashnikova E, et al. Circulating tumor DNA based molecular residual disease detection for treatment monitoring in advanced melanoma patients. *Cancer* (2023). <https://doi.org/10.1002/cncr.34716>
- Eroglu Z, Krinshpun S, Martin J, et al. Molecular residual disease detection and circulating tumor DNA dynamics during treatment in patients with advanced melanoma. Abstract presented at: 10th World Congress of Melanoma; April 15-17, 2021; virtual meeting. #P-096. <https://worldmelanoma2021.com/>

ORIGINAL ARTICLE

Depression, anxiety, and the risk of cancer: An individual participant data meta-analysis

Lonneke A. van Tuijl PhD^{1,2}  | Maartje Basten PhD³ | Kuan-Yu Pan PhD^{4,5,6} |
 Roel Vermeulen PhD⁷ | Lützen Portengen PhD⁷ | Alexander de Graeff MD, PhD⁸ |
 Joost Dekker PhD^{4,5}  | Mirjam I. Geerlings PhD^{3,9,10,11} |
 Adriaan Hoogendoorn PhD^{4,12} | Femke Lamers PhD^{4,5} | Adri C. Voogd PhD^{13,14} |
 Jessica Abell PhD¹⁵ | Philip Awadalla PhD^{16,17,18} | Aartjan T. F. Beekman PhD⁴ |
 Ottar Bjerkeset PhD^{19,20} | Andy Boyd BA²¹ | Yunsong Cui MSc²² |
 Philipp Frank MSc¹⁵ | Henrike Galenkamp PhD²³ | Bert Garssen PhD¹ |
 Sean Hellingman MSc²⁴ | Martijn Huisman PhD^{25,26} | Anke Huss PhD⁷ |
 Trynke R. de Jong PhD²⁷ | Melanie R. Keats PhD²⁸ | Almar A. L. Kok PhD^{4,25} |
 Steinar Krokstad PhD^{29,30} | Flora E. van Leeuwen PhD³¹ | Annemarie I. Luik PhD³² |
 Nolwenn Noisel PhD³³ | N. Charlotte Onland-Moret PhD³ | Yves Payette MSc³³ |
 Brenda W. J. H. Penninx PhD⁴ | Ina Rissanen MD, PhD³ | Annelieke M. Roest PhD³⁴ |
 Rikje Rutter MD, PhD^{32,35} | Robert A. Schoevers MD, PhD³⁶ | David Soave PhD^{16,24} |
 Mandy Spaan PhD³¹ | Andrew Steptoe DSc¹⁵ | Karien Stronks PhD^{23,37} |
 Erik R. Sund PhD^{19,29,30} | Ellen Sweeney PhD²² | Emma L. Twait MSc^{3,9,10} |
 Alison Teyhan PhD²¹ | W. M. Monique Verschuren PhD^{3,38} |
 Kimberly D. van der Willik PhD^{31,32}  | Judith G. M. Rosmalen PhD^{36,39} |
 Adelita V. Ranchor PhD¹

Correspondence

Lonneke A. van Tuijl, Health Psychology
 Section, HPC FA12, University Medical Center
 Groningen, Hanzeplein 1, Postbus 30 001,
 Groningen 9700 RB, the Netherlands.
 Email: l.a.van.tuijl@umcg.nl

Funding information

Dutch Cancer Society, Grant/Award Number:
 VU2017-8288; UK Medical Research Council
 and Wellcome, Grant/Award Number:
 217065/Z/19/Z; University of Bristol;
 Netherlands Organisation for Health Research
 and Development, Grant/Award Number:
 10-000-1002; National Institute on Aging,
 Grant/Award Number: R01AG017644;

Abstract

Background: Depression and anxiety have long been hypothesized to be related to an increased cancer risk. Despite the great amount of research that has been conducted, findings are inconclusive. To provide a stronger basis for addressing the associations between depression, anxiety, and the incidence of various cancer types (overall, breast, lung, prostate, colorectal, alcohol-related, and smoking-related cancers), individual participant data (IPD) meta-analyses were performed within the Psychosocial Factors and Cancer Incidence (PSY-CA) consortium.

Methods: The PSY-CA consortium includes data from 18 cohorts with measures of depression or anxiety (up to $N = 319,613$; cancer incidences, 25,803; person-years

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. Cancer published by Wiley Periodicals LLC on behalf of American Cancer Society.

National Institute for Health and Care
Research, Grant/Award Number: 198/1074-
02

of follow-up, 3,254,714). Both symptoms and a diagnosis of depression and anxiety were examined as predictors of future cancer risk. Two-stage IPD meta-analyses were run, first by using Cox regression models in each cohort (stage 1), and then by aggregating the results in random-effects meta-analyses (stage 2).

Results: No associations were found between depression or anxiety and overall, breast, prostate, colorectal, and alcohol-related cancers. Depression and anxiety (symptoms and diagnoses) were associated with the incidence of lung cancer and smoking-related cancers (hazard ratios [HRs], 1.06–1.60). However, these associations were substantially attenuated when additionally adjusting for known risk factors including smoking, alcohol use, and body mass index (HRs, 1.04–1.23).

Conclusions: Depression and anxiety are not related to increased risk for most cancer outcomes, except for lung and smoking-related cancers. This study shows that key covariates are likely to explain the relationship between depression, anxiety, and lung and smoking-related cancers.

Preregistration number: https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=157677.

KEYWORDS

anxiety, cancer, depression, meta-analysis, risk

INTRODUCTION

Depression and anxiety have long been hypothesized to increase the risk for cancer. It is thought that the increased cancer risk can occur via several pathways, including health behaviors, or by influencing mutation, viral oncogenes, cell proliferation, or DNA repair.¹ Conclusions drawn in meta-analyses vary greatly, with some supporting an association between depression, anxiety, and cancer incidence^{2–4} and others finding no or a negligible association.^{5–7} These meta-analyses are limited by one or more of the following: inclusion of only published studies, which results in possible underrepresentation of null findings; inclusion of studies that do not use prospective designs; combination of different cancer outcomes, which may hide effects for specific cancer types; limited adjustment for potential confounding factors; or lack of consistency in measurements of depression and anxiety.

Individual participant data (IPD) meta-analysis uses crude data from studies. This has several key advantages that lead to more reliable results than traditional meta-analyses.⁸ First, via the harmonization of data, constructs across studies are, conceptually, as similar as possible. Second, there is more consistency across studies with regard to the statistical model tested and the covariates adjusted for.⁹ Third, data can be included that have not previously been published or used to test the research hypothesis, which reduces the risk of overestimating effects due to publication bias. In the present study, we tested whether depression and anxiety were associated with increased cancer risk by means of IPD meta-analysis. In addition to looking at overall cancers, we also consider the four most prevalent cancers (breast, lung, prostate, and colorectal) and

cancers with established common lifestyle risk factors (smoking-related cancers and alcohol-related cancers). We hypothesized that both depression and anxiety are associated with increased risk for overall cancers and breast, prostate, lung, colorectal, smoking-related, and alcohol-related cancers.

MATERIALS AND METHODS

PSY-CA consortium

The international Psychosocial Factors and Cancer Incidence (PSY-CA) consortium was established to investigate whether psychosocial factors are associated with an increased risk of cancer. The present study focuses specifically on depression and anxiety. An overview of the various research questions can be found in the preregistration in PROSPERO (<http://www.crd.york.ac.uk/PROSPERO>) with the following IDs: CRD42020157677, CRD42020181623, and CRD42020193716.

Inclusion/exclusion criteria

Fourteen cohorts and four subcohorts (10 [sub]cohorts in the Netherlands and eight [sub]cohorts¹ in the United Kingdom,

¹In some cohorts, inclusion criteria or depression/anxiety instruments changed over time or there was a long recruitment period over many years. In these instances, the cohorts divided their data into subcohorts to increase homogeneity within cohorts.

Norway, and Canada) were included on the basis of the following criteria: (1) a valid and reliable measure of depression or anxiety (or another psychosocial factor¹⁰; in the present article, we focus on depression and anxiety); (2) an objective measure of cancer diagnosis during follow-up is available or possible to attain; (3) an assessment of smoking behavior, alcohol use, sex, and age; and (4) a prospective design (i.e., depression, anxiety, or other psychosocial factors were measured before cancer incidence). A thorough overview of the design of PSY-CA is provided in the protocol paper.¹⁰

Study design

A two-stage approach to IPD meta-analysis was used in PSY-CA. During the first stage, cohort-specific data harmonization instructions, based on Maelstrom guidelines, were provided to each cohort.¹¹ Subsequently, local researchers from each cohort ran standardized R scripts over their harmonized data sets and sent the output to the central researcher (L.A.v.T.) who, during the second stage, pooled the estimated effects from stage 1 using random-effects meta-analyses.

Variables

Assessment of depression and anxiety

Depression and anxiety were conceptualized and analyzed in two ways: diagnoses and symptoms. Diagnoses were based on clinical interviews (if available) or scoring above a clinically validated cutoff on self-report questionnaires.² Depression diagnosis was based on meeting the *Diagnostic and Statistical Manual of Mental Disorders (DSM)* criteria for a current major depressive disorder or dysthymia. Anxiety diagnosis was based on the presence of a current generalized anxiety disorder, social anxiety, panic disorder (with or without agoraphobia), and/or agoraphobia, according to *DSM* criteria. Depression and anxiety symptoms were assessed by using self-report questionnaires that covered symptoms in the previous week(s). Continuous sum scores were converted to z scores in each cohort, with higher scores indicative of more severe symptoms (see Supplementary Materials 1).

Cancer incidence

Cohorts based in the Netherlands obtained information about cancer incidence including site, morphology, and diagnosis date from the Netherlands Cancer Registry. Cohorts outside of the

Netherlands obtained information from country/province-specific registries. In four cohorts (RS1, RS2, RS3, and CARTa-GENE), information from registries was supplemented with data on hospital visits, insurance claims, and general practitioner records.

Participants who had a history of cancer or cancer at baseline were excluded, except when the cancer was (nonmelanoma) skin cancer. Participants diagnosed with any cancer (including carcinoma in situ and tumors where malignancy could not be determined) within 1 year from baseline were also excluded to reduce the likelihood of reverse causality.

In the main analyses, cancer incidence included malignant tumors, tumors where malignancy could not be determined (i.e., borderline), and carcinoma in situ. Seven cancer outcomes were analyzed: overall cancers and breast, lung, prostate, colorectal, smoking-related, and alcohol-related cancers as listed by the International Agency for Research on Cancer¹² (see Supplementary Materials 2 for *International Classification of Diseases* codes).

Covariates

In the minimally adjusted models, we adjusted for birth year (age-period cohort effects), sex, country of birth, and education level across all cohorts. In the maximally adjusted models, we further adjusted for a number of health-related covariates depending on cancer outcome and availability within the cohort:

- all outcomes: weekly alcohol intake, current antidepressant use, body mass index, hours of physical activity per week (metabolic equivalent, if available), smoking status (former, current, or never), pack-years (or current number smoked), and family history of cancer;
- breast cancer: parity (0 pregnancies, 1 or 2 pregnancies, or 3+ pregnancies), (hormonal) contraceptive use, age at menarche, menopause status, and family history of breast cancer;
- colorectal cancer: sedentary behavior and family history of colorectal cancer; and
- lung and prostate cancer: family history of lung or prostate cancer, respectively.

A full overview of covariates in the maximally adjusted models per cohort is given in Supplementary Materials 3.

Statistical analysis

Stage 1: Local analyses

Cox regression models (minimally or maximally adjusted; see the Covariates section) were used in all analyses to calculate hazard ratios (HRs) and 95% confidence intervals (CIs), with entry age (age

²Although diagnoses cannot be made on the basis of self-report questionnaires alone, we apply validated cutoffs that have been found to show a sufficient degree of sensitivity and specificity. We refer to scores above the cutoff as "diagnosis."

at baseline) and exit age (age at diagnosis, death, or dropout/study end) as timescale.

Only participants without missing information on depression or anxiety were included in the analyses. Covariates were included if they were available for at least 60% of the sample. Extreme values for continuous covariates were truncated to three times the interquartile range above the third quartile or below the first quartile. For the covariates related to physical activity, alcohol use, and smoking, only upper extreme outliers were truncated because a substantial number reported 0 (e.g., never smokers).

Stage 2: Meta-analyses

For each meta-analysis, we selected models (from stage 1) that included at least 10 cancer events and, for depression and anxiety diagnosis, at least five expected events (cancer incidences) in the smallest category of depression/anxiety diagnosis (based on the observed cohort-specific cancer incidence) (Table 2).

Beta coefficients and robust standard errors from stage 1 were entered into a random-effects meta-analysis separately for each exposure and cancer outcome. Between-study heterogeneity was estimated by using restricted maximum likelihood (REML). The Hartung–Knapp–Sidik–Jonkman approach was used to adjust test statistics and CIs to more fully account for the estimation uncertainty.¹³ To examine between-cohort heterogeneity, we report I^2 using the following general rule¹⁴: 0%–25%, low heterogeneity; 26%–50%, moderate heterogeneity; and 51%–75%, substantial heterogeneity. Because the value of I^2 depends critically on the size of the studies included,¹⁵ we also looked at Tau^2 (estimated by using REML) and Cochrane's Q . For ease of reading, only I^2 is reported, as Tau^2 and Cochrane's Q were essentially similar to I^2 . Given the number of meta-analyses conducted that increases the likelihood of type I errors, we focused our conclusions on consistent patterns of results, as stated in the PSY-CA protocol.¹⁰

Additional analyses

Several additional analyses were conducted in both stages of the analyses to explore the robustness of our findings.

Stage 1. First, to explore the effect of follow-up duration, analyses were rerun when capping the follow-up duration at four cutoff points: 5, 10, 15, and 20 years. Second, the main analyses were rerun in which carcinoma in situ or borderline tumors were not considered a cancer event and were censored at the time of diagnosis. Third, the minimally adjusted models were rerun with the sample for the corresponding maximally adjusted model to test whether change in effects from the minimally adjusted to the maximally adjusted models could be explained by sample-size reduction because of missing values on covariates. Fourth, given

the correlation between anxiety and depression, when both depression and anxiety showed a statistically significant association with a cancer outcome in the maximally adjusted model, a final model was run that included both depression and anxiety. Finally, analyses were repeated within males and females separately when sex modified the effects of depression or anxiety on the basis of the relative excess risk due to interaction, a measure for additive interaction.

Stage 2. First, to explore the effect of measurement type in depression and anxiety diagnoses, subgroup analyses were conducted that compared the effects observed between cohorts using self-report questionnaires and cohorts using clinical interviews. Second, to examine influential cohorts, we conducted leave-one-out analyses across all models and present the range of pooled effects and I^2 when excluding each cohort in turn. We looked at the effect of influential cohorts in a particular model only when there was at least substantial heterogeneity (>50% for I^2). In these cases, influential studies were identified with the R package Meta¹⁶ on the basis of the criteria outlined by Viechtbauer and Cheung¹⁷ and by considering the change in I^2 as reported in the range.

Changes from protocol

Initially, we had planned to report univariable models.¹⁰ However, the conclusions were very similar to those of the minimally adjusted models and therefore we do not report these results. Furthermore, we had initially planned to run subgroup analyses that looked at differences between cancer stages at diagnosis. This turned out not to be possible because few cohorts had sufficient information. Finally, we applied extra exclusion criteria at stage 2 on the basis of the number of expected cases (see the Stage 2: Meta-analyses section) because models with few cancer incidences resulted in convergence issues.

RESULTS

Descriptive statistics

The percentage of participants with a diagnosis of depression or anxiety in each cohort, along with other descriptives, is displayed in Table 1. Table 2 conveys the number of cancer diagnoses per cohort.

Main results

We found no evidence for an association between depression or anxiety and overall cancer incidence at follow-up (HRs, 0.98–1.05; Figure 1). When looking at specific cancer types, we found that depression and anxiety were associated with an increased risk of lung cancer in the minimally adjusted models (HRs, 1.12–1.60). In

TABLE 1 Prevalence of anxiety and depression along with descriptives of each cohort.

| Cohorts | Total N ^a | Age, mean (SD), years | Female, % | Low education, % | High education, % | Nonnative, % | Depression diagnosis clinical interview, % | Depression diagnosis questionnaire, % | Anxiety diagnosis clinical interview, % | Anxiety diagnosis questionnaire, % |
|--------------------------------|----------------------|-----------------------|-----------|------------------|-------------------|--------------|--|---------------------------------------|---|------------------------------------|
| ALSPAC ^{18,19} | 10,276 | 27.6 (5.0) | 100 | 18 | 17.6 | 2 | — | 23 | — | 22 |
| Atlantic PATH ^{20,21} | 2141 | 50.2 (8.8) | 67 | 3 | 38 | 2 | — | 7 | — | 5 |
| CARTaGENE ^{20,22,23} | 32,093 | 52.9 (7.8) | 53 | 22 | 57 | 15 | — | 5 | — | 4 |
| ELSA ²⁴ | 9248 | 63.4 (10.6) | 55 | 40 | 12 | 6 | — | 24 | — | — |
| HELIUS ^{25,26} | 19,103 | 44.7 (13.2) | 57 | 44 | 26 | 79 | — | 15 | — | — |
| HUNT ²⁷ | 54,697 | 48.3 (16.5) | 53 | 67 | 20 | 1 | — | 3 | — | 5 |
| HUNT3 ²⁷ | 8602 | 40.2 (14.6) | 56 | 28 | 40 | 5 | — | 2 | — | 6 |
| LASA ^{28,29} | 3506 | 67.8 (9.0) | 49 | 66 | 17 | 1 | 3 | 14 | 3 | 3 |
| Lifelines ³⁰ | 141,134 | 44.5 (12.7) | 58 | 29 | 30 | 6 | 3 | 3 | 8 | — |
| NESDA ³¹ | 2296 | 41.7 (13.1) | 66 | 30 | 36 | 7 | 36 | 52 | 42 | 22 |
| OHS ^{20,22} | 14,384 | 49.7 (14.8) | 64 | 1 | 71 | 49 | — | 10 | — | 11 |
| RS1 ³² | 2812 | 75.7 (6.4) | 61 | 59 | 11 | 2 | 4 | 12 | 7 | — |
| RS2 ³² | 1981 | 67.9 (7.3) | 57 | 52 | 17 | 3 | 2 | 9 | 9 | — |
| RS3 ³² | 2968 | 56.7 (6.9) | 56 | 46 | 27 | 3 | 2 | 10 | — | 7 |
| UCC-SMART-2 ³³ | 1810 | 64.8 (10.0) | 25 | 26 | 27 | 10 | — | 7 | — | — |
| UHP1 ³⁴ | 4385 | 38.6 (12.1) | 55 | 30 | 37 | 22 | — | No validated cutoff | — | No validated cutoff |
| UHP2 ³⁴ | 2660 | 39.0 (11.6) | 54 | 17 | 55 | 21 | — | 4 | — | 2 |
| Whitehall II ³⁵ | 5517 | 60.9 (5.9) | 28 | 12 | 35 | 9 | — | 15 | — | — |

Abbreviations: ALSPAC, Avon Longitudinal Study of Parents and Children (mothers cohort); Atlantic PATH, Atlantic Partnership for Tomorrow's Health; ELSA, English Longitudinal Study of Ageing; HELIUS, Healthy Life in an Urban Setting; HUNT, Nord-Trøndelag Health Study; LASA, Longitudinal Aging Study Amsterdam; NESDA, Netherlands Study of Depression and Anxiety; OHS, Ontario Health Study; RS, Rotterdam Study; UCC-SMART-2, Utrecht Cardiovascular Cohort—Second Manifestations of Arterial Disease 2; UHP, Utrecht Health Project.

^aBased on the maximum N in a minimally adjusted model. The number of missing participants per cohort varies per model, and the final number in a particular model may be lower than the frequencies reported. The final N and number of incidences per cohort in the meta-analysis can be seen in the individual forest plots in Supplementary Materials 4.

TABLE 2 Follow-up duration and number of cancer events per type and cohort.

| Cohort (country) | Follow-up duration, years | All cancers ^a | Breast cancer ^a | Lung cancer ^a | Prostate cancer ^a | Colorectal cancer | Smoking-related cancers ^a | Alcohol-related cancers ^a |
|----------------------------|---------------------------|--------------------------|----------------------------|--------------------------|------------------------------|--------------------|--------------------------------------|--------------------------------------|
| ALSPAC (UK) | 20 | 307 | 150 | 8 ^{b,c} | 0 | 8 ^{b,c} | 36 | 160 |
| Atlantic PATH (CAN) | 10 | 67 ^c | 18 ^{b,c} | 5 ^{b,c} | 7 ^{b,c} | 7 ^{b,c} | 24 ^{b,c} | 26 ^{b,c} |
| CARTaGENE (CAN) | 10 | 3875 | 528 | 393 | 429 | 297 | 1336 | 1040 |
| ELSA (UK) | 16 | 2038 | 190 | 173 | 249 | 201 | 705 | 466 |
| HELIUS (NL) | 8 | 421 | 80 | 31 | 51 | 32 | 143 | 125 |
| HUNT2 (NOR) | 24 | 8998 | 1046 | 778 | 1581 | 1292 | 3640 | 2555 |
| HUNT3 (NOR) | 13 | 452 | 79 ^b | 34 ^{b,c} | 78 ^b | 54 ^{b,c} | 153 ^b | 141 ^b |
| LASA (NL) | 26 | 874 | 80 ^{b,c} | 104 ^{b,c} | 100 ^{b,c} | 111 ^{b,c} | 420 | 236 |
| Lifelines (NL) | 13 | 5587 | 1332 | 274 | 503 | 549 | 1569 | 2058 |
| NESDA (NL) | 15 | 223 | 35 | 19 ^b | 10 ^{b,c} | 20 | 85 | 63 |
| Ontario Health Study (CAN) | 10 | 482 | 103 | 33 ^{b,c} | 73 | 28 ^b | 163 | 151 |
| RS1 (NL) | 13 | 507 | 47 ^{b,c} | 87 ^b | 53 ^{b,c} | 83 ^b | 306 | 157 |
| RS2 (NL) | 13 | 293 | 46 ^{b,c} | 37 ^{b,c} | 57 ^b | 38 ^{b,c} | 140 ^b | 96 ^b |
| RS3 (NL) | 9 | 183 | 34 ^{b,c} | 18 ^{b,c} | 20 ^{b,c} | 26 ^{b,c} | 89 ^b | 76 ^b |
| UCC-SMART-2 (NL) | 12 | 207 | 4 ^b | 22 ^b | 25 ^b | 20 ^b | 86 | 39 ^b |
| UHP1 (NL) | 19 | 277 | 61 | 18 | 28 | 29 | 96 | 98 |
| UHP2 (NL) | 16 | 104 ^c | 27 ^{b,c} | 6 ^{b,c} | 10 ^{b,c} | 6 ^{b,c} | 30 ^{b,c} | 36 ^{b,c} |
| Whitehall II (UK) | 13 | 908 | 66 | 30 | 221 | 92 | 245 | 181 |
| Total | — | 25,803 | 3926 | 2070 | 3495 | 2893 | 9266 | 7704 |

Abbreviations: ALSPAC, Avon Longitudinal Study of Parents and Children (mothers cohort); Atlantic PATH, Atlantic Partnership for Tomorrow's Health; CAN, Canada; ELSA, English Longitudinal Study of Ageing; HELIUS, Healthy Life in an Urban Setting; HUNT, Nord-Trøndelag Health Study; LASA, Longitudinal Aging Study Amsterdam; NESDA, Netherlands Study of Depression and Anxiety; NL, the Netherlands; NOR, Norway; OHS, Ontario Health Study; RS, Rotterdam Study; UCC-SMART-2, Utrecht Cardiovascular Cohort–Second Manifestations of Arterial Disease 2; UHP, Utrecht Health Project; UK, United Kingdom.

^aBased on the maximum number of incidences in a minimally adjusted model.

^bFewer than five expected cancer cases in participants with a depression diagnosis.

^cFewer than five expected cancer cases in participants with an anxiety diagnosis.

maximally adjusted models, all effect estimates were attenuated (HRs, 1.07–1.23; Figure 2). Similarly, for smoking-related cancers, depression diagnosis, anxiety diagnosis, and anxiety symptoms were associated with an increased risk of incidence (HRs, 1.06–1.24). Again, these effect estimates were attenuated in the maximally adjusted models (HRs, 1.04–1.08; Figure 3). We found no evidence for an association between depression or anxiety and the incidence of colorectal cancer (HRs, 0.88–1.13; Figure 4), prostate cancer (HRs, 0.97–1.17; Figure 5), or alcohol-related cancers (HRs, 0.97–1.06; Figure 6). For breast cancer, all pooled HRs were consistently negative but mean pooled HRs were close to 1 (HRs, 0.92–0.98; Figure 7) and the upper limit of the 95% CIs all exceeded 1 (with the exception of anxiety symptoms). Across all outcomes, conclusions were similar for diagnoses and symptoms unless indicated otherwise. Forest plots for each meta-analysis are presented in Supplementary Materials 4.

Additional analyses

Analyses with different times of follow-up showed similar HRs for overall, smoking-related, and alcohol-related cancer incidence when capping at 10 and 15 years (Supplementary Materials 5). Other capping points and cancer forms could not be analyzed because of too few cancer incidences and limited follow-up. Analyses excluding carcinomas in situ and borderline tumors did not change the previously observed associations regarding lung cancer and smoking-related cancers (Supplementary Materials 6). The reduction in HRs from the minimally adjusted to maximally adjusted models was not explained by reduced power because conclusions did not change when rerunning minimally adjusted models with the samples used in maximally adjusted models (Supplementary Materials 7). Because neither depression nor anxiety was significantly related to cancer incidence in maximally adjusted models, we did not run meta-analyses with both anxiety and depression in the models. No sex

differences were identified in the associations between depression/anxiety and cancer incidence (Supplementary Materials 8).

The analysis for anxiety and depression diagnosis by measurement type (clinical interview vs. self-report questionnaire) indicated in most cases no differences related to measurement type (Supplementary Materials 9). The exception to this was the association between depression diagnosis and breast cancer, which was associated with increased risk when measured by clinical interview (two studies) compared to decreased risk when measured by self-report questionnaire (seven studies; Figures S9.21 and S9.22). Leave-one-out ranges shown in the figures indicated consistent HRs when excluding each cohort in turn. Further leave-one-out analysis in meta-analyses where $I^2 > 50\%$ did not reveal any influential cohorts.

DISCUSSION

The current study is the first IPD meta-analysis regarding the association between depression and anxiety and cancer risk. Methodological strengths are the use of validated measures of depression or anxiety, the harmonization of data to reach conceptually similar variables, the use of the same statistical procedure across all cohorts, and the control of key confounders in the Cox regression models. In our IPD meta-analyses, on the basis of 18 prospective cohorts, we observed that (1) anxiety and depression were not associated with an increased risk for overall cancers, nor for breast, prostate, colorectal, or alcohol related-cancers; and (2) anxiety and depression were associated with an increased risk for lung cancer and smoking-related cancers after adjustment for sociodemographic factors only but not after additionally adjusting for health-related covariates and other potential confounders. Furthermore, our findings were robust because conclusions did not change when accounting for the reduced power in the maximally adjusted models or the type of measurement for anxiety and depression diagnoses, nor when excluding in situ or borderline tumors, nor when comparing 10- to 15-year follow-up. Finally, we observed no evidence of sex differences in any associations.

The lack of association between anxiety and depression and all cancers is not consistent with several previous meta-analyses.²⁻⁴ These differences are likely to be explained by the higher consistency in the covariates included in our models across all cohorts via data harmonization, the unlikely presence of publication bias in our meta-analysis because inclusion was not restricted to published studies, and the large sample size in our study. Furthermore, cancer diagnosis was based on registry information in all cohorts. Therefore, data on malignancy, site, and time of diagnosis are less influenced by recall bias.³⁶ Also, we used age as the timescale in the Cox regression model whereas previous studies tended to use time on study, which can lead to bias given the nonparametric confounding effect of age in cancer.³⁷

For lung cancer, in minimally adjusted models, depression and anxiety diagnoses were related to increased risk (58% and 60% increased risk, respectively). Furthermore, depression symptoms and anxiety symptoms (for each SD increase) were related to 15% and

12% increased risk, respectively. Similar patterns were seen for smoking-related cancers, although the effect estimates were smaller. Because effects attenuated considerably when further adjusting for health behaviors and other potential confounders (including smoking-related variables), it is possible that tobacco smoking or other unhealthy behaviors play a more crucial role, via mediation or interaction. Indeed, previous studies have attributed the association between depression and smoking-related cancers to increased prevalence of smoking among depressed persons.³⁸⁻⁴⁰ To understand the link between depression, anxiety, and the incidence of lung and smoking-related cancers, further research on the mediating and interactive effects of health behaviors is needed. The residual effect in maximum models may be explained by imperfect adjustment for unhealthy behaviors. Another possibility is a common Ras proto-oncogene that increases the risk for both depression and cancers related to the Ras oncogene family such as lung cancer.⁴¹

A few limitations exist in the current study. We did not use multiple imputation to deal with missing data⁴² because it was not feasible to do this in a standardized way in all 18 cohorts. Furthermore, we did not consider specific subtypes of cancer (e.g., non-small cell lung cancer vs. other types of lung cancer) that may be differentially related to depression and anxiety.⁴¹ Other cohorts may have met the inclusion and exclusion criteria but were neither known to the consortium nor came up in a literature search. PSY-CA is set up in such a way that information from additional cohorts can be added in the future. To reduce the likelihood of type I errors, we focused our conclusions on consistent findings, but it is possible that small, sporadic effects relate to true effects that require even more power to detect. Finally, we looked at current depression or anxiety at baseline, whereas individuals with a history of depression and anxiety, or more chronic forms of depression and anxiety, may be at a higher risk of cancer.⁴³

In conclusion, our study does not support the hypothesis that depression and anxiety are related to an increased risk for overall cancers nor breast, prostate, colorectal, or alcohol-related cancers. Although we found support for an association with smoking-related cancers such as lung cancer, when we adjusted for additional covariates the effect was substantially reduced. The results from the current study may help health professionals to alleviate feelings of guilt and self-blame in patients with cancer who attribute their diagnosis to previous depression or anxiety. However, further research is required to test whether depression and anxiety interact with or moderate the effects of health behaviors on the incidence of lung and smoking-related cancers.

AUTHOR CONTRIBUTIONS

Lonneke A. van Tuijl: Conceptualization, data curation, formal analysis, methodology, project administration, visualization, writing—original draft, and writing—review and editing. **Maartje Basten:** Conceptualization, data curation, formal analysis, methodology, project administration, writing—original draft, validation, and writing—review and editing. **Kuan-Yu Pan:** Conceptualization, data curation, formal analysis, methodology, project administration, writing—original




FIGURE 1 Association between depression/anxiety and all cancer types. Error bars convey the 95% confidence interval around the pooled HR. HR indicates hazard ratio; max., maximally adjusted (additionally adjusting for health-related covariates and other potential risk factors); min., minimally adjusted (demographic variables).




FIGURE 2 Association between depression/anxiety and lung cancer. Error bars convey the 95% confidence interval around the pooled HR. HR indicates hazard ratio; max., maximally adjusted (additionally adjusting for health-related covariates and other potential risk factors); min., minimally adjusted (demographic variables).




FIGURE 3 Association between depression/anxiety and smoking-related cancers. Error bars convey the 95% confidence interval around the pooled HR. HR indicates hazard ratio; max., maximally adjusted (additionally adjusting for health-related covariates and other potential risk factors); min., minimally adjusted (demographic variables).




FIGURE 4 Association between depression/anxiety and colorectal cancer. Error bars convey the 95% confidence interval around the pooled HR. HR indicates hazard ratio; max., maximally adjusted (additionally adjusting for health-related covariates and other potential risk factors); min., minimally adjusted (demographic variables).




FIGURE 5 Association between depression/anxiety and prostate cancer. Error bars convey the 95% confidence interval around the pooled HR. HR indicates hazard ratio; max., maximally adjusted (additionally adjusting for health-related covariates and other potential risk factors); min., minimally adjusted (demographic variables).




FIGURE 6 Association between depression/anxiety and alcohol-related cancers. Error bars convey the 95% confidence interval around the pooled HR. HR indicates hazard ratio; max., maximally adjusted (additionally adjusting for health-related covariates and other potential risk factors); min., minimally adjusted (demographic variables).




FIGURE 7 Association between depression/anxiety and breast cancer. Error bars convey the 95% confidence interval around the pooled HR. HR indicates hazard ratio; max., maximally adjusted (additionally adjusting for health-related covariates and other potential risk factors); min., minimally adjusted (demographic variables).

draft, and writing–review and editing. **Roel Vermeulen:** Conceptualization, formal analysis, methodology, project administration, funding acquisition, formal analysis, supervision, and writing–review and editing. **Lützen Portengen:** Formal analysis, methodology, and writing–review and editing. **Alexander de Graeff:** Conceptualization, funding acquisition, methodology, project administration, supervision, and writing–review and editing. **Joost Dekker:** Conceptualization, data curation, funding acquisition, methodology, project administration, supervision, writing–original draft, and writing–review and editing. **Mirjam I. Geerlings:** Conceptualization, data curation, formal analysis, funding acquisition, methodology, project administration, supervision, writing–original draft, and writing–review and editing. **Adriaan Hoogendoorn:** Conceptualization, data curation, formal analysis, funding acquisition, methodology, project administration, supervision, and writing–review and editing. **Femke Lamers:** Conceptualization, data curation, formal analysis, funding acquisition, methodology, project administration, supervision, writing–original draft, and writing–review and editing. **Adri C. Voogd:** Conceptualization, funding acquisition, methodology, project administration, supervision, and writing–review and editing. **Jessica Abell:** Formal analysis and writing–review and editing. **Philip Awadalla:** Formal analysis and writing–review and editing. **Aartjan T. F. Beekman:** Conceptualization, methodology, and writing–review and editing. **Ottar Bjerkeset:** Formal analysis and writing–review and editing. **Andy Boyd:** Formal analysis and writing–review and editing. **Yunsong Cui:** Formal analysis and writing–review and editing. **Philipp Frank:** Formal analysis and writing–review and editing. **Henrike Galenkamp:** Formal analysis and writing–review and editing. **Bert Garssen:** Conceptualization, methodology, and writing–review and editing. **Sean Hellingman:** Formal analysis and writing–review and editing. **Martijn Huisman:** Conceptualization, formal analysis, methodology, and writing–review and editing. **Anke Huss:** Writing–review and editing. **Trynke R. de Jong:** Writing–review and editing. **Melanie R. Keats:** Formal analysis and writing–review and editing. **Almar A. L. Kok:** Conceptualization, formal analysis, methodology, and writing–review and editing. **Steinar Krokstad:** Formal analysis and writing–

review and editing. **Flora E. van Leeuwen:** Conceptualization, formal analysis, methodology, and writing–review and editing. **Annemarie I. Luik:** Conceptualization, formal analysis, methodology, and writing–review and editing. **Nolwenn Noisel:** Formal analysis and writing–review and editing. **N. Charlotte Onland-Moret:** Formal analysis and writing–review and editing. **Yves Payette:** Formal analysis and writing–review and editing. **Brenda W. J. H. Penninx:** Conceptualization, formal analysis, methodology, and writing–review and editing. **Ina Rissanen:** Formal analysis and writing–review and editing. **Annelieke M. Roest:** Conceptualization, funding acquisition, methodology, and writing–review and editing. **Rikje Ruiter:** Writing–review and editing. **Robert A. Schoevers:** Conceptualization, methodology, and writing–review and editing. **David Soave:** Formal analysis and writing–review and editing. **Mandy Spaan:** Conceptualization, formal analysis, methodology, and writing–review and editing. **Andrew Steptoe:** Formal analysis and writing–review and editing. **Karien Stronks:** Conceptualization, formal analysis, methodology, and writing–review and editing. **Erik R. Sund:** Formal analysis and writing–review and editing. **Ellen Sweeney:** Formal analysis and writing–review and editing. **Emma L. Twait:** Writing–review and editing. **Alison Teyhan:** Formal analysis and writing–review and editing. **W. M. Monique Verschuren:** Formal analysis and writing–review and editing. **Kimberly D. van der Willik:** Formal analysis and writing–review and editing. **Judith G. M. Rosmalen:** Conceptualization, formal analysis, methodology, supervision, writing–original draft, and writing–review and editing. **Adelita V. Ranchor:** Conceptualization, data curation, formal analysis, methodology, project administration, supervision, writing–original draft, and writing–review and editing.

AFFILIATIONS

¹Health Psychology Section, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

²Department of Clinical Psychology, Utrecht University, Utrecht, the Netherlands

³Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht and Utrecht University, Utrecht, the Netherlands

- ⁴Department of Psychiatry, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands
- ⁵Mental Health Program, Amsterdam Public Health, Amsterdam, the Netherlands
- ⁶Unit of Occupational Medicine, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden
- ⁷Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands
- ⁸Department of Medical Oncology, University Medical Center Utrecht, Utrecht, the Netherlands
- ⁹Department of General Practice, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands
- ¹⁰Aging & Later Life and Personalized Medicine, Amsterdam Public Health, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands
- ¹¹Neurodegeneration and Mood, Anxiety, Psychosis, Stress, and Sleep, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands
- ¹²GGZ inGeest Specialized Mental Health Care, Amsterdam, the Netherlands
- ¹³Department of Epidemiology, Maastricht University, Maastricht, the Netherlands
- ¹⁴Department of Research and Development, Netherlands Comprehensive Cancer Organization, Utrecht, the Netherlands
- ¹⁵Department of Behavioural Science and Health, University College London, London, UK
- ¹⁶Ontario Institute for Cancer Research, Toronto, Ontario, Canada
- ¹⁷Department of Molecular Genetics, University of Toronto, Toronto, Ontario, Canada
- ¹⁸Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada
- ¹⁹Faculty of Nursing and Health Sciences, Nord University, Levanger, Norway
- ²⁰Department of Mental Health, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway
- ²¹Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK
- ²²Atlantic Partnership for Tomorrow's Health, Faculty of Medicine, Dalhousie University, Halifax, Nova Scotia, Canada
- ²³Department of Public and Occupational Health, Amsterdam UMC, and Amsterdam Public Health Research Institute, University of Amsterdam, Amsterdam, the Netherlands
- ²⁴Department of Mathematics, Wilfrid Laurier University, Waterloo, Ontario, Canada
- ²⁵Department of Epidemiology and Data Science, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands
- ²⁶Department of Sociology, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands
- ²⁷Lifelines Cohort and Biobank, Roden, the Netherlands
- ²⁸School of Health and Human Performance, Faculty of Health, Dalhousie University, Halifax, Nova Scotia, Canada
- ²⁹Department of Public Health and Nursing, Trøndelag Health Study Research Centre, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway
- ³⁰Levanger Hospital, Nord-Trøndelag Hospital Trust, Levanger, Norway
- ³¹Division of Psychosocial Research and Epidemiology, The Netherlands Cancer Institute, Amsterdam, the Netherlands
- ³²Department of Epidemiology, Erasmus MC–University Medical Center, Rotterdam, the Netherlands
- ³³CARTaGENE, CHU Sainte-Justine, Montreal, Quebec, Canada

³⁴Department of Developmental Psychology, University of Groningen, Groningen, the Netherlands

³⁵Department of Internal Medicine, Maastrad, Rotterdam, the Netherlands

³⁶Department of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

³⁷Centre for Urban Mental Health, University of Amsterdam, Amsterdam, the Netherlands

³⁸Centre for Nutrition, Prevention and Health Services, National Institute for Public Health and the Environment, Utrecht Bilthoven, the Netherlands

³⁹Department of Internal Medicine, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

ACKNOWLEDGMENTS

We thank the Dutch Cancer Society (grant VU2017-8288). The UK Medical Research Council and Wellcome (reference 217065/Z/19/Z) and the University of Bristol provide core support for the Avon Longitudinal Study of Parents and Children (ALSPAC). The linkage of ALSPAC to the cancer registry was funded by Wellcome (reference 086118). The infrastructure of the Netherlands Study of Depression and Anxiety study (<http://www.nesda.nl>) is funded via the Geestkracht program of the Netherlands Organisation for Health Research and Development (grant 10-000-1002) and financial contributions by the participating universities and mental health care organizations (VU University Medical Center, GGZ inGeest, Leiden University Medical Center, Leiden University, GGZ Rivierduinen, University Medical Center Groningen, University of Groningen, Lentis, GGZ Friesland, GGZ Drenthe, Dimence, and Rob Giel Onderzoekscentrum). The English Longitudinal Study of Ageing is funded by the National Institute on Aging (R01AG017644) and the National Institute for Health and Care Research (198/1074-02). The funding organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

The Trøndelag Health Study (HUNT) is a collaboration between the HUNT Research Centre (Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology), Trøndelag County Council, Central Norway Regional Health Authority, and Norwegian Institute of Public Health.

This research has been conducted using Atlantic Partnership for Tomorrow's Health (Atlantic PATH) data under application 2019-103. The data used in this research were made available by the Atlantic PATH study, which is the Atlantic Canada regional component of the Canadian Partnership for Tomorrow's Health funded by the Canadian Partnership Against Cancer and Health Canada. The views expressed herein represent the views of the authors and do not necessarily represent the views of Health Canada.

The Healthy Life in an Urban Setting (HELIUS) study is conducted by the Amsterdam University medical centers, location AMC, and the Public Health Service of Amsterdam. Both organizations provide core support for HELIUS. The HELIUS study is also funded by the Dutch Heart Foundation, the Netherlands Organisation for Health Research and Development (ZonMw), the European Union (FP-7), and the European Fund for the Integration of non-EU immigrants (EIF).

The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam, Netherlands Organisation for Health Research and Development (ZonMw), the Research Institute for Diseases in the Elderly, the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam. We are grateful to the study participants, the staff from the Rotterdam Study, and the participating general practitioners and pharmacists.

We thank the whole CARTaGENE team (<https://cartagene.qc.ca/en/about>), represented by the authors N.N. and Y.P., for their contribution.

Lifelines is a multidisciplinary prospective population-based cohort study examining in a unique three-generation design the health and health-related behaviors of 167,729 persons living in the north of the Netherlands. It employs a broad range of investigative procedures in assessing the biomedical, sociodemographic, behavioral, physical, and psychological factors that contribute to the health and disease of the general population, with a special focus on multimorbidity and complex genetics. The Lifelines initiative has been made possible by a subsidy from the Dutch Ministry of Health, Welfare and Sport, the Dutch Ministry of Economic Affairs, University Medical Center Groningen, Groningen University, and the provinces in the north of the Netherlands (Drenthe, Friesland, Groningen).

We are extremely grateful to all the families who took part in the ALSPAC study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses.

We gratefully acknowledge the contribution of the research nurses: R. van Petersen (data manager); A. Vandersteen (study manager); and the members of the Utrecht Cardiovascular Cohort–Second Manifestations of Arterial Disease study group (UCC-SMART): F. W. Asselbergs and H. M. Nathoe, Department of Cardiology; G. J. de Borst, Department of Vascular Surgery; M. L. Bots and M. I. Geerlings, Julius Center for Health Sciences and Primary Care; M. H. Emmelot-Vonk, Department of Geriatrics; P. A. de Jong and T. Leiner, Department of Radiology; A. T. Lely, Department of Gynecology and Obstetrics; N. P. van der Kaaij, Department of Cardiothoracic Surgery; L. J. Kappelle and Y. M. Ruigrok, Department of Neurology and Hypertension; M. C. Verhaar, Department of Nephrology and Hypertension; and F. L. J. Visseren (chair), Department of Vascular Medicine, University Medical Center Utrecht and Utrecht University.

This study has used data from the Cancer Registry of Norway. The interpretation and reporting of these data are the sole responsibility of the authors, and no endorsement by the Cancer Registry of Norway is intended nor should it be inferred.

CONFLICT OF INTEREST STATEMENT

Lonneke A. van Tuijl has received grants and travel support from the Dutch Cancer Society (KWF). Maartje Basten has received grants from the KWF. Kuan-Yu Pan has received grants from the Dutch

Cancer Foundation. Joost Dekker has received grants from the KWF. Femke Lamers has received grants from the KWF. Ottar Bjerkeset has been a consultant for Nord University. Andy Boyd has received grants from the Medical Research Council. Philipp Frank has received grants from the Economic and Social Research Council. Trynke R. de Jong has been a consultant for Medische Biobank Noord–Nederland. Ina Rissanen has received grants from Orionin Tutkimussäätiö. Annelieke M. Roest has received grants and travel support from the KWF. Mandy Spaan has received grants from the Netherlands Cancer Institute. Kimberly D. van der Willik has been a consultant for Erasmus Medisch Centrum. Adelita V. Ranchor has received grants from the KWF. The other authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are owned by the participating cohort studies. Data are not publicly available but may be shared upon reasonable request at each cohort depending on cohort-specific regulations.

ORCID

Lonneke A. van Tuijl  <https://orcid.org/0000-0001-9031-0886>

Joost Dekker  <https://orcid.org/0000-0003-2027-0027>

Kimberly D. van der Willik  <https://orcid.org/0000-0002-4479-2914>

REFERENCES

1. Antoni MH, Lutgendorf SK, Cole SW, et al. The influence of bio-behavioural factors on tumour biology: pathways and mechanisms. *Nat Rev Cancer*. 2006;6(3):240-248. doi:10.1038/nrc1820
2. Chida Y, Hamer M, Wardle J, Steptoe A. Do stress-related psychosocial factors contribute to cancer incidence and survival? *Nat Clin Pract Oncol*. 2008;5(8):466-475. doi:10.1038/npcnc1134
3. Jia Y, Li F, Liu YF, Zhao JP, Leng MM, Chen L. Depression and cancer risk: a systematic review and meta-analysis. *Public Health*. 2017; 149:138-148. doi:10.1016/j.puhe.2017.04.026
4. Wang YH, Li JQ, Shi JF, et al. Depression and anxiety in relation to cancer incidence and mortality: a systematic review and meta-analysis of cohort studies. *Mol Psychiatr*. 2020;25(7):1487-1499. doi:10.1038/s41380-019-0595-x
5. McGee R, Williams S, Elwood M. Depression and the development of cancer: a meta-analysis. *Soc Sci Med*. 1994;38(1):187-192. doi:10.1016/0277-9536(94)90314-x
6. Oerlemans ME, van den Akker M, Schuurman AG, Kellen E, Buntinx F. A meta-analysis on depression and subsequent cancer risk. *Clin Pract Epidemiol Ment Health*. 2007;3(1):29. doi:10.1186/1745-0179-3-29
7. Ahn HK, Bae JH, Ahn HY, Hwang IC. Risk of cancer among patients with depressive disorder: a meta-analysis and implications. *Psychooncology*. 2016;25(12):1393-1399. doi:10.1002/pon.4084
8. Stewart LA, Tierney JF. To IPD or not to IPD?: Advantages and disadvantages of systematic reviews using individual patient data. *Eval Health Prof*. 2002;25(1):76-97. doi:10.1177/0163278702025001006
9. Tierney JF, Vale C, Riley R, et al. Individual participant data (IPD) meta-analyses of randomised controlled trials: guidance on their use. *PLoS Med*. 2015;12(7):e1001855. doi:10.1371/journal.pmed.1001855
10. van Tuijl LA, Voogd AC, de Graeff A, et al. Psychosocial Factors and Cancer Incidence (PSY-CA): protocol for individual participant data meta-analyses. *Brain Behav*. 2021;11(10):e2340. doi:10.1002/brb3.2340

11. Fortier I, Raina P, Van den Heuvel ER, et al. Maelstrom Research guidelines for rigorous retrospective data harmonization. *Int J Epidemiol.* 2017;46(1):103-105. doi:10.1093/ije/dyw075
12. IARC Monographs on the Identification of Carcinogenic Hazards to Humans and Handbooks of Cancer Prevention. International Agency for Research on Cancer. Published 2019. Accessed October 14, 2019. <https://monographs.iarc.who.int/wp-content/uploads/2019/12/OrganSitePoster.PlusHandbooks.pdf>
13. IntHout J, Ioannidis JP, Borm GF. The Hartung-Knapp-Sidik-Jonkman method for random effects meta-analysis is straightforward and considerably outperforms the standard DerSimonian-Laird method. *BMC Med Res Methodol.* 2014;14(1):25. doi:10.1186/1471-2288-14-25
14. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327(7414):557-560. doi:10.1136/bmj.327.7414.557
15. Rücker G, Schwarzer G, Carpenter JR, Schumacher M. Undue reliance on I^2 in assessing heterogeneity may mislead. *BMC Med Res Methodol.* 2008;8(1):79. doi:10.1186/1471-2288-8-79
16. Schwarzer G, Carpenter JR, Rücker G. *Meta-Analysis with R*. Springer International Publishing; 2015. doi:10.1007/978-3-319-21416-0
17. Viechtbauer W, Cheung MWL. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods.* 2010;1(2):112-125. doi:10.1002/jrsm.11
18. Boyd A, Golding J, Macleod J, et al. Cohort profile: the 'children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol.* 2013;42(1):111-127. doi:10.1093/ije/dys064
19. Fraser A, Macdonald-Wallis C, Tilling K, et al. Cohort profile: The Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol.* 2013;42(1):97-110. doi:10.1093/ije/dys066
20. Borugian MJ, Robson P, Fortier I, et al. The Canadian Partnership for Tomorrow Project: building a pan-Canadian research platform for disease prevention. *CMAJ.* 2010;182(11):1197-1201. doi:10.1503/cmaj.091540
21. Sweeney E, Cui Y, DeClercq V, et al. Cohort profile: The Atlantic Partnership for Tomorrow's Health (Atlantic PATH) study. *Int J Epidemiol.* 2017;46(6):1762-1763. doi:10.1093/ije/dyx124
22. Dummer TJB, Awadalla P, Boileau C, et al. The Canadian Partnership for Tomorrow project: a pan-Canadian platform for research on chronic disease prevention. *CMAJ.* 2018;190(23):E710-E717. doi:10.1503/cmaj.170292
23. Awadalla P, Boileau C, Payette Y, et al. Cohort profile of the CAR-TaGENE study: Quebec's population-based biobank for public health and personalized genomics. *Int J Epidemiol.* 2013;42(5):1285-1299. doi:10.1093/ije/dys160
24. Steptoe A, Breeze E, Banks J, Nazroo J. Cohort profile: The English Longitudinal Study of Ageing. *Int J Epidemiol.* 2013;42(6):1640-1648. doi:10.1093/ije/dys168
25. Snijder MB, Galenkamp H, Prins M, et al. Cohort profile: the Healthy Life in an Urban Setting (HELIUS) study in Amsterdam, the Netherlands. *BMJ Open.* 2017;7(12):e017873. doi:10.1136/bmjopen-2017-017873
26. Stronks K, Snijder MB, Peters RJ, Prins M, Schene AH, Zwinderman AH. Unravelling the impact of ethnicity on health in Europe: the HELIUS study. *BMC Public Health.* 2013;13(1):402. doi:10.1186/1471-2458-13-402
27. Krokstad S, Langhammer A, Hveem K, et al. Cohort profile: The HUNT study, Norway. *Int J Epidemiol.* 2013;42(4):968-977. doi:10.1093/ije/dys095
28. Hoogendijk EO, Deeg DJH, Poppelaars J, et al. The Longitudinal Aging Study Amsterdam: cohort update 2016 and major findings. *Eur J Epidemiol.* 2016;31(9):927-945. doi:10.1007/s10654-016-0192-0
29. Huisman M, Poppelaars J, van der Horst M, et al. Cohort profile: The Longitudinal Aging Study Amsterdam. *Int J Epidemiol.* 2011;40(4):868-876. doi:10.1093/ije/dyq219
30. Scholtens S, Smidt N, Swertz MA, et al. Cohort profile: LifeLines, a three-generation cohort study and biobank. *Int J Epidemiol.* 2015;44(4):1172-1180. doi:10.1093/ije/dyu229
31. Penninx BWJH, Beekman ATF, Johannes HS, et al. The Netherlands Study of Depression and Anxiety (NESDA): rationale, objectives and methods. *Int J Methods Psychiatr Res.* 2008;17(3):121-140. doi:10.1002/mpr.256
32. Hofman A, Brusselle GGO, Murad SD, et al. The Rotterdam Study: 2016 objectives and design update. *Eur J Epidemiol.* 2015;30(8):661-708. doi:10.1007/s10654-015-0082-x
33. Simons PCG, Algra A, van de Laak MF, Grobbee DE, van der Graaf Y. Second Manifestations of Arterial Disease (SMART) study: rationale and design. *Eur J Epidemiol.* 1999;15(9):773-781. doi:10.1023/a:1007621514757
34. Grobbee DE, Hoes AW, Verheij TJM, Schrijvers AJP, van Ameijden EJC, Numans ME. The Utrecht Health Project: optimization of routine healthcare data for research. *Eur J Epidemiol.* 2005;20(3):285-287. doi:10.1007/s10654-004-5689-2
35. Marmot M, Brunner E. Cohort profile: The Whitehall II study. *Int J Epidemiol.* 2005;34(2):251-256. doi:10.1093/ije/dyh372
36. Cho S, Shin A, Song D, et al. Validity of self-reported cancer history in the Health Examinees (HEXA) study: a comparison of self-report and cancer registry records. *Cancer Epidemiol.* 2017;50:16-21. doi:10.1016/j.canep.2017.07.010
37. Thiébaud ACM, Bénichou J. Choice of time-scale in Cox's model analysis of epidemiologic cohort data: a simulation study. *Stat Med.* 2004;23(24):3803-3820. doi:10.1002/sim.2098
38. Dalton SO, Møller-Jensen L, Olsen JH, Mortensen PB, Johansen C. Depression and cancer risk: a register-based study of patients hospitalized with affective disorders, Denmark, 1969-1993. *Am J Epidemiol.* 2002;155(12):1088-1095. doi:10.1093/aje/k155.12.1088
39. Knekt P, Raitasalo R, Heliövaara M, et al. Elevated lung cancer risk among persons with depressed mood. *Am J Epidemiol.* 1996;144(12):1096-1103. doi:10.1093/oxfordjournals.aje.a008887
40. Linkins RW, Comstock GW. Depressed mood and development of cancer. *Am J Epidemiol.* 1990;132(5):962-972. doi:10.1093/oxfordjournals.aje.a115739
41. Brewer JK. Behavioral genetics of the depression/cancer correlation: a look at the Ras oncogene family and the 'cerebral diabetes paradigm.' *J Mol Neurosci.* 2008;35(3):307-322. doi:10.1007/s12031-008-9078-2
42. Knol MJ, Janssen KJM, Donders ART, et al. Unpredictable bias when using the missing indicator method or complete case analysis for missing confounder values: an empirical example. *J Clin Epidemiol.* 2010;63(7):728-736. doi:10.1016/j.jclinepi.2009.08.028
43. Penninx BWJH, Guralnik JM, Pahor M, et al. Chronically depressed mood and cancer risk in older persons. *J Natl Cancer Inst.* 1998;90(24):1888-1893. doi:10.1093/jnci/90.24.1888

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: van Tuijl LA, Basten M, Pan K-Y, et al. Depression, anxiety, and the risk of cancer: An individual participant data meta-analysis. *Cancer.* 2023;1-13. doi:10.1002/cncr.34853