


















Article

Independent and Joined Association between Socioeconomic Indicators and Pediatric Obesity in Spain: The PASOS Study

Clara Homs ^{1,2}, Paula Berruezo ¹, Albert Arcarons ^{3,4} , Julia Wärnberg ^{5,6} , Maddi Osés ⁷,
Marcela González-Gross ^{5,8} , Narcis Gusi ⁹ , Susana Aznar ^{10,11} , Elena Marín-Cascales ^{12,13,14},
Miguel Ángel González-Valeiro ¹⁵ , Lluís Serra-Majem ^{5,16} , Nicolás Terrados ¹⁷ , Josep A. Tur ^{5,18} ,
Marta Segú ¹⁹, Montserrat Fitó ^{5,20}, Juan Carlos Benavente-Marín ^{5,6}, Idoia Labayen ^{5,7}, Augusto G. Zapico ^{8,21} ,
Jesús Sánchez-Gómez ⁹, Fabio Jiménez-Zazo ¹⁰ , Pedro E. Alcaraz ^{12,13,14} , Marta Sevilla-Sánchez ¹⁵ ,
Estefanía Herrera-Ramos ²², Susana Pulgar-Muñoz ²³, Cristina Bouzas ^{5,18} , Raimon Milà ² ,
Helmut Schröder ^{20,24,*}  and Santiago F. Gómez ^{1,20,24,25,*} 

- ¹ Gasol Foundation Europe, 08830 Sant Boi de Llobregat, Spain
- ² Global Research on Wellbeing (GroW), Faculty of Health Sciences, Blanquerna Ramon Llull University, 08025 Barcelona, Spain
- ³ Office of the High Commissioner against Child Poverty, 28079 Madrid, Spain
- ⁴ Department of Sociology, National Distance Education University (UNED), 28012 Madrid, Spain
- ⁵ Physiopathology of Obesity and Nutrition Networking Biomedical Research Center (CIBEROBN), Institute of Health Carlos III, 28029 Madrid, Spain
- ⁶ EpiPHAAN Research Group, School of Health Sciences, Instituto de Investigación Biomédica en Málaga (IBIMA), University of Málaga, 29590 Málaga, Spain
- ⁷ IS-FOOD—Institute for Sustainability & Food Chain Innovation, Universidad Pública de Navarra (UPNA), IDISNA, 31006 Pamplona, Spain
- ⁸ ImFINE Research Group, Department of Health and Human Performance, Universidad Politécnica de Madrid, 28003 Madrid, Spain
- ⁹ Physical Activity and Quality of Life Research Group (AFYCAV), Faculty of Sport Sciences, University of Extremadura, 10003 Cáceres, Spain
- ¹⁰ PAFS Research Group, Faculty of Sports Sciences, University of Castilla-La Mancha-Toledo Campus, 45071 Toledo, Spain
- ¹¹ Biomedical Research Networking Center on Frailty and Healthy Aging (CIBERFES), 28029 Madrid, Spain
- ¹² UCAM Research Center for High Performance Sport, UCAM Universidad Católica de Murcia, 30107 Murcia, Spain
- ¹³ Facultad de Deporte, UCAM Universidad Católica de Murcia, 30107 Murcia, Spain
- ¹⁴ Strength & Conditioning Society, 30008 Murcia, Spain
- ¹⁵ Faculty of Sports Sciences and Physical Education, Universidad de la Coruña, 15001 A Coruña, Spain
- ¹⁶ Preventive Medicine Service, Canarian Health Service, Centro Hospitalario Universitario Insular Materno Infantil (CHUIMI), 35016 Las Palmas de Gran Canaria, Spain
- ¹⁷ Regional Unit of Sports Medicine-Municipal Sports Foundation of Avilés, 33402 Avilés, Spain
- ¹⁸ Research Group of Community Nutrition & Oxidative Stress, University of the Balearic Islands-IUNICS & Health Research Institute of the Balearic Islands (IDISBA), 07122 Palma de Mallorca, Spain
- ¹⁹ Barça Foundation, 08028 Barcelona, Spain
- ²⁰ Cardiovascular Risk and Nutrition Research Group (CARIN), Hospital del Mar Medical Research Institute (IMIM), 08003 Barcelona, Spain
- ²¹ Department of Language, Arts and Physical Education, Universidad Complutense de Madrid, 28040 Madrid, Spain
- ²² Research Institute of Biomedical and Health Sciences (IUIBS), University of Las Palmas de Gran Canaria, 35016 Las Palmas de Gran Canaria, Spain
- ²³ Health Research Institute of the Principality of Asturias (ISPA), 33011 Oviedo, Spain
- ²⁴ CIBER Epidemiology and Public Health (CIBERESP), Carlos III Health Institute, 28029 Madrid, Spain
- ²⁵ Nursing and Physiotherapy Department, University of Lleida, 25198 Lleida, Spain
- * Correspondence: hschroder@imim.es (H.S.); sgomez@gasolfoundation.org (S.F.G.)
- † These authors contributed equally to this work.



Citation: Homs, C.; Berruezo, P.; Arcarons, A.; Wärnberg, J.; Osés, M.; González-Gross, M.; Gusi, N.; Aznar, S.; Marín-Cascales, E.; González-Valeiro, M.Á.; et al. Independent and Joined Association between Socioeconomic Indicators and Pediatric Obesity in Spain: The PASOS Study. *Nutrients* **2023**, *15*, 1987. <https://doi.org/10.3390/nu15081987>

Academic Editor: Cherubini Valentino

Received: 7 March 2023

Revised: 14 April 2023

Accepted: 18 April 2023

Published: 20 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Childhood obesity is a public health problem worldwide. An important determinant of child and adolescent obesity is socioeconomic status (SES). However, the magnitude of the impact of different SES indicators on pediatric obesity on the Spanish population scale is unclear. The aim of this study was to assess the association between three SES indicators and obesity in a nationwide,

representative sample of Spanish children and adolescents. A total of 2791 boys and girls aged 8 to 16 years old were included. Their weight, height, and waist circumference were measured. SES was assessed using two parent/legal guardian self-reported indicators (educational level -University/non-University- and labor market status -Employed/Unemployed-). As a third SES indicator, the annual mean income per person was obtained from the census section where the participating schools were located ($\geq 12.731\text{€}$ / $< 12.731\text{€}$). The prevalence of obesity, severe obesity, and abdominal obesity was 11.5%, 1.4%, and 22.3%, respectively. Logistic regression models showed an inverse association of both education and labor market status with obesity, severe obesity, and abdominal obesity (all $p < 0.001$). Income was also inversely associated with obesity ($p < 0.01$) and abdominal obesity ($p < 0.001$). Finally, the highest composite SES category (University/Employed/ $\geq 12.731\text{€}$ $n = 517$) showed a robust and inverse association with obesity (OR = 0.28; 95% CI: 0.16–0.48), severe obesity (OR = 0.20; 95% CI: 0.05–0.81), and abdominal obesity (OR = 0.36; 95% CI: 0.23–0.54) in comparison with the lowest composite SES category (Less than University/Unemployed/ $< 12.731\text{€}$; $n = 164$). No significant interaction between composite SES categories and age and gender was found. SES is strongly associated with pediatric obesity in Spain.

Keywords: obesity; child; adolescent; socioeconomic status; cross-sectional study

1. Introduction

Pediatric obesity is a public health problem worldwide [1]. In Spain, the prevalence of excess body weight in children aged 7 to 13 increased from 32.3% (95% CI: 29.1–35.6%) during 1999–2010 to 35.3% (95% CI, 32.9–37.7%) during 2011–2021 [2]. The obesity epidemic had short-, mid-, and long-term consequences on the physical, psychological, and social spheres of health [3].

Obesity is the result of complex interactions of multiple determinants, such as nutrition [4], physical activity and sedentary behaviors [5], sleep [6], psychological factors [7], and genetic predisposition [8]. Evidence from high-income countries (HICs) showed an inverted association between socioeconomic status (SES) and pediatric obesity [9–12]. Nonetheless, a recent review describes the complexity of studying the association between SES and childhood obesity and concludes that more research is needed [11].

A European study showed that in 7 out of the 13 HICs, the prevalence of overweightness was double among children whose parents had a low educational level than among those whose parents had a high educational level [13]. Another study showed that, in the United Kingdom, adolescents from the poorest backgrounds had a two-fold risk of obesity than those from the richest ones [14].

In Spain, the impact of different SES indicators on pediatric obesity in a nationwide, representative sample remains unclear [15] since the evidence is only available for the 6- to 9-year-old population [15]. Three studies that used self-reported anthropometric data from the Spanish National Health Survey (Encuesta Nacional de Salud de España, ENS) that collected information on children aged 0–15 years found that (i) obesity and overweightness showed an inverse association with parental occupation [16]; (ii) there is a growing tendency of obesity and overweightness among adolescents from families with a low educational level [17]; and (iii) there is a higher tendency of obesity and overweightness among adolescents whose parents have a lower economic situation [18]. However, self-reported data tend to overestimate height while underestimating weight [16–18]. Moreover, contrasting results were obtained in studies in different regions of Spain. In particular, a longitudinal study from 2006 to 2016 in Catalonia [19] indicated a socioeconomic gradient for childhood obesity: The prevalence was higher both in the most deprived areas and among children with non-Spanish nationalities, especially Africans and Asians. Conversely, a cross-sectional study carried out in 2018 with a representative sample in Andalusia [20] reported that the youngest population had no significant social gradient for overweightness and obesity prevalence.

The World Health Organization (WHO) highlighted the urgency of reducing socioeconomic inequalities to prevent childhood obesity [21,22]. In Spain, structural measures to tackle childhood obesity that strongly consider health inequalities are being implemented in the framework of the National Plan for the Reduction of Childhood Obesity for 2030 [23] presented in 2022. Other initiatives especially benefitting the young population from low SES backgrounds include applying taxes on sugar-sweetened beverages in Catalonia [24] and regulating the advertising of unhealthy food and beverages to children under the age of 16 [25,26]. Finally, it is crucial to understand the association between SES and weight status among Spanish children and adolescents and to monitor the trend of this association over a prolonged time to apply the appropriate measures to fight childhood obesity.

The present study aimed to assess the association between three SES indicators (educational level, labor market status, and income) and weight status (obesity, severe obesity, and abdominal obesity) in a nationwide, representative sample of Spanish children and adolescents. We also assessed the joined association between combinations of the three SES indicators and weight status.

2. Materials and Methods

2.1. Study Design

This study is a cross-sectional analysis within the framework of the PASOS study (Physical Activity, Sedentarism, and Obesity in Spanish Youth), a nationwide, representative, observational, and multicenter research study. Such project aimed to determine the levels of physical activity, sedentarism, lifestyle factors, and weight status of the Spanish youth population. Details of the PASOS protocol have been fully described [27]. A STROBE list is attached (see Supplementary File S1).

2.2. Participants and Recruitment

The participants of the PASOS study were children and adolescents aged 8 to 16 enrolled through randomly selected primary and secondary schools. Data were collected from March 2019 to February 2020 from 244 primary and secondary schools across the 17 Spanish autonomous communities. Informed consent was obtained for each participant, reaching a total of 4025 parents or legal guardians. Data on children's and adolescent's lifestyles were collected with self-reported online questionnaires with the support of trained professionals in each school. Additionally, anthropometric measurements (body weight, height, and waist circumference) of the participants were taken by trained professionals during school time.

2.3. Anthropometric Variables

Anthropometric variables were taken according to the WHO standardized protocol [28]. Body weight, height, and waist circumference were gathered using an electronic SECA 899 scale, a portable SECA 217 stadiometer, and a flexible non-stretch SECA 201 metric tape, respectively. The body mass index (BMI) (kg/m^2) was calculated using weight and height measures according to WHO growth charts [29]. Obesity and severe obesity were defined as ">WHO growth reference median + 2 standard deviation (SD)" and ">WHO growth reference median + 3 SD", respectively [29]. The waist-to-height ratio (WHtR) was calculated by dividing waist circumferences (cm) by height (cm), and abdominal obesity was defined as $\text{WHtR} \geq 0.5$ [30].

2.4. Assessment of Socioeconomic Status

Three SES indicators were gathered: The educational level and labor market status of parents or legal guardians were obtained through a self-reported paper questionnaire; the income was obtained through the database of the Spanish National Statistics Institute on the annual mean income per person in 2019 [31].

Each participant's family received two copies of the adult questionnaire. To Designated Adult 1, parents or legal guardians were asked who spent more time with the

child or adolescent. If a second parent or legal guardian answered the second copy of the questionnaire, they were designated as Adult 2. Data from mothers/female legal guardians designated as Adult 1 were prioritized (76.4%) followed by data from mothers/female legal guardians designated as Adult 2 (17.1%) if the parent/legal guardian designated as Adult 1 was male or absent; finally, data from fathers/male legal guardians designated as Adult 1 (6.1%) or Adult 2 (0.4%) were used if data were still lacking.

Parental educational level was categorized into “university” and “less than university”. The latter included general certificate of education, vocational education and training, general certificate of secondary education, primary education, and no education. Parental labor market status was designated as “employed” and “unemployed” excluding other labor market categories, such as household work, student, retirement, and permanent disability.

The annual mean income per person was assigned to each participant according to the corresponding value for the census section where their school was located. Two categories were defined: (i) above the total sample mean value for this variable ($\geq 12.731\text{€}$) and (ii) below the mean ($< 12.731\text{€}$).

Composite SES categories were created by combining each indicator and obtaining the less favorable category (Less than University/Unemployed/ $< 12.731\text{€}$; $n = 164$), the most favorable one (University/Employed/ $\geq 12.731\text{€}$; $n = 517$), and the intermediate ones.

2.5. Other Variables

The data on gender and age were obtained from the informed consent. Child and adolescent adherence to the Mediterranean diet and minutes dedicated to moderated/vigorous physical activity (MVPA) were self-reported by the KidMed index [32] and PAU7-S questionnaires [33], respectively.

2.6. Statistical Analysis

Logistic regression analysis was performed to determine the association between each of the 3 SES indicators and the prevalence of obesity, severe obesity, and abdominal obesity. Furthermore, a composite SES category was calculated including educational level (University/Less than University), labor market status (Employed/Unemployed), and annual mean income per person (Above/Below the studied population mean). The final composite variable of 9 categories was used for analysis after excluding 2 categories (University/Unemployed/ $< 12.731\text{€}$ and University/Unemployed/ $\geq 12.731\text{€}$) with less than 1% of the study population. The 6 composite SES categories were included as independent variables in logistic regression models with 3 anthropometric outcomes (obesity, severe obesity, and abdominal obesity) being the dependent variables. A first model was adjusted for gender and age and a second model also for school, adherence to the Mediterranean diet, and minutes of MVPA. Additionally, we tested also for interaction between indicators of SES with adherence to the Mediterranean diet and MVPA.

Moreover, the association of the composite SES categories with age and gender was tested. The associations were considered significant if $p < 0.05$. A flowchart of case selection included in this analysis is attached (see Supplementary File S2). The distribution by regions of the final analyzed sample is also declared and compared with the originally expected distribution (see Supplementary File S3). All statistical analyses were performed using SPSS for Windows version 22 (SPSS, Inc., Chicago, IL, USA).

2.7. Ethic Aspects

The PASOS study was performed according to the guidelines of the Declaration of Helsinki and approved by the Ethics Committee of the Fundació Sant Joan de Déu, Barcelona, Spain, on 17 December 2018 and Reference PIC-179-18. The trial was registered in 2019 at the International Standard Randomized Controlled Trial (ISRCT; <https://www.isrctn.com/ISRCTN34251612>, accessed on 1 March 2023).

3. Results

Table 1 shows that most parents had an educational level lower than a university degree (66.0%) and were employed (90.0%). Regarding income, 45.3% of the participants attended schools located in a census section with an annual mean income per person higher or equal to 12.731€. The prevalence of obesity, severe obesity, and abdominal obesity in children and adolescents was 11.5%, 1.4%, and 22.3%, respectively.

Table 1. Characteristics of the study population ($n = 2791$)—PASOS study. Spain, 2019–2020.

Study Sample Characteristics	Values (n ; % or Mean)
Girls	1449 (51.9%)
Age	12.59 (12.51–12.68)
BMI (kg/m ²)	20.29 (20.15–20.44)
Waist (cm)	70.70 (70.30–71.10)
Obesity	321 (11.5%)
Severe obesity	39 (1.4%)
Abdominal obesity	623 (22.3%)
Adherence to the Mediterranean diet (unit)	6.82 (6.73–6.91)
MVPA (min/d)	93.01 (91.75–94.28)
Parental SES indicators	
University degree	949 (34.0%)
Employed	2512 (90.0%)
Annual Income \geq 12.731€	1264 (45.3%)

Values are expressed as mean (95% CI) and number (%) for continuous variables and proportions, respectively. Body Mass Index (BMI). Moderated/Vigorous Physical Activity (MVPA).

Figure 1 shows the individual association of each of the three SES indicators with the anthropometric outcomes of the participants. Logistic regression analysis adjusted for gender and age revealed an inverse association of parental educational level with the prevalence of obesity, severe obesity, and abdominal obesity of adolescents (Figure 2). Being employed was also inversely associated with the three anthropometric outcomes. The magnitude of these associations was strongest for high educational levels and slightly attenuated after controlling for adherence to the Mediterranean diet, minutes of MVPA, and school. An inverse association was also found between parental income above the median and the prevalence of obesity and abdominal obesity. This association was not statistically significant for severe obesity ($p > 0.05$). Among the three indicators of SES, income shows the weakest associations with the anthropometric outcomes. The magnitude of these associations was lower in comparison to the other two indicators of SES, educational level, and labor market status. Parental education was the only indicator of SES that showed a significant interaction with adherence to the Mediterranean diet ($p = 0.024$) and MVPA ($p = 0.016$). However, there were no changes in the direction between the exposures and outcomes.

Figure 2 shows the association of the composite SES categories and the prevalence of obesity, severe obesity, and abdominal obesity in children and adolescents participating in this study. The direction and magnitude of the associations evidenced by Model 1 and Model 2 are very similar. According to Model 2, the combination of the most favorable SES indicators (University/Employed/Income above the media) was the least associated with obesity, severe obesity, and abdominal obesity. In particular, in comparison with the combination of the most unfavorable SES indicators (Less than University/Unemployed/Income below the media), the combination of the most favorable SES indicators was associated with a 72%, 80%, and 65% lower likelihood of obesity, severe obesity, and abdominal obesity, respectively. There was no statistically significant interaction between the composite SES categories and children's and adolescents' age and gender ($p > 0.05$).




Figure 1. Association between socioeconomic status and general, severe, and abdominal obesity ($n = 2791$)—PASOS study. Spain, 2019–2020. ¹ Performed by logistic regression analysis. ² Model 1: Adjusted for gender and age. ³ Model 2: Adjusted for gender, age, adherence to the Mediterranean diet, minutes of MVPA, and school. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.




Figure 2. Association between composite socioeconomic status and general, severe, and abdominal obesity ($n = 2791$)—PASOS study. Spain, 2019–2020. ¹ Performed by logistic regression analysis. ² Model 1: Adjusted for gender and age. ³ Model 2: Adjusted for gender, age, school, adherence to the Mediterranean diet, and minutes of MVPA. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

4. Discussion

The aim of this study was to evaluate the association between three parental SES indicators (individually and in combination) and weight status in a nationwide, representative sample of the Spanish child and adolescent population. Parental educational level, labor market status, and annual income per person were inversely related to obesity and abdominal obesity; moreover, parental educational level and labor market status also showed a negative association with severe obesity. Income showed a weaker association with weight status in comparison to educational level and labor market status indicators, but all three SES indicators should be considered when studying pediatric obesity. Finally, children and adolescents living exposed to the most advantageous SES factors were less likely to present obesity, severe obesity, or abdominal obesity than those living exposed to the least favorable SES factors.

Different studies led to opposite conclusions: Miqueleiz et al. reported a statistically significant upward trend in the prevalence of overweightness and obesity between 1987 and 2007 in Spanish girls and boys aged 10 to 15 years [17] but not in 5- to 9-year-old children from families with a lower educational level [17]. Another cross-sectional study with a population aged 2 to 15 years showed a lower prevalence of youth overweightness and obesity within families with a higher SES. This association was reported for all gender and age ranges except for girls under 12 [16]. Data from a longitudinal study that included a large young population from north–east Spain revealed the highest prevalence of overweightness and obesity in all gender and age categories for the most deprived participants [19]. Conversely, another cross-sectional study on the prevalence trend in childhood obesity and SES with a representative sample of a southern Spanish region [20] reported no significant social gradient for the prevalence of excess weight. Although a tendency towards an inverse association was found for the 2011–2012 data, it was less obvious for the 2015–2016 data. The present paper contributes to clarifying the heterogeneous results found by the original studies published so far in Spain. Moreover, it solves previous limitations, such as using anthropometric self-reported data and recruiting regional samples.

A global systematic review published in 2021 [34] found that in HICs, there was a higher risk for children from disadvantaged backgrounds to have a higher fat mass. Similarly, the present study found an inverse association between parental higher education and being employed with obesity, severe obesity, and abdominal obesity among Spanish children and adolescents. The magnitude of the association was higher for chil-

dren and adolescents whose parents had a university degree and slightly attenuated by adjusting for adherence to the Mediterranean diet, minutes of MVPA, and school. A recent systematic review identified parental education as the most commonly used SES measure [34] and the one that is more often significantly and inversely associated with childhood overweight/obesity in HICs. The second most commonly used SES indicator is parental income [11]. Our findings are consistent with the above-mentioned ones from European countries. However, in contrast to them, we showed that parental employment status was associated with the prevalence of being overweight [35].

The negative effects of severe childhood obesity on health [36] and its rising prevalence in many HICs are alarming [37]. Although emergent scientific evidence is showing that severe obesity is in general associated with lower socioeconomic status [38], more in-depth analyses are needed to study its association with specific SES indicators [39]. Data from the WHO European Childhood Obesity Surveillance Initiative (COSI) found that in European countries, severe obesity was more common among children whose mothers had a lower educational level. Southern European countries had the highest levels of severe obesity with a prevalence of 4% [40]. The prevalence of severe obesity was 5.1% among children whose mothers had a lower educational level and 2.9% among children whose mothers had a higher educational level [40]. In the present study, a strong and negative association of severe obesity with parental educational level was also obtained. Specifically, children and adolescents whose parents had a university degree were 34% less likely to present severe obesity in comparison with those whose parents had a lower educational level.

To the best of our knowledge, few studies have reported the association between SES indicators and abdominal obesity in young populations. A cross-sectional study among European adolescents [41] indicates an inverse association, which is consistent with the findings of this study. The magnitude of the association between abdominal obesity and SES indicators reported by Costa de Oliveira Forkert et al. [41] is comparable to the one found in the present study. Moreover, a longitudinal study showed that abdominal obesity was two-folds higher in Spanish children aged 4 to 9 years with parents of low SES in comparison to those with parents of high SES [42]. In our study, children and adolescents exposed to the most favorable SES factors were less likely to present abdominal obesity.

SES inequalities are an important determinant of obesity [11]. The pathways through which inequalities are associated with weight status in the young population are complex and interrelated [43]: structural, community, and individual factors influencing the reaction to food advertising and marketing [44]; food environment [45,46]; access to sports facilities [47] and safe [48] and green spaces [49]; parental role [50,51]; and individual lifestyles [52]. For example, children from socioeconomically disadvantaged backgrounds are disproportionately exposed to unhealthy food advertising [44]. Moreover, those attending schools located in more deprived neighborhoods had fewer opportunities to find establishments selling healthy food products, and they were more exposed to fast food establishments, convenience stores, supermarkets, and grocery stores offering unhealthy foods [45,46]. Additionally, young people in low-SES communities have larger obstacles in participating in sports and physical activities because their neighborhoods might not be safe enough [48]. As for green spaces, in southern European cities, many of the neighborhoods had access to them, but the mean distance was higher for deprived neighborhoods [49]. Additionally, green spaces in the most deprived neighborhoods presented significantly more safety concerns, signs of damage, and a lack of equipment to engage in active leisure activities and had significantly fewer amenities, such as seating or public toilets [49]. Finally, in Madrid, although disadvantaged areas had a shorter distance to the closest sports facility, especially for public and low-cost facilities, the overall number of such facilities was lower in low SES areas [47].

Scientific evidence highlights that families with both low educational levels and income have a greater predisposition to experience psychological distress [53], and children are more likely to be exposed to multiple adversities [54]. Moreover, several studies found that parental stress was associated with childhood obesity [51,55]. Stress and adversity

affect parents' abilities to provide a safe, stable, responsive, and nurturing environment for children [51] and are associated with unhealthy practices, such as fast food consumption by children [55]. Indeed, a higher SES is associated with better health in children, fewer behavioral difficulties, a better quality of life, fewer critical life events, and a healthier lifestyle (less time spent watching television and more practicing physical activity) [52]. In conclusion, children who grow up in circumstances with more socioeconomic risk factors will have a predisposition towards sustained weight growth, whereas those who grow up with more protective factors will better maintain a healthy weight [56]. Lastly, different studies reported that children living in lower socioeconomic backgrounds are more vulnerable to overweightness and obesity later on during childhood or adulthood even if their economic environment improves [57,58].

The main strength of this study is the nationwide, representative design. Furthermore, body weight, height, and waist circumference were measured by trained professionals. Another strength is the use of three different standardized SES indicators to examine the individual and joined association between SES and weight status. Vazquez and Cubbin suggested using multiple measures of SES because one variable cannot capture all aspects embedded in SES [11]. The present analysis includes fewer participants than the total recruited due to missing the different interest variables. However, the cases studied were from all Spanish regions following a very similar percentual distribution than the total invited eligible participants (see Supplementary File S3), which is a strength of this study.

The main limitation of the present study is that the annual mean income per person is based on the census section where the participant schools were located, and using aggregated data is less reliable than using individual data. However, it is known that families commonly choose a school located in their neighborhood [59,60]. Finally, even if some schools are situated at the boundary between two census sections and may enroll students from each of them, we assumed that children and adolescents were living in the same census section of their school [59,60].

Previous scientific evidence highlighted the need to find common indicators to better approximate the SES of an individual [61], and our study represents a step forward in this direction. Future research is needed to harmonize the use of indicators and to enhance comparability between results from different regions and countries [12].

The results included in this paper correspond to the first edition of the PASOS project carried out in 2019. Data collection from a second edition will be completed in 2023. It will include a new cross-sectional study and a follow-up of participants from the first edition, allowing for a comparative analysis of the association between SES and weight status over an extended period. Moreover, the longitudinal follow-up of the participants from the first edition will enhance the analysis of the association of obesity, severe obesity, and abdominal obesity with SES. Finally, emerging evidence suggests the need to study the impact of the COVID-19 pandemic on the aforementioned association because a higher increase in weight was observed among young people living in unfavorable socioeconomic conditions in comparison to people living in more favorable socioeconomic conditions [62].

Previous research suggested the need to explore the factors mediating socioeconomic differences in adiposity among children and adolescents to define youth policy development [63]. In a review, Gebremariam et al. reported several factors that could be targeted in interventions aimed at reducing socioeconomic differences in excess weight among young people. These include early life experience (particularly breastfeeding, early weaning, and maternal smoking in pregnancy); child dietary behavior (particularly consumption of sugar-sweetened beverages and breakfast eating patterns); child sedentary activity (particularly watching television and using computers); and maternal BMI [63]. Understanding the mechanisms behind the association of these factors with socioeconomic differences is crucial to tackling childhood obesity by reducing SES inequalities. Moreover, research indicated that programs to prevent childhood obesity tended to reach more people with higher SES levels who often benefit more from innovative initiatives [64,65]. In this regard,

it would be interesting to better understand how to target obesity prevention programs for the most disadvantaged population.

A recent report by the Spanish Agency for Food Safety and Nutrition and High Commissioner Against Child Poverty [15] suggests implementing initiatives oriented at facilitating healthy eating in a sustained manner throughout the year at school. This way, it would be possible to address social inequalities; promote daily physical activity and active leisure among school children and reduce sedentary lifestyles; reinforce education and awareness-raising among parents and schoolchildren; and continue to develop epidemiological surveillance of childhood obesity and its determinants, paying particular attention to lowering inequities.

In this sense, the National Strategic Plan for the Reduction of Childhood Obesity (2022–2030) is starting its implementation in Spain [23] and should guide the efforts addressed to prevent childhood obesity overall with a special focus on children and adolescents living exposed to disadvantaged factors.

5. Conclusions

This study demonstrated that Spanish children and adolescents with a lower SES are more likely to present obesity, severe obesity, and abdominal obesity. Therefore, to tackle the obesity epidemic, it is essential to consider the associated social health inequalities.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/nu15081987/s1>, File S1: STROBE Statement—Checklist of items that should be included in reports of cross-sectional studies; File S2: Flowchart; File S3: Comparative of expected study sample size and final sample analyzed by country region.

Author Contributions: Conceptualization, C.H., A.A., M.S., H.S. and S.F.G.; Methodology, C.H., R.M., H.S. and S.F.G.; Validation, C.H.; Formal analysis, C.H., H.S. and S.F.G.; Investigation, C.H., P.B., J.W., M.O., M.G.-G., N.G., S.A., E.M.-C., M.Á.G.-V., L.S.-M., N.T., J.A.T., M.F., J.C.B.-M., I.L., A.G.Z., J.S.-G., F.J.-Z., P.E.A., M.S.-S., E.H.-R., S.P.-M., C.B., H.S. and S.F.G.; Resources, S.F.G.; Data curation, C.H., H.S. and S.F.G.; Writing—original draft, C.H., H.S. and S.F.G.; Writing—review and editing, C.H., P.B., A.A., J.W., M.O., M.G.-G., N.G., S.A., E.M.-C., M.Á.G.-V., L.S.-M., N.T., J.A.T., M.S., M.F., J.C.B.-M., I.L., A.G.Z., J.S.-G., F.J.-Z., P.E.A., M.S.-S., E.H.-R., S.P.-M., C.B., R.M., H.S. and S.F.G.; Visualization, C.H., H.S. and S.F.G.; Supervision, H.S. and S.F.G.; Project administration, S.F.G.; Funding acquisition, S.F.G. All authors have read and agreed to the published version of the manuscript.

Funding: The PASOS study was funded by Fundación PROBITAS (2019) and the Gasol Foundation (2019–2020). Additional funds were received from the Barça Foundation (2019–2020), Banco Santander (2019), IFA (2019–2020), Vienna (2019), and the Fundación Deporte Joven (2019) (no references are applicable). J.A.T., M.G.-G. and C.B. are funding by Instituto de Salud Carlos III through the CIBEROBN CB12/03/30038, which are co-funded by the European Regional Development Fund.

Institutional Review Board Statement: The PASOS study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Ethics Committee of the Fundació Sant Joan de Déu, Barcelona, Spain, on 17 December 2018 and Reference PIC-179-18. (Please see our paper on the protocol of the study: [27].)

Informed Consent Statement: Informed consent has been obtained from parents or legal guardians of each participant.

Data Availability Statement: There are restrictions on the availability of the data for this trial because of the signed consent agreements and concerns regarding data sharing, which only allow access to external researchers for studies following the project purposes. Requestors wishing to access the trial data used in this study can make a request to Santi F. Gómez Santos, PhD (sgomez@gasolfoundation.org) principal researcher of the PASOS study.

Acknowledgments: We thank the staff, pupils, parents, schools, and municipalities for their participation, enthusiasm, and support. PASOS has the institutional support of Spain's Ministry of Education and Vocational Training, the Ministry of Health, Consumption and Social Welfare through the Spanish Agency for Food Safety and Nutrition (ASEAN), the High Commission Against Child

Poverty, the High Sports Council, the General College of Professional Associations of Physical Education and Sports, and the Departments of Education and/or Health and/or Sports of Spain's 17 autonomous regions.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the analysis or interpretation of the data in this study.

References

- Bentham, J.; Di Cesare, M.; Bilano, V.; Bixby, H.; Zhou, B.; Stevens, G.A.; Riley, L.M.; Taddei, C.; Hajifathalian, K.; Lu, Y.; et al. Worldwide Trends in Body-Mass Index, Underweight, Overweight, and Obesity from 1975 to 2016: A Pooled Analysis of 2416 Population-Based Measurement Studies in 128.9 Million Children, Adolescents, and Adults. *Lancet* **2017**, *390*, 2627–2642. [[CrossRef](#)]
- Bravo-Saquicela, D.M.; Sabag, A.; Rezende, L.F.M.; Rey-Lopez, J.P. Has the Prevalence of Childhood Obesity in Spain Plateaued? A Systematic Review and Meta-Analysis. *Int. J. Environ. Res. Public Health* **2022**, *19*, 5240. [[CrossRef](#)] [[PubMed](#)]
- Kumar, S.; Kelly, A.S. Review of Childhood Obesity: From Epidemiology, Etiology and Comorbidities to Clinical Assessment and Treatment. *Mayo Clin. Proc.* **2017**, *92*, 251–265. [[CrossRef](#)] [[PubMed](#)]
- Neri, D.; Steele, E.M.; Khandpur, N.; Cediel, G.; Zapata, M.E.; Rauber, F.; Marrón-Ponce, J.A.; Machado, P.; Louzada, M.L.C.; Andrade, G.C.; et al. Ultraprocessed food consumption and dietary nutrient profiles associated with obesity: A multicountry study of children and adolescents. *Obes. Rev.* **2022**, *23*, e13387. [[CrossRef](#)]
- Kelley, G.A.; Kelley, K.S.; Pate, R.R. Exercise and BMI in Overweight and Obese Children and Adolescents: A Systematic Review and Trial Sequential Meta-Analysis. *BioMed. Res. Int.* **2015**, *2015*, 704539. [[CrossRef](#)]
- Morrissey, B.; Taveras, E.; Allender, S.; Strugnell, C. Sleep and obesity among children: A systematic review of multiple sleep dimensions. *Pediatr. Obes.* **2020**, *15*, e12619. [[CrossRef](#)]
- Moradi, M.; Mozaffari, H.; Askari, M.; Azadbakht, L. Association between overweight/obesity with depression, anxiety, low self-esteem, and body dissatisfaction in children and adolescents: A systematic review and meta-analysis of observational studies. *Crit. Rev. Food Sci. Nutr.* **2022**, *62*, 555–570. [[CrossRef](#)]
- Littleton, S.; Berkowitz, R.I.; Grant, S.F.A. Genetic Determinants of Childhood Obesity. *Mol. Diagn. Ther.* **2020**, *24*, 653–663. [[CrossRef](#)]
- Barriuso, L.; Miqueleiz, E.; Albaladejo, R.; Villanueva, R.; Santos, J.M.; Regidor, E. Socioeconomic position and childhood-adolescent weight status in rich countries: A systematic review, 1990–2013. *BMC Pediatr.* **2015**, *15*, 129. [[CrossRef](#)]
- Chung, A.; Backholer, K.; Wong, E.; Palermo, C.; Keating, C.; Peeters, A. Trends in child and adolescent obesity prevalence in economically advanced countries according to socioeconomic position: A systematic review. *Obes. Rev.* **2016**, *17*, 276–295. [[CrossRef](#)]
- Vazquez, C.E.; Cubbin, C. Socioeconomic Status and Childhood Obesity: A Review of Literature from the Past Decade to Inform Intervention Research. *Curr. Obes. Rep.* **2020**, *9*, 562–570. [[CrossRef](#)] [[PubMed](#)]
- Sares-Jäske, L.; Grönqvist, A.; Mäki, P.; Tolonen, H.; Laatikainen, T. Family socioeconomic status and childhood adiposity in Europe—A scoping review. *Prev. Med.* **2022**, *160*, 107095. [[CrossRef](#)] [[PubMed](#)]
- Buonocristiano, M.; Williams, J.; Simmonds, P.; Nurk, E.; Ahrens, W.; Nardone, P.; Rito, A.I.; Rutter, H.; Bergh, I.H.; Starc, G.; et al. Socioeconomic inequalities in overweight and obesity among 6- to 9-year-old children in 24 countries for the World Health Organization European region. *Obes. Rev.* **2021**, *22*, e13213. [[CrossRef](#)] [[PubMed](#)]
- Fitzsimons, E.; Bann, D. *Obesity Prevalence and Its Inequality from Childhood to Adolescence: Initial Findings from the Millennium Cohort Study Age 17 Survey*; Center for Longitudinal Studies: London, UK, 2020; p. 4.
- García Solano, M.; Gutiérrez González, E.; Peña-Rey Lorenzo, I.; Arcarons Feixas, A.; Arisas Lera, A.; Jorquera Rojas, G. *Obesidad y Pobreza Infantil: Radiografía de Una Doble Desigualdad. Estudio Del Rol de Los Factores Socioeconómicos En La Obesidad de Los Escolares En España*; Agencia Española de Seguridad Alimentaria y Nutrición, Ministerio de Consumo, Alto Comisionado contra la Pobreza Infantil: Madrid, Spain, 2022; p. 30.
- Pizarro, J.V. Prevalence of Childhood Obesity in Spain; National Health Survey 2006–2007. *Nutr. Hosp.* **2012**, *27*, 154–160. [[CrossRef](#)]
- Miqueleiz, E.E.; Lostao, L.; Ortega, P.; Santos, J.M.; Astasio, P.; Regidor, E.E. Trends in the prevalence of childhood overweight and obesity according to socioeconomic status: Spain, 1987–2007. *Eur. J. Clin. Nutr.* **2014**, *68*, 209–214. [[CrossRef](#)]
- Gil, J.M.; Takourab, S. Socio-economics, food habits and the prevalence of childhood obesity in Spain. *Child Care Health Dev.* **2017**, *43*, 250–258. [[CrossRef](#)]
- De Bont, J.; Díaz, Y.; Casas, M.; García-Gil, M.; Vrijheid, M.; Duarte-Salles, T. Time Trends and Sociodemographic Factors Associated with Overweight and Obesity in Children and Adolescents in Spain. *JAMA Netw. Open* **2020**, *3*, e201171. [[CrossRef](#)]
- Sánchez-Cruz, J.-J.; De Ruiter, I.; Jiménez-Moleón, J.J.; García, L.; Sánchez, M.-J. Stabilization and reversal of child obesity in Andalusia using objective anthropometric measures by socioeconomic status. *BMC Pediatr.* **2018**, *18*, 322. [[CrossRef](#)]
- Loring, B.; Robertson, A. *Obesity and Inequities: Guidance for Addressing Inequities in Overweight and Obesity*; World Health Organization: Geneva, Switzerland, 2014; ISBN 978 92 890 5048 7.

22. WHO. *Regional Office for Europe WHO European Regional Obesity Report*; World Health Organization: Copenhagen, Denmark, 2022; p. 220.
23. High Commissioner against Child Poverty. *National Strategic Plan for The Reduction of Childhood Obesity (2022–2030) Spain. Executive Summary*; Alto Comisionado contra la Pobreza Infantil, Presidencia del Gobierno: Madrid, Spain, 2022; p. 23.
24. Royo-Bordonada, M.; Fernández-Escobar, C.; Gil-Bellosta, C.J.; Ordaz, E. Effect of excise tax on sugar-sweetened beverages in Catalonia, Spain, three and a half years after its introduction. *Int. J. Behav. Nutr. Phys. Act.* **2022**, *19*, 24. [CrossRef]
25. Gómez, S.F.; Rajmil, L. Advertising, obesity and child health: The case of Spain. *BMJ Paediatr. Open* **2022**, *6*, e001482. [CrossRef]
26. Borrador de Real Decreto Sobre Regulación de La Publicidad de Alimentos y Bebidas Dirigida al Público Infantil | Ministerio de Consumo. Available online: <https://www.consumo.gob.es/es/consulta-p-blica/borrador-de-real-decreto-sobre-regulacion-de-la-publicidad-de-alimentos-y-bebidas> (accessed on 6 February 2023).
27. Gómez, S.F.; Homs, C.; Wärnberg, J.; Medrano, M.; Gonzalez-Gross, M.; Gusi, N.; Aznar, S.; Cascales, E.M.; González-Valeiro, M.; Serra-Majem, L.; et al. Study protocol of a population-based cohort investigating Physical Activity, Sedentarism, lifestyles and Obesity in Spanish youth: The PASOS study. *BMJ Open* **2020**, *10*, e036210. [CrossRef] [PubMed]
28. World Health Organization (WHO). Weighing and Measuring. *Child Growth Standards*. Available online: <https://www.who.int/tools/child-growth-standards> (accessed on 6 February 2023).
29. De Onis, M.; Onyango, A.W.; Borghi, E.; Siyam, A.; Nishida, C.; Siekmann, J. Development of a WHO growth reference for school-aged children and adolescents. *Bull. World Health Organ.* **2007**, *85*, 660–667. [CrossRef]
30. Taylor, R.W.; Jones, I.E.; Williams, S.M.; Goulding, A. Evaluation of waist circumference, waist-to-hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by dual-energy X-ray absorptiometry, in children aged 3–19 y. *Am. J. Clin. Nutr.* **2000**, *72*, 490–495. [CrossRef] [PubMed]
31. INE: Indicadores de Renta Media y Mediana. Available online: <https://www.ine.es/jaxiT3/Tabla.htm?t=31097> (accessed on 6 February 2023).
32. Serra-Majem, L.; Ribas, L.; Ngo, J.; Ortega, R.M.; García, A.; Pérez-Rodrigo, C.; Aranceta, J. Food, youth and the Mediterranean diet in Spain. Development of KIDMED, Mediterranean Diet Quality Index in children and adolescents. *Public Health Nutr.* **2004**, *7*, 931–935. [CrossRef] [PubMed]
33. Schröder, H.; Subirana, I.; Wärnberg, J.; Medrano, M.; González-Gross, M.; Gusi, N.; Aznar, S.; Alcaraz, P.E.; González-Valeiro, M.A.; Serra-Majem, L.; et al. Validity, reliability, and calibration of the physical activity unit 7 item screener (PAU-7S) at population scale. *Int. J. Behav. Nutr. Phys. Act.* **2021**, *18*, 98. [CrossRef] [PubMed]
34. Staatz, C.B.; Kelly, Y.; Lacey, R.E.; Blodgett, J.M.; George, A.; Arnot, M.; Walker, E.; Hardy, R. Socioeconomic position and body composition in childhood in high- and middle-income countries: A systematic review and narrative synthesis. *Int. J. Obes.* **2021**, *45*, 2316–2334. [CrossRef]
35. Buoncristiano, M.; Spinelli, A.; Williams, J.; Nardone, P.; Rito, A.I.; García-Solano, M.; Grøholt, E.K.; Gutiérrez-González, E.; Klepp, K.I.; Starc, G.; et al. Childhood overweight and obesity in Europe: Changes from 2007 to 2017. *Obes. Rev.* **2021**, *22*, e13226. [CrossRef]
36. Chung, Y.L.; Rhie, Y.-J. Severe Obesity in Children and Adolescents: Metabolic Effects, Assessment, and Treatment. *J. Obes. Metab. Syndr.* **2021**, *30*, 326–335. [CrossRef]
37. Jebeile, H.; Kelly, A.S.; O'Malley, G.; Baur, L.A. Obesity in children and adolescents: Epidemiology, causes, assessment, and management. *Lancet Diabetes Endocrinol.* **2022**, *10*, 351–365. [CrossRef]
38. Pinhas-Hamiel, O.; Hamiel, U.; Bendor, C.D.; Bardugo, A.; Twig, G.; Cukierman-Yaffe, T. The Global Spread of Severe Obesity in Toddlers, Children, and Adolescents: A Systematic Review and Meta-Analysis. *Obes. Facts* **2022**, *15*, 118–134. [CrossRef]
39. Kelly, A.S.; Barlow, S.E.; Rao, G.; Inge, T.H.; Hayman, L.L.; Steinberger, J.; Urbina, E.M.; Ewing, L.J.; Daniels, S.R. Severe Obesity in Children and Adolescents: Identification, Associated Health Risks, and Treatment Approaches: A Scientific Statement from the American Heart Association. *Circulation* **2013**, *128*, 1689–1712. [CrossRef] [PubMed]
40. Spinelli, A.; Buoncristiano, M.; Kovacs, V.A.; Yngve, A.; Spiroski, I.; Obreja, G.; Starc, G.; Pérez, N.; Rito, A.I.; Kunešová, M.; et al. Prevalence of Severe Obesity among Primary School Children in 21 European Countries. *Obes. Facts* **2019**, *12*, 244–258. [CrossRef] [PubMed]
41. Forkert, E.C.D.O.; de Moraes, A.C.F.; Carvalho, H.B.; Kafatos, A.; Manios, Y.; Sjöström, M.; González-Gross, M.; Gottrand, F.; Beghin, L.; Censi, L.; et al. Abdominal obesity and its association with socioeconomic factors among adolescents from different living environments. *Pediatr. Obes.* **2017**, *12*, 110–119. [CrossRef] [PubMed]
42. Ortiz-Marrón, H.; Ortiz-Pinto, M.A.; Pujadas, G.C.; Mosquera, J.G.M.; Miñarro, M.L.; Pinos, F.M.; Gavín, M.O.; Galán, I. Tracking and risk of abdominal and general obesity in children between 4 and 9 years of age. The Longitudinal Childhood Obesity Study (ELOIN). *BMC Pediatr.* **2022**, *22*, 198. [CrossRef]
43. Pearce, A.; Dundas, R.; Whitehead, M.; Taylor-Robinson, D. Pathways to inequalities in child health. *Arch. Dis. Child.* **2019**, *104*, 998–1003. [CrossRef]
44. Backholer, K.; Gupta, A.; Zorbas, C.; Bennett, R.; Huse, O.; Chung, A.; Isaacs, A.; Golds, G.; Kelly, B.; Peeters, A. Differential exposure to, and potential impact of, unhealthy advertising to children by socio-economic and ethnic groups: A systematic review of the evidence. *Obes. Rev.* **2021**, *22*, e13144. [CrossRef]

45. Londoño-Cañola, C.; Serral, G.; Díez, J.; Martínez-García, A.; Franco, M.; Artazcoz, L.; Ariza, C. Retail Food Environment around Schools in Barcelona by Neighborhood Socioeconomic Status: Implications for Local Food Policy. *Int. J. Environ. Res. Public Health* **2023**, *20*, 649. [[CrossRef](#)]
46. De França, F.C.O.; Andrade, I.D.S.; Zandonadi, R.P.; Sávio, K.E.; Akutsu, R.D.C.C.D.A. Food Environment around Schools: A Systematic Scope Review. *Nutrients* **2022**, *14*, 5090. [[CrossRef](#)]
47. Cereijo, L.; Gullón, P.; Cebrecos, A.; Bilal, U.; Santacruz, J.A.; Badland, H.; Franco, M. Access to and availability of exercise facilities in Madrid: An equity perspective. *Int. J. Health Geogr.* **2019**, *18*, 15. [[CrossRef](#)]
48. Mohammed, S.H.; Habtewold, T.D.; Birhanu, M.M.; Sissay, T.A.; Tegegne, B.S.; Abuzerr, S.; Esmailzadeh, A. Neighbourhood socioeconomic status and overweight/obesity: A systematic review and meta-analysis of epidemiological studies. *BMJ Open* **2019**, *9*, e028238. [[CrossRef](#)]
49. Hoffmann, E.; Barros, H.; Ribeiro, A.I. Socioeconomic Inequalities in Green Space Quality and Accessibility—Evidence from a Southern European City. *Int. J. Environ. Res. Public Health* **2017**, *14*, 916. [[CrossRef](#)] [[PubMed](#)]
50. Danford, C.A.; Schultz, C.; Marvicsin, D. Parental roles in the development of obesity in children: Challenges and opportunities. *Res. Rep. Biol.* **2015**, *6*, 39–53. [[CrossRef](#)]
51. Jang, M.; Owen, B.; Lauver, D.R. Different types of parental stress and childhood obesity: A systematic review of observational studies. *Obes. Rev.* **2019**, *20*, 1740–1758. [[CrossRef](#)] [[PubMed](#)]
52. Poulain, T.; Vogel, M.; Sobek, C.; Hilbert, A.; Körner, A.; Kiess, W. Associations Between Socio-Economic Status and Child Health: Findings of a Large German Cohort Study. *Int. J. Environ. Res. Public Health* **2019**, *16*, 677. [[CrossRef](#)]
53. Hemmingsson, E. A new model of the role of psychological and emotional distress in promoting obesity: Conceptual review with implications for treatment and prevention. *Obes. Rev.* **2014**, *15*, 769–779. [[CrossRef](#)]
54. Francis, L.; DePriest, K.; Wilson, M.; Gross, D. Child Poverty, Toxic Stress, and Social Determinants of Health: Screening and Care Coordination. *Online J. Issues Nurs.* **2018**, *23*, 2. [[CrossRef](#)]
55. Parks, E.P.; Kumanyika, S.; Moore, R.H.; Stettler, N.; Wrotniak, B.H.; Kazak, A. Influence of Stress in Parents on Child Obesity and Related Behaviors. *Pediatrics* **2012**, *130*, e1096–e1104. [[CrossRef](#)]
56. Kim, T.J.; Von Dem Knesebeck, O. Income and Obesity: What Is the Direction of the Relationship? A Systematic Review and Meta-Analysis. *BMJ Open* **2018**, *8*, e019862. [[CrossRef](#)]
57. Kakinami, L.; Séguin, L.; Lambert, M.; Gauvin, L.; Nikiema, B.; Paradis, G. Poverty's latent effect on adiposity during childhood: Evidence from a Québec birth cohort. *J. Epidemiol. Community Health* **2014**, *68*, 239–245. [[CrossRef](#)]
58. Li, M.; Mustillo, S.; Anderson, J. Childhood poverty dynamics and adulthood overweight/obesity: Unpacking the black box of childhood. *Soc. Sci. Res.* **2018**, *76*, 92–104. [[CrossRef](#)]
59. Calsamiglia, C.; Güell, M. *¿Cómo Escogen Los Padres La Escuela de Sus Hijos? Teoría y Evidencia Para España*; FEDE: Barcelona, Spain, 2013; p. 27.
60. Murillo, F.J.; Almazán, A.; Garrido, C.M. La elección de centro educativo en un sistema de cuasi-mercado escolar mediado por el programa de bilingüismo. *Rev. Complut. De Educ.* **2021**, *32*, 89–97. [[CrossRef](#)]
61. Darin-Mattsson, A.; Fors, S.; Kåreholt, I. Different indicators of socioeconomic status and their relative importance as determinants of health in old age. *Int. J. Equity Health* **2017**, *16*, 173. [[CrossRef](#)] [[PubMed](#)]
62. La Fauci, G.; Montalti, M.; Di Valerio, Z.; Gori, D.; Salomoni, M.G.; Salussolia, A.; Soldà, G.; Guaraldi, F. Obesity and COVID-19 in Children and Adolescents: Reciprocal Detrimental Influence—Systematic Literature Review and Meta-Analysis. *Int. J. Environ. Res. Public Health* **2022**, *19*, 7603. [[CrossRef](#)] [[PubMed](#)]
63. Gebremariam, M.K.; Lien, N.; Nianogo, R.A.; Arah, O.A. Mediators of socioeconomic differences in adiposity among youth: A systematic review. *Obes. Rev.* **2017**, *18*, 880–898. [[CrossRef](#)] [[PubMed](#)]
64. Peeters, A.; Backholer, K. Prioritising and tackling socio-economic inequalities in obesity. *BMC Obes.* **2014**, *1*, 16. [[CrossRef](#)]
65. Beauchamp, A.; Backholer, K.; Magliano, D.; Peeters, A. The effect of obesity prevention interventions according to socioeconomic position: A systematic review. *Obes. Rev.* **2014**, *15*, 541–554. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.