

## Laser cooling of stored relativistic ion beams with large momentum spreads using a laser system with a wide scanning range

Weiqliang Wen<sup>1</sup>, Danyal Winters<sup>2</sup>, Tobias Beck<sup>3</sup>, Benjamin Rein<sup>3</sup>, Thomas Walther<sup>3</sup>, Sascha Tichelmann<sup>3</sup>, Gerhard Birkel<sup>3</sup>, Rodolfo Sanchez-Alarcon<sup>2</sup>, Johannes Ullmann<sup>2</sup>, Matthias Lochmann<sup>2,4</sup>, Wilfried Nörtershäuser<sup>2,4,5</sup>, Colin Clark<sup>2</sup>, Christophor Kozhuharov<sup>2</sup>, Thomas Kühl<sup>2,4,6</sup>, Shahab Sanjari<sup>2</sup>, Yuri Litvinov<sup>2</sup>, Tino Giacomini<sup>2</sup>, Markus Steck<sup>2</sup>, Christina Dimopoulou<sup>2</sup>, Fritz Nolden<sup>2</sup>, Thomas Stöhlker<sup>2,6,7</sup>, Jie Yang<sup>1</sup>, Dacheng Zhang<sup>1</sup>, Xinwen Ma<sup>1</sup>, Michael Seltmann<sup>8</sup>, Matthias Siebold<sup>8</sup>, Ulrich Schramm<sup>8</sup> and Michael Bussmann<sup>8,\*</sup>

<sup>1</sup> Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou, 730000, Gansu, China

<sup>2</sup> GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany

<sup>3</sup> Institut für Angewandte Physik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

<sup>4</sup> Institut für Kernchemie, Johannes Gutenberg Universität Mainz, 55099 Mainz, Germany.

<sup>5</sup> Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

<sup>6</sup> Helmholtz-Institut Jena, 07743 Jena, Germany


<sup>7</sup> Institut für Optik und Quantenelektronik, Universität Jena, 07737 Jena, Germany.

<sup>8</sup> Helmholtz-Zentrum Dresden-Rossendorf, 01328 Dresden, Germany

**Synopsis** New results on laser cooling of stored, bunched, relativistic ion beams are presented. For the first time it has been possible to cool an ion beam with large momentum spread without initial electron cooling or scanning of the bunching frequency by using a single cw laser system.

We present results on a recent laser cooling experiment on bunched  $C^{3+}$  ion beams at relativistic energies at the Experimental Storage Ring, GSI Darmstadt, Germany. In order to directly cool all ions with an initial momentum spread of the RF-bunched beam, a scanning cw laser system has been applied during the experiment, the frequency of this laser system at the UV range (257.3 nm) can be scanned more than 12 GHz corresponding to the longitudinal momentum spread about  $\sim 10^{-5}$  of the ion beam. A schematic view of the laser cooling of a coasting ion beam and a bunched ion beam is shown in Figure 1(a) and Figure 1(b), respectively.

Our results prove that the initially large momentum spread of bunched ion beams can be efficiently cooled using a single cw laser by scanning the laser frequency over a broad frequency range. The cooling rate is mainly limited by the speed of the frequency scan. In comparison to previous laser cooling experiments at ESR [1, 2], this new cooling scheme does not require initial electron cooling of the ion beam or scanning of the bunching frequency. This very promising scheme can be directly adopted to future high-energy ion storage rings. In order to further improve the technique, follow up experiments at the CSRe in Lanzhou are planned for this year.



**Figure 1.** Schematic view of the cooling scheme for a coasting ion beam (a) and a bunched ion beam (b). The Schottky noise signal strength is plotted logarithmically.

### References

- [1] U Schramm *et al* 2006 *AIP Conf. Proc.* **821** 501
- [2] M Bussmann *et al* 2007 *J. Phys.: Conf.* **88** 01204

\* E-mail: [m.bussmann@hzdr.de](mailto:m.bussmann@hzdr.de)