

The m -Order Linear Recursive Quaternions

Orhan Dişkaya and Hamza Menken

Abstract. This study considers the m -order linear recursive sequences yielding some well-known sequences (such as the Fibonacci, Lucas, Pell, Jacobsthal, Padovan, and Perrin sequences). Also, the Binet-like formulas and generating functions of the m -order linear recursive sequences have been derived. Then, we define the m -order linear recursive quaternions, and give the Binet-like formulas and generating functions for them.

Key Words and Phrases: Linear Recursive, Fibonacci numbers, quaternions, generating functions.

2010 Mathematics Subject Classifications: Primary 11B39, 11R52, 05A15

1. Introduction

Primarily, we will consider a linear recursion sequence that gives us some special sequences such as Fibonacci, Lucas, Pell, Jacobsthal, Padovan, Perrin, and Tribonacci with certain initial conditions and coefficients. Then, we obtain the Binet-like formula and generating functions of the linear recursive sequence to find the Binet-like formulas and the generating functions of some special sequences by choosing certain initial conditions and coefficients. Thus, we will make it easier for us to prove the Binet-like formulas and generating functions of some special sequences as a result of this study. The m -order linear recursive sequence definition given below is given by Matyas and Szakacs in [1, 4]. Now, let's examine some identities by reminding this definition again.

For $a_0, a_1, \dots, a_{m-1} \in \mathbb{Z}$ with $a_{m-1} \neq 0$ and $m \in \mathbb{Z}^+$, the m -order linear recursive sequence $\{S_n\}_{n \geq 0}$ are defined by recurrence relation

$$S_{n+m} = \sum_{k=0}^{m-1} a_k S_{n+k} \quad (1)$$

where the initial conditions S_0, S_1, \dots, S_{m-1} with $|S_0| + |S_1| + \dots + |S_{m-1}| \neq 0$. The recurrence relation (1) involves the characteristic equation

$$a_0 + a_1x + a_2x^2 + \dots + a_{m-1}x^{m-1} - x^m = 0.$$

By the complex numbers q_1, q_2, \dots, q_m , we denote the roots of the characteristic equation. Assume that the numbers a_i 's are chosen such that the roots of the characteristic equation are distinct.

Linear recursive sequences have been studied by many authors [1, 2, 3, 4, 5, 42]. Matyas investigated some sequence transformations of $\{G_{n+d}/G_n\}_{n=0}^{\infty}$ of linear recursive sequences and linear recurrences and roots-finding methods in [1, 5] where $\{G_n\}$ is a linear sequence with m -order. Gatta and D'amito studied sequences H_n for which H_{n+1}/H_n approaches the golden ratio in [2] where $\{H_n\}$ is a third order linear sequence. Komatsu continued the work of Gatta and D'amito, and examined the sequence H_n for which H_{n+1}/H_n approaches an irrational number in [3]. Szakacs investigated sequence $\{G_{n+1}/G_n\}_{n=1}^{\infty}$ which are approaching the Golden Ratio, in case $\{G_n\}_{n=0}^{\infty}$ is defined the k -order linear recursive sequence of real numbers [4]. In the present work, we derive the Binet-like Formula and generating functions in the general case.

In [41, 43], the Binet-like formula of the m -order linear recursive sequences is

$$S_n = \sum_{r=1}^m p_r q_r^n, \quad (n \geq 0) \quad (2)$$

where

$$p_1 = \frac{\begin{vmatrix} S_0 & 1 & \dots & 1 \\ S_1 & q_2 & \dots & q_m \\ \dots & \dots & \dots & \dots \\ S_{m-1} & q_2^{m-1} & \dots & q_m^{m-1} \end{vmatrix}}{\prod_{1 \leq j < i \leq m} (q_i - q_j)},$$

$$p_2 = \frac{\begin{vmatrix} 1 & S_0 & \dots & 1 \\ q_1 & S_1 & \dots & q_m \\ \dots & \dots & \dots & \dots \\ q_1^{m-1} & S_{m-1} & \dots & q_m^{m-1} \end{vmatrix}}{\prod_{1 \leq j < i \leq m} (q_i - q_j)},$$

...

$$p_m = \frac{\begin{vmatrix} 1 & 1 & \dots & S_0 \\ q_1 & q_2 & \dots & S_1 \\ \dots & \dots & \dots & \dots \\ q_1^{m-1} & q_2^{m-1} & \dots & S_{m-1} \end{vmatrix}}{\prod_{1 \leq j < i \leq m} (q_i - q_j)}$$

By choosing suitable initial conditions and coefficients we obtain the Binet-like formulas for the well-known sequences as follows:

If the terms of the sequence (1) take $m = 2$, $S_0 = 0, S_1 = 1$ and $a_0 = 1, a_1 = 1$, the Binet-like formula for the Fibonacci numbers will be denoted by

$$S_n = \frac{q_2^n - q_1^n}{q_2 - q_1}.$$

where q_1 and q_2 are the roots of the characteristic equation $x^2 - x - 1 = 0$ of the Fibonacci sequence $S_{n+2} = S_{n+1} + S_n$.

If the terms of the sequence (1) take $m = 2$, $S_0 = 2$, $S_1 = 1$ and $a_0 = 1$, $a_1 = 1$, the Binet-like formula for the Lucas numbers will be denoted by

$$S_n = q_2^n + q_1^n.$$

where q_1 and q_2 are the roots of the characteristic equation $x^2 - x - 1 = 0$ of the Lucas sequence $S_{n+2} = S_{n+1} + S_n$.

If the terms of the sequence (1) take $m = 2$, $S_0 = 0$, $S_1 = 1$ and $a_0 = 1$, $a_1 = 2$, the Binet-like formula for the Pell numbers will be denoted by

$$S_n = \frac{q_2^n - q_1^n}{q_2 - q_1}.$$

where q_1 and q_2 are the roots of the characteristic equation $x^2 - 2x - 1 = 0$ of the Pell sequence $S_{n+2} = 2S_{n+1} + S_n$.

If the terms of the sequence (1) take $m = 2$, $S_0 = 0$, $S_1 = 1$ and $a_0 = 2$, $a_1 = 1$, the Binet-like formula for the Jacobsthal numbers will be denoted by

$$S_n = \frac{q_2^n - q_1^n}{q_2 - q_1}.$$

where q_1 and q_2 are the roots of the characteristic equation $x^2 - x - 2 = 0$ of the Jacobsthal sequence $S_{n+2} = S_{n+1} + 2S_n$.

If the terms of the sequence (1) take $m = 2$, $S_0 = a$, $S_1 = b$ and $a_0 = -q$, $a_1 = p$, the Binet-like formula for the Horadam numbers will be denoted by

$$S_n = \frac{(aq_1 - b)q_2^n - (aq_2 - b)q_1^n}{q_2 - q_1}.$$

where q_1 and q_2 are the roots of the characteristic equation $x^2 - px + q = 0$ of the Horadam sequence $S_{n+2} = pS_{n+1} - qS_n$.

If the terms of the sequence (1) take $m = 2$, $S_0 = 1$, $S_1 = t$ and $a_0 = -1$, $a_1 = 2t$, the Binet-like formula for the Chebyshev polynomials will be denoted by

$$S_n = \frac{(t - q_1)q_2^n - (t - q_2)q_1^n}{q_2 - q_1}.$$

where q_1 and q_2 are the roots of the characteristic equation $x^2 - 2tx + 1 = 0$ of the Chebyshev polynomial sequence $S_{n+2} = 2tS_{n+1} - S_n$.

If the terms of the sequence (1) take $m = 3$, $S_0 = 1$, $S_1 = 1$, $S_2 = 1$ and $a_0 = 1$, $a_1 = 1$, $a_2 = 0$, the Binet-like formula for the Padovan numbers will be denoted by

$$S_n = p_1q_1^n + p_2q_2^n + p_3q_3^n,$$

where $p_1 = \frac{(q_2 - 1)(q_3 - 1)}{(q_1 - q_2)(q_1 - q_3)}$, $p_2 = \frac{(q_1 - 1)(q_3 - 1)}{(q_2 - q_1)(q_2 - q_3)}$, $p_3 = \frac{(q_1 - 1)(q_2 - 1)}{(q_3 - q_1)(q_3 - q_2)}$ and q_1, q_2, q_3 are the roots of the characteristic equation $x^3 - x - 1 = 0$ of the Padovan sequence $S_{n+3} = S_{n+1} + S_n$.

If the terms of the sequence (1) take $m = 3, S_0 = 3, S_1 = 0, S_2 = 2$ and $a_0 = 1, a_1 = 1, a_2 = 0$, the Binet-like formula for the Perrin numbers will be denoted by

$$S_n = q_1^n + q_2^n + q_3^n.$$

where q_1, q_2, q_3 are the roots of the characteristic equation $x^3 - x - 1 = 0$ of the Perrin sequence $S_{n+3} = S_{n+1} + S_n$.

If the terms of the sequence (1) take $m = 3, S_0 = 0, S_1 = 1, S_2 = 1$ and $a_0 = 1, a_1 = 1, a_2 = 1$, the Binet-like formula for the Tribonacci numbers will be denoted by

$$S_n = p_1 q_1^n + p_2 q_2^n + p_3 q_3^n,$$

where $p_1 = \frac{q_1^{n+2}}{(q_1 - q_2)(q_1 - q_3)}$, $p_2 = \frac{q_2^{n+2}}{(q_2 - q_1)(q_2 - q_3)}$, $p_3 = \frac{q_3^{n+2}}{(q_3 - q_1)(q_3 - q_2)}$ and q_1, q_2, q_3 are the roots of the characteristic equation $x^3 - x^2 - x - 1 = 0$ of the Tribonacci sequence $S_{n+3} = S_{n+2} + S_{n+1} + S_n$.

The Binet-like formulas and generating functions of some special sequences are available in the studies in [25, 27, 7, 20, 10, 15, 12, 23, 38, 17, 14, 31, 11, 9, 13, 33, 18, 8, 16, 30, 32, 6, 34, 36, 35, 39, 26, 19, 37, 21, 22, 28, 29, 24, 40]. Now we give generating function for the m -order linear recursive sequences.

In [43], the generating function of the m -order linear recursive sequences is

$$\sum_{n=0}^{\infty} S_n x^n = \frac{\sum_{i=0}^{m-1} S_i x^i \left(1 - \sum_{j=1}^{m-i-1} a_{m-j} x^j\right)}{1 - \sum_{k=0}^{m-1} a_k x^{m-k}}.$$

By choosing suitable initial conditions and coefficients we obtain the generating functions for the well-known sequences as follows:

m	S_0, S_1, \dots, S_{m-1}	a_0, a_1, \dots, a_{m-1}	Generating Functions	Names of sequence
2	$S_0 = 0, S_1 = 1$	$a_0 = 1, a_1 = 1$	$\frac{x}{1-x-x^2}$	Fibonacci
2	$S_0 = 2, S_1 = 1$	$a_0 = 1, a_1 = 1$	$\frac{2-x}{1-x-x^2}$	Lucas
2	$S_0 = 0, S_1 = 1$	$a_0 = 1, a_1 = 2$	$\frac{x}{1-2x-x^2}$	Pell
2	$S_0 = 0, S_1 = 1$	$a_0 = 2, a_1 = 1$	$\frac{1-x-2x^2}{1-2x-x^2}$	Jacobsthal
2	$S_0 = a, S_1 = b$	$a_0 = q, a_1 = p$	$\frac{a+(b-ap)x}{1-px-qx^2}$	Horadam
2	$S_0 = 1, S_1 = t$	$a_0 = -1, a_1 = 2t$	$\frac{1-tx}{1-2tx+x^2}$	Chebyshev polynomials
3	$S_0 = 1, S_1 = 1, S_2 = 1$	$a_0 = 1, a_1 = 1, a_2 = 0$	$\frac{x+1}{1-x^2-x^3}$	Padovan
3	$S_0 = 3, S_1 = 0, S_2 = 2$	$a_0 = 1, a_1 = 1, a_2 = 0$	$\frac{3-x^2}{1-x^2-x^3}$	Perrin
3	$S_0 = 0, S_1 = 1, S_2 = 1$	$a_0 = 1, a_1 = 1, a_2 = 1$	$\frac{x}{1-x-x^2-x^3}$	Tribonacci

Now we give exponential generating function for the m -order linear recursive sequences.

The exponential generating function of the m -order linear recursive sequences is

$$\sum_{n=0}^{\infty} S_n \frac{x^n}{n!} = \sum_{r=1}^m p_r e^{q_r x}.$$

By choosing suitable initial conditions and coefficients we obtain the generating functions for the well-known sequences as follows:

m	S_0, S_1, \dots, S_{m-1}	a_0, a_1, \dots, a_{m-1}	Exponential Generating Functions	Names of sequence
2	$S_0 = 0, S_1 = 1$	$a_0 = 1, a_1 = 1$	$\frac{e^{q_2 x} - e^{q_1 x}}{q_2 - q_1}$	Fibonacci
2	$S_0 = 2, S_1 = 1$	$a_0 = 1, a_1 = 1$	$\frac{e^{q_2 x} - e^{q_1 x}}{q_2 - q_1} + e^{q_1 x}$	Lucas
2	$S_0 = 0, S_1 = 1$	$a_0 = 1, a_1 = 2$	$\frac{e^{q_2 x} - e^{q_1 x}}{q_2 - q_1}$	Pell
2	$S_0 = 0, S_1 = 1$	$a_0 = 2, a_1 = 1$	$\frac{e^{q_2 x} - e^{q_1 x}}{q_2 - q_1}$	Jacobsthal
2	$S_0 = a, S_1 = b$	$a_0 = q, a_1 = p$	$\frac{(aq_1 - b)e^{q_2 x} - (aq_2 - b)e^{q_1 x}}{q_2 - q_1}$	Horadam
2	$S_0 = 1, S_1 = t$	$a_0 = -1, a_1 = 2t$	$\frac{(t - q_1)e^{q_2 x} - (t - q_2)e^{q_1 x}}{q_2 - q_1}$	Chebyshev polynomials
3	$S_0 = 1, S_1 = 1, S_2 = 1$	$a_0 = 1, a_1 = 1, a_2 = 0$	$\frac{e^{q_1 x} + e^{q_2 x} + e^{q_3 x}}{q_1 + q_2 + q_3}$	Padovan
3	$S_0 = 3, S_1 = 0, S_2 = 2$	$a_0 = 1, a_1 = 1, a_2 = 0$	$\frac{3e^{q_1 x} + e^{q_2 x} + e^{q_3 x}}{q_1 + q_2 + q_3}$	Perrin
3	$S_0 = 0, S_1 = 1, S_2 = 1$	$a_0 = 1, a_1 = 1, a_2 = 1$	$\frac{e^{q_1 x} + e^{q_2 x} + e^{q_3 x}}{q_1 + q_2 + q_3}$	Tribonacci

1.1. m -Order Linear Recursive Quaternions

A quaternion is defined by

$$q = a_0 e_0 + a_1 e_1 + a_2 e_2 + a_3 e_3$$

where a_0, a_1, a_2 and a_3 are real numbers and $e_0 = 1, e_1 = i, e_2 = j$ and $e_3 = k$ are the standart basis in \mathbb{R}^4 .

The quaternion multiplication is defined using the rules:

$$e_0^2 = 1, \quad e_1^2 = e_2^2 = e_3^2 = -1$$

$$e_1e_2 = -e_2e_1 = e_3, \quad e_2e_3 = -e_3e_2 = e_1 \quad \text{and} \quad e_3e_1 = -e_1e_3 = e_2.$$

This algebra is associative and non-commutative.

Let $q = a_0e_0 + a_1e_1 + a_2e_2 + a_3e_3$ and $p = b_0e_0 + b_1e_1 + b_2e_2 + b_3e_3$ be any two quaternions. Then the addition and subtraction of them is

$$q \mp p = (a_0 \mp b_0)e_0 + (a_1 \mp b_1)e_1 + (a_2 \mp b_2)e_2 + (a_3 \mp b_3)e_3$$

and for $k \in \mathbb{R}$, the multiplication by scalar is

$$kq = ka_0e_0 + ka_1e_1 + ka_2e_2 + ka_3e_3$$

and the conjugate and norm of a quaterion are

$$\bar{q} = a_0e_0 - a_1e_1 - a_2e_2 - a_3e_3$$

and

$$N(q) = q\bar{q} = a_0^2 + a_1^2 + a_2^2 + a_3^2.$$

Addition, equality and multiplication by scalar of two quaternions can be found [1, 2, 5].

Definition 1.1. The m -order linear recursive quaternion $\{QS_n\}_{n \geq 0}$ is defined by

$$QS_n = S_n e_0 + S_{n+1} e_1 + S_{n+2} e_2 + S_{n+3} e_3 \quad (3)$$

where S_n is the m -order linear recursive numbers.

Theorem 1.2. The Binet-like formula for the m -order linear recursive quaternion $\{QS_n\}_{n \geq 0}$ is

$$QS_n = \sum_{r=1}^m p_r \hat{q}_r q_r^n \quad (4)$$

where $\hat{q}_r = e_0 + q_r e_1 + q_r^2 e_2 + q_r^3 e_3$.

Proof. From the definition of the m -order linear recursive quaternion QS_n in (3) and Binet-like formula for the m -order linear recursive number S_n , we write

$$\begin{aligned} QS_n &= S_n e_0 + S_{n+1} e_1 + S_{n+2} e_2 + S_{n+3} e_3 \\ &= \sum_{r=1}^m p_r q_r^n e_0 + \sum_{r=1}^m p_r q_r^{n+1} e_1 + \sum_{r=1}^m p_r q_r^{n+2} e_2 + \sum_{r=1}^m p_r q_r^{n+3} e_3 \\ &= \sum_{r=1}^m p_r (e_0 + q_r^1 e_1 + q_r^2 e_2 + q_r^3 e_3) q_r^n \\ &= \sum_{r=1}^m p_r \hat{q}_r q_r^n \end{aligned}$$

□

As a special case of the equality (4), the Binet-like formula of Fibonacci quaternions can be given as follows:

For $m = 2$, the Binet-like formula for the Fibonacci quaternions will be denoted by

$$\mathcal{Q}S_n = \frac{\hat{q}_2 q_2^n - \hat{q}_1 q_1^n}{q_2 - q_1}.$$

where q_1 and q_2 are the roots of the characteristic equation $x^2 - x - 1 = 0$ of the Fibonacci sequence $S_{n+2} = S_{n+1} + S_n$, and $\hat{q}_1 = e_0 + q_1^1 e_1 + q_1^2 e_2 + q_1^3 e_3$, $\hat{q}_2 = e_0 + q_2^1 e_1 + q_2^2 e_2 + q_2^3 e_3$. Binet-like formulas of other special quaternion sequences can be obtained in a similar way using (4).

Theorem 1.3. *The generating function for m -order linear recursive quaternion $\{\mathcal{Q}S_n\}_{n \geq 0}$ is*

$$G_{\mathcal{Q}S}(x) = \frac{(e_0 x^3 + e_1 x^2 + e_2 x + e_3) G_S(x) - (S_0(e_1 x^2 + e_2 x + e_3) + S_1(e_2 x^2 + e_3 x) + S_2(e_3 x^2))}{x^3}$$

where $G_S(x)$ is the generating function of the m -order linear recursive sequences

Proof. Let

$$G_{\mathcal{Q}S}(x) = \sum_{n=0}^{\infty} \mathcal{Q}S_n x^n \quad (5)$$

be generating function of the m -order linear recursive quaternion. We have

$$\begin{aligned} G_{\mathcal{Q}S}(x) &= \sum_{n=0}^{\infty} (S_n e_0 + S_{n+1} e_1 + S_{n+2} e_2 + S_{n+3} e_3) x^n \\ &= e_0 \sum_{n=0}^{\infty} S_n x^n + e_1 \sum_{n=0}^{\infty} S_{n+1} x^n + e_2 \sum_{n=0}^{\infty} S_{n+2} x^n + e_3 \sum_{n=0}^{\infty} S_{n+3} x^n \\ &= e_0 G_S(x) + e_1 \left(G_S(x) \frac{1}{x} - \frac{S_0}{x} \right) + e_2 \left(G_S(x) \frac{1}{x^2} - \frac{S_0}{x^2} - \frac{S_1}{x} \right) \\ &\quad + e_3 \left(G_S(x) \frac{1}{x^3} - \frac{S_0}{x^3} - \frac{S_1}{x^2} - \frac{S_2}{x} \right) \\ &= \frac{(e_0 x^3 + e_1 x^2 + e_2 x + e_3) G_S(x) - (S_0(e_1 x^2 + e_2 x + e_3) + S_1(e_2 x^2 + e_3 x) + S_2(e_3 x^2))}{x^3} \end{aligned}$$

□

As a special case of the equality (5), the Binet-like formula of Fibonacci quaternions can be given as follows:

For $m = 2$, the generating function for the Fibonacci quaternions will be denoted by

$$G_{\mathcal{Q}S}(x) = \frac{x + e_1 + e_2(x + 1) + e_3(x + 2)}{1 - x - x^2}.$$

Generating functions of other special quaternion sequences can be obtained in a similar way using (5).

Theorem 1.4. *The exponential generating function of the m -order linear recursive quaternions is*

$$E_S(x) = \sum_{r=1}^m p_r \hat{q}_r e^{q_r x} \quad (6)$$

Proof. Let

$$E_S(x) = \sum_{n=0}^{\infty} Q S_n \frac{x^n}{n!}$$

Using the identity (2), we get

$$E_S(x) = \sum_{n=0}^{\infty} Q S_n \frac{x^n}{n!} = \sum_{n=0}^{\infty} \sum_{r=1}^m p_r \hat{q}_r q_r^n \frac{x^n}{n!} = \sum_{r=1}^m p_r \hat{q}_r \sum_{n=0}^{\infty} \frac{(q_r x)^n}{n!} = \sum_{r=1}^m p_r \hat{q}_r e^{q_r x}$$

□

For $m = 2$, the exponential generating function for the Fibonacci quaternions will be denoted by

$$E_S(x) = \frac{\hat{q}_2 e^{q_2 x} - \hat{q}_1 e^{q_1 x}}{q_2 - q_1}$$

Exponential generating functions of other special quaternion sequences can be obtained in a similar way using (6).

References

- [1] F. Matyas, *Sequence transformations and linear recurrences of higher order*. Acta Mathematica et Informatica Universitatis Ostraviensis, 2001, v. 9, no 1, 45-51.
- [2] F. Gatta, A. D'amico, *Sequences H_n for which H_{n+1}/H_n Approaches the Golden Ratio*, 2008.
- [3] T. Komatsu, *Sequences H_n for which H_{n+1}/H_n approaches an irrational number*, Fibonacci Quaterly, 2010, v. 48, no 3, 265-275.
- [4] T. Szakács, *K -order Linear Recursive Sequences and the Golden Ratio*, Fibonacci Quarterly, 2017, v. 55, 186-191.
- [5] F. Matyas, *Linear recurrences and rootfinding methods*. Acta Academiae Paedagogicae Agriensis, Sectio Mathematicae, 2001, v. 28, 27-34.
- [6] G. Lee, M. Asci, *Some properties of the (p, q) -Fibonacci and (p, q) -Lucas polynomials*, Journal of applied mathematics, 2012.
- [7] A. Suvarnamani, M. Tatong, *Some properties of (p, q) -Fibonacci numbers*. Progress in Applied Science and Technology, 2015, v. 5, no 2, 17-21.

- [8] A. Ipek, *On (p, q) -Fibonacci quaternions and their Binet formulas, generating functions and certain binomial sums*. Advances in Applied Clifford Algebras, 2015, v. 27, no 2, 1343-1351.
- [9] S. Falcon, A. Plaza, *The k -Fibonacci sequence and the Pascal 2-triangle*. Chaos, Solitons & Fractals, 2007, v. 33, no 1, 38-49.
- [10] C. Bolat, H. Köse, *On the properties of k -Fibonacci numbers*, Int. J. Contemp. Math. Sciences, 2010, v. 5, no 22, 1097-1105.
- [11] M. El-Mikkawy, T. Sogabe, *A new family of k -Fibonacci numbers*. Applied Mathematics and Computation, 2010, v. 215, no 12, 4456-4461.
- [12] P. Catarino, *On some identities for k -Fibonacci sequence*. Int. J. Contemp. Math. Sci, 2014, v. 9, no 1, 37-42.
- [13] A. D. Godase, M. B. Dhakne, *On the properties of k -Fibonacci and k -Lucas numbers*. International Journal of Advances in Applied Mathematics and Mechanics, 2014, v. 2, no 1, 100-106.
- [14] O. Deveci, E. Karaduman, *The Pell sequences in finite groups*. Util. Math, 2015, v. 96, 263-276.
- [15] J. J. Bravo, J. L. Herrera, F. Luca, *On a generalization of the Pell sequence*. Mathematica Bohemica, 2021, v. 146, no 2, 199-213.
- [16] S. H. Jafari-Petroudia, B. Pirouzb, *On some properties of (k, h) -Pell sequence and (k, h) -Pell-Lucas sequence*. Int. J. Adv. Appl. Math. and Mech, 2015, v. 3, no 1, 98-101.
- [17] A. Dasdemir, *On the Pell, Pell-Lucas and modified Pell numbers by matrix method*. Applied Mathematical Sciences, 2011, v. 5, no 64, 3173-3181.
- [18] A. F. Horadam, *Jacobsthal representation numbers*. significance, 1996, v. 2, 2-8.
- [19] M. Tastan, E. Özkan, *Catalan transform of the k -jacobsthal sequence*. Electronic Journal of Mathematical Analysis and Applications, 2020, v. 8, no 2, 70-74.
- [20] F. T. Aydin, *On generalizations of the Jacobsthal sequence*. Notes on number theory and discrete mathematics, 2018, v. 24, no 1, 120-135.
- [21] S. Uygun, *The (s, t) -Jacobsthal and (s, t) -Jacobsthal Lucas sequences*. Applied Mathematical Sciences, 2015, v. 70, no 9, 3467-3476.
- [22] S. Uygun, H. Eldogan, *Properties of k -Jacobsthal and k -Jacobsthal Lucas sequences*. General Mathematics Notes, 2016, v. 36, no 1, 34pp.
- [23] S. Celik, İ. Durukan, E. Özkan, *New recurrences on Pell numbers, Pell-Lucas numbers, Jacobsthal numbers, and Jacobsthal-Lucas numbers*. Chaos, Solitons & Fractals, 2021, 150, 111173pp.

- [24] N. Yilmaz, N. Taskara, *Matrix sequences in terms of Padovan and Perrin numbers*. Journal of Applied Mathematics, 2013.
- [25] A. Faisant, A. *On the Padovan sequence*. 2013, arXiv preprint arXiv:1905.07702.
- [26] D. Tasci, *Gaussian padovan and gaussian pell-padovan sequences*. Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics, 2018, v. 67, no 2, 82-88.
- [27] R. Sivaraman, *Properties of Padovan Sequence*. Turkish Journal of Computer and Mathematics Education, 2021, v. 12, no 2, 3098-3101.
- [28] R. P. M. Vieira, F. R. V. Alves, P. M. M. C. Catarino, *A Historical Analysis of The Padovan Sequence*. International Journal of Trends in Mathematics Education Research, 2020, v. 3, no 1, 8-12.
- [29] E. W. Weisstein, *Perrin sequence*. 2021, <https://mathworld.wolfram.com/>.
- [30] K. Khompungson, B. Rodjanadid, S. Sompong, *Some matrices in term of Perrin and Padovan sequences*. Thai Journal of Mathematics, 2019, v. 17, no 3, 767-774.
- [31] O. Diskaya, H. Menken, *On the Split (s, t) -Padovan and (s, t) -Perrin Quaternions*. International Journal of Applied Mathematics and Informatics, 2019, v. 13, 25-28.
- [32] P. J. Larcombe, O. D. Bagdasar, E. J. Fennessey, *Horadam sequences: a survey*. Bulletin of the Institute of Combinatorics and its, 2017.
- [33] P. Haukkanen, *A note on Horadam's sequence*. Fibonacci Quarterly, 2002, v. 40, no 4, 358-361.
- [34] J. C. Mason, D. C. Handscomb, *Chebyshev polynomials*. Chapman and Hall/CRC, 2002.
- [35] T. J. Rivlin, *Chebyshev polynomials*. Courier Dover Publications, 2020.
- [36] H. Merzouk, A. Boussayoud, M. Chelgham, *Generating Functions of Generalized Tribonacci and Tricobsthal Polynomials*. Montes Taurus Journal of Pure and Applied Mathematics, 2020, v. 2, no 2, 7-37.
- [37] Y. Taşyurdu, *Tribonacci and Tribonacci-Lucas hybrid numbers*. International Journal of Contemporary Mathematical Sciences, 2019, v. 14, no 4, 245-254.
- [38] J. L. Cereceda, *Binet's formula for generalized tribonacci numbers*. International journal of mathematical education in science and technology, 2015, v. 46, no 8, 1235-1243.
- [39] Y. Soykan, *Tribonacci and tribonacci-Lucas sedenions*. Mathematics, 2019, v. 7, no 1, 74.

- [40] N. Yilmaz, N. Taskara, *Incomplete Tribonacci–Lucas Numbers and Polynomials*. Advances in Applied Clifford Algebras, 2015, v. 25, no 3, 741-753.
- [41] L. M. Milne-Thomson, *The calculus of finite differences*. American Mathematical Soc., 2000.
- [42] T. N. Shorey, R. Tijdeman, *Exponential diophantine equations*, 1986.
- [43] E. Kiliç, P. Stanica, *A matrix approach for general higher order linear recurrences*. Bull. Malays. Math. Sci. Soc., 2011, v. 34, no 1, 51-67.

Orhan Dişkaya
Mersin University, Mersin, Turkey
E-mail: orhandiskaya@mersin.edu.tr

Hamza Menken
Mersin University, Mersin, Turkey
E-mail: orhandiskaya@mersin.edu.tr

Received 30 July 20..

Accepted 18 January 20..