

Effects of selenium supplementation on pregnancy outcome and disease progression in HIV-infected pregnant women in Lagos, Nigeria

Study protocol for a randomised, double-blind, placebo-controlled trial

Kehinde S. Okunade, FWACS, FMCOG^{a,d,*}, Sarah John-Olabode, FMCPATH^b, Oluwatosin J. Akinsola, PhD^c, Opeyemi Akinajo, FWACS^d, Sulaimon A. Akanmu, FMCPATH^b, Phyllis J. Kanki, DVM, DSc^e

Abstract

Background: Micronutrient deficiencies are common during pregnancy, especially in pregnant women from economically disadvantaged settings where diets with low content of minerals and vitamins are consumed. Selenium is a non-metallic chemical element of great importance to human health. This study will assess the effect of selenium supplementation on major pregnancy outcomes and disease progression among HIV-infected pregnant women in Lagos, Nigeria.

Methods: A randomized, double-blind, placebo-controlled trial involving confirmed HIV-positive pregnant women at the Lagos University Teaching Hospital (LUTH) between September 2018 and February 2019. Eligible participants are HIV-infected pregnant women aged 15 to 49 years and have a singleton gestation at 14 to 27 weeks' gestation. At enrolment, 90 women will be randomly assigned into each intervention arm to receive either a daily tablet of 200 µg elemental selenium or placebo. Relevant participants' data will be collected at enrolment and at delivery. Statistical analyses will be carried out using SPSS version 23.0 for Windows. The associations between any 2 groups of continuous variables will be tested using the *t* test or the Mann-Whitney *U* test and that of 2 groups of categorical variables with chi-square or Fishers exact test where appropriate. A series of multivariable analyses will also be carried out to identify and control for several possible confounders of the major pregnancy outcomes and HIV disease progression. Statistical significance will be defined as $P < .05$. Ethical approval for the study was obtained from the LUTH's Health Research and Ethics Committee (Approval number: ADM/DCST/HREC/APP/2438; 30th August 2018).

Discussion: This trial will assess the effect of selenium supplementation on pregnancy outcome and HIV disease progression among HIV-infected pregnant women in Lagos. This will help to determine if routine selenium supplementation in HIV-infected pregnant women will contribute to the improvement in the major adverse pregnancy outcomes such as preterm birth and low birth weight and the HIV disease surrogate markers such as CD4+ cells count and viral load.

Trial registration: PACTR, PACTR201809756724274. Registered on 3rd September 2018, <https://pactr.samrc.ac.za/TrialDisplay.aspx?TrialID=3571>.

Ethical approval for this proposed trial was obtained the Health Research and Ethics Committee of the Lagos University Teaching Hospital on September 30th, 2018 with approval number ADM/DCST/HREC/APP/2438. Ethics principles of the Helsinki Declaration will be applied throughout the course of the study. All participants will be counselled prior to enrolment and read and signed an informed consent form. The investigators will ensure strict confidentiality of participant information.

The authors certify that they will obtain all appropriate patient consent forms. In the form, the patients will give their consent for their clinical information to be reported in the journal. The patients understand that their names and initials will not be published, and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

The datasets that will be generated or analysed during the current study will be available from the corresponding author upon reasonable request.

The work reported in this publication was supported by the Fogarty International Center and National Institute of Mental Health, of the National Institutes of Health under Award Number D43TW010543. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Supplemental Digital Content is available for this article.

The authors declare that they have no competing interests.

^a Department of Obstetrics & Gynaecology, ^b Department of Haematology & Blood Transfusion, ^c Epidemiology and Biostatistics unit, Department of Community Health & Primary Health Care, College of Medicine, University of Lagos, ^d Department of Obstetrics & Gynaecology, Lagos University Teaching Hospital, Nigeria, ^e Department of Immunology & Infectious Diseases, Harvard School of Public Health, USA.

* Correspondence: Kehinde S. Okunade, Department of Obstetrics and Gynaecology, College of Medicine, University of Lagos, Lagos, Nigeria (e-mail: sokunade@unilag.edu.ng).

Copyright © 2019 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Medicine (2019) 98:3(e12735)

Received: 14 September 2018 / Accepted: 17 September 2018

<http://dx.doi.org/10.1097/MD.00000000000012735>

Abbreviations: AE = adverse events, ART = antiretroviral treatment, CRF = case report form, GPX = glutathione peroxidase, HIV = human immune deficiency virus, LUTH = Lagos university teaching hospital, PI = principal investigator, SP = sulfadoxine-pyrimethamine, SPSS = statistical package for social science, WHO = World Health Organization.

Keywords: CD4+ cells count, lagos, low birth weight, LUTH, preterm birth, selenium, viral load

1. Introduction

Micronutrient deficiencies are common during pregnancy, especially in pregnant women from economically disadvantaged settings where diets with low content of minerals and vitamins are consumed.^[1] Selenium is a non-metallic chemical element of great importance to human health.^[2,3] It is a key component of several human seleno-proteins and mostly involved in redox reactions.^[4] It acts as an antioxidant and helps protect the body against the damaging effects of free radicals.^[5] Selenium is essential for the activity of the enzyme glutathione peroxidase (GPX), a main intracellular antioxidant that protects against reactive oxygen species and subsequent oxidation-induced cellular damage.^[5,6] Biochemical selenium deficiency has been associated with increased mortality among those infected with HIV^[7] and with accelerated HIV disease progression through increased viral load.^[8] Selenium's role in antioxidant defence^[9] and immunity^[10] may be the underlying mechanism. There is, however, limited data on the selenium status of Nigerian pregnant women with HIV-infection, with a recent study conducted by us showing a prevalence rate of selenium deficiency to be 20.4% and we reported significantly increased rates of preterm birth and delivery low birth weight babies among the HIV infected participants with selenium deficiency.^[11] Several other studies have also reported low selenium status in HIV-infected individuals, and that serum selenium concentration declines with disease progression.^[12,13]

Selenium is found in cereals, meat, poultry, seafood, and eggs.^[6] Cereals may provide about 50% of dietary selenium, however data on the selenium content of Nigerian or African foods is also limited. The selenium content of plant foods varies with the selenium content of the soil.^[6] The Recommended Dietary Intake of selenium for pregnant women is 60 mg/day^[6] and most interventional studies make use of daily doses of up to 200 µg selenium in its elemental form (selenomethionine).^[14-17] Supplemental doses up to 200 µg of the mineral should not cause adverse effects, even if temporarily exceeded, since the selenium upper tolerable intake (UL) for this group is established as 400 µg (5.1 µmol) selenium/day.^[18] Given the high UL and the regulation of body homeostasis through urinary excretion, selenium supplementation can be regarded as a safe intervention.^[18,19] There is still limited data about selenium toxicity in humans, but the most common side effects include: hair and nail brittleness and loss, gastrointestinal disturbances, skin rash, fatigue, irritability and nervous system disturbances.^[6,19] Selenium containing supplements are not recommended for children under the age of 15 years.^[6] There is currently paucity of studies that examined the effects of selenium supplementation among pregnant HIV-infected black African women. This study will therefore attempt to assess the effect of antenatal selenium supplementation on major pregnancy outcomes (preterm birth and low birth weight) and HIV disease progression rate among pregnant women at the Lagos University Teaching Hospital (LUTH), Lagos, Nigeria.

2. Methods and design

2.1. Study design and setting

The study is designed as a prospective, randomized, double-blind, placebo-controlled, single-center trial involving confirmed HIV-seropositive pregnant women who are seeking care at the antenatal clinics of the Lagos University Teaching Hospital (LUTH). LUTH is the teaching hospital of the College of Medicine, University of Lagos. It has more than 1000 beds and is located in the metropolis of Lagos in Southwest Nigeria. The hospital provides services to patients from the neighbouring states in Southwest Nigeria. It is the largest hospital in Lagos State and mainly offers clinical services, including prenatal, intra-partum, and postnatal care. Pregnant women with HIV infections are jointly managed by haematologists and HIV physicians at the hospital's AIDS Prevention Initiative Nigeria (APIN) clinic and undergo regular monitoring of their disease with 3-monthly CD4 + cells count and viral load. Participants' enrolment will start in September 2018, and the last woman is expected to be included in the trial in February 2019. The Health Research and Ethics Committee of LUTH's Institutional Review Board has granted ethics approval for this study (Approval Number ADM/DCST/HREC/APP/2438; 30th August 2018).

2.2. Study population/ participant enrolment

Eligible participants are confirmed HIV-seropositive pregnant women aged 15 to 49 years who have a singleton gestation at 14 to 27 weeks' gestation. The exclusion criteria at enrolment are multiple gestations, significant renal and hepatic impairment, an expected delivery date beyond March 2019, receipt of a long course of mineral supplement containing selenium during the 6 months prior to enrolment, refusal of consent at enrolment or withdrawal of consent during the study.

2.3. Randomisation and blinding schedule

Eligible women will be invited to give consent for participation in the trial within 2 weeks of screening. At enrolment, participants will be randomly assigned by a trained research assistant into 2 arms to receive daily oral tablet of 200 µg elemental selenium (arm I) or placebo (arm II) till their delivery using a 1:1 block randomisation code to be generated from Random Allocation software version 1.0 (May 2004) by the study Statistician. The proposed dose of selenium is one-half of the safe and tolerable upper intake level of 400 µg/day for pregnant women.^[18] The active drug (selenium) and placebo tablets will be indistinguishable in shape, size, and colour and will be packed in identical coded transparent dispensing sachets by the onsite pharmacy technician who will then store the coded randomization list in a sealed opaque envelope that will be kept in a locked file cabinet at the study site until the end of the study. Twelve (12) dispensing sachets with identical numeric regimen code, containing the same intervention drug will be prepared for each of the participants. At

every 2-weekly follow-up visit, a new dispensing sachet with the same numeric code, containing 16 tablets of the same intervention drugs will be given to each woman and pills remaining in used sachets will be counted to assess compliance. In addition, all the women will receive the standard prenatal care (see below). Study investigators and participants will be blinded to the treatment arms until after the final analysis. The onsite pharmacy technician is not involved in the design, conduct, statistical analysis, and reporting of study findings.

2.4. Standard prenatal care

All the women will receive their routine prenatal care which includes daily doses of ferrous gluconate (300mg, equivalent to 60 mg of elemental iron) and folic acid (5 mg), and 4-weekly doses of malarial chemoprophylaxis in the form of Sulfadoxine-Pyrimethamine (SP) tablets (from enrolment to 36 weeks of gestation) or instead daily doses of Cotrimoxazole 980mg for women with CD4+ cell counts <200 cells/mm³ according to established protocol for *Pneumocystis carinii* prophylaxis^[20] irrespective of treatment assignments. Alternatively, daily proguanil tablet (100mg) is given to women with known or recent allergy to Sulphonamides from enrolment to delivery. All women will also be commenced on ART at some point during their follow-up according to the National HIV prevention and treatment guidelines.^[21]

2.5. General procedure and monitoring

2.5.1. Data collection and management. At enrolment, the investigators will collect data on the participants' socioeconomic status, education, and obstetric history, duration of HIV diagnosis (in months), WHO clinical stage of the disease,^[22] type and duration of ART use (in months), baseline Haemoglobin (Hb) levels (in g/dL), CD4+ cells count (in cells/mm³) and viral load (in copies/mL), and BMI (actual maternal weight is calculated by subtracting the estimated fetal weight at enrolment using a normogram chart from the measured maternal weight) (in kg/m²) using a structured interviewer-administered questionnaire. Gestational age is based on the date of the participant's last menstrual period, which will be obtained at the time of random assignment. Data collection begins on the day a participant is allocated into an intervention arm and continues until the termination of the trial or until the participant withdraws from the trial at any time for any reason. At delivery, data will be collected from the participants' case notes on the latest CD4+ cells count and viral load (done within the last 6 weeks of delivery), WHO stage of the disease, and latest Hb levels. The total duration of supplement intake – selenium or placebo (in weeks), level of compliance to use of the supplement (good compliance level is defined as usage of at least 10 of the 16, 2-weekly prescribed tablets), mode of delivery, gestational age at delivery (in weeks) and the baby's birth weight (in grams) will also be recorded in the questionnaire. Original data will be




Figure 1. Trial flow chart.

to guarantee consistent practice. The training will include understanding of inclusion/exclusion/exit criteria, follow-up procedures, and completion of questionnaire and CRF. The trial will be monitored by quality assurance personnel from the research management office of the College of Medicine, University of Lagos, who will be independent from the study team, and an independent steering committee. There will be a periodic monitoring to guarantee accuracy and quality throughout the study period. The essential documents (consent information, enrolment, protocol deviations, number and proportion of missed visits, and losses to follow-up) will be monitored and checked for accuracy and completeness by the monitors. The principal investigator (PI) is responsible for the overall project and for organizing steering committee meetings. An independent steering committee will be responsible for ensuring the overall safety of participants, coordinating study meetings, supervising the study, monitoring data safety, and overseeing quality control.

2.11. Ethics and dissemination

The study protocol was approved by the Health Research and Ethics Committee of the Lagos University Teaching Hospital on September 30th, 2018 with approval number ADM/DCST/HREC/APP/2438 [Additional file 1, <http://links.lww.com/MD/C761>]. Participants will provide written informed consent [Additional file 2, <http://links.lww.com/MD/C761>] prior to participation in the trial and the participant will keep a signed copy of the consent form for future references. Participants below the age of 18 years will be regarded as emancipated minors. ‘Emancipated minor’ is a person that is not of legal age to give consent (below 18 years of age in Nigeria) for a research study but who by marriage, pregnancy, being the mother of a child whether married or not or has left home and is self-sufficient can be allowed to do so legally. The trial will be reported in line with the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) checklist [Additional file 3, <http://links.lww.com/MD/C761>] and the results will be disseminated in peer review scientific journals.

3. Discussion

Selenium is a non-metallic chemical element of great important to human health. Biochemical selenium deficiency has been associated with increased mortality among those infected with HIV and with accelerated HIV disease progression. This study aims to assess the effects of giving routine antenatal selenium supplementation to HIV-infected pregnant women on their pregnancy outcome and HIV-disease progression. If it is found to be significantly effective and safe, this trial will provide a basis for prescribing this antioxidant to HIV-seropositive women in pregnancy.

3.1. Trial status

At the time of initial manuscript submission, recruitment is yet to commence into the trial. Participants’ enrolment will start in September 2018, and the last woman is expected to be included in the trial in February 2019. The manuscript reports protocol version 2.0 (30th June 2018).

Acknowledgments

The authors would like to thank the HBNU Global Health Fellows Consortium for providing the funding for this work. We

also appreciate the efforts of Prof. Folasade Ogunsola (Deputy Vice-Chancellor, Development Services of the University of Lagos, Lagos, Nigeria) in making the application for the HBNU Fellowship a success through her very valuable criticisms.

Author contributions

KSO, SAA, and PJK participated in the conception, design, and writing of the study protocol. SJ-O contributed to the revision and editing of the study protocol. KSO, SJ-O, OJA and OA will be involved in the recruitment of participants and the acquisition of data. KS wrote the first draft of this manuscript. SJ-O, OJA, OA, SA, PK were all involved in the revision of the manuscript. All authors approved the final version of the manuscript to be submitted.

Conceptualization: Kehinde Sharafadeen Okunade, Sulaimon A Akanmu, Phyllis J Kanki.

Formal analysis: Oluwatosin J Akinsola.

Funding acquisition: Kehinde Sharafadeen Okunade, Sulaimon A Akanmu, Phyllis J Kanki.

Methodology: Kehinde Sharafadeen Okunade, Sarah John-Olabode, Oluwatosin J Akinsola, Opeyemi Akinajo, Sulaimon A Akanmu, Phyllis J Kanki.

Project administration: Kehinde Sharafadeen Okunade.

Resources: Kehinde Sharafadeen Okunade.

Software: Kehinde Sharafadeen Okunade, Oluwatosin J Akinsola.

Supervision: Kehinde Sharafadeen Okunade, Sarah John-Olabode, Opeyemi Akinajo, Sulaimon A Akanmu.

Validation: Kehinde Sharafadeen Okunade.

Visualization: Kehinde Sharafadeen Okunade.

Writing – original draft: Kehinde Sharafadeen Okunade, Sarah John-Olabode, Opeyemi Akinajo.

Writing – review & editing: Kehinde Sharafadeen Okunade, Sarah John-Olabode, Oluwatosin J Akinsola, Opeyemi Akinajo, Sulaimon A Akanmu, Phyllis J Kanki.

Kehinde Sharafadeen Okunade orcid: 0000-0002-0957-7389.

References

- [1] World Health Organisation. Zinc. In: Trace elements in human nutrition and health. WHO, Geneva, Switzerland;1996:72–104.
- [2] Fawzi WW, Msamanga GI, Spiegelman D, et al. A randomized trial of multivitamin supplements and HIV disease progression and mortality. *N Engl J Med* 2004;351:23–32.
- [3] Mehta S, Fawzi WW. Micronutrient supplementation as adjunct treatment for HIV-infected patients. *Clin Infect Dis* 2010;50:1661–3.
- [4] Bellinger FP, Raman AV, Reeves MA, et al. Regulation and function of selenoproteins in human disease. *Biochem J* 2009;422:11–22.
- [5] Rayman MP. The importance of selenium to human health. *Lancet* 2000;356:233–41.
- [6] National health medical research council nutrient reference values for australia and new zealand including recommended dietary intakes. Canberra, Australia. Commonwealth Department of Health and Ageing 2006;202–8.
- [7] Kupka R, Msamanga GI, Spiegelman D, et al. Selenium status is associated with accelerated HIV disease progression among HIV-1-infected pregnant women in Tanzania. *J Nutr* 2004;134:2556–60.
- [8] Hurwitz BE, Klaus JR, Llabre MM, et al. Suppression of human immunodeficiency virus type 1 viral load with selenium supplementation: a randomized controlled trial. *Arch Intern Med* 2007;167:148–54.
- [9] Flohe L. Glutathione peroxidase. *Basic Life Sci* 1988;49:663–8.
- [10] Spallholz JE, Boylan LM, Larsen HS. Advances in understanding selenium’s role in the immune system. *Ann N Y Acad Sci* 1990;587:123–39.
- [11] Okunade KS, Olowoselu OF, Osanyin GE, et al. Selenium deficiency and pregnancy outcome in pregnant women with HIV in Lagos, Nigeria *Int J Gynaecol Obstet* 2018;142:207–13.

- [12] Dworkin BM, Rosenthal WS, Wormser GP, et al. Abnormalities of blood selenium and glutathione peroxidase activity in patients with acquired immunodeficiency syndrome and aids-related complex. *Biol Trace Elem Res* 1988;15:167–77.
- [13] Beck KW, Schramel P, Hedl A, et al. Serum trace element levels in HIV-infected subjects. *Biol Trace Elem Res* 1990;25:89–96.
- [14] Kupka R, Mugusi F, Aboud S, et al. Randomized, double-blind, placebo-controlled trial of selenium supplements among HIV infected pregnant women in Tanzania: effects on maternal and child outcomes. *Am J Clin Nutr* 2008;87:1802–8.
- [15] Kupka K, Mugusi F, Aboud S, et al. Effect of selenium supplements on hemoglobin concentration and morbidity among HIV-1-infected Tanzanian women. *Clin Infect Dis* 2009;48:1475–8.
- [16] Moore MA, Wander RC, Xia YM, et al. Selenium supplementation of Chinese women with habitually low selenium intake increases plasma selenium, plasma glutathione peroxidase activity, and milk selenium, but not milk glutathione peroxidase activity. *J Nutr Biochem* 2000;11:341–7.
- [17] Negro R, Greco G, Mangieri T, et al. The influence of selenium supplementation on postpartum thyroid women with thyroid peroxidase autoantibodies. *J Clin Endocrinol Metab* 2007;92:1263–8.
- [18] Institute of Medicine, Food and Nutrition Board. Dietary Reference Intakes: Vitamin C, Vitamin E, Selenium, and Carotenoids. National Academy Press, Washington, DC. 2000.
- [19] Bergamaschi DP, Mariath AB, Abbade JF, et al. Selenium supplementation during pregnancy for improving maternal and newborn outcomes. *Cochrane Datab Syst Rev* 2012;3: Art. No.: CD009673.
- [20] Connelly RT, Lourwood DL. *Pneumocystis carinii* pneumonia prophylaxis during pregnancy. *Pharmacotherapy* 1994;14:424–9.
- [21] Federal Ministry of Health (Nigeria). Integrated national guidelines for HIV prevention, treatment, and care: Chapters 5 and 6. *J HIV Hum Reprod*. 2015; 3: 47–55.
- [22] Interim WHO Clinical Staging of HIV/AIDS and HIV/AIDS Case Definitions for Surveillance (African Region). WHO 2005, Geneva, Switzerland: 20–28.