



## King's Research Portal

*Document Version*  
Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Knowles, G., Gayer-Anderson, C., Beards, S., Blakey, R., Davis, S., Lewis, K., Stanyon, D., Ofori, A., Turner, A., Pinfold, V., Bakolis, I., Reininghaus, U., Harding, S., & Morgan, C. (Accepted/In press). Mental Distress among Young People in Inner-Cities: the Resilience, Ethnicity and AdolesCent Mental Health (REACH) study. *Journal of Epidemiology and Community Health*.

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Mental Distress among Young People in Inner-Cities:  
the Resilience, Ethnicity and AdolesCent Mental Health (REACH) study**

**Authors:** Gemma Knowles<sup>1,2</sup>, PhD; Charlotte Gayer-Anderson<sup>1,2</sup>, PhD; Stephanie Beards<sup>1,3</sup>, PhD; Rachel Blakey<sup>1,2</sup>, MSc; Sam Davis<sup>1,2</sup>, BA; Katie Lowis<sup>1</sup>, BSc; Daniel Stanyon<sup>1</sup>, MSc; Aisha Ofori<sup>1</sup>, MSc; Alice Turner<sup>1,2</sup>, MSc; Schools Working Group; Vanessa Pinfold<sup>4</sup>, PhD; Ioannis Bakolis<sup>5,6</sup>, PhD; Ulrich Reininghaus<sup>7</sup>, PhD; Seeromanie Harding<sup>8</sup>, PhD; Craig Morgan<sup>1,2</sup>, PhD.

CONFIDENTIAL

## **ABSTRACT**

### **Background**

Recent estimates suggest around 14% of 11-16-year-olds in England have a mental health problem. However, we know very little about the extent and nature of mental health problems among diverse groups in densely populated inner cities, where contexts and experiences may differ from the national average.

### **Aims**

To estimate the extent and nature of mental health problems in inner-city London, overall and by social group, using data from our school-based accelerated cohort study of adolescent mental health, REACH (Resilience, Ethnicity and AdolesCent Mental Health).

### **Methods**

Self-report data on mental health (general mental health, depression, anxiety, self-harm) were analysed (n, 4353; 11-14yrs, 85% BAME). Mixed models were used to estimate weighted prevalences and adjusted risks of each type of problem, overall and by gender, cohort, ethnic group, and free school meals (FSM) status.

### **Results**

The weighted prevalence of mental health problems was 18.6% (95% CI 16.4-20.8). Each type of mental health problem was more common among girls compared with boys (adjusted risk ratios [adjRR]: mental health problems, 1.33 [95% CI: 1.18-1.48]; depression, 1.52 [1.30-1.73]; anxiety, 2.09 [1.58-2.59], self-harm, 1.40 [1.06-1.75]). Gender differences were more pronounced in older cohorts compared with the youngest. Mental health problems (1.28 [1.05-1.51]) and self-harm (1.29 [1.02-1.56]) – but not depression or anxiety – were more common among those receiving (vs. not receiving) FSM. There were many similarities, with some variations, by ethnic group.

### **Conclusions**

Adolescent mental health problems and self-harm are common in inner-city London. Gender differences in mental health problems may emerge during early adolescence.

### **What is already known on this subject?**

Recent estimates from national surveys in England suggest that around 1-in-7-8 adolescents have a mental health problem. While national surveys provide useful information on the prevalence of mental health problems nationally, they tend not to provide detailed information on specific groups in which prevalence (and change in prevalence over time) may differ from the national average, e.g., minority ethnic and socially disadvantaged groups living in densely populated inner-city areas. At present, there is very little up-to-date information on the extent and nature of problems among young people from diverse groups and inner-city areas in the UK.

### **What this study adds?**

REACH – Resilience, Ethnicity and AdolesCent Mental Health – is an accelerated cohort study of 4,353 young people (age 11-14 years at baseline; 85% BAME, 1-in-4 receiving free school meals) attending secondary schools in inner-city London. In analyses of baseline data, we found that around 1-in-5 (19%) experienced recognisable mental health issues in the 6 months prior to assessment, a notably higher prevalence than in recent national estimates (10-14%) and estimates for inner city London from around 15-20 years ago (10-12%). Around 1-in-7 (14%) reported lifetime self-harm. Mental health problems were more common among girls than boys, a difference that was more pronounced in older cohorts, and among those from economically disadvantaged backgrounds. We observed many similarities – with some variations – in prevalence of mental health problems across ethnic groups, with highest prevalence (around 1 in 4) being among those of mixed ethnic backgrounds (25%). These findings underscore the importance of local population-based studies to more fully characterise and understand variations in prevalence, and in the distribution of risk and protective factors, by area and social group and to better inform local strategies for prevention and intervention. The many similarities in prevalence of mental health problems across ethnic and socioeconomic groups are striking and understanding these similarities should be a priority for future research; it will advance our understanding of resilience and protective factors, and therefore how to promote mental health and prevent problems, in young people from all backgrounds.

## **INTRODUCTION**

Mental ill-health is a major public health concern.<sup>1</sup> Adolescence is a critical period of emotional and behavioural development and the period during which most adult mental health problems first emerge.<sup>2</sup> Adolescents who develop mental health problems, particularly persistent or recurring problems, are at increased risk of wide-ranging adverse outcomes in adulthood.<sup>3-5</sup> The costs are vast. In 2009/10, the estimated economic and social costs of mental ill-health in England was £105.2 billion.<sup>6</sup>

The most recent national survey of the Mental Health of Children and Young People in England (MHCYP) suggests around 14% of 11-16-year olds have a mental health problem.<sup>7</sup> This is broadly in

line with other national surveys using self-report measures, e.g., 12% in Understanding Society, 2011/12.<sup>8</sup> Despite widespread perceptions that mental health problems have risen among young people in recent years, the MHCYP surveys suggest little change in the overall prevalence of mental health problems among 5-15 year olds over the past 15-20 years. Notable increases in emotional problems were evident among older (16-19 year old) females, but among younger adolescents (11-16 year olds) there was at most a small increase in emotional problems between 1999 and 2017 and little change for other types of mental health problems.<sup>7</sup> This is somewhat surprising in the context of profound social change in the UK during this period (e.g., recession, austerity, changes in migration). However, the effects of such changes are unlikely to be uniform across social groups and geographical areas. While national surveys provide useful information on the prevalence of mental health problems nationally, they do not provide detailed information on specific groups in which prevalence may differ from the national average, e.g., minority ethnic and socially disadvantaged groups in densely-populated inner-cities. Moreover, national surveys may obscure changes over time in some groups. At present, there is very little up-to-date information on the extent and nature of problems among young people from diverse groups and inner-city areas in the UK. Limited evidence suggests young people from UK minority ethnic groups generally have similar or better mental health than their white British peers, but findings are inconsistent, the most robust data are now almost two decades old,<sup>9,10</sup> and much of the recent evidence is based on small, unrepresentative samples and data aggregated over broad, heterogeneous groups (e.g., 'Black', 'Asian').<sup>7,11</sup> One of the more consistent findings is that young people from Indian or, in some studies, 'Asian' backgrounds tend to have better mental health than their white British peers.<sup>11,12</sup>

In this paper, we present baseline data from a prospective study of adolescent mental health in inner-city London – Resilience, Ethnicity and AdolesCent Mental Health (REACH) – on the extent and nature of mental health problems among adolescents from diverse backgrounds. REACH is the largest localised contemporary study of its kind in the UK. We hypothesised that the prevalence of mental health problems would be higher in inner-city London compared with nationally; higher among girls compared with boys, particularly depression; higher among those from economically disadvantaged compared with more affluent backgrounds; and lower among those from Indian/Pakistani/Bangladeshi backgrounds but similar across other ethnic groups.

## **METHODS**

### **Study design**

REACH is an accelerated cohort study of adolescent mental health in two inner-city London boroughs, Lambeth and Southwark. Three cohorts – age 11-12 (Cohort 1; school year 7), 12-13 (Cohort 2; school

year 8), and 13-14 (Cohort 3; school year 9) at baseline – are completing annual questionnaires. A subset (around 20%) are also completing face-to-face interviews and cognitive assessments.

### **Setting, participants**

Participants were recruited from twelve mainstream secondary schools in two inner-city London boroughs, Lambeth and Southwark. These boroughs are among the most densely-populated, socio-economically, and ethnically diverse areas in England,<sup>13-15</sup> and have high rates of adult mental health problems.<sup>16</sup> Schools were selected to be representative of mainstream secondary schools within the two boroughs, based on the proportion of pupils receiving free school meals (FSM) and the proportion in minority ethnic groups. At each school, all students in Years 7-9 were invited to participate (n, 4945).

### **Procedures**

Informed consent was obtained for all participants. Eligible young people attended an in-school talk about REACH, delivered by a researcher, and received written information for themselves and their parents/carers. Parents were asked to return a form, or contact the school or research team, if they did not want their child to take part. On the day of assessment, students received further verbal and written information from researchers and provided written assent before completing a battery of validated questionnaires, in-class, on study tablet computers. The baseline assessment took around 1-hour to complete and collected detailed information on mental health and putative risk and protective factors. Trained researchers were present throughout all sessions to explain procedures and answer questions. All baseline questionnaires were administered between February 2016 and January 2018.

### **Ethical approval**

All study procedures were approved by the Psychiatry, Nursing and Midwifery Research Ethics Subcommittee (PNM-RESC), King's College London (ref:15/162320).

### **Mental health**

General mental health was assessed using the 25-item self-report Strength and Difficulties Questionnaire (SDQ) for 11-17 year olds,<sup>17</sup> a widely-used and validated<sup>17</sup> measure of emotional and behavioural problems during the previous 6 months. Scores  $\geq 18$  indicate high-to-very high risk of mental health problems (hereafter, 'mental health problem'). Internalising and externalising scores were calculated using established procedures.<sup>18</sup> Higher scores indicate more difficulties.

### **Depression**

Depression was assessed using the 13-item Short Mood and Feelings Questionnaire (SMFQ),<sup>19</sup> a self-report measure of depressive symptoms over the previous two weeks. Scores  $\geq 12$  indicate high risk of depression (hereafter, 'depression'). The SMFQ has high internal consistency and convergent validity and moderate diagnostic accuracy among adolescents.<sup>19,20</sup>

### **Anxiety**

Generalised anxiety was assessed using the 7-item self-report Generalised Anxiety Disorder Scale (GAD-7), which measures experiences of generalised anxiety over the preceding two weeks.<sup>21</sup> Scores  $\geq 10$  indicate moderate-to-severe anxiety (hereafter, 'anxiety').<sup>21</sup> Recent community-based research supports the validity and reliability of the GAD-7 among adolescents.<sup>22</sup>

### **Self-harm**

Lifetime self-harm (yes/no) was self-reported using an item from the Development and Adolescent Wellbeing Assessment.<sup>23</sup>

### **Demographic information**

Participants were asked to describe their ethnic group based on eighteen categories used in the 2011 ONS census.<sup>24</sup> We combined some smaller groups (e.g., Arab, Chinese) and used ten ethnic groups in this analysis (see Table 1). FSM status, a marker of household socioeconomic status, was self-reported.

### **Statistical analysis**

Frequencies and proportions were calculated for demographic data. Multilevel linear and logistic regression models (melogit and mixed commands) were used to estimate weighted prevalence (margins command) of each type of mental health problem, overall and by group (cohort, gender, ethnic group, FSM), with school fitted as a second-level variable in all models to account for clustering. The marginalised delta method was used to calculate risk ratios (RR) and 95% confidence intervals (CI) from odds ratios.<sup>25</sup> Gender differences in mental health were examined in models stratified by cohort. In the absence of a clear reference group in comparisons of mental health problems by ethnic group (due to the diversity of the sample), the whole sample prevalence was used as the reference. Adjusted RRs are presented throughout. Unadjusted RRs are presented in Supplement S1. Sample weights were calculated using 2016/17 data from the National Pupil Database. Weighted estimates are presented throughout. Partially observed variables (SDQ [missing n, 106], SMFQ [missing n, 396], GAD-7 [missing n, 243], self-harm [missing n, 474], FSM [missing n, 240]) were multiply imputed in a multilevel model, under the missing at random assumption, using REALCOM<sup>26</sup>. *Sensitivity analyses*: (1) depression, anxiety, and self-harm questions were not administered at two (originally pilot) schools. To assess any

impact of including these two schools in analyses of SDQ (but not other outcome) data, we repeated analyses excluding participants from these two schools (n=818); (2) we conducted a complete case analysis to assess the impact of missing data.

## **RESULTS**

### **Participant characteristics**

Of 4945 eligible students, 4353 (88.0%) participated at baseline and are included in this analysis. Reasons for non-participation were: persistent absence despite repeated visits by researchers (n 354, 7.2%); did not have parental consent (n 167, 3.4%); did not assent (n 57, 1.2%); insufficient data due to technical issues (n 15, 0.03%). Just under a quarter (23.7%) of participants received FSM and 85% were from minority ethnic groups. Demographic characteristics were similar across cohorts, except for a higher proportion of British white students in Cohort 1 (17.9%) compared with Cohorts 2 (13.0%) and 3 (14.7%) ( $\chi^2=40.3$ ,  $df=18$ ,  $p=0.002$ ). The REACH sample is highly representative of the target population (Year 7-9 pupils attending mainstream schools in the boroughs); weighting made little difference to our estimates (Supplement S2).

### **Mental health problems, overall**

The weighted prevalence was 18.6% (95% CI 16.4-20.8) for mental health problems, 14.5% (11.8-17.2) for depression, 13.7% (10.9-16.6) for anxiety, and 14.5% (12.4-16.6) for lifetime self-harm (Table 2). Of those with mental health problems, 69.0% were also categorised as having problems on at least one other mental health measure.

### **Mental health problems, by group**

#### **Gender**

Overall, prevalence was higher among girls than among boys for mental health problems (aRR 1.33 [95% CI 1.18-1.48]), depression (1.52 [1.30-1.73]), moderate-to-severe anxiety (2.09 [1.58-2.59]), and lifetime self-harm (1.40 [1.06-1.75]) (Table 2). Mean internalising score was also higher among girls (6.1 [95% CI 5.9-6.4]) compared with boys (5.0 [4.8-5.3]), as were mean GAD (5.1 [4.7-5.5] vs. 3.3 [2.9-3.7]) and SMFQ (6.0 [5.4-6.6] vs. 4.4 [4.0-4.9]) scores (Table 3).

#### **Cohort**

There was no clear evidence for variations in prevalence of mental health problems by cohort.

#### **Gender and cohort**



In models stratified by cohort (Figure 1), clear patterns emerged of variation in gender differences by cohort. Prevalence of mental health problems was similar for boys and girls in Cohort 1 (Year 7), but higher among girls in Cohort 2 (Year 8) and higher still in Cohort 3 (Year 9). For example, the relative risks of mental health problems among girls (vs. boys) were 1.09 (95% CI 0.85-1.34), 1.27 (0.99-1.55), and 1.74 (1.38-2.10) in Cohorts 1, 2 and 3, respectively (Figure 1). A similar pattern was observed for depression, anxiety, lifetime self-harm (Supplement S3), and internalising scores (Supplement S4). By contrast, mean externalising scores were similar between boys and girls in each cohort (Supplement S4).

### **Socioeconomic Disadvantage**

Among those receiving (vs. not receiving) FSM, risks of mental health problems (aRR 1.28 [95% CI 1.05-1.51]) and of lifetime self-harm (1.29 [1.02-1.56]) were around 30% higher (Table 2). However, risks of anxiety and depression were similar between the two groups.

### **Ethnic group**

Overall, there were many similarities, with some variations, in prevalence of mental health problems by ethnic group (Table 2). For example, compared with the overall sample, mental health problems were more common among those from other mixed ethnic backgrounds (aRR 1.34 [95% CI 1.02-1.74]) and less common among those from Indian/Pakistani/Bangladeshi backgrounds (0.67 [0.42-1.01]), but otherwise similar across ethnic groups. A similar pattern was observed for anxiety, although there was also some evidence for higher prevalence of anxiety among those from Latin American (1.35 [0.95-1.86]) and non-British white (1.20 [0.97-1.35]) backgrounds compared with the overall sample. We found no evidence of differences in prevalence of depression by ethnic group (14-16% across all groups). For self-harm, risk was lower among those from black African backgrounds compared with the overall sample (0.84 [0.70-0.99]), and there was some evidence for increased risk among those from other mixed ethnic backgrounds (1.25 [0.90-1.69]), but otherwise there was little difference across groups. Overall, prevalence of mental health problems was generally similar between those from black African, black Caribbean, and white British backgrounds (Table 2), but there was evidence for variations in externalising scores between these groups. For example, mean externalising scores were higher among black Caribbean (7.4 [7.0-7.8]), mixed black and white (7.2 [6.6-7.8]), and other Black (7.0 [6.5-7.5]), and lower among British white (6.1 [5.8-6.4]) (Table 3).

### **Sensitivity analyses**

The findings were not substantively changed in sensitivity analyses.

## **DISCUSSION**

In the most recent and comprehensive study of adolescent mental health in an inner-city area in the UK, we found that mental health problems were notably more common than reported nationally; around 1-in-5 had a mental health problem compared with around 1-in-7-8 nationally. This equates to, on average, 6 pupils in a class of 30 in inner-city schools. We also found that differences between boys and girls emerge in older cohorts, suggesting this may be a critical point at which lifelong differences emerge, and we found interesting similarities, with some variations, by ethnic group and socioeconomic status.

### **Methodological Considerations**

Several methodological issues should be noted when considering our findings. First, in this part of REACH we used self-report measures of mental health. These measures are not designed to provide clinical diagnoses; they are screening tools designed to identify those likely to have a diagnosable problem. It is likely, then, that some young people in our study are misclassified. Nonetheless, the measures used in our study were developed/validated for this age group, are widely used in epidemiological studies of adolescent mental health, and our use of multiple screening measures (SDQ, SMFQ, GAD-7) provides more detailed information on participants' mental health compared with many other studies.<sup>8,11</sup> Collection of more in-depth information via interviews with a nested subsample is ongoing and will enable triangulation of data.

Second, those who did not participate were primarily those who were persistently absent from school. This is a potential source of bias, but the impact, if any, would be conservative estimates – i.e., underestimation of the prevalence of, and many subgroup differences in, mental health problems – because those who did not participate may be more likely to have problems and disadvantaged backgrounds compared with those who did. Moreover, our cohorts are highly representative of the target population, weighting made little difference to our estimates, and we achieved high participation rates: 88% in REACH compared with, for example, 52% in the MHCYP.<sup>7</sup>

### **The Nature and Extent of Adolescent Mental Health Problems**

Estimates of the prevalence of mental health problems among adolescents vary across studies. Previous national studies in the UK estimate a prevalence of around 10-14%, with, overall, little change among secondary school-aged pupils over time.<sup>7,8,27</sup> Some of this variation may be due to the methods used. For instance, in the MHCYP (2017, 11-16-years), the estimated prevalence was 14% using multi-informant interviews.<sup>7</sup> Understanding Society (2009/10 and 2011/12, 10-15-years) estimated 12% using similar methods to ours.<sup>8</sup> Nonetheless, these differences are small. Our estimate, 1-in-5 (19%), is notably higher, and strongly suggests higher prevalence in inner-cities. Further, it may be that in

diverse inner-city contexts prevalence has increased, more so than reported nationally, over the past 15-20 years. Our estimates (16% among boys, 21% among girls; 11-14-years) are notably higher than observed in two similar multi-ethnic inner-city London studies conducted in 2001 (boys, 10%; girls 12%; 11-13-years<sup>9</sup>) and 2003 (boys, 14%; girls, 17%; 11-14-years<sup>10</sup>). Taken together – while noting the limitations of directly comparing these data – the evidence suggests that (a) problems are more common among adolescents in inner-city London compared with national samples, and (b) within inner-city London, prevalence has increased over the last 15-20 years.

However, while our data suggest the prevalence of mental health problems may, overall, be elevated in inner-city London compared with nationally, prevalence of (self-report) depression and self-harm appear to be similar to, or slightly lower than, recent national estimates (around 15% for lifetime self-harm and 16% for depression<sup>27,28</sup>). It is possible, then, that elevated risk in inner-cities is driven by externalising/behavioural problems rather than internalising/emotional problems. Directly comparable data on externalising problems in this age group are scarce and further research is required, but this observation may reflect substantive differences in the experiences of young people growing up in diverse inner-city areas, and/or variations in the manifestation of distress among diverse groups. These findings are important and reinforce the need for localised studies of mental health.<sup>16</sup> Nationally, there is a strong focus on addressing rising emotional problems among girls and young women; much less attention is directed at externalising problems, or indeed emotional problems and self-harm among boys. That 12% of boys in inner-city London have self-harmed by age 11-14 is concerning and there is an urgent need for greater understanding and preventive strategies.

Consistent with existing evidence,<sup>7,29</sup> mental health problems were more common among girls than among boys. However, our data – which permit more fine-grained analyses than is often possible in other surveys – suggest risk of problems is similar between boys and girls in Year 7 and becomes more pronounced with age. These data, at this stage, are cross-sectional, so cohort effects cannot be ruled out, but this observation mirrors findings from previous research in London<sup>10</sup> and internationally,<sup>30</sup> and, if replicated with longitudinal REACH data, may have significant implications. Irrespective of whether gender differences at this age reflect genuine differences in risk, or under-reporting of distress among boys, these findings point to early adolescence as the possible point at which life-long gender inequalities in (reported) mental health problems first emerge.

Our data – the most recent and detailed information on adolescent mental health among diverse groups and one of few studies with sufficient power for ethnic group comparisons – suggests prevalence of mental health problems is lower among Indian/Bangladeshi/Pakistani adolescents, and higher among those from other mixed ethnic backgrounds (a group that is growing in size in the UK),

but otherwise similar across ethnic groups. This observation is broadly consistent with earlier findings.<sup>9-11</sup> Given that most UK minority ethnic groups experience disproportionately high levels of adversity (e.g., discrimination, poverty), these observations are striking.

### **Implications**

Our findings suggest important implications for research, prevention/intervention, and local services and policy. First, national data is important, but of limited use in understanding the extent and nature of distress in local communities; localised studies are needed to inform prevention strategies and local service provision.<sup>16</sup> Second, the widely reported gender difference in mental health likely emerges during the first few years of secondary school; this may be a critical period for prevention and intervention. Third, the many similarities in prevalence of mental health problems across ethnic and socioeconomic groups are striking and understanding these similarities should be a priority for future research; it will advance our understanding of resilience and protective factors, and therefore how to promote mental health and prevent problems, in young people from all backgrounds.

CONFIDENTIAL

**Author affiliations:** <sup>1</sup>Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK; <sup>2</sup>ESRC Centre for Society and Mental Health, King's College London; <sup>3</sup>National Children's Bureau, London, UK; <sup>4</sup>The McPin Foundation, London, UK; <sup>5</sup>Centre for Implementation Science, Health Services and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK; <sup>6</sup>Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK; <sup>7</sup>Department of Public Mental Health, Central Institute of Mental Health, Medical Faculty Mannheim, University of Heidelberg, Mannheim, Germany; <sup>8</sup>Department of Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London, UK.

**Correspondence to:** Professor Craig Morgan, ESRC Centre for Society and Mental Health, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, SE5 8AF. Tel: 0207 848 0351. Email: [craig.morgan@kcl.ac.uk](mailto:craig.morgan@kcl.ac.uk).

**Funding:** REACH is supported by funding from the European Research Council (Ref: REACH 648837). The funder had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

**Acknowledgments:** We would like to thank all participating schools, teachers, young people, and parents, and our Young Persons Advisory Group, for their invaluable contributions to REACH. We would also like to thank the research assistants and students who have contributed to data collection.

**Declaration of Interest:** none.

**Author contributions:** **GK:** drafted and revised the manuscript; approved the final manuscript; collected, analysed, and interpreted the data; contributed to the design of the study; coordinated the study. **CGA:** revised the manuscript; approved the final manuscript; collected, analysed, and interpreted the data; contributed to the design of the study. **SB, RB, SD, KL, DS, AO, AT:** collected, cleaned, and interpreted the data; critically revised the manuscript; approved the final manuscript. **SWG:** made substantial contributions to the acquisition and interpretation of data and critically reviewed and approved the final manuscript. **VP:** contributed to the conceptualisation and design of the work; interpreted the data; revised the manuscript; approved the final manuscript. **IB:** contributed to data analysis and interpretation; critically revised the manuscript; approved the final manuscript. **UR:** contributed to the conceptualisation and design of the work; interpreted the data; critically revised the manuscript; approved the final manuscript. **SH:** contributed to the conceptualisation and design of the work; interpreted the data; critically revised the manuscript; approved the final manuscript. **CM:** conceptualised and designed the work; collected, analysed, and interpreted the data; critically revised the manuscript; approved the final manuscript.

## References

- 1 Steel Z, Marnane C, Iranpour C, et al. The global prevalence of common mental disorders: a systematic review and meta-analysis 1980-2013. *Int J Epidemiol* 2014; **43**(2): 476-93.
- 2 Kessler RC, Amminger GP, Aguilar-Gaxiola S, Alonso J, Lee S, Ustun TB. Age of onset of mental disorders: a review of recent literature. *Curr Opin Psychiatry* 2007; **20**(4): 359-64.
- 3 Fergusson DM, Horwood LJ, Ridder EM. Show me the child at seven: the consequences of conduct problems in childhood for psychosocial functioning in adulthood. *J Child Psychol Psychiatry* 2005; **46**(8): 837-49.
- 4 Kessler RC, Foster CL, Saunders WB, Stang PE. Social consequences of psychiatric disorders, I: Educational attainment. *Am J Psychiatry* 1995; **152**(7): 1026-32.
- 5 Fergusson DM, Boden JM, Horwood LJ. Recurrence of major depression in adolescence and early adulthood, and later mental health, educational and economic outcomes. *Br J Psychiatry* 2007; **191**: 335-42.
- 6 Centre for Mental Health, 2010. The economic and social costs of mental health problems in 2009/10. [https://www.centreformentalhealth.org.uk/sites/default/files/2018-10/Economic\\_and\\_social\\_costs\\_2010.pdf](https://www.centreformentalhealth.org.uk/sites/default/files/2018-10/Economic_and_social_costs_2010.pdf) (accessed: 26/04/2019).
- 7 Sadler K, Vizard T, Ford T, et al. Mental Health of Children and Young People in England, 2017. Office for National Statistics: NHS Digital, 2018.
- 8 Office for National Statistics, 2015. Measuring National Well-being: Insights into children's mental health and well-being. <https://www.ons.gov.uk/peoplepopulationandcommunity/wellbeing/articles/measuringnationalwellbeing/2015-10-20#what-do-the-data-tell-us> (accessed: 20/08/2019).
- 9 Maynard MJ, Harding S, Minnis H. Psychological well-being in Black Caribbean, Black African, and White adolescents in the UK Medical Research Council DASH study. *Social Psychiatry and Psychiatric Epidemiology* 2007; **42**(9): 759-69.
- 10 Stansfeld S, Haines M, Booy R, et al., 2003. The health of young people in East London: the RELACHS study, 2001. [http://archive.wolfson.qmul.ac.uk/relachs/academics/publications/Relachs\\_report.pdf](http://archive.wolfson.qmul.ac.uk/relachs/academics/publications/Relachs_report.pdf) (accessed: 20/08/2019).
- 11 Goodman A, Patel V, Leon DA. Child mental health differences amongst ethnic groups in Britain: a systematic review. *BMC Public Health* 2008; **8**(1): 258.
- 12 Smith NR, Lewis DJ, Fahy A, et al. Individual socio-demographic factors and perceptions of the environment as determinants of inequalities in adolescent physical and psychological health: the Olympic Regeneration in East London (ORIEL) study. *BMC Public Health* 2015; **15**: 150.
- 13 Office for National Statistics. 2011 Census: Key Statistics for local authorities in England and Wales. Table KS201EW: Ethnic group. 2011.
- 14 Lambeth Council, 2016. State of the Borough 2016: Lambeth. <https://www.lambeth.gov.uk/sites/default/files/State%20of%20Borough%202016%20-%20v3.pdf> (accessed: 18/06/2019).

- 15 Southwark Council, 2018. Southwark Joint Strategic Needs Assessment 2018-19: Demography. <https://www.southwark.gov.uk/assets/attach/7379/JSNA%20Factsheet%202018-19%20-%20Demography%2020180725.pdf> (accessed: 18/06/2019).
- 16 Hatch SL, Woodhead C, Frissa S, et al. Importance of thinking locally for mental health: data from cross-sectional surveys representing South East London and England. *PLoS One* 2012; **7**(12): e48012.
- 17 Goodman R, Meltzer H, Bailey V. The Strengths and Difficulties Questionnaire: a pilot study on the validity of the self-report version. *Eur Child Adolesc Psychiatry* 1998; **7**(3): 125-30.
- 18 Goodman A, Lamping DL, Ploubidis GB. When to use broader internalising and externalising subscales instead of the hypothesised five subscales on the Strengths and Difficulties Questionnaire (SDQ): data from British parents, teachers and children. *J Abnorm Child Psychol* 2010; **38**(8): 1179-91.
- 19 Angold A, Costello EJ, Messer SC, Pickles A. Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *International Journal of Methods in Psychiatric Research* 1995; **5**(4): 237-49.
- 20 Thabrew H, Stasiak K, Bavin LM, Frampton C, Merry S. Validation of the Mood and Feelings Questionnaire (MFQ) and Short Mood and Feelings Questionnaire (SMFQ) in New Zealand help-seeking adolescents. *Int J Methods Psychiatr Res* 2018; **27**(3): e1610.
- 21 Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006; **166**(10): 1092-7.
- 22 Tiirikainen K, Haravuori H, Ranta K, Kaltiala-Heino R, Marttunen M. Psychometric properties of the 7-item Generalized Anxiety Disorder Scale (GAD-7) in a large representative sample of Finnish adolescents. *Psychiatry Res* 2019; **272**: 30-5.
- 23 Goodman R, Ford T, Richards H, Gatward R, Meltzer H. The Development and Well-Being Assessment: description and initial validation of an integrated assessment of child and adolescent psychopathology. *J Child Psychol Psychiatry* 2000; **41**(5): 645-55.
- 24 Office for National Statistics, 2012. Ethnicity and National Identity in England and Wales: 2011. <https://www.ons.gov.uk/peoplepopulationandcommunity/culturalidentity/ethnicity/articles/ethnicityandnationalidentityinenglandandwales/2012-12-11> (accessed: 20/08/2019).
- 25 Localio AR, Margolis DJ, Berlin JA. Relative risks and confidence intervals were easily computed indirectly from multivariable logistic regression. *J Clin Epidemiol* 2007; **60**(9): 874-82.
- 26 Carpenter JR, Goldstein H, Kenward MG. REALCOM-IMPUTE Software for Multilevel Multiple Imputation with Mixed Response Types. *Journal of Statistical Software* 2011; **45**(5).
- 27 Patalay P, Gage SH. Changes in millennial adolescent mental health and health-related behaviours over 10 years: a population cohort comparison study. *Int J Epidemiol* 2019.
- 28 Morey Y, Mellon D, Dailami N, Verne J, Tapp A. Adolescent self-harm in the community: an update on prevalence using a self-report survey of adolescents aged 13–18 in England. *Journal of Public Health* 2016; **39**(1): 58-64.

- 29 McManus S BP, Jenkins R, Brugha T., 2016. Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014. Leeds: NHS Digital. <https://digital.nhs.uk/data-and-information/publications/statistical/adult-psychiatric-morbidity-survey/adult-psychiatric-morbidity-survey-survey-of-mental-health-and-wellbeing-england-2014> (accessed: 20/08/2019).
- 30 Wade TJ, Cairney J, Pevalin DJ. Emergence of gender differences in depression during adolescence: national panel results from three countries. *J Am Acad Child Adolesc Psychiatry* 2002; **41**(2): 190-8.

CONFIDENTIAL



Tables, Figures

**Table 1. Participant characteristics**

	All (n, 4353)		Boys (n, 2138)		Girls (n, 2215)		Cohort 1 (n, 1593)		Cohort 2 (n, 1421)		Cohort 3 (n, 1339)		
	n	%	n	%	n	%	n	%	n	%	n	%	
Gender (n, 4,353)													
Boys	2,138	49.1	..	..	..	..	778	48.8	701	49.3	659	49.2	
Girls	2,215	50.9	..	..	..	..	815	51.2	720	50.7	680	50.8	
Cohort (school year group) (n, 4,353)													
1 (Y7)	1,593	36.6	778	36.4	815	36.8	..	..	..	..	..	..	
2 (Y8)	1,421	32.6	701	32.8	720	32.5	..	..	..	..	..	..	
3 (Y9)	1,339	30.8	659	30.8	680	30.7	..	..	..	..	..	..	
Eligible for free school meals (n, 4,113)													
No	3,137	76.3	1,563	77.2	1,574	75.4	1,142	76.5	1,015	75.4	980	76.9	
Yes	976	23.7	461	22.8	515	24.7	351	23.5	331	24.6	294	23.1	
Ethnic group (n, 4,353)													
Black African	1,113	25.6	545	25.5	568	25.6	383	24.0	374	26.3	356	26.6	
Black Caribbean	719	16.5	350	16.4	369	16.7	234	14.7	257	18.1	228	17.0	
Other black	127	2.9	51	2.4	76	3.4	49	3.1	46	3.2	32	2.4	
Mixed white and black	380	8.7	182	8.5	198	8.9	143	9.0	117	8.2	120	9.0	
Other mixed ethnic groups	237	5.4	105	4.9	132	6.0	87	5.5	82	5.8	68	5.1	
Indian, Pakistani, Bangladeshi	181	4.2	83	3.9	98	4.4	79	5.0	57	4.0	45	3.4	
Latin American	217	5.0	110	5.1	107	4.8	58	3.6	79	5.6	80	6.0	
White British	667	15.3	337	15.8	330	14.9	285	17.9	185	13.0	197	14.7	
Non-British white	409	9.4	213	10.0	196	8.9	166	10.4	121	8.5	122	9.1	
Any other/Unknown	303	7.0	162	7.6	141	6.4	109	6.8	103	7.3	91	6.8	

The proportion of white British students was slightly higher in Cohort 1 compared with Cohorts 2 and 3 ( $\chi^2= 40.3223$ ,  $df=18$ ,  $p=0.002$ ).

**Table 2. Weighted prevalence of mental health problems (and 95% confidence intervals), overall and by group**


	Probable mental health problems		Probable depression		Moderate-to-severe anxiety		Lifetime self-harm		
	% (95% CI)	Adj. RR (95% CI)	% (95% CI)	Adj. RR (95% CI)	% (95% CI)	Adj. RR (95% CI)	% (95% CI)	Adj. RR (95% CI)	
<b>Gender</b>	All	18.6 (16.4, 20.8)	-	14.5 (11.8, 17.2)	-	13.7 (10.9, 16.6)	-	14.5 (12.4, 16.6)	-
	Boys	16.0 (14.1, 17.8)	1	11.5 (9.3, 13.8)	1	8.8 (6.3, 11.3)	1	12.2 (10.0, 14.3)	1
	Girls	21.2 (18.7, 23.8)	1.33 (1.18, 1.48)	17.5 (14.3, 20.7)	1.52 (1.30, 1.73)	18.5 (16.2, 20.8)	2.09 (1.58, 2.59)	16.9 (13.9, 19.8)	1.40 (1.06, 1.75)
<b>Cohort (school year group)</b>	1 (Y7)	19.2 (16.6, 21.7)	1	13.5 (10.2, 16.7)	1	13.0 (9.8, 16.1)	1	12.7 (9.6, 15.8)	1
	2 (Y8)	18.8 (15.9, 21.6)	0.97 (0.79, 1.14)	16.4 (13.0, 19.9)	1.22 (0.92, 1.53)	14.7 (9.6, 19.9)	1.12 (0.71, 1.53)	16.4 (13.0, 19.8)	1.29 (0.90, 1.68)
	3 (Y9)	17.7 (15.1, 20.3)	0.91 (0.80, 1.02)	13.7 (10.3, 17.0)	1.02 (0.75, 1.28)	13.6 (10.5, 16.8)	1.03 (0.71, 1.35)	14.8 (10.6, 19.0)	1.17 (0.73, 1.61)
<b>Eligible for free school meals</b>	No	17.4 (14.9, 19.9)	1	14.1 (11.1, 17.0)	1	14.0 (11.1, 16.9)	1	13.8 (11.4, 16.1)	1
	Yes	22.5 (19.8, 25.2)	1.28 (1.05, 1.51)	16.1 (12.3, 19.8)	1.15 (0.84, 1.46)	12.6 (8.6, 16.5)	0.87 (0.64, 1.10)	17.4 (14.2, 20.5)	1.29 (1.02, 1.56)
<b>Ethnic group</b>	All	18.6 (16.4, 20.8)	1	14.5 (11.8, 17.2)	1	13.7 (10.9, 16.6)	1	14.5 (12.4, 16.6)	1
	Black African	17.2 (14.1, 20.4)	0.93 (0.80, 1.07)	13.7 (10.7, 16.7)	0.96 (0.81, 1.12)	11.9 (9.1, 14.7)	0.88 (0.73, 1.05)	12.1 (7.9, 16.2)	0.84 (0.70, 0.99)
	Black Caribbean	20.8 (16.8, 24.8)	1.12 (0.94, 1.31)	15.8 (10.3, 21.2)	1.08 (0.89, 1.30)	12.5 (8.0, 16.9)	0.89 (0.71, 1.11)	15.2 (11.9, 18.5)	1.04 (0.85, 1.25)
	Other black	19.4 (15.1, 23.7)	1.04 (0.67, 1.54)	14.3 (9.8, 18.8)	0.94 (0.55, 1.51)	14.1 (8.1, 20.1)	1.00 (0.57, 1.62)	16.6 (9.2, 24.0)	1.16 (0.72, 1.78)
	Mixed white and black	22.4 (16.6, 28.3)	1.19 (0.94, 1.47)	14.0 (8.8, 19.3)	0.96 (0.72, 1.26)	13.7 (8.5, 19.0)	1.02 (0.75, 1.35)	15.3 (9.9, 20.7)	1.04 (0.78, 1.35)
	Other mixed ethnic groups	25.4 (19.3, 31.5)	1.34 (1.02, 1.74)	15.7 (11.5, 20.0)	1.07 (0.75, 1.48)	21.2 (15.6, 26.9)	1.57 (1.15, 2.08)	18.4 (13.9, 22.9)	1.25 (0.90, 1.69)
	Indian, Pakistani, Bangladeshi	12.3 (6.8, 17.8)	0.67 (0.42, 1.01)	14.5 (7.7, 21.4)	1.01 (0.66, 1.48)	10.6 (5.6, 15.6)	0.74 (0.43, 1.19)	13.3 (6.9, 19.6)	0.93 (0.60, 1.39)
	Latin American	18.8 (14.2, 23.4)	1.04 (0.74, 1.41)	16.1 (8.2, 24.1)	1.10 (0.76, 1.54)	18.2 (11.1, 25.2)	1.35 (0.95, 1.86)	15.7 (9.8, 21.7)	1.07 (0.74, 1.50)
	White British	17.9 (13.2, 22.7)	0.98 (0.81, 1.17)	15.6 (12.0, 19.3)	1.09 (0.89, 1.32)	16.2 (10.9, 21.5)	1.20 (0.97, 1.45)	14.9 (12.6, 17.1)	1.05 (0.85, 1.27)
	Non-British white	17.4 (13.3, 21.5)	0.97 (0.76, 1.22)	13.7 (9.4, 18.0)	0.95 (0.71, 1.23)	13.1 (9.1, 17.1)	0.95 (0.70, 1.25)	17.2 (12.3, 22.1)	1.22 (0.95, 1.54)
	Any other/Unknown	16.9 (12.4, 21.5)	0.93 (0.69, 1.22)	12.2 (7.6, 16.7)	0.86 (0.61, 1.19)	12.6 (8.1, 17.1)	0.99 (0.70, 1.36)	13.6 (8.1, 19.2)	0.93 (0.66, 1.27)

RR, risk ratio; CI, confidence interval; %, weighted prevalence; Y7, school year 7 (age 11-12y); Y8, school year 8 (age 12-13y); Y9, school year 9 (age 13-14y). Risk ratios adjusted for gender, free school meal eligibility, gender and ethnic group, as applicable. All estimates adjusted for clustering at school level.

**Table 3. Weighted mean mental health scores (and 95% confidence intervals), overall and by group**

		SDQ total difficulties score	SDQ internalising score	SDQ externalising score	SMFQ score	GAD-7 score
		mean (95% CI)	mean (95% CI)	mean (95% CI)	mean (95% CI)	mean (95% CI)
Gender	All	12.1 (11.7, 12.5)	5.6 (5.3, 5.8)	6.6 (6.3, 6.8)	5.2 (4.7, 5.7)	4.2 (3.7, 4.7)
	Boys	11.7 (11.4, 12.1)	5.0 (4.8, 5.3)	6.8 (6.5, 7.0)	4.4 (4.0, 4.9)	3.3 (2.9, 3.7)
	Girls	12.5 (11.9, 13.0)	6.1 (5.9, 6.4)	6.3 (6.0, 6.7)	6.0 (5.4, 6.6)	5.1 (4.7, 5.5)
Cohort (school year group)	1 (Y7)	12.1 (11.6, 12.5)	5.6 (5.3, 5.8)	6.5 (6.2, 6.8)	5.0 (4.4, 5.5)	4.0 (3.6, 4.5)
	2 (Y8)	12.2 (11.8, 12.6)	5.7 (5.4, 6.0)	6.6 (6.4, 6.8)	5.7 (5.1, 6.2)	4.4 (3.7, 5.1)
	3 (Y9)	12.0 (11.5, 12.5)	5.5 (5.2, 5.8)	6.6 (6.3, 6.8)	5.1 (4.5, 5.7)	4.2 (3.6, 4.8)
Eligible for free school meals	No	12.0 (11.5, 12.4)	5.5 (5.2, 5.8)	6.5 (6.3, 6.7)	5.1 (4.6, 5.7)	4.2 (3.7, 4.7)
	Yes	12.6 (12.0, 13.1)	5.8 (5.5, 6.1)	6.8 (6.5, 7.2)	5.6 (4.9, 6.3)	4.2 (3.6, 4.9)
Ethnic group	Black African	11.6 (11.1, 12.2)	5.2 (4.9, 5.6)	6.4 (6.1, 6.8)	5.0 (4.4, 5.5)	3.8 (3.3, 4.3)
	Black Caribbean	12.8 (12.1, 13.5)	5.3 (4.9, 5.7)	7.4 (7.0, 7.8)	5.6 (4.6, 6.5)	4.0 (3.1, 5.0)
	Other black	12.3 (11.4, 13.2)	5.3 (4.7, 6.0)	7.0 (6.5, 7.5)	4.6 (3.9, 5.3)	3.9 (3.0, 4.7)
	Mixed white and black	12.7 (11.8, 13.7)	5.5 (5.0, 6.1)	7.2 (6.6, 7.8)	5.0 (4.3, 5.8)	4.0 (3.4, 4.7)
	Other mixed ethnic groups	13.2 (12.4, 13.9)	6.3 (5.8, 6.7)	6.9 (6.5, 7.3)	5.7 (4.9, 6.4)	5.3 (4.4, 6.2)
	Indian, Pakistani, Bangladeshi	10.9 (10.1, 11.8)	5.2 (4.7, 5.8)	5.7 (5.2, 6.1)	5.1 (4.1, 6.0)	3.9 (3.0, 4.7)
	Latin American	12.8 (12.2, 13.4)	6.2 (5.7, 6.7)	6.6 (5.9, 7.4)	5.7 (4.2, 7.1)	4.6 (3.6, 5.5)
	White British	12.0 (11.5, 12.6)	6.0 (5.6, 6.3)	6.1 (5.8, 6.4)	5.6 (5.0, 6.2)	4.8 (4.1, 5.6)
	non-British white	11.9 (11.2, 12.6)	5.7 (5.2, 6.3)	6.2 (5.7, 6.6)	5.0 (4.2, 5.9)	4.3 (3.7, 4.9)
	Any other/unknown	11.9 (11.1, 12.7)	5.7 (5.3, 6.2)	6.2 (5.7, 6.8)	4.9 (4.2, 5.7)	4.0 (3.3, 4.8)

Figure 1. Prevalence of probable mental health problems (and 95% CIs), by gender and cohort.



## Supplement S1

**Table S1. Unadjusted risk ratios (and 95% confidence intervals) for mental health problems**

	Probable mental health problems	Probable depression	Moderate-to- severe anxiety	Lifetime self- harm
	RR (95% CI)	RR (95% CI)	RR (95% CI)	RR (95% CI)
<b>Gender</b>				
Boys	1	1	1	1
Girls	<b>1.33 (1.18, 1.48)</b>	<b>1.52 (1.32, 1.73)</b>	<b>2.11 (1.61, 2.61)</b>	<b>1.39 (1.06, 1.72)</b>
<b>Cohort (school year group)</b>				
1 (Y7)	1	1	1	1
2 (Y8)	0.98 (0.81, 1.15)	1.22 (0.93, 1.52)	1.14 (0.72, 1.56)	1.29 (0.91, 1.67)
3 (Y9)	0.92 (0.82, 1.03)	1.02 (0.75, 1.28)	1.05 (0.73, 1.38)	1.16 (0.74, 1.58)
<b>Eligible for free school meals</b>				
No	1	1	1	1
Yes	<b>1.30 (1.07, 1.53)</b>	1.14 (0.85, 1.44)	0.89 (0.66, 1.13)	<b>1.27 (1.01, 1.52)</b>
<b>Ethnic group</b>				
All	1	1	1	1
Black African	0.93 (0.80, 1.07)	0.94 (0.80, 1.11)	0.86 (0.72, 1.02)	<b>0.83 (0.70, 0.98)</b>
Black Caribbean	1.12 (0.95, 1.32)	1.09 (0.90, 1.31)	0.91 (0.73, 1.12)	1.05 (0.86, 1.26)
Other black	1.06 (0.69, 1.56)	0.98 (0.58, 1.55)	1.03 (0.61, 1.63)	1.14 (0.71, 1.74)
Mixed white and black	1.20 (0.96, 1.49)	0.97 (0.72, 1.26)	1.00 (0.74, 1.31)	1.05 (0.80, 1.36)
Other mixed ethnic groups	<b>1.36 (1.04, 1.75)</b>	1.08 (0.76, 1.49)	<b>1.54 (1.14, 2.03)</b>	1.28 (0.93, 1.72)
Indian, Pakistani, Bangladeshi	<b>0.65 (0.41, 0.99)</b>	0.99 (0.65, 1.46)	0.77 (0.46, 1.19)	0.92 (0.59, 1.36)
Latin American	1.02 (0.73, 1.38)	1.12 (0.78, 1.55)	1.31 (0.93, 1.79)	1.08 (0.75, 1.51)
White British	0.97 (0.80, 1.16)	1.08 (0.88, 1.31)	1.18 (0.97, 1.42)	1.02 (0.83, 1.25)
Non-British white	0.93 (0.73, 1.18)	0.95 (0.72, 1.23)	0.96 (0.72, 1.26)	1.18 (0.92, 1.49)
Any other/Unknown	0.91 (0.67, 1.19)	0.84 (0.59, 1.16)	0.91 (0.65, 1.25)	0.93 (0.67, 1.27)


## Supplement S2

**Table S2. Unweighted prevalence (and 95% confidence intervals) of mental health problems, overall and by group.**


	Probable mental health problems	Probable depression	Moderate-to-severe anxiety	Lifetime self-harm
	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
<b>Gender</b>				
All	18.8 (16.6, 20.9)	14.5 (11.9, 17.2)	13.9 (10.6, 17.2)	14.4 (12.2, 16.5)
Boys	16.1 (14.0, 18.2)	11.6 (9.2, 14.1)	9.0 (7.1, 10.9)	12.3 (10.1, 14.5)
Girls	21.4 (18.9, 23.8)	17.3 (14.1, 20.5)	18.3 (15.3, 21.2)	16.3 (13.8, 18.9)
<b>Cohort (school year group)</b>				
1 (Y7)	19.4 (16.7, 22.1)	13.5 (10.5, 16.4)	13.1 (9.6, 16.5)	12.6 (10.1, 15.1)
2 (Y8)	18.7 (16.0, 21.4)	16.6 (13.0, 20.1)	15.2 (11.2, 19.1)	16.3 (13.2, 19.5)
3 (Y9)	18.0 (15.3, 20.8)	13.8 (10.6, 17.0)	13.7 (9.9, 17.4)	14.8 (11.8, 17.7)
<b>Eligible for free school meals</b>				
No	17.6 (15.5, 19.7)	14.1 (11.4, 16.8)	14.2 (10.8, 17.5)	13.6 (11.4, 15.8)
Yes	22.4 (19.1, 25.6)	16.0 (12.2, 19.8)	12.8 (9.0, 16.6)	17.3 (13.8, 20.8)
<b>Ethnic group</b>				
Black African	17.3 (14.5, 20.1)	13.6 (10.4, 16.9)	11.9 (8.5, 15.3)	12.0 (9.3, 14.7)
Black Caribbean	20.9 (17.3, 24.5)	15.9 (11.6, 20.3)	12.8 (8.6, 16.9)	15.3 (11.5, 19.1)
Other black	20.1 (12.9, 27.3)	14.4 (7.2, 21.5)	14.5 (7.3, 21.8)	17.5 (10.0, 25.0)
Mixed white and black	22.8 (18.1, 27.5)	14.1 (9.4, 18.7)	14.0 (9.0, 19.0)	15.5 (10.9, 20.1)
Other mixed ethnic groups	25.8 (19.7, 31.8)	15.9 (10.2, 21.6)	21.0 (14.1, 27.9)	17.9 (12.2, 23.6)
Indian, Pakistani, Bangladeshi	12.5 (7.5, 17.6)	15.6 (8.8, 22.3)	10.9 (5.3, 16.6)	13.5 (7.6, 19.4)
Latino/Latina	18.7 (13.2, 24.1)	16.2 (9.7, 22.7)	18.0 (10.8, 25.1)	16.3 (10.0, 22.6)
British white	17.3 (13.8, 20.8)	15.4 (11.4, 19.3)	15.7 (11.2, 20.3)	14.1 (10.6, 17.5)
Non-British white	17.5 (13.4, 21.6)	13.6 (9.0, 18.2)	13.1 (8.3, 17.9)	16.8 (12.0, 21.5)
Any other/Unknown	17.1 (12.5, 21.7)	12.4 (7.6, 17.1)	13.1 (8.0, 18.3)	13.4 (8.7, 18.0)

Supplement S3


S3(a). Prevalence of probable depression, by gender and cohort.



S3(b). Prevalence of moderate-to-severe anxiety, by gender and cohort.



S3(c). Prevalence of lifetime self-harm, by gender and cohort.




CONFIDENTIAL



## Supplement S4

S4(a). Weighted mean internalising scores (and 95% CIs), by gender and cohort



S4(b). Weighted mean externalising scores (and 95% CIs), by gender and cohort

