



## King's Research Portal

*Document Version*  
Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Weber, C., Lupo, C., Tse, T., & Jamet, F. (2021). Maximally Localized Dynamical Quantum Embedding for Solving Many-Body Correlated Systems. *Nature Computational Science*.

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# 1 Maximally Localized Dynamical Quantum 2 Embedding for Solving Many-Body Correlated 3 Systems

4 Carla Lupo<sup>1,2,\*</sup>, François Jamet<sup>2,1,\*\*</sup>, Terence Tse<sup>1</sup>, Ivan Rungger<sup>2,++</sup>, and Cedric Weber<sup>1,+</sup>

5 <sup>1</sup>King's College London, Theory and Simulation of Condensed Matter, The Strand, WC2R 2LS London, UK

6 <sup>2</sup>National Physical Laboratory, Teddington, TW11 0LW, United Kingdom

7 \*carla.lupo@kcl.ac.uk

8 \*\*francois.jamet@npl.co.uk

9 ++ivan.rungger@npl.co.uk

10 +cedric.weber@kcl.ac.uk

11

12

## 13 ABSTRACT

14 Quantum computing opens new avenues for modelling correlated materials, notoriously challenging to solve due to the presence  
15 of large electronic correlations. Quantum embedding approaches, such as the dynamical mean-field theory, provide corrections  
16 to first-principles calculations for strongly correlated materials, which are poorly described at lower levels of theory. Such  
17 embedding approaches are computationally demanding on classical computing architectures, and hence remain restricted  
18 to small systems, which limits the scope of applicability. Hitherto, implementations on quantum computers are limited by  
19 hardware constraints. Here, we derive a compact representation, where the number of quantum states is reduced for a given  
20 system, while retaining a high level of accuracy. We benchmark our method for archetypal quantum states of matter that emerge  
21 due to electronic correlations, such as Kondo and Mott physics, both at equilibrium and for quenched systems. We implement  
22 this approach on a quantum emulator demonstrating a reduction of the required number of qubits.

## 23 Introduction

24 Correlated materials have attracted a widespread interest due to their broad range of complex properties and possible avenues  
25 for technological applications. Modelling correlated transition metal systems remains a challenge for standard approaches, such  
26 as first-principle based methods. With the discovery of high-temperature superconductivity, a large effort has been devoted  
27 to the study of archetypal theories for describing Mott insulators, leading to developments of theoretical tools for solving  
28 correlated materials accurately. An exact solution of the Hubbard model, which describes such correlated materials, is lacking  
29 in two or three dimensions. However, embedding approaches based on controllable approximations, such as the dynamical  
30 mean-field theory (DMFT)<sup>1,2</sup>, have provided accurate predictions<sup>3-5</sup>. DMFT is a non-perturbative approach that allowed  
31 to obtain the phase diagram of the Mott-Hubbard transition. The marriage of density functional theory (DFT) and DMFT  
32 (DFT+DMFT) provides the work-horse for studying correlated materials.

33 DMFT is based on a self-consistent mapping between local properties of a given material onto a so-called Anderson  
34 impurity model (AIM), a correlated impurity embedded in an infinite non-interacting bath. The AIM can be solved by high-level  
35 many-body methods, such as the continuous-time Monte Carlo approach (CTQMC). The latter provides an exact solution within  
36 statistical error bars, but is limited to the imaginary time representation. The evaluation of real-frequency spectral quantities  
37 requires the ill-defined analytical continuation<sup>6,7</sup>. Cluster extensions of DMFT suffer from the *fermionic sign-problem* when  
38 inter-orbital hybridizations are present. Other solvers are based on the numerical renormalization group (NRG)<sup>8</sup>, which allow  
39 real axis calculations and access to Kondo physics, but remain challenging to extend for multi-orbital systems. Another method  
40 that provides solutions for real frequencies is the exact diagonalisation (ED) approach, where a finite size discretization of  
41 the AIM is used, through representation of the infinite bath in terms of a small number of effective *bath-sites*. In typical  
42 implementations, the bath size ( $N_b$ ) is restricted because of the exponential growth of the Hilbert space with the total number of  
43 sites  $N_s$  (bath sites and impurity orbitals). Nonetheless, Lanczos-based algorithms allows to deal with large Hilbert spaces,  
44 where the discretization at low temperature<sup>9,10</sup> is fine enough to compute observables accurately. Some success has been

45 achieved by ED for multi-orbital systems with three or five orbitals<sup>11–13</sup>, but in general the limitation in the bath size limits the  
46 scope of applicability for realistically describing transition metal oxides. Towards the achievement of the largest number of bath  
47 sites, ED calculations have been extended to handle the single impurity embedding problem, allowing up to  $O(100)$ <sup>14–17</sup> and  
48  $O(300)$ <sup>18</sup> uncorrelated bath sites. Although the latter approaches allow the obtaining of approximations of the zero temperature  
49 Green’s function, the building of systematically high energy excited states remains challenging.

50 Moreover, scaling the precision of the latter approach on classical computers is not feasible due to the exponential increase  
51 of the Hilbert space with the number of bath sites. In view of the recent progress in the implementation of ED on quantum  
52 computers<sup>19</sup>, the latter approach has gained interest due to the possibility of scaling linearly on quantum computers. However,  
53 due to current hardware limitations (noise and decoherence issues), such algorithms are limited to a small number of qubits and  
54 short quantum circuits on the currently available noisy intermediate-scale quantum (NISQ) computers. Hence it is important to  
55 develop an ED based solver that can obtain precise results with a reduced number of bath sites, since the required number of  
56 qubits is proportional to this.

57 In this light, here we present the maximally localized dynamical embedding (MLDE) methodology to solve  
58 the AIM for DMFT, which extends the ED method by adding electron-electron interactions also in the bath. Our method  
59 provides a maximally localized quantum impurity model, where the non-local self-energy component of the correlation due to  
60 interactions in the bath remains minimal, and hence the AIM minimally breaks locality (DMFT is a purely local theory). MLDE  
61 provides the benefit that the environment used in the quantum embedding approach is described by propagating correlated  
62 electrons (instead of free electrons in DMFT). This firstly improves the hybridization fitting procedure compared to ED due to  
63 the polynomial increase of the number of fitting parameters with number of bath sites. Secondly, the number of poles in the  
64 MLDE hybridization function increases exponentially with the number of bath sites, compared to a linear increase for ED. This  
65 allows to obtain converged system properties with a much smaller number of bath sites than the one required by ED. This is  
66 reminiscent of the representation of correlated electrons by a Green’s function embedding approach, where correlations are  
67 described by hidden fictitious additional fermionic degrees of freedom<sup>20,21</sup>.

68 This representation has hence the potential to improve the scope of applicability of the quantum embedding approach, whilst  
69 limiting the small number of bath sites. We report that quantum impurity models with as few as 3 bath sites can reproduce  
70 both the Kondo regime and the Mott transition and obtain good agreement for dynamical magnetic susceptibilities, poising  
71 this approach as a candidate to describe 2-particle excitations such as excitons in correlated systems, with applications for  
72 high-temperature superconductors<sup>22,23</sup>. We also present a quantum computing algorithm for MLDE, and show that with a  
73 number of qubits, available on current hardware, one can achieve a fine description of correlations in materials.

## 74 Results

### 75 From the Anderson impurity model to maximally localized dynamical embedding

76 Within DMFT, the lattice model is mapped to an effective Anderson impurity model (AIM) where a correlated atom is connected  
77 to a non-interacting bath with the hybridization function  $\Delta(\omega)$  as shown in Fig 1.a.

78 In the ED approximation, the continuum bath is represented with a finite number of effective sites (Fig.1.b). Typically, for a  
79 fixed set of the Hamiltonian parameters (see Eq.1 Methods section), the AIM is solved (1) by using a Lanczos algorithm to  
80 converge the ground state and excited states<sup>9,12</sup> which contribute to the thermal average. Once the eigenstates are obtained, the  
81 dynamical and static observables are computed. As previously stated, the number of bath sites is severely limited because of  
82 the exponential scaling of the Hilbert space with the number of sites. To improve the performance of the algorithm, without  
83 increasing the number of bath sites, we introduced a two-body interaction between the bath electrons (Fig.1.c). The resulting  
84 approximation is represented by an extended ED solver where the non-local component of the correlation potential remain  
85 minimal. Thus the ED method has been extended to a maximally localized dynamical embedding (MLDE) model where the  
86 same accuracy is obtained with a reduced number of bath sites. (see Methods section for the formal mathematical definition of  
87 the model).

### 88 Comparison between MLDE and CTQMC solver

89 We now turn to a simple test of the MLDE equations for a typical AIM. The bath hybridization used as a test case is obtained  
90 from a typical correlated material (The hybridization function is available in Extended Data Figure 1), and the impurity  
91 energy is set to  $\epsilon_f = -U/2$  to stay at half-filling. We perform a benchmark of the MLDE approach with respect to both a  
92 continuous-time Monte Carlo and Lanczos solver, which both provide in this case the same answer used as a reference. Both  
93 the imaginary part (Fig.2.a) and real part (Fig.2.b) obtained by MLDE with as few as  $N_b = 3$  bath sites provide a very good  
94 agreement with the exact solution. We considered both the nearly free electron (NFE) limit ( $U < 10$ ) and the atomic limit  
95 ( $U > 10$ ), and in both cases the MLDE solution is consistently in agreement with the exact answer. Noteworthy, the strength  
96 of correlation and overall physical properties are also well captured by MLDE with  $N_b = 2$ , whereas the ED solver with the  
97 same number of bath sites largely overestimates the strength of correlation (see Extended Data Figure 2). MLDE captures well

98 the delocalization-localization transition (see Fig.2.b-c), where the MLDE with  $N_b = 2$  only differs near the transition while  
99 MLDE with  $N_b = 3$  is exact (Fig. 2.c).

## 100 Kondo physics

101 We extend further the application of MLDE to a realistic study case of the deposition of a correlated Kondo molecule on a gold  
102 surface (see Fig.3.a). Stable organic radical molecules exhibit a Kondo peak in the low-temperature experimental conductance,  
103 which is due to the presence of a single unpaired electron in the highest occupied molecular orbital<sup>24–26</sup>. In the gas phase  
104 these molecules are paramagnetic. Due to their low spin-orbit interaction and small hyperfine splitting, they are expected to  
105 exhibit long spin-coherence times, and therefore have potential as building blocks of molecular spintronics applications<sup>26</sup>.  
106 When brought in contact with a metal surface, the system corresponds to a single impurity Anderson model (SIAM) and has  
107 been modeled in the past using CTQMC or NRG as impurity solvers<sup>27,28</sup>. To demonstrate the capability of MLDE to describe  
108 Kondo physics, we choose the 1,3,5-triphenyl-6-oxoverdazyl (TOV) organic radical molecule, which, when deposited on a  
109 Au substrate has been shown experimentally to exhibit a Kondo temperature of about  $\approx 37\text{K}$ <sup>24</sup>. Compared to other radical  
110 molecules on surfaces, this molecule has the advantage to have a well-defined contact geometry, where the molecule lies  
111 flat on the surface. We use the same simulation setup and parameters as in a previous study<sup>27</sup> so that also the hybridization  
112 function of the SIAM is the same (see Extended Data Figure 5). Describing Kondo physics at low temperature is a notoriously  
113 difficult problem for quantum impurity solvers. In particular, the collapse of energy scales in the Kondo limit prevents typical  
114 Lanczos or ED solvers from capturing the Kondo resonance, as the finite discretization tends to introduce fictitious gaped states  
115 at very low temperature. We performed calculation at  $T = 5\text{K}$ . In the Kondo regime (see Fig.3.b) the hybridization remains  
116 constant up to the lowest frequency. We note that  $\Delta_1(i\omega_n) = \Delta_2(i\omega_n)$ , which confirms that the embedding source field  $\delta\Delta(i\omega_n)$   
117 remains negligible. Furthermore, MLDE fares better with  $N_b = 4$  than the best possible fit obtained by the usual discretized  
118 Lanczos approach ( $N_b = 12$ , triangles in Fig.3.b), as the hybridization vanishes at small frequencies (in the molecule, it remains  
119 constant). The imaginary part of the self-energy obtained by MLDE shows a Fermi liquid type behavior at small frequencies  
120 (see Fig.3.c), whereas the self-energy obtained by ED shows an artificial Mott singularity, due to the bath discretization. We  
121 note that this low temperature was beyond the reach of our CTQMC solver ( $\beta = 35000 \text{ Ryd}^{-1}$ ). Below the Kondo temperature,  
122 we recover the Fermi liquid behavior of the self-energy (see Fig.3.d). As the MLDE representation is compact, it opens a large  
123 degree of possible manipulation once the AIM is established. In particular, we extended the calculation to the time dynamics of  
124 the Kondo molecule with the Keldysh formalism<sup>29</sup> after a magnetic quench, where at time  $t_0 = 0$  the molecule is magnetically  
125 polarized along the  $e_z$  axis, and the external magnetic field released for  $t > t_0$ . The magnetic moment enters in a precession  
126 dynamics with the frequency equal to the Kondo temperature (see Fig.3.e). The estimated Kondo temperature( $T_K$ ) which we  
127 achieved with different methods, e.g. from the time dynamics ( $T_K \approx 33.64$ ) and from perturbation theory<sup>30,31</sup> ( $T_K \approx 33$ , see  
128 subsection MLDE for Kondo molecule in the Methods section), which provides a further check of the validity of the MLDE  
129 approach. The dynamics can also be resolved below  $T_K$ , where we observe additional harmonics, reminiscent from the Kondo  
130 Zeeman splitting at  $t = t_0$ <sup>32</sup>.

## 131 Mott transition

132 We have so far focused on simple AIM systems in the absence of mean-field corrections to the hybridization. We now turn to  
133 the dynamical mean-field MLDE approach, applied to the Mott transition where the local Green's function is defined with  
134 the density of state of the two-dimensional square-lattice Hubbard model (see Fig.4.a). Using the MLDE as a solver for  
135 DMFT, we recover the well-known metal-insulator transition (MIT) associated with the charge localization induced by the local  
136 Hubbard repulsion  $U$ . Here, the DMFT self-consistency is enforced in Matsubara frequency. We recover with a simple MLDE  
137 and  $N_b = 3$  bath sites the spectral function of the Hubbard model, with the usual features (lower and upper Hubbard bands,  
138 quasi-particle peak below the transition  $U_c \approx 9$ , Mott gap for  $U > U_c$ ). We note that interestingly the spectral function obtained  
139 by MLDE also shows the satellite peaks at the gap edge for  $U > U_c$ . This feature is typically difficult to obtain with analytically  
140 continued spectral function from Matsubara quantity. Indeed the inner peak near the Mott gap edge are associated with the real  
141 part of the self-energy, and in particular, is derived from an actual quasi-particle solution<sup>33</sup>.

142 The MLDE charge gap  $\Delta$  reproduces the known trend  $\Delta = U - W$ , which provides a further test of the theory. The benchmark  
143 with the converged solution obtained by CTQMC is accurate (see Fig.4.b), and the MLDE solution essentially within the error  
144 bars of the CTQMC for  $U/t = 12$ . Across the Mott transition, the agreement between the Green's functions remains good (see  
145 Fig.4.c).

## 146 Bethe-Salpeter Equation

147 We extended the calculations to the Bethe-Salpeter Equation (BSE) formalism, applied to the MLDE solution. In particular we  
148 calculate the local irreducible vertex  $\Gamma$  (see Section Vertex calculation in MLDE in the Methods section), which enables the  
149 calculation of the non-local dynamical magnetic susceptibility  $\chi^{mag}(\omega)$  within MLDE. We performed calculations for  $U/t = 12$   
150 at various temperatures, to explore the behavior of magnetic excitations across the Mott gap melting. Within the Mott gap

151 phase at temperature  $T/t = 0.025$  (Fig.5.a), the Neel fluctuations are local in momentum at  $Q_{\text{Neel}} = (\pi, \pi)$ . As expected, the  
152 spin fluctuations are largely located at  $Q_{\text{Neel}}$  in the Mott phase ( $T = 0.025t$ ). We observe that the antiferromagnetic magnetic  
153 fluctuations become gradually more incoherent as the temperature is increased throughout (Fig.5.b). Interestingly, we observe a  
154 large degree of incoherence at all considered temperature, which is reminiscent of the large scattering rate of the archetypal  
155 Mott system La2CuO4. At very large temperature (Fig.5.c), the fluctuations are fully incoherent, and the spectrum uniform  
156 across the Brillouin zone.

157 We note that a known challenge for the BSE approach is the extraction of the local irreducible vertex  $\Gamma$ , which is obtained  
158 by calculating the two-electron response function  $G^{(2)}$ . Such a quantity requires traditionally computationally demanding  
159 collection of statistics by CTQMC and few other alternatives exist. The vertex can be calculated in the Lehman representation  
160 (see Section Vertex calculation in MLDE in the Methods section), but requires sampling over the whole Hilbert space, which  
161 is not possible for AIM with more than few bath sites. In MLDE this task is largely simplified as the Hilbert space remains  
162 compact. We provide a benchmark of the local vertex  $\Gamma$  with CTQMC for both  $U/t = 6$  and  $U/t = 12$  (see respectively Figs.5.e  
163 and 5.g), in both cases the agreement is quantitative. This agreement is also obtained in the fermionic representation of the local  
164 magnetic susceptibility  $\chi_{iv,iv}$  (see Figs.5.d and 5.f). In conclusion, our MLDE solver can be integrated in ab-initio DFT+DMFT,  
165 where the irreducible vertex and the self-energy are the key quantities which are ultimately used to compute any observable.  
166 We ensure correctness of the approximation up to the two particle response function, while for higher order response functions  
167 the technique is limited as compared as ED.

### 168 MLDE for multiorbital system

169 So far, we have considered a single orbital case as a proof of concept of the formalism. To highlight the advantage of the  
170 MLDE solver, we now turn to the application to a multi-orbital case. The ED solver has been notoriously limited for the case of  
171 multi-orbital systems<sup>34</sup> as it would require a large number of bath sites, whereas CTQMC solver would be affected by the sign  
172 problem triggered by the non-zero off-diagonal terms of the hybridization between the orbitals. Therefore, in this section, we  
173 demonstrate the ability to treat multi-orbital systems within MLDE framework. As a test system, we applied DFT+DMFT on  
174 LaNiO<sub>3</sub>, whose nontrivial contributions stemming from electronic correlations have been studied using CTQMC solver<sup>35</sup>. In  
175 particular, The crystal field splits the Ni  $d$  orbital into two manifolds  $t_{2g}$  (composed of  $d_{xy}, d_{xz}, d_{yz}$  orbital) and  $e_g$  ( $d_{z^2}, d_{x^2-y^2}$ );  
176 here, the  $e_g$  are active orbitals and pose this compound as our ideal candidate of two orbitals system.

177 Our main focus is the comparison between a standard DFT+DMFT implementation using CTQMC and ED impurity solver  
178 with our MLDE environment with 3 bath sites. As shown in Fig 6a, a good agreement is achieved between CTQMC and MLDE  
179 when considering the local self energy for the first iteration of the DMFT loop.

180 We also show the local self energy obtained with ED using 3 bath sites. Notice that, differently from the results obtained by  
181 MLDE, the ED solver does not well capture the correct self energy behavior. A better agreement between the ED solver and the  
182 CTQMC would be achieved at the cost of adding more bath sites to increase the number of parameters and density of poles in  
183 the hybridization function required for the fit of the hybridization, as it is done in MLDE.

184 Lastly, we consider the local Green's function for the final iteration of the self-consistent DMFT loop. Notice that while we  
185 only include 3 sites in the bath, the MLDE Green's function is in a good agreement compared to CTQMC results.

### 186 MLDE on a quantum computer

187 Since MLDE reduces the number of bath sites compare to ED, it opens the opportunity to run Hamiltonian simulation on Noisy  
188 Intermediate-Scale Quantum computer(NISQ) with a reduced number of qubits. Here we develop a quantum algorithm to  
189 run MLDE on NISQ devices and demonstrate it using a quantum simulator on the benchmark system considered in Fig. 2.  
190 To calculate the Green's function on a quantum computer, we use the variational quantum eigensolver (VQE) based method  
191 presented by Rungger et al<sup>36</sup>, which was demonstrated to run on currently available hardware as it is rather resilient to noise.  
192 Details of the method are presented in the Sec. 5 of the method. Here we show results obtained with our implementation within  
193 the Quest quantum simulator<sup>37</sup>. To demonstrate that the quantum MLDE algorithm gives the same accuracy as the classical  
194 MLDE algorithm, we apply it on the same Anderson impurity model shown in Fig. 2. The system has one impurity site and  
195 two bath sites, which we map to a 6 qubit system using a Jordan-Wigner transform<sup>38</sup>. In Fig. 2.d we show that the self-energy  
196 for  $U = 6$  computed with the quantum computing algorithm agrees very very well with the result obtained using the classical  
197 computing algorithm. We have verified that the same is true for all values of  $U$ . This, therefore, demonstrates the functionality  
198 of our quantum algorithm for MLDE.

## 199 Discussion

200 We now turn to the discussion of MLDE and its advantages and limitations with respect to either other methods or known limits.  
201 Indeed, the MLDE non-local self energy vanishes, as expected, when: the local impurity correlation is naught ( $U_{imp} = 0$ ), the  
202 bath correlations are naught, or when the impurity-bath coupling is naught ( $V = 0$ ). In between the latter limits, we note that



203 MLDE is an approximation of the exact AIM self-energy, on the same ground as the standard ED technique. Although MLDE  
 204 mitigates to a large extent the bath discretisation error present in the standard ED approach, it introduces another error due to  
 205 non-local components in the self-energy between the impurity and the bath orbitals. MLDE and ED converge to the same exact  
 206 solution in the limit where the bath discretisation involves a large number of bath orbitals.

207 In this regard, we showed in this work that MLDE also performs optimally in the intermediate regime, where typically  
 208 emergent quantum state of matters occur, i.e. Kondo physics, Mott physics, and correlated Fermi liquids. MLDE always  
 209 performs better than ED (for the same number of bath sites) when the non local components in the self-energy are minimised.  
 210 The reason for the better performance are twofold: MLDE improves the hybridization fitting procedure compared to ED due to  
 211 the polynomial increase of the number of fitting parameters with number of bath sites, and the number of poles in the MLDE  
 212 hybridization function increases exponentially with the number of bath sites, compared to a linear increase for ED. This allows  
 213 to obtain converged system properties with a much smaller number of bath sites than the one required by ED. In MLDE,  
 214 the non-local self-energy provides a figure of merit of the obtained solution. A quantitative way to check that the MLDE is  
 215 converged to a good set of parameters is to systematically evaluate the self-energy for increasing number of bath sites, in the  
 216 same way as it is done for ED.

217 A further advantage is that MLDE can be applied in the realm of quantum computing on NISQ devices, where, with a  
 218 reduced number of required qubits, MLDE can allow the simulation of transition metal systems on a quantum computer, with  
 219 minor errors induced by the bath discretization. Note that on classical computers the computational cost increases exponentially  
 220 with system size, both for ED and MLDE, while quantum computers can potentially reduce this to a polynomial scaling<sup>39,40</sup>.  
 221 On near term quantum computers the noise limits the size of calculations that can be done in practice, so that only very small  
 222 proof of concept DMFT systems have been demonstrated on hardware<sup>36</sup>. Since MLDE significantly reduces the number of  
 223 sites needed to describe a given physical system, it allows to reduce the required number of qubits as well as the depth of the  
 224 quantum circuits. Therefore, in the near term, it is a potential framework to run larger physical systems on NISQ devices, while  
 225 in the long term, the reduction of the required number of sites will be beneficial also in the fault-tolerant quantum computing  
 226 regime.

## 227 Methods

### 228 Overview of the MLDE method

229 Within DMFT, the effective AIM is subject to a self-consistency condition which relates the Green's function of the impurity  
 230 model  $G(i\omega_n)$  to the so-called Weiss-field  $\mathcal{G}_0^{-1}(i\omega_n)$ , which completely characterizes the AIM. For the single-band case, the  
 231 Hamiltonian of the AIM reads:

$$H = \sum_{ij\sigma} \varepsilon_{ij\sigma} \hat{d}_{i\sigma}^\dagger \hat{d}_{j\sigma} + \sum_{i\sigma} V_i (\hat{d}_{i\sigma}^\dagger \hat{f}_\sigma + hc) + U \hat{n}_{f\uparrow} \hat{n}_{f\downarrow} + \sum_{\sigma} \varepsilon_f \hat{f}_{\sigma}^\dagger \hat{f}_{\sigma} \quad (1)$$

232 where  $\hat{d}_{p\sigma}^\dagger$  ( $\hat{d}_{p\sigma}$ ) creates (destroys) a particle with spin  $\sigma$  in the d-orbitals of the uncorrelated bath ( $p \in [1, N_b]$ ) and  $\hat{f}_{\sigma}^\dagger$  ( $\hat{f}_{\sigma}$ )  
 233 creates (destroys) a spin  $\sigma$  particle on the impurity,  $U$  is the static Coulomb repulsion on the impurity and  $V$  is the tunneling  
 234 amplitude between the impurity and the bath.

235 In the MLDE approach we add an additional general two-electron interaction to the bath sites, and the interaction vertex  
 236 reads:

$$H_{\text{int}} = \sum_{\sigma_1, \sigma_2} \sum_{i, j, k, l} U_{i, j, k, l}^{(2)} \hat{d}_{i\sigma_1}^\dagger \hat{d}_{j\sigma_2}^\dagger \hat{d}_{k\sigma_2} \hat{d}_{l\sigma_1} \quad (2)$$

The  $U^{(2)}$  tensor is a fictitious two-body interaction between bath electrons introduced to enlarge the number of fitting parameters  
 for a given number of bath sites, and also to enlarge the number of poles in the hybridization function. Although the free  
 electron propagator takes a simple polynomial form, e.g.  $G_0^{-1} = i\omega_n - \varepsilon_f - \sum_i V_i^2 / (i\omega_n - \varepsilon_i)$ , a correlated Green's function is  
 instead described by a mapping to an exponentially long one-dimensional chain within the Krylov space<sup>41</sup>. In particular, when  
 the bath is correlated, the Weiss field incorporates the dressed propagator of the bath electrons  $\tilde{G}_d^{(1)}$ :

$$G_0^{-1} = i\omega_n - \varepsilon_f - \underbrace{\mathbf{V}^\dagger \tilde{\mathbf{G}}_d^{(1)} \mathbf{V}}_{\Delta^{(1)}}, \quad (3)$$

where the hybridization to the bath is denoted as  $\Delta^{(1)}$ . We emphasize that at this level of the theory, the self energy of  
 the impurity  $\Sigma_f$  remains naught, and hence we have a fully local theory from the perspective of the impurity. However, as

we introduce a local correlation  $U$  on the impurity, two effects occur: i) the propagator of the bath  $\tilde{G}_d^{(1)}$  is dependent on the correlation on the impurity, and hence changes to  $\tilde{G}_d^{(2)}$  and ii) a non-local part of the self-energy  $\Sigma_{fd}$  between the impurity and the bath emerges:

$$G_{imp}^{-1} = i\omega_n - \varepsilon_f - \underbrace{(\mathbf{V} + \Sigma_{fd})^\dagger \tilde{\mathbf{G}}_d^{(2)} (\mathbf{V} + \Sigma_{fd})}_{\Delta^{(2)}} - \Sigma_f \quad (4)$$

We have now a set of embedding equations, which leads to a generalized Dyson equation

$$\mathbf{G}^{-1} - \mathbf{G}_0^{-1} = \delta\Delta - \Sigma_f, \quad (5)$$

where the  $\delta\Delta(z)$  is a source field that stems from the non-local correlations. Indeed, the latter term can be rationalized following a simple argument via the Migdal energy functional, which in MLDE reads

$$U\langle\hat{n}_\uparrow\hat{n}_\downarrow\rangle = \frac{1}{2} \left( Tr(\Sigma_f G_f) + \underbrace{Tr(\Sigma_{fd} \mathbf{G}_{fd})}_{\bar{Z}} \right) \quad (6)$$

233 Part of the correlation energy spills effectively on the bath<sup>42</sup>, leading to a correlation leakage term  $\bar{Z}$ .

234 The DMFT equations can be recovered when the leakage potential and leakage correlation energy are small. Note that  
 235 the obtained DMFT equations are for a self-energy with interactions in the bath, which differs from the one obtained without  
 236 interactions in the bath, and which only becomes equal to the exact self-energy for an infinitely large number of bath sites. As  
 237 the tuning of the bath propagator allows for a very large set of parameters, these constraints can be successfully enforced via  
 238 Lagrange parameters (see the cost function defined in Eq. 19) in the fit of the DMFT hybridization, to maximize the locality  
 239 of the embedding, with a concomitant exponential improvement of the bath discretization errors. It is known that in high  
 240 dimension, or when a system is strongly correlated, the electron self-energy is well separable into a local dynamical part and a  
 241 static non-local contribution<sup>43</sup>. In this respect, we reabsorb the embedding potential into a shift of the static part of the MLDE  
 242 self energy, which ensures that the Migdal energy remains exact.

## 243 Details of the MLDE derivation

244 In this section, we provide an extended derivation of the MLDE formalism briefly introduced in the previous section. To this  
 245 end, we start with reviewing the case of a non-correlated bath. We then proceed to introduce the quantities of interest in the  
 246 MLDE formalism, where bath interactions are included. We also provide the pseudo-code where the algorithm is described in  
 247 more detail.

### 248 Case of non correlated bath

In the case of the non correlated bath, shown in Fig. 1.a, the AIM is described by the Hamiltonian of Eq. 1. Then the free-electron propagator is

$$G_0^{-1}(i\omega_n) = i\omega_n - \varepsilon_f - \Delta(i\omega_n). \quad (7)$$

In the ED approximation, the continuum bath is represented with a finite number of effective sites. Hereby, we consider and AIM in a start geometry, where the hybridization  $\Delta(i\omega_n)$  is discretized as follows:

$$\Delta^{ED}(i\omega_n) = \sum_d \frac{V_d^2}{i\omega_n - \varepsilon_d} \quad (8)$$

In the case of correlated impurity,  $U \neq 0$ , the problems acquire a many-body nature, and the electron propagation is affected by the scattering events, therefore correlations between the electrons in the system. We then introduce the interacting propagator of the impurity of the AIM in the function of the hopping terms  $V_d$  between impurity and bath sites and onsite energies  $\varepsilon_d$  of the bath sites:

$$(G^{ED})_{imp}^{-1}(i\omega_n) = i\omega_n - \varepsilon_f - \Delta^{ED}(i\omega_n) - \Sigma^{ED}(i\omega_n) \quad (9)$$

The Dyson equation for the impurity reads:

$$(G^{ED})_{imp}^{-1}(z) = G_0^{-1}(z) - \Sigma_f^{ED}(z) \quad (10)$$

For our convenience, we report the Green's function and the self-energy  $\Sigma$  of the problem, in a matrix formalism Hence:

$$\Sigma^{ED} = \left( \begin{array}{c|c} \Sigma_f^{ED} & 0 \\ \hline 0 & 0 \end{array} \right) \quad \text{and} \quad (\mathbf{G}^{ED})^{-1} = \left( \begin{array}{c|c} (i\omega_n - \varepsilon_f - \Sigma_f^{ED}) & -\mathbf{V} \\ \hline -\mathbf{V} & (i\omega_n - \varepsilon_d) \end{array} \right) \quad (11)$$

249 Notice that their dimension is  $N_b + 1$ , with  $N_b$  being the number of bath sites.

250 **Case of correlated bath**

In the maximally localized dynamical embedding approach (MLDE), we extend the formalism of the ED approximation to include correlations in the bath sites, adding a two-body interaction term of Eq. 2 in the Hamiltonian. Hence, the MLDE Hamiltonian reads as:

$$H = \sum_{ij\sigma} \varepsilon_{ij\sigma} \hat{d}_{i\sigma}^\dagger \hat{d}_{j\sigma} + \sum_{i\sigma} V_i (\hat{d}_{i\sigma}^\dagger \hat{f}_\sigma + hc) + U \hat{n}_{f\uparrow} \hat{n}_{f\downarrow} + \sum_{\sigma} \varepsilon_f \hat{f}_\sigma^\dagger \hat{f}_\sigma + \sum_{\sigma_1, \sigma_2} \sum_{i,j,k,l} U_{i,j,k,l}^{(2)} \hat{d}_{i\sigma_1}^\dagger \hat{d}_{j\sigma_2}^\dagger \hat{d}_{k\sigma_2} \hat{d}_{l\sigma_1} \quad (12)$$

251 We emphasize that this formalism can be extended further by the addition of higher order terms (e.g. if a three-rank tensor is  
 252 introduced, the number of parameters scales at most as  $N_b^6$ ) being limited only by the dimension of the Hilbert space. Therefore  
 253 our method introduces a great advantage over the ED solver, where the number of fitting parameters scales only as  $N_b^2$ , and  
 254 therefore a larger number of bath sites is needed for the increase of the accuracy.

255 Consistent with the previous section, we start from the definition of the free-electron propagator in Eq. 3, where the  
 256 hybridisation to the bath is denoted as  $\Delta^{(1)}$  and includes a Green's function for the bath  $\tilde{G}_d^{(1)}$  which is interacting rather than  
 257 being a free-electron propagator as in Eq. 8 (more details in the pseudocode **Algorithm 1**). We emphasise that in case of  
 258 non correlated bath the number of poles in the hybridization scales linearly with  $N_b$ , while the inclusion of a correlated bath  
 259 introduces an exponential scaling of the number of poles in the hybridization function with  $N_b$ . Notice that at this level of the  
 260 theory, the self-energy of the impurity  $\Sigma_f$  remains nought, and hence we have a fully local theory from the perspective of the  
 261 impurity.

262 We now turn to the case of non-zero correlation on the impurity where the interacting propagator on the bath  $G_d^{(2)}$  is affected by  
 263 the correlation in the impurity and a non-local part of the self-energy  $\Sigma_{fd}$  between the impurity and the bath emerges.

264 To better clarify the formalism introduced in Eq.(4) so far, we recall the Green's function and the self-energy in their matrix  
 265 form. Hence:

$$\mathbf{G}(i\omega_n) = \left( \begin{array}{c|c} G_f(i\omega_n) & \mathbf{G}_{fd}(i\omega_n) \\ \hline \mathbf{G}_{df}(i\omega_n) & \mathbf{G}_d(i\omega_n) \end{array} \right) \quad \text{and} \quad \mathbf{\Sigma}(i\omega_n) = \left( \begin{array}{c|c} \Sigma_f(i\omega_n) & \mathbf{\Sigma}_{fd}(i\omega_n) \\ \hline \mathbf{\Sigma}_{df}(i\omega_n) & \mathbf{\Sigma}_d(i\omega_n) \end{array} \right) \quad (13)$$

Notice that to simplify the notation, the repeated indices are dropped. Therefore  $\mathbf{G}_{dd} = \mathbf{G}_d$ . A comparison with the matrices defined in the case of non-correlated bath, provides further clarification on the consequences of introducing interactions in the bath sites. The block of the self-energy matrix related to the bath has now an non-zero contribution by definition of our model. A less trivial observation concern the off-diagonal terms which are non zero, due to the interaction leaked in the bonds between impurity and bath sites. Therefore we can distinguish different contribution to the correlation energy, and in terms of Migdal energy functional, it reads:

$$E_U^{tot} = \frac{2}{\beta} \sum_{i\omega_n} \text{Tr} \left[ \mathbf{G}^\dagger(i\omega_n) \left( \mathbf{\Sigma}^\dagger(i\omega_n) - \mathbf{\Sigma}^\dagger(\infty) \right) \right] + \frac{2}{\beta} \text{Tr} \left( \boldsymbol{\rho}^\dagger \mathbf{\Sigma}^\dagger(\infty) \right) \quad (14)$$

$$E_U^{bath} = \frac{2}{\beta} \sum_{i\omega_n} \text{Tr} \left[ \mathbf{G}_d^\dagger(i\omega_n) \left( \mathbf{\Sigma}^\dagger(i\omega_n) - \mathbf{\Sigma}^\dagger(\infty) \right)_d \right] + \frac{2}{\beta} \text{Tr} \left( \boldsymbol{\rho}_d^\dagger \mathbf{\Sigma}_d^\dagger(\infty) \right) \quad (15)$$

$$E_U^{imp} = \frac{2}{\beta} \sum_{i\omega_n} G_f^\dagger(i\omega_n) \left( \Sigma_f^\dagger(i\omega_n) - \Sigma_f^\dagger(\infty) \right) + \frac{2}{\beta} \boldsymbol{\rho}_f^\dagger \Sigma_f^\dagger(\infty) \quad (16)$$

where  $\boldsymbol{\rho}$  is the density matrix. Part of the correlation energy spills effectively on the bath<sup>44</sup>, leading to a correlation leakage term, which is defined dynamically

$$z_U^{int}(i\omega_n) = \frac{(\mathbf{\Sigma}_{fd}(i\omega_n) \cdot \mathbf{G}_{fd}(i\omega_n))^2}{|i\omega_n|} \quad (17)$$

and statically as a result of the average over matsubara frequencies:

$$E_U^{leakage} = \frac{2}{\rho_f} \left( E_U^{tot} - E_U^{bath} - E_U^{imp} \right) \quad (18)$$

The locality principle behind the DMFT equations can be recovered when the leakage potential  $\delta\Delta(z)$  and leakage correlation energy are small. We clarify that the leakage field  $\tilde{Z}$  defined in Eq. 6 is  $\tilde{Z} = E_U^{leakage} R_f/2$ . As the tuning of the bath propagator allows for a very large set of parameters, these constraints can be successfully enforced via Lagrange parameters in the fit of the



DMFT hybridization, to maximize the locality of the embedding, with a concomitant exponential improvement of the bath discretization errors. To that end, the cost function of the minimization procedure is defined as:

$$d = \sum_{i\omega_n} \frac{|\Delta^{(1)} - \Delta_{target}|^2 + |\Delta^{(2)} - \Delta_{target}|^2 + \alpha_3 |\Delta^{(1)} - \Delta^{(2)}|^2}{|i\omega_n|^{\alpha_1}} + \lambda_s |E_U^{leakage}| + \lambda_d \sum_{i\omega_n} z_U^{int}(i\omega_n) \quad (19)$$

where the parameters  $\alpha_1, \alpha_3, \lambda_d, \lambda_s$  are properly tuned according to the case of study to improve the minimisation procedure and  $\Delta_{target}$  is the hybridisation of the continuous bath. We noticed that in most of the cases, the leakage field  $E_U^{leakage}$  turns out to be naturally small. Indeed this follows from the constraint that  $|\Delta^{(1)} - \Delta^{(2)}|$  has to be minimal. Therefore the fitting process tries to reduce the leakage by construction. It is worth it to notice, however, that all the terms in the cost function are important to give more flexibility to the minimisation process and avoid it to get stuck into local minima. Moreover, the first and the last term allow to have a better control on the low frequencies contributions and increase the precision of the description of phenomena occurring at a low energy scale, e.g. Kondo physics.

To ensure that the Migdal energy remains exact, we reabsorb the leakage field into a shift of the static part of the MLDE self-energy. This is an allowed procedure as it is known that in high dimension, or when a system is strongly correlated, the electron self-energy is well separable into a local dynamical part and a static non-local contribution<sup>43</sup>.

To summarise, in the MLDE formalism, non-local correlations are introduced to enlarge the parameters for the fitting of the parameters. This represents an improvement over existent ED formalism, as the increased number of fitting parameters allow the use of less bath sites. The non-locality is kept small through a Lagrange parameter in the cost function. To recover consistency with MLDE, the leaked (non-local) correlated energy is reabsorbed in the impurity self-energy. More details on the different steps of the algorithm are shown in the pseudo-code (**Algorithm 1**) reported below.

---

#### Algorithm 1 MLDE solver

---

inputs:  $\Delta_{target}, \epsilon_f, \alpha, \alpha_3, \lambda_s, \lambda_d, U, N_b$

**Require:**  $U_{imp} = 0$

**call solver:** input:  $\Delta_{target}$ , output  $\epsilon^{(1)}, \mathbf{V}^{(1)}, \mathbf{G}^{(1)}$

$$\Sigma^{(1)} \leftarrow i\omega_n \mathbf{I} - \epsilon^{(1)} - \left(\mathbf{G}^{(1)}\right)^{-1} \quad (\text{Dyson Equation})$$

$$\tilde{\mathbf{G}}_d^{(1)} \leftarrow (i\omega_n \mathbf{I})^{-1} - \epsilon_d^{(1)} - \Sigma_d^{(1)}$$

$$\Delta^{(1)} \leftarrow \mathbf{V}^{(1)\dagger} \tilde{\mathbf{G}}_d^{(1)} \mathbf{V}^{(1)}$$

$$d_1 \leftarrow \sum_n \frac{|\Delta^{(1)} - \Delta_{target}|^2}{|i\omega_n|^{\alpha}}$$

**Require:**  $U_{imp} \neq 0$

**call solver:** input:  $\Delta_{target}$ , output  $\epsilon^{(2)}, \mathbf{V}^{(2)}, \mathbf{G}^{(2)}$

$$\Sigma^{(2)} \leftarrow i\omega_n \mathbf{I} - \epsilon^{(2)} - \left(\mathbf{G}^{(2)}\right)^{-1} \quad (\text{Dyson Equation})$$

$$\Sigma_{int}^{(2)} \leftarrow -\mathbf{V}^{(2)} - \Sigma_{fd}^{(2)}$$

$$\tilde{\mathbf{G}}_d^{(2)} \leftarrow \left(-\epsilon_d^{(2)} - \Sigma_d^{(2)} + i\omega_n \mathbf{I}\right)^{-1}$$

$$\Delta^{(2)} \leftarrow \Sigma_{int}^{(2)\dagger} \tilde{\mathbf{G}}_d^{(2)} \Sigma_{int}^{(2)}$$

compute  $E_U^{tot}, E_U^{bath}, E_U^{imp}$  and  $E_U^{leakage}$

$$d_2 \leftarrow \sum_n \frac{|\Delta^{(2)} - \Delta_{target}|^2}{|i\omega_n|^{\alpha}}$$

$$d_3 \leftarrow \sum_n \frac{|\Delta^{(2)} - \Delta^{(1)}|^2}{|i\omega_n|^{\alpha}}$$

$$d_4 \leftarrow |E_U^{leakage}|$$

$$d_5 \leftarrow \sum_{i\omega_n} |z_U^{int}(i\omega_n)|$$

$$d \leftarrow d_1 + d_2 + \alpha_3 d_3 + d_4 \lambda_s + d_5 \lambda_d$$

$$\Sigma_f \leftarrow \Sigma_f + E_U^{leakage}$$


---

#### MLDE error scaling

As we stated previously, we find that in our calculations the non-local self energy contribution remains small. Additionally, we note that the non-local self energy vanishes in known limits of the theory: when the local impurity correlation is naught

284 ( $U_{imp} = 0$ ), when the bath correlations are naught ( $U_b = 0$ ) (Here for simplicity we denote as  $U_b$  the bath fictitious interactions),  
 285 or when the impurity-bath coupling is naught ( $V = 0$ ). In any of these limits, the non-local self energy vanishes and the MLDE  
 286 distance measure is optimal. The first two cases are identical in nature, as they are mirror of each others, and both corresponds  
 287 to the usual DMFT implementation where a correlated sub-system is embedded within a non-correlated one. We now turn the  
 288 discussion to the case of small  $V$  coupling within MLDE. To address this scenario, we consider the smallest possible example  
 289 system, where the local density of states of the impurity is identical in a free electron bath and in a correlated bath:

- 290 • **First system (with bath interactions):** the impurity is connected to a single interacting bath orbital with local interaction  
 291  $U_b$ . The green's function of the interacting bath has poles at  $-U_b/2$  and  $U_b/2$  (we stay at half-filling).
- 292 • **Second system (no bath interactions):** the impurity is coupled to two non-interacting bath orbitals with energies  
 293  $-U_b/2, U_b/2$

294 In both cases, the local density of states of the bath is set to half-filling (see Extended Data Figure 3.a). In Fig. S1b, we  
 295 report the MLDE scaling error as a function of leakage for different impurity to bath coupling  $V$ . As mentioned above and  
 296 expected, for  $V = 0$  MLDE is exact and the self energies of both approaches identical. Upon increasing  $V$ , we compare the  
 297 self energies obtained by both methods, as MLDE is an approximation in this limit, and compare the error obtained in the self  
 298 energy with the leakage spilling. As shown in Extended Data Figure 3.b, in this simple limit the MLDE error obtained in the  
 299 self energy correlates with the leakage. This is an illustration of the scaling of the MLDE error in a simple known limit, near  
 300 the limit where MLDE is exact (weak bath correlations or weak impurity-bath coupling).

### 301 Benchmark in real frequency

302 Previously, we preferred to benchmark the MLDE with ED and CTQMC, in the Matsubara formalism, rather than in real  
 303 frequency. Indeed, the spectral functions are notoriously known for being a poor benchmark when using the continuous fraction  
 304 method, used for the benchmark ED solver (as the number of bath sites is large, Lehmann representation of the GF is not  
 305 achievable). In particular, reliable benchmarks can be obtained via the Keldysh formalism and Fourier transforming the time  
 306 evolution to real frequency<sup>45</sup>, but require extensive computing resources for the large systems used as the benchmark. In this  
 307 regard, we compared the spectral functions obtained by the MLDE for the Mott transitions with the one obtained by ED (see  
 308 Extended Data Figure 4). However, a comparison in the real axis is less reliable due to the continuous fraction. Moreover, the  
 309 ill-conditioned maxent approach prevents us from detailed comparison on the real axis between CTQMC and MLDE. Therefore,  
 310 we have benchmarked the different solvers in the Matsubara formalism.

### 311 Vertex calculations in MLDE

In this section, we provide further insights on the application of MLDE to the calculation of two particle Green's function  
 quantities. In particular, the dynamical susceptibility fermionic matrix is computed by inverting the Bethe-Salpeter Equation  
 (BSE)

$$\chi(q, i\Omega_n)_{(iv\sigma)(iv'\sigma')} = [(\chi^0(q, i\Omega_n)^{-1} + \Gamma^{imp}(i\Omega_n))]_{(iv\sigma)(iv'\sigma')}^{-1}, \quad (20)$$

where :

$$\chi(q, i\Omega)_{(iv\sigma)(iv'\sigma')}^0 = -\delta_{iviv'} \delta_{\sigma\sigma'} \sum_k g_{k\sigma}(iv) g_{k+q, \sigma'}(iv + i\Omega), \quad (21)$$

and  $\Gamma^{imp}$  is the impurity vertex obtained by inverting the BSE for the MLDE impurity problem. The impurity susceptibility  
 reads:

$$G_{(iv\sigma)(iv'\sigma')}^{(2)imp}(i\Omega) = \int_0^\beta d\tau_1 d\tau_2 d\tau_3 d\tau_4 e^{iv(\tau_1 - \tau_2) + iv'(\tau_3 - \tau_4) + i\Omega(\tau_1 - \tau_4)} \left\langle \mathcal{T}_\tau c_\sigma^\dagger(\tau_1) c_\sigma(\tau_2) c_{\sigma'}^\dagger(\tau_3) c_{\sigma'}(\tau_4) \right\rangle \quad (22)$$

$$\chi_{(iv\sigma)(iv'\sigma')}^{imp}(i\Omega) = G_{(iv\sigma)(iv'\sigma')}^{(2)imp}(i\Omega) - \beta \delta_{0\Omega} G(iv) G(iv'). \quad (23)$$

In our work, we compute the two-particle response function  $G^2$  via the Lehman representation<sup>46,47</sup>:

$$G_{iviv'}^{(2)imp}(i\Omega) = \frac{1}{Z} \sum_{ijkl} \sum_{\Pi} \Phi(E_i, E_j, E_k, E_l; \omega_{\Pi_1}, \omega_{\Pi_2}, \omega_{\Pi_3}) \text{sign}(\Pi) \langle i | \mathcal{O}_{\Pi_1} | j \rangle \langle j | \mathcal{O}_{\Pi_2} | k \rangle \langle k | \mathcal{O}_{\Pi_3} | l \rangle \langle l | c_\sigma | i \rangle \quad (24)$$

$$\Phi(E_i, E_j, E_k, E_l; \omega_1, \omega_2, \omega_3) = \int_0^\beta d\tau_1 \int_0^{\tau_1} d\tau_2 \int_0^{\tau_2} d\tau_3 e^{-\beta E_i + \tau_1(E_j - E_i) + \tau_2(E_k - E_j) + \tau_3(E_l - E_k)} e^{i(\omega_1 \tau_1 + \omega_2 \tau_2 + \omega_3 \tau_3)}, \quad (25)$$

312 where we have introduced the closure relation, and define  $O_1 = c_\sigma$ ,  $O_2 = c_\sigma^\dagger$  and  $O_3 = c_{\sigma'}$ .  $E_i$  are the eigenvalues of the MLDE  
 313 hamiltonian,  $\Pi$  are the triplet permutations, and the summation holds over all states of the Hilbert space.

The magnetic susceptibility is obtained as

$$\chi^{mag}(q, i\Omega_n) = \frac{1}{2} \sum_{\sigma} \chi(q, i\Omega_n)_{(iv, \sigma)(iv', \sigma)} - \chi(q, i\Omega_n)_{(iv, \sigma)(iv', -\sigma)}. \quad (26)$$

314 In principle, the minimization of the single particle leakage does not guarantee the minimization of the leakage for the two  
 315 particle irreducible vertex. For the case we have studied, even without imposing a constraint on the two particle leakage, the  
 316 agreement with CTQMC is good. We propose here a diagrammatic argument to support our findings, in particular, that MLDE  
 317 also provides a good estimate of higher order response functions.

Firstly, the lowest order diagram contributing to the irreducible vertex is given by (see Ref.<sup>48</sup>):

$$\Gamma_{abcd}^{(1)} = U_{abcd}. \quad (27)$$

Since in MLDE, there are no direct many-body interaction terms in the Hamiltonian between the impurity and the bath,  
 this diagram is equal to zero by construction for the cross term contribution. Furthermore, the MLDE non-local self-energy  
 connecting the bath and the impurity can be expressed in term of the reducible vertex<sup>49</sup>:

$$\Sigma(14) = -U(12'3'1')G(1'4')G(23')G(2'3)F(4'234) \quad (28)$$

where  $F$  is the reducible vertex, which is related to the irreducible vertex by the Bethe-Salpeter equation

$$F(1234) = \Gamma(1234) + \Gamma(122'3')\chi^0(3'2'4'3')F(3'4'34). \quad (29)$$

318 Here we use the usual short-handed notations  $1 = (\tau_1, l_1, \sigma_1)$  and assume sum over repeated indices. We now consider the  
 319 case of  $\Sigma(14)$  corresponding to the non-local self-energy that connects the impurity to the bath degrees of freedom. When the  
 320 left-hand side of Eq. 28 is minimized in MLDE (reduction of the one-particle leakage), it naturally imposes that  $F$  remains  
 321 small (for non-local contributions). Moreover, we see from Eq. 29 that minimizing  $F$  also imposes a minimization of the  
 322 non-local terms in  $\Gamma$ , that connects the impurity to the bath. In summary, when the non-local self-energy is minimal, MLDE  
 323 provides a good approximation of the irreducible vertex used throughout to calculate the dynamical magnetic susceptibility.

324 Finally, we emphasize that it is usually difficult to provide energy cutoffs on the low energy converged states of the  
 325 Hamiltonian (typically in Lanczos or other iterative approaches), as the Boltzmann statistics only truncates one of the sums (on  
 326 index  $i$ ), whereas all eigenstates are required for the other summation. MLDE provides an ideal candidate for such calculations,  
 327 as the number of bath sites can be kept small with very little cost in accuracy. We note indeed that the previous equation can  
 328 be arranged to show a computational complexity growing with the cube of the number of eigenstates  $n$ , where  $n$  itself grows  
 329 exponentially with the number of bath sites.

### 330 MLDE for Kondo molecule

The density of states (DOS) for different temperatures is shown in Extended Data Figure 6a. The fingerprint Kondo peak around  
 zero energy can be seen at low temperatures. We emphasize that this feature is notoriously hard to obtain when using an ED  
 solver as the formalism is affected by the small broadening factor needed to correctly capture this Kondo physics fingerprint.  
 The Kondo temperature can be obtained from the relation<sup>30,31</sup> obtained from perturbation theory, which reads

$$T_K = -\frac{\pi Z}{4} \Im \Delta(i\omega_n) |_{\omega_n \rightarrow 0} \quad (30)$$

331 With the computed quasi-particle weight ( $Z = 0.026$ ) and value of the hybridization in the limit of zero frequency (see Extended  
 332 Data Figure 6.b), we obtain a Kondo temperature of  $T = 33.8$  K, which is comparable with the experimental one at  $\approx 37$ K<sup>24</sup>.  
 333 We notice that temperature effect corrections to the Fermi liquid regime can be more easily studied in the Matsubara self-energy  
 334 (see Extended Data Figure 6.b). In this representation, the quadratic behavior of the self-energy is apparent at low energy scales.

### 335 DFT+DMFT for LaNiO<sub>2</sub> : Extension of MLDE to multiorbital

336 DFT+DMFT calculation is performed within Questaal package<sup>50</sup>. We used the local density approximation (LDA) functional  
 337 with a k-mesh 10x10x10 points. The DMFT subspace is defined by the projection on Ni-d orbitals. The filled  $t_{2g}$  manifold is  
 338 treated at the Hartree Fock level, and the  $e_g$  is treated using DMFT. In order to solve the Anderson impurity model composed  
 339 of  $e_g$  orbital, we use MLDE with 2 impurity sites and 3 sites in the bath. The comparison with an exact solver is done using

340 CTQMC solver in TRIQS Package<sup>51</sup>. The interaction between the 2  $e_g$  orbitals is given by the rotationally invariant Slater  
 341 Hamiltonian

$$H_{int} = \frac{1}{2} \sum_{ijkl, \sigma\sigma'} U_{ijkl} c_{i\sigma}^\dagger c_{j\sigma'}^\dagger c_{k\sigma'} c_{l\sigma} \quad (31)$$

The tensor  $U_{ijkl}$  is given in term of Slater integral

$$U_{ijkl} = \sum_{k=0}^2 \alpha_k^{ijkl} F^{2k} \quad (32)$$

342 Where  $\alpha_k^{ijkl}$  are the Racah-Wigner numbers and  $F_k$  are the Slater integrals.

343 In the literature, we often use the parameter corresponding to the Hubbard interaction  $U$  and Hund's coupling  $J$  to express  
 344  $F^k$  with the relations

$$U = F^0 \quad (33)$$

$$J = \frac{F^2 + F^4}{14} \quad (34)$$

$$\frac{F^4}{F^2} = 10/16 \quad (35)$$

345 In this letter, we use the parameters  $U = 7$  and  $J = 1$  as suggested in<sup>35</sup>.

### 346 MLDE quantum computing algorithm

347 In this section, we present a quantum algorithm to run MLDE on noisy intermediate-scale quantum (NISQ) computers. We  
 348 perform a Jordan-Wigner (JW) transformation<sup>38</sup> to map the MLDE Hamiltonian (Eq. 12) to a qubit representation. After  
 349 the transformation, the Hamiltonian is in the form of a linear combination of Pauli tensors. With the JW transformation, the  
 350 required number of qubits is two times the number of sites in the system, the factor two being due to spin.

To compute the Green's function, we use a variational quantum eigensolver (VQE) based approach as presented in<sup>36</sup> since  
 it is rather resilient to noise. The quality of the quantum computing solution corresponds largely to the ability of the quantum  
 circuit to represent the eigenstates of  $H$ , and hence obtain accurate energies and amplitudes of the peaks in the Green's function.  
 One, therefore, needs a state preparation ansatz able to accurately represent the ground state as well as the excited states with  
 one electron added or removed<sup>36</sup>. For the results presented in the Results section, we use a so-called Hamiltonian variational  
 ansatz (HVA)<sup>52</sup> to represent the eigenstates of  $H$ . Note that we obtain a similar accuracy also with different types of circuit  
 ansatz, such as a so-called hardware efficient ansatz<sup>53</sup>. A general quantum state  $|\psi(\boldsymbol{\theta})\rangle$  is prepared with a circuit ansatz as  
 $|\psi(\boldsymbol{\theta})\rangle = \hat{U}(\boldsymbol{\theta}) |0\rangle$ , where  $|0\rangle$  is the initial state corresponding to all qubits set to the zero state, and  $\hat{U}(\boldsymbol{\theta})$  is the unitary generated  
 by the state preparation quantum circuit. Here  $\boldsymbol{\theta}$  is a vector of parameters, which determine the specific state generated by the  
 unitary. For the HVA  $U(\boldsymbol{\theta})$  has the form

$$\hat{U}(\boldsymbol{\theta}) = \prod_k^{n_{\text{layers}}} \prod_j e^{-i\theta_j^k \hat{P}_j} \quad (36)$$

351 where  $P_j$  is a Pauli tensor in  $H$  after JW transformation and  $\theta_j^k$  a real-valued parameter. The multiplication over  $j$  goes over  
 352 all Pauli terms in the Hamiltonian. The ansatz corresponds to  $n_{\text{layers}}$  repetitions of individual blocks. In a given block, each  
 353 Pauli tensor term in  $H$ ,  $P_j$ , is included in the ansatz as imaginary exponentiation with the parameter  $\theta_j^k$ , which can be readily  
 354 implemented on a quantum computer<sup>52</sup>. The accuracy of the ansatz generally improves with increasing  $n_{\text{layers}}$ . We verified that  
 355 for the system considered in the section MLDE on a quantum computer,  $n_{\text{layers}} = 2$  gives good accuracy.

To obtain the state preparation circuit parameters for the ground state, we minimize the cost function corresponding to the  
 ground state energy

$$E_{\text{GS}} = \min_{\boldsymbol{\theta}} E(\boldsymbol{\theta}) = \min_{\boldsymbol{\theta}} \langle \psi(\boldsymbol{\theta}) | H | \psi(\boldsymbol{\theta}) \rangle. \quad (37)$$

To obtain the state preparation circuit parameters for the excited states there are several possibilities<sup>54-56</sup>. Here we use the  
 method proposed in<sup>55</sup>, where a penalty term is added to the Hamiltonian to impose orthogonality of each excited state to the

lower energy states. To calculate the  $l^{\text{th}}$  excited state,  $|\psi_l\rangle = |\psi(\boldsymbol{\theta}_l)\rangle$ , with energy  $E_l$ , one therefore minimises

$$E_l = \min_{\boldsymbol{\theta}} \left[ \langle \psi(\boldsymbol{\theta}) | H | \psi(\boldsymbol{\theta}) \rangle + \sum_{j=0}^{l-1} \alpha_j \langle \psi_j | \psi(\boldsymbol{\theta}) \rangle^2 \right]. \quad (38)$$

356 Here the positive-valued  $\alpha_j$  are arbitrary parameters, set in a way to optimize the convergence of the algorithm. Once all  
357 required state preparation parameters  $\boldsymbol{\theta}_l$  are computed, the Green's function is obtained using the Lehman representation, and  
358 with it, the self-energy<sup>36</sup>.

## 359 Data availability

360 Source data and code to generate the data for all the figures are available via Code Ocean<sup>57</sup>.

## 361 Code availability

362 The code is available via Code Ocean<sup>57</sup>.

## 363 Acknowledgements

364 CW acknowledges insightful and stimulating discussions with Andrew Mitchell. CW is supported by grant EP/R02992X/1 from  
365 the UK Engineering and Physical Sciences Research Council (EPSRC). CL is supported by the EPSRC Centre for Doctoral  
366 Training in Cross-Disciplinary Approaches to Non-Equilibrium Systems (CANES, EP/L015854/1). IR and FJ acknowledge  
367 the support of the UK government department for Business, Energy and Industrial Strategy through the UK national quantum  
368 technologies programme (InnovateUK Industrial Strategy Challenge Fund (ISCF) QUANTIFI project). FJ was also partly  
369 supported by the Simons Many-Electron Collaboration and by the EPSRC Centre for Doctoral Training in Cross-Disciplinary  
370 Approaches to Non-Equilibrium Systems (CANES, EP/L015854/1). This work was performed using resources provided by the  
371 ARCHER UK National Supercomputing Service and the Cambridge Service for Data Driven Discovery (CSD3) operated by the  
372 University of Cambridge Research Computing Service (www.csd3.cam.ac.uk), provided by Dell EMC and Intel using Tier-2  
373 funding from the Engineering and Physical Sciences Research Council (capital grant EP/P020259/1), and DiRAC funding from  
374 the Science and Technology Facilities Council (www.dirac.ac.uk).

## 375 Author contributions statement

376 CW designed the research. CL and FJ equally contributed to the work. IR and FJ developed the quantum computing algorithm.  
377 All authors contributed to the code design, results production and analysis. All authors reviewed the manuscript.

## 378 Competing Interests Statement

379 The authors declare no competing interests.

## 380 Corresponding authors

381 carla.lupo@kcl.ac.uk  
382 francois.jamet@npl.co.uk  
383 ivan.rungger@npl.co.uk  
384 cedric.weber@kcl.ac.uk

**Figure 1. From AIM to MLDE.** (a) Mapping of the lattice model (with inter-site hopping parameter  $t$  and on-site correlation  $U$ ) onto a local impurity model, with a correlated atom (pink sphere) embedded in a non-interacting bath (blue shaded area) with energy  $\varepsilon$  as for the Anderson impurity model (AIM). The four possible configurations describe the quantum evolution of the atom. Electrons may hop from the atom to the bath via the frequency dependent hybridization function  $\Delta(\omega)$ , which plays the role of a dynamical mean field. b) Discretization of the continuum bath in non-interacting bath sites coupled to the interacting impurity through the hopping function  $V_i$ . This is the picture related to the ED impurity solver. c) Cartoon related to the Maximally Localised Dynamical Embedding (MLDE) solver where an interacting impurity is coupled to interacting bath sites.



**Figure 2. MLDE benchmark and application on quantum computer.** (a) Imaginary and (b) real part of the MLDE with  $N_b = 3$  self-energy (dashed lines) obtained from on-site correlation  $U$  in range  $2 - 15$  for a test hybridisation function for a half-filled impurity (units are arbitrary) and compared with continuous time Monte Carlo (continuous line). The scales are in arbitrary units. (c) Quasi-particle weight obtained by MLDE with  $N_b = 2$  and  $N_b = 3$  compared to the continuous time Monte Carlo. The scales are in arbitrary units. (d) Imaginary part of the self-energy as function of the Matsubara frequencies for the MLDE system presented in panel (b) and  $U = 6$  ( $N_b = 2$ ), computed using the quantum computing algorithm (“MLDE-VQE”, blue solid curve), and compared to the results obtained with the conventional computing algorithm (“MLDE-exact”, red dashed curves).  $\Im$  and  $\Re$  stand respectively for the imaginary and real part of the complex functions. The latter notations are used throughout the manuscript.

**Figure 3. Kondo physics.** (a) TOV organic molecule deposited on a gold substrate at  $T = 5\text{K}$ . (b) Imaginary parts of the molecule’s hybridisation function (AIM), represented by MLDE  $\Delta_{1,2}(i\omega_n)$ , and as obtained by the ED solver with  $N_b = 12$ . (c) Imaginary part of the self-energy obtained by MLDE (squares) and ED (triangles). (d) Imaginary part of the self-energy in real axis frequencies obtained with MLDE and ED. (e) The inset shows the time dynamics of the magnetically quenched system, at a temperature  $T = 100\text{K}$  ( $T = 20\text{K}$ ) above (below) the Kondo temperature ( $T_K \approx 37\text{K}$ ). The main panel outlines the Fourier transform of the dynamics at  $T = 100\text{K}$  showing a peak at  $\omega = 0.0029\text{eV}$  ( $= 33.64\text{K}$ ).

**Figure 4. Mott transition.** (a) Spectral function obtained for the square lattice Hubbard model solved by MLDE ( $N_b = 3$  for all calculations) with increasing values of the on-site dimensionless correlation ratio  $U/t$ . (b) Imaginary part of the self-energy obtained for  $U/t = 12$  at different MLDE iteration of the self-consistent cycle compared with CQMC solutions. (c) Converged imaginary part of the MLDE Green’s function for different values of  $U/t$  (symbols) compared with the DMFT CTQMC solution (dashed lines).

**Figure 5. Dynamical susceptibility.** Momentum resolved spin susceptibility obtained by Bethe Salpeter with the vertex calculated with MLDE in the Hubbard model with  $U/t = 12$ , at increasing temperature  $T/t = 0.025$  (a),  $T/t = 1$  (b),  $T/t = 10$  (c). (d) Magnetic susceptibility  $\chi$  and (e) irreducible vertex  $\Gamma$  resolved in fermionic frequency  $i\nu$  obtained by MLDE (continuous line) and compared with the exact vertex (dashed line) at temperature  $T/t = 0.025$  and  $U/t = 6$ .

**Figure 6. MLDE multiorbital correlated  $\text{LaNiO}_3$ :** Self energy of the first iteration of the self-consistent DMFT loop (panel a) and the Green’s function for the final iteration (panel b) for the  $e_g$  orbitals of the Ni ion in the  $\text{LaNiO}_3$  ( $d_{z^2}$  in red and  $d_{x^2-y^2}$  in blue). In both case, we compare the quantities obtained with MLDE with 3 sites in the bath (solid line) with the quantities obtained with CTQMC (dashed line).

## References

- 385
- 386 1. Kotliar, G. *et al.* Electronic structure calculations with dynamical mean-field theory. *Rev. Mod. Phys.* **78**, 865, DOI:  
387 [10.1103/RevModPhys.78.865](https://doi.org/10.1103/RevModPhys.78.865) (2006).
- 388 2. Georges, A., Kotliar, G., Krauth, W. & Rozenberg, M. J. Dynamical mean-field theory of strongly correlated fermion  
389 systems and the limit of infinite dimensions. *Rev. Mod. Phys.* **68**, 13, DOI: [10.1103/RevModPhys.68.13](https://doi.org/10.1103/RevModPhys.68.13) (1996).
- 390 3. Zgid, D. & Gull, E. Finite temperature quantum embedding theories for correlated systems. *New J. Phys.* **19**, 023047, DOI:  
391 [10.1088/1367-2630/aa5d34](https://doi.org/10.1088/1367-2630/aa5d34) (2017).
- 392 4. Fertitta, E. & Booth, G. H. Rigorous wave function embedding with dynamical fluctuations. *Phys. Rev. B* **98**, 235132,  
393 DOI: [10.1103/PhysRevB.98.235132](https://doi.org/10.1103/PhysRevB.98.235132) (2018).
- 394 5. Sun, Q. & Chan, G. K.-L. Quantum embedding theories. *Accounts Chem. Res.* **49**, 2705–2712, DOI: [10.1021/acs.accounts.](https://doi.org/10.1021/acs.accounts.6b00356)  
395 [6b00356](https://doi.org/10.1021/acs.accounts.6b00356) (2016). PMID: 27993005, <https://doi.org/10.1021/acs.accounts.6b00356>.
- 396 6. Gubernatis, J. E., Jarrell, M., Silver, R. N. & Sivia, D. S. Quantum Monte Carlo simulations and maximum entropy:  
397 Dynamics from imaginary-time data. *Phys. Rev. B* **44**, 6011–6029, DOI: [10.1103/PhysRevB.44.6011](https://doi.org/10.1103/PhysRevB.44.6011) (1991).
- 398 7. Gunnarsson, O., Haverkort, M. W. & Sangiovanni, G. Analytical continuation of imaginary axis data for optical conductivity.  
399 *Phys. Rev. B* **82**, 165125, DOI: [10.1103/PhysRevB.82.165125](https://doi.org/10.1103/PhysRevB.82.165125) (2010).
- 400 8. Mitchell, A. K. & Fritz, L. Kondo effect with diverging hybridization: Possible realization in graphene with vacancies.  
401 *Phys. Rev. B* **88**, 075104, DOI: [10.1103/PhysRevB.88.075104](https://doi.org/10.1103/PhysRevB.88.075104) (2013).
- 402 9. Capone, M., de' Medici, L. & Georges, A. Solving the dynamical mean-field theory at very low temperatures using the  
403 Lanczos exact diagonalization. *Phys. Rev. B* **76**, 245116, DOI: [10.1103/PhysRevB.76.245116](https://doi.org/10.1103/PhysRevB.76.245116) (2007).
- 404 10. Perroni, C. A., Ishida, H. & Liebsch, A. Exact diagonalization dynamical mean-field theory for multiband materials: Effect  
405 of coulomb correlations on the fermi surface of *na0.3coo2*. *Phys. Rev. B* **75**, 045125, DOI: [10.1103/PhysRevB.75.045125](https://doi.org/10.1103/PhysRevB.75.045125)  
406 (2007).
- 407 11. Capone, M., Fabrizio, M., Castellani, C. & Tosatti, E. Strongly correlated superconductivity. *Science* **296**, 2364–2366,  
408 DOI: [10.1126/science.1071122](https://doi.org/10.1126/science.1071122) (2002).
- 409 12. Liebsch, A. & Ishida, H. Temperature and bath size in exact diagonalization dynamical mean field theory. *J. Physics:*  
410 *Condens. Matter* **24**, 053201 (2012).
- 411 13. Ishida, H. & Liebsch, A. Fermi-liquid, non-fermi-liquid, and mott phases in iron pnictides and cuprates. *Phys. Rev. B* **81**,  
412 054513, DOI: [10.1103/PhysRevB.81.054513](https://doi.org/10.1103/PhysRevB.81.054513) (2010).
- 413 14. Arrigoni, E., Knap, M. & von der Linden, W. Nonequilibrium dynamical mean-field theory: An auxiliary quantum master  
414 equation approach. *Phys. Rev. Lett.* **110**, 086403, DOI: [10.1103/PhysRevLett.110.086403](https://doi.org/10.1103/PhysRevLett.110.086403) (2013).
- 415 15. Lu, Y., Höppner, M., Gunnarsson, O. & Haverkort, M. W. Efficient real-frequency solver for dynamical mean-field theory.  
416 *Phys. Rev. B* **90**, 085102, DOI: [10.1103/PhysRevB.90.085102](https://doi.org/10.1103/PhysRevB.90.085102) (2014).
- 417 16. Ganahl, M. *et al.* Efficient dmft impurity solver using real-time dynamics with matrix product states. *Phys. Rev. B* **92**,  
418 155132, DOI: [10.1103/PhysRevB.92.155132](https://doi.org/10.1103/PhysRevB.92.155132) (2015).
- 419 17. Bauernfeind, D., Zingl, M., Triebel, R., Aichhorn, M. & Evertz, H. G. Fork tensor-product states: Efficient multiorbital  
420 real-time dmft solver. *Phys. Rev. X* **7**, 031013, DOI: [10.1103/PhysRevX.7.031013](https://doi.org/10.1103/PhysRevX.7.031013) (2017).
- 421 18. Lu, Y., Höppner, M., Gunnarsson, O. & Haverkort, M. W. Efficient real-frequency solver for dynamical mean-field theory.  
422 *Phys. Rev. B* **90**, 085102, DOI: [10.1103/PhysRevB.90.085102](https://doi.org/10.1103/PhysRevB.90.085102) (2014).
- 423 19. Rungger, I. *et al.* Dynamical mean field theory algorithm and experiment on quantum computers (2019). [1910.04735](https://arxiv.org/abs/1910.04735).
- 424 20. Sakai, S., Civelli, M. & Imada, M. Hidden fermionic excitation boosting high-temperature superconductivity in cuprates.  
425 *Phys. Rev. Lett.* **116**, 057003, DOI: [10.1103/PhysRevLett.116.057003](https://doi.org/10.1103/PhysRevLett.116.057003) (2016).
- 426 21. Sen, S., Wong, P. J. & Mitchell, A. K. The mott transition as a topological phase transition (2020). [2001.10526](https://arxiv.org/abs/2001.10526).
- 427 22. Baldini, E. *et al.* Electron–phonon-driven three-dimensional metallicity in an insulating cuprate. *Proc. Natl. Acad. Sci.*  
428 **117**, 6409–6416, DOI: [10.1073/pnas.1919451117](https://doi.org/10.1073/pnas.1919451117) (2020). <https://www.pnas.org/content/117/12/6409.full.pdf>.
- 429 23. Acharya, S. *et al.* Metal-insulator transition in copper oxides induced by apex displacements. *Phys. Rev. X* **8**, 021038, DOI:  
430 [10.1103/PhysRevX.8.021038](https://doi.org/10.1103/PhysRevX.8.021038) (2018).
- 431 24. Liu, J. *et al.* First observation of a kondo resonance for a stable neutral pure organic radical, 1,3,5-triphenyl-6-oxoverdazyl,  
432 adsorbed on the au(111) surface. *J. Am. Chem. Soc.* **135**, 651 (2013).

- 433 **25.** Zhang, Y.-h. *et al.* Temperature and magnetic field dependence of a kondo system in the weak coupling regime. *Nat.*  
434 *Commun.* **4**, 2110 (2013).
- 435 **26.** Frisenda, R. *et al.* Kondo effect in a neutral and stable all organic radical single molecule break junction. *Nano letters* **15**,  
436 3109–3114 (2015).
- 437 **27.** Droghetti, A. & Rungger, I. Quantum transport simulation scheme including strong correlations and its application to  
438 organic radicals adsorbed on gold. *Phys. Rev. B* **95**, 085131, DOI: [10.1103/PhysRevB.95.085131](https://doi.org/10.1103/PhysRevB.95.085131) (2017).
- 439 **28.** Appelt, W. H. *et al.* Predicting the conductance of strongly correlated molecules: the kondo effect in perchlorotriphenyl-  
440 methyl/au junctions. *Nanoscale* **10**, 17738–17750, DOI: [10.1039/C8NR03991G](https://doi.org/10.1039/C8NR03991G) (2018).
- 441 **29.** Balzer, M., Gdaniec, N. & Potthoff, M. Krylov-space approach to the equilibrium and nonequilibrium single-particle  
442 green's function. *J. Phys.: Condens. Matter* **24**, 035603 (2012).
- 443 **30.** Lee, H., Plekhanov, E., Blackburn, D., Acharya, S. & Weber, C. The mott to kondo transition in diluted kondo superlattices.  
444 *Commun. Phys.* **2**, 49 (2019).
- 445 **31.** Surer, B. *et al.* Multiorbital kondo physics of co in cu hosts. *Phys. Rev. B* **85**, 085114, DOI: [10.1103/PhysRevB.85.085114](https://doi.org/10.1103/PhysRevB.85.085114)  
446 (2012).
- 447 **32.** Žitko, R., Peters, R. & Pruschke, T. Splitting of the kondo resonance in anisotropic magnetic impurities on surfaces. *New J.*  
448 *Phys.* **11**, 053003, DOI: [10.1088/1367-2630/11/5/053003](https://doi.org/10.1088/1367-2630/11/5/053003) (2009).
- 449 **33.** Granath, M. & Schött, J. Signatures of coherent electronic quasiparticles in the paramagnetic mott insulator. *Phys. Rev. B*  
450 **90**, 235129, DOI: [10.1103/PhysRevB.90.235129](https://doi.org/10.1103/PhysRevB.90.235129) (2014).
- 451 **34.** Liebsch, A. & Ishida, H. Temperature and bath size in exact diagonalization dynamical mean field theory. *J. Physics:*  
452 *Condens. Matter* **24**, 053201, DOI: [10.1088/0953-8984/24/5/053201](https://doi.org/10.1088/0953-8984/24/5/053201) (2011).
- 453 **35.** Nowadnick, E. A. *et al.* Quantifying electronic correlation strength in a complex oxide: A combined dmft and arpes study  
454 of lanio<sub>3</sub>. *Phys. Rev. B* **92**, 245109, DOI: [10.1103/PhysRevB.92.245109](https://doi.org/10.1103/PhysRevB.92.245109) (2015).
- 455 **36.** Rungger, I. *et al.* Dynamical mean field theory algorithm and experiment on quantum computers (2020). [1910.04735](https://arxiv.org/abs/1910.04735).
- 456 **37.** Jones, T., Brown, A., Bush, I. & Benjamin, S. C. Quest and high performance simulation of quantum computers. *Sci.*  
457 *Reports* **9**, 10736, DOI: [10.1038/s41598-019-47174-9](https://doi.org/10.1038/s41598-019-47174-9) (2019).
- 458 **38.** Fradkin, E. Jordan-wigner transformation for quantum-spin systems in two dimensions and fractional statistics. *Phys. Rev.*  
459 *Lett.* **63**, 322–325, DOI: [10.1103/PhysRevLett.63.322](https://doi.org/10.1103/PhysRevLett.63.322) (1989).
- 460 **39.** Wecker, D. *et al.* Solving strongly correlated electron models on a quantum computer. *Phys. Rev. A* **92**, 062318, DOI:  
461 [10.1103/PhysRevA.92.062318](https://doi.org/10.1103/PhysRevA.92.062318) (2015).
- 462 **40.** Bauer, B., Wecker, D., Millis, A. J., Hastings, M. B. & Troyer, M. Hybrid quantum-classical approach to correlated  
463 materials. *Phys. Rev. X* **6**, 031045, DOI: [10.1103/PhysRevX.6.031045](https://doi.org/10.1103/PhysRevX.6.031045) (2016).
- 464 **41.** Cini, M. Topics and methods in condensed matter theory. *From Basic Quantum Mech. to Front. Res.* Springer DOI:  
465 [ISBN978-3-540-70727-1](https://doi.org/10.1007/978-3-540-70727-1) (2007).
- 466 **42.** Zgid, D. & Gull, E. Finite temperature quantum embedding theories for correlated systems. *New J. Phys.* **19**, 023047, DOI:  
467 [10.1088/1367-2630/aa5d34](https://doi.org/10.1088/1367-2630/aa5d34) (2017).
- 468 **43.** Schäfer, T., Toschi, A. & Tomczak, J. M. Separability of dynamical and nonlocal correlations in three dimensions. *Phys.*  
469 *Rev. B* **91**, 121107, DOI: [10.1103/PhysRevB.91.121107](https://doi.org/10.1103/PhysRevB.91.121107) (2015).
- 470 **44.** Neuhauser, D., Baer, R. & Zgid, D. Stochastic self-consistent second-order green's function method for correlation energies  
471 of large electronic systems. *J. Chem. Theory Comput.* **13**, 5396–5403, DOI: [10.1021/acs.jctc.7b00792](https://doi.org/10.1021/acs.jctc.7b00792) (2017). PMID:  
472 28961398, <https://doi.org/10.1021/acs.jctc.7b00792>.
- 473 **45.** Wolf, F. A., Go, A., McCulloch, I. P., Millis, A. J. & Schollwöck, U. Imaginary-time matrix product state impurity solver  
474 for dynamical mean-field theory. *Phys. Rev. X* **5**, 041032, DOI: [10.1103/PhysRevX.5.041032](https://doi.org/10.1103/PhysRevX.5.041032) (2015).
- 475 **46.** Toschi, A., Katanin, A. A. & Held, K. Dynamical vertex approximation: A step beyond dynamical mean-field theory. *Phys.*  
476 *Rev. B* **75**, 045118, DOI: [10.1103/PhysRevB.75.045118](https://doi.org/10.1103/PhysRevB.75.045118) (2007).
- 477 **47.** Hafermann, H. *et al.* Superperturbation solver for quantum impurity models. *EPL (Europhysics Lett.)* **85**, 27007, DOI:  
478 [10.1209/0295-5075/85/27007](https://doi.org/10.1209/0295-5075/85/27007) (2009).
- 479 **48.** Rohringer, G., Valli, A. & Toschi, A. Local electronic correlation at the two-particle level. *Phys. Rev. B* **86**, 125114, DOI:  
480 [10.1103/PhysRevB.86.125114](https://doi.org/10.1103/PhysRevB.86.125114) (2012).

- 481 **49.** Rohringer, G. *et al.* Diagrammatic routes to nonlocal correlations beyond dynamical mean field theory. *Rev. Mod. Phys.*  
482 **90**, 025003, DOI: [10.1103/RevModPhys.90.025003](https://doi.org/10.1103/RevModPhys.90.025003) (2018).
- 483 **50.** Pashov, D. *et al.* Questaal: A package of electronic structure methods based on the linear muffin-tin orbital technique.  
484 *Comput. Phys. Commun.* **249**, 107065, DOI: [10.1016/j.cpc.2019.107065](https://doi.org/10.1016/j.cpc.2019.107065) (2020).
- 485 **51.** Seth, P., Krivenko, I., Ferrero, M. & Parcollet, O. Triqs/cthyb: A continuous-time quantum monte carlo hybridisation  
486 expansion solver for quantum impurity problems. *Comput. Phys. Commun.* **200**, 274 – 284, DOI: [http://dx.doi.org/10.](http://dx.doi.org/10.1016/j.cpc.2015.10.023)  
487 [1016/j.cpc.2015.10.023](http://dx.doi.org/10.1016/j.cpc.2015.10.023) (2016).
- 488 **52.** Wiersema, R. *et al.* Exploring entanglement and optimization within the hamiltonian variational ansatz. *PRX Quantum* **1**,  
489 DOI: [10.1103/prxquantum.1.020319](https://doi.org/10.1103/prxquantum.1.020319) (2020).
- 490 **53.** Kandala, A. *et al.* Hardware-efficient variational quantum eigensolver for small molecules and quantum magnets. *Nature*  
491 **549**, 242 (2017).
- 492 **54.** Nakanishi, K. M., Mitarai, K. & Fujii, K. Subspace-search variational quantum eigensolver for excited states. *Phys. Rev.*  
493 *Res.* **1**, 033062, DOI: [10.1103/PhysRevResearch.1.033062](https://doi.org/10.1103/PhysRevResearch.1.033062) (2019).
- 494 **55.** Higgott, O., Wang, D. & Brierley, S. Variational Quantum Computation of Excited States. *Quantum* **3**, 156, DOI:  
495 [10.22331/q-2019-07-01-156](https://doi.org/10.22331/q-2019-07-01-156) (2019).
- 496 **56.** Jones, T., Endo, S., McArdle, S., Yuan, X. & Benjamin, S. C. Variational quantum algorithms for discovering hamiltonian  
497 spectra. *Phys. Rev. A* **99**, 062304, DOI: [10.1103/PhysRevA.99.062304](https://doi.org/10.1103/PhysRevA.99.062304) (2019).
- 498 **57.** Lupo, C., Tse, T., Jamet, F., Rungger, I. & Weber, C. Maximally localized dynamical embedding (mlde). [https:](https://www.codeocean.com/)  
499 [/www.codeocean.com/](https://www.codeocean.com/), DOI: [10.24433/CO.9420336.v4](https://doi.org/10.24433/CO.9420336.v4) (2021).