

***FMNS 2023***  
*JUNE 14-18*  
*BLAGOEVGRAD, BULGARIA*

***PRODUCTS OF DISTRIBUTIONS IN  
COLOMBEAU ALGEBRA***

**Marija Miteva, Limonka Koceva Lazarova, Biljana  
Zlatanovska, Natasha Stojkovikj**  
Goce Delcev University – Stip, Macedonia

---

# INTRODUCTION TO THE THEORY OF DISTRIBUTIONS

---

- Theory of distributions 1950s
  - Mathematical meaning of many concepts in the science that were described heuristically
  - Dirac  $\delta$  - function and its derivatives
    - Concepts and their properties were defined heuristically, to be appropriate to the experimental results and to be adequate for analysis and solving the problems that they characterize
  - The operations with those concepts remained mathematically unsupported

$$\delta(x) = \begin{cases} 0, & x \neq 0 \\ \infty & x = 0 \end{cases} \quad \int_{-\infty}^{\infty} \delta(x) dx = 1 \quad (1)$$

# INTRODUCTION TO THE THEORY OF DISTRIBUTIONS

---

- Necessity for generalizing the concept of function
- Distribution (generalized function)
- Laurent Schwartz, '*Theory of Distributions*' (1950)
  - Many concepts that can not be described with functions can be described applying distributions
  - Concepts that can be described with functions can also be described using distributions

# INTRODUCTION TO THE THEORY OF DISTRIBUTIONS

---

- ✗  $\mathcal{D} = \mathcal{D}(\mathbf{R}^n)$  - space of smooth functions with compact support (test functions)
- ✗ *Generalized function (distribution) is continuous linear mapping  $f : \mathcal{D} \rightarrow \mathbf{C}$*

$$f(\varphi) = \langle f, \varphi \rangle \quad (2)$$

$\mathcal{D}' = \mathcal{D}'(\mathbf{R}^n)$  - the space of distributions with domain  $\mathcal{D}$

# INTRODUCTION TO THE THEORY OF DISTRIBUTIONS

---

- $f$  - locally integrable function

$$f(\varphi) = \langle f, \varphi \rangle = \int_{\mathbf{R}^n} f(x) \varphi(x) dx \quad (3)$$

- $f$  - regular distribution

- *Singular distributions*

- Dirac  $\delta$  distribution:

$$\langle \delta, \varphi \rangle = \varphi(0) \quad (4)$$

# INTRODUCTION TO THE THEORY OF DISTRIBUTIONS

---

## ➤ Differentiation of distributions

$$f \in \mathcal{D}'(\mathbf{R}) \quad \varphi \in \mathcal{D}(\mathbf{R})$$

$$\left\langle f^{(n)}(x), \varphi(x) \right\rangle = (-1)^n \left\langle f(x), \varphi^{(n)}(x) \right\rangle \quad (5)$$

$$f \in \mathcal{D}'(\mathbf{R}^n) \quad \varphi \in \mathcal{D}(\mathbf{R}^n)$$

$$D^k = \prod_{i=1}^n \left( \frac{\partial}{\partial x_i} \right)^{k_i} \quad k_i \in \mathbf{N}_0 \quad k = \sum_{i=1}^n k_i$$

$$\left\langle D^k f, \varphi \right\rangle = (-1)^k \left\langle f, D^k \varphi \right\rangle \quad (6)$$

# INTRODUCTION TO THE THEORY OF DISTRIBUTIONS

---

- ❖ **A derivative of function, with arbitrary order, will always exist if we consider that function as generalized function (distribution)**
- ❖ **Two main problems for the theory of distributions:**
  - **Product of distributions: two arbitrary distributions can not always be multiplied**
    - the product of distributions is not an associative operation
  - **Differentiation of the product of distributions** (the product of distributions not always satisfy the Leibniz rule)

# INTRODUCTION TO THE THEORY OF DISTRIBUTIONS

- ▶ The application of distributions in non-linear systems needs products of singular distributions
- ▶ Attempts for defining product of distributions that will be generalization of the existing products

## ▶ **Regularization method**

$\varphi_n \rightarrow \delta(x)$  - delta sequence;  $f$  - distribution

$$f_n(x) = (f * \varphi_n)(x) = \langle f(y), \varphi_n(x-y) \rangle \quad (7)$$

- ▶ Sequence of smooth functions  $(f_n)$ ;  $f_n \rightarrow f$
- ▶  $(f_n)$  - regularization of the distribution  $f$

- ▶ 
$$fg = \lim_{n \rightarrow \infty} (f * \varphi_n) \cdot (g * \varphi_n) \quad (8)$$

# INTRODUCTION TO THE THEORY OF DISTRIBUTIONS

---

➤ with regularization method:

➤  $\frac{1}{x} \cdot \delta = -\frac{1}{2} \delta'$  (9)

➤ but,  $\delta \cdot \delta = \delta^2$  is not defined neither with the regularization method

# COLOMBEAU ALGEBRA

---

- ❖ Overcoming the problem with product of distributions
- ❖ Construction of algebra  $A$  with properties:
  - 1) Contains the space of distributions  $\mathcal{D}'(\mathbf{R}^n)$  and  $f(x) \equiv 1$  is neutral element in  $A$
  - 2) There exist linear differential operators  $\partial_i : A \rightarrow A$  which satisfy the Leibnitz's rule
  - 3)  $\partial_i$  generalizes the derivation on the space of distributions
  - 4) The product in  $A$  generalizes the product of continuous functions

## ❖ Schwartz's impossibility result

- ▶ the functions  $f(x) = x$  и  $g(x) = |x|$  are considered
- ▶ derivative of their classical product:

$$\partial(x|x|) = 2|x| \quad (10)$$

$$\partial^2(x|x|) = 2\partial(|x|) \quad (11)$$

- ▶ Derivative of their product in  $A$

$$\partial(x \cdot |x|) = |x| + x \cdot \partial(|x|) \quad (12)$$

$$\partial^2(x \cdot |x|) = 2\partial(|x|) + x \cdot \partial^2(|x|) \quad (13)$$

$$\partial^2(x \cdot |x|) = 2\partial(|x|) + 2x \cdot \delta \quad (14)$$

## ❖ Schwartz's impossibility result

- From (11) and (14) it follows:  $x \cdot \delta = 0$  (15)
- *Theorem:* In  $A$ , if  $x \cdot a = 0$  then  $a = 0$ .
- From (15)  $\Rightarrow \delta = 0$

# COLOMBEAU ALGEBRA

---

- ❑ New theory of generalized functions, more general than the theory of distributions
- ❑ **Jean-Francois Colombeau**
  - ❑ *New generalized function and multiplication of distributions (1984)*
  - ❑ *Elementary introduction to new generalized functions (1985)*
- ❑ **Colombeau algebra**
  - ❑ The product in the algebra generalizes classical product of  $C^\infty$  - functions

# CONSTRUCTION OF THE COLOMBEAU ALGEBRA

- $\mathbf{N}_0 = \mathbf{N} \cup \{0\}$  - non negative integers
- $\mathcal{D}(\mathbf{R}^n)$  - the space of  $C^\infty$  - functions  $\varphi: \mathbf{R}^n \rightarrow \mathbf{C}$  with compact support

- for  $j \in \mathbf{N}_0$  and  $q \in \mathbf{N}_0$  the next sets are defined:

$$A_0(\mathbf{R}^n) = \left\{ \varphi(x) \in \mathcal{D}(\mathbf{R}^n) \left| \int_{\mathbf{R}^n} \varphi(x) dx = 1 \right. \right\}$$

$$A_q(\mathbf{R}^n) = \left\{ \varphi(x) \in \mathcal{D}(\mathbf{R}^n) \left| \int_{\mathbf{R}^n} \varphi(x) dx = 1, \int_{\mathbf{R}^n} x^j \varphi(x) dx = 0; 1 \leq |j| \leq q \right. \right\}$$

$$q \geq 1 \quad x = (x_1, x_2, \dots, x_n) \in \mathbf{R}^n \quad j = (j_1, j_2, \dots, j_n) \in \mathbf{N}^n$$
$$|j| = j_1 + j_2 + \dots + j_n \quad x^j = (x_1)^{j_1} (x_2)^{j_2} \dots (x_n)^{j_n}$$

# CONSTRUCTION OF THE COLOMBEAU ALGEBRA

---

$$\diamond A_0 \supset A_1 \supset A_2 \supset A_3 \dots$$

$\diamond$  **Theorem:** The sets  $A_q$  are non empty sets.

Proof: J.F.Colombeau, *Elementary Introduction to New Generalized Functions* (1985)

$\diamond$  For  $\varphi \in A_q(\mathbf{R}^n)$  and  $\varepsilon > 0$  we denote:

$$\varphi_\varepsilon(x) = \frac{1}{\varepsilon^n} \varphi\left(\frac{x}{\varepsilon}\right) \tag{16}$$

$$\check{\varphi}(x) = \varphi(-x) \tag{17}$$

# CONSTRUCTION OF THE COLOMBEAU ALGEBRA

---

- ❖  $\mathcal{E}(\mathbf{R}^n)$  - an algebra of functions  $f(\varphi, x): A_0(\mathbf{R}^n) \times \mathbf{R}^n \rightarrow \mathbf{C}$  that are infinitely differentiable regarding the second variable  $x$  (with fixed test function  $\varphi$ )
- ❖ The space  $C^\infty(\mathbf{R}^n)$  is subalgebra of  $\mathcal{E}(\mathbf{R}^n)$  (those functions that don't depend of  $\varphi$ )
- ❖ Embedding of the distributions in  $\mathcal{E}(\mathbf{R}^n)$  is such that embedding of  $C^\infty(\mathbf{R}^n)$  - functions is identity

# CONSTRUCTION OF THE COLOMBEAU ALGEBRA

---

- $\mathcal{E}_M[\mathbf{R}^n]$  - subalgebra of  $\mathcal{E}(\mathbf{R}^n)$  with elements such that for every compact subset  $K$  from  $\mathbf{R}^n$  and every  $p \in \mathbf{N}_0$  there exists  $q \in \mathbf{N}$  such that for arbitrary  $\varphi \in A_q(\mathbf{R}^n)$  there exist  $c > 0$ ,  $\eta > 0$  and the relation holds:

$$\sup_{x \in K} |\partial^p f(\varphi_\varepsilon, x)| \leq c\varepsilon^{-q} \quad (18)$$

for  $0 < \varepsilon < \eta$ .

- Functions in  $\mathcal{E}(\mathbf{R}^n)$  which derivatives on compact sets are bounded with negative powers of  $\varepsilon$

# CONSTRUCTION OF THE COLOMBEAU ALGEBRA

$$f \in C(\mathbf{R}^n)$$

✘ With the mapping:

$$F(\varphi, x) = \int_{\mathbf{R}^n} f(y) \varphi(y-x) dy = \int_{\mathbf{R}^n} f(x+t) \varphi(t) dt \quad (19)$$

the space  $C(\mathbf{R}^n)$  is embedded in  $\mathcal{E}_M[\mathbf{R}^n]$ .

□  $C^\infty(\mathbf{R}^n)$  is contained in  $\mathcal{E}_M[\mathbf{R}^n]$  in a way that  $f(x)$  are those functions  $f(\varphi, x)$  that don't depend of  $\varphi$ .

□  $C^\infty(\mathbf{R}^n) \subset C(\mathbf{R}^n)$  are embedded in  $\mathcal{E}_M[\mathbf{R}^n]$  with (19).

$$f(x) \neq \int_{\mathbf{R}^n} f(x+\varepsilon t) \varphi(t) dt \quad (20)$$

□ Ideal such that the difference in (20) will vanish

# CONSTRUCTION OF THE COLOMBEAU ALGEBRA

- $\mathcal{I}[\mathbf{R}^n]$  is an ideal in  $\mathcal{E}_M[\mathbf{R}^n]$  consisting of functions  $f(\varphi, x)$  such that for every compact subset  $K$  of  $\mathbf{R}^n$  and each  $p \in \mathbf{N}_0$  there exist  $q \in \mathbf{N}$  such that for any  $r \geq q$  and each  $\varphi \in A_r(\mathbf{R}^n)$  there exist  $c > 0, \eta > 0$  and it holds:

$$\sup_{x \in K} \left| \partial^p f(\varphi_\varepsilon, x) \right| \leq c \varepsilon^{r-q} \quad (21)$$

For  $0 < \varepsilon < \eta$  .

- The elements of  $\mathcal{I}[\mathbf{R}^n]$  are called *null functions*

# CONSTRUCTION OF THE COLOMBEAU ALGEBRA

❖ **Generalized functions in Colombeau theory are elements of quotient algebra**

$$\mathcal{G} \equiv \mathcal{G}(\mathbf{R}^n) = \frac{\mathcal{E}_M[\mathbf{R}^n]}{\mathcal{I}[\mathbf{R}^n]} \quad (22)$$

❖ In  $\mathcal{E}_M[\mathbf{R}^n]$  the equivalence relation is defined '  $\sim$  ' :

$$F_1 \sim F_2 \Leftrightarrow F_1 - F_2 \in \mathcal{I}[\mathbf{R}^n] \quad (23)$$

❖ **The generalized functions in Colombeau theory are an equivalence classes of smooth functions**

❖ New generalized functions (Colombeau generalized functions)

## EMBEDDING OF DISTRIBUTIONS IN COLOMBEAU ALGEBRA

$$\square \quad f \in C^\infty(\mathbf{R}^n) \quad f \rightarrow f(\varphi, x) \in \mathcal{G}(\mathbf{R}^n)$$

$$f(\varphi, x) = f(x) \tag{24}$$

$$\square \quad f \in C(\mathbf{R}^n) \quad f \rightarrow f(\varphi, x) \in \mathcal{G}(\mathbf{R}^n)$$

$$f(\varphi, x) = \int_{\mathbf{R}^n} f(y) \varphi(y-x) dy = \int_{\mathbf{R}^n} f(x+y) \varphi(y) dy \tag{25}$$

$$\square \quad f \in \mathcal{D}'(\mathbf{R}^n) \quad f \rightarrow f(\varphi, x) \in \mathcal{G}(\mathbf{R}^n)$$

$$f(\varphi, x) = \left( f * \overset{\vee}{\varphi} \right)(x) = \langle f(y), \varphi(y-x) \rangle = \int_{\mathbf{R}^n} f(y) \varphi(y-x) dy \tag{26}$$

# ASSOCIATION IN COLOMBEAU ALGEBRA

---

- ✓ The space of smooth functions  $C^\infty(\mathbf{R}^n)$  is subalgebra of the Colombeau algebra  $\mathcal{G}(\mathbf{R}^n)$
- ✓ The space of continuous functions  $C(\mathbf{R}^n)$  and the space of distributions  $\mathcal{D}'(\mathbf{R}^n)$  are not subalgebras of Colombeau algebra  $\mathcal{G}(\mathbf{R}^n)$ 
  - ✓ If  $f, g$  are two continuous functions (or distributions which classical product exists), the embedding of their classical product,  $fg$ , and the product of their embeddings  $f \cdot g$  in  $\mathcal{G}$  may not coincide
  - ✓ This difference of the products has been overcome introducing the concept of „association“ in  $\mathcal{G}$

# ASSOCIATION IN COLOMBEAU ALGEBRA

- ❖ Generalized functions  $F, G \in \mathcal{G}(\mathbf{R}^n)$  are said to be **associated** ( $F \approx G$ ) if for each representatives  $f(\varphi_\varepsilon, x)$  and  $g(\varphi_\varepsilon, x)$  and each  $\psi(x) \in \mathcal{D}(\mathbf{R}^n)$ , there exists  $q \in \mathbf{N}_0$  such that for any  $\varphi(x) \in A_q(\mathbf{R}^n)$  holds:

$$\lim_{\varepsilon \rightarrow 0_+} \int_{\mathbf{R}^n} |f(\varphi_\varepsilon, x) - g(\varphi_\varepsilon, x)| \psi(x) dx = 0 \quad (27)$$

- ❖ Generalized function  $F \in \mathcal{G}$  is associated with the distribution  $u \in \mathcal{D}'$  ( $F \approx u$ ) if for each representative of that generalized function  $f(\varphi_\varepsilon, x)$  and each  $\psi(x) \in \mathcal{D}(\mathbf{R}^n)$ , there exist  $q \in \mathbf{N}_0$  such that for any  $\varphi(x) \in A_q(\mathbf{R}^n)$  holds:

$$\lim_{\varepsilon \rightarrow 0_+} \int_{\mathbf{R}^n} f(\varphi_\varepsilon, x) \psi(x) dx = \langle u, \psi \rangle \quad (28)$$

# ASSOCIATION IN COLOMBEAU ALGEBRA

---

- ✓ Previous definitions are independent of the representatives chosen
- ✓ The distribution associated, if it exists, is unique
- ✓ To an element of Colombeau algebra, with this process of association, is associated element in  $\mathcal{D}'$ , which allows us to consider obtained results in the sense of distribution.
  - ✓ Not any element in Colombeau algebra has an associated distribution!

# ASSOCIATION IN COLOMBEAU ALGEBRA

- *Theorem:* If  $f, g \in C(\mathbf{R}^n)$  are two continuous functions, their product  $f \cdot g$  in  $\mathcal{G}(\mathbf{R})$  is associated with their classical product  $fg$  in  $C(\mathbf{R}^n)$ .
- *Theorem:* If  $f \in C^\infty(\mathbf{R}^n)$  and  $T \in \mathcal{D}'(\mathbf{R}^n)$ , the product  $f \cdot T$  in  $\mathcal{G}(\mathbf{R}^n)$  is associated with the classical product  $fT$  in  $\mathcal{D}'(\mathbf{R}^n)$ .
- *Theorem:* If  $S$  and  $T$  are two distributions in  $\mathcal{D}'(\mathbf{R}^n)$  and their classical product  $ST$  in  $\mathcal{D}'(\mathbf{R}^n)$  exists, then the product of these two distributions  $S \cdot T$  in  $\mathcal{G}(\mathbf{R}^n)$  is associated with their classical product  $ST$ .

# ASSOCIATION IN COLOMBEAU ALGEBRA

---

- ✓ Two distributions embedded in Colombeau algebra are new (Colombeau) generalized functions
- ✓ Product of two distributions in  $\mathcal{G}$  is in general new (Colombeau) generalized function (for which there may not exist associated distribution)
- ✓ *If for the product of two distributions in  $\mathcal{G}$  there exists an associated distribution, we say that there exists **the Colombeau product of those two distributions***
- ✓ **If the classical product of two distributions exists, then their Colombeau product also exists and is the same with the first one**

# RESULTS ON PRODUCTS OF DISTRIBUTIONS IN COLOMBEAU ALGEBRA

- $\ln|x| \cdot \delta^{(s-1)}(x) \approx \frac{-1}{s} \delta^{(s-1)}(x) \quad s = 1, 2, \dots \quad (29)$
- $x_+^{-k} \cdot \delta^{(p)}(x) \approx \frac{(-1)^k k \cdot p!}{(p+k+1)!} \delta^{(k+p)}(x) \quad k = 1, 2, \dots \quad p = 0, 1, 2, \dots$
- $x_+^{-r-1/2} \cdot x_-^{-k-1/2} \approx \frac{(-1)^{r+k} \pi}{2(r+k)!} \delta^{(r+k)}(x) \quad r = 0, 1, 2, \dots \quad k = 0, 1, 2, \dots$

# RESULTS ON PRODUCTS OF DISTRIBUTIONS IN COLOMBEAU ALGEBRA

$$\square \quad x_+^{-r-1/2} \cdot x_-^{k-1/2} \approx C_{r,k} \delta^{(r-k)}(x) \quad r = 0, 1, 2, \dots \quad k = 0, 1, 2, \dots \quad r \geq k$$

$$C_{r,k} = \frac{(-1)^r (2k-1)!! k! r! \pi}{2(4k-1)!! (2r-1)!! (r-k)! (r+k)!} \sum_{q=0}^{2k} (-1)^q \binom{2k}{q} \binom{r-k}{k-q} (2(r+q)-1)!! (2(k-q)-1)!!$$

$$x_-^{-k} \cdot \delta^{(p)}(x) \approx \frac{k \cdot p!}{(p+k+1)!} \delta^{(k+p)}(x) \quad k = 1, 2, \dots \quad p = 0, 1, 2, \dots$$

# NEW RESULTS ON PRODUCTS OF DISTRIBUTIONS IN COLOMBEAU ALGEBRA

---

$$\square \quad \ln^2 |x| \cdot \delta^{(s-1)}(x) \approx \frac{2}{s^2} \delta^{(s-1)}(x) \quad s = 1, 2, \dots \quad (30)$$

$$\square \quad \ln^3 |x| \cdot \delta^{(s-1)}(x) \approx \frac{-3!}{s^3} \delta^{(s-1)}(x) \quad s = 1, 2, \dots \quad (31)$$

$$\square \quad \ln^r |x| \cdot \delta^{(s-1)}(x) \approx \frac{(-1)^r r!}{s^r} \delta^{(s-1)}(x) \quad s = 1, 2, \dots \quad r = 0, 1, 2, \dots \quad (32)$$

---

***THANK YOU FOR YOUR  
ATTENTION***