



TITLE:

Finite time blow up and concentration phenomena for a solution to drift-diffusion equations in higher dimensions

AUTHOR(S):

Ogawa, Takayoshi; Suguro, Takeshi; Wakui, Hiroshi

CITATION:

Ogawa, Takayoshi ...[et al]. Finite time blow up and concentration phenomena for a solution to drift-diffusion equations in higher dimensions. *Calculus of Variations and Partial Differential Equations* 2023, 62(2): 47.

ISSUE DATE:

2023-03

URL:

<http://hdl.handle.net/2433/284972>

RIGHT:

© The Author(s) 2022; This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.



Finite time blow up and concentration phenomena for a solution to drift-diffusion equations in higher dimensions

Takayoshi Ogawa¹ · Takeshi Suguro²  · Hiroshi Wakui³Received: 24 July 2022 / Accepted: 1 October 2022 / Published online: 24 December 2022
© The Author(s) 2022

Abstract

We show the finite time blow up of a solution to the Cauchy problem of a drift-diffusion equation of a parabolic-elliptic type in higher space dimensions. If the initial data satisfies a certain condition involving the entropy functional, then the corresponding solution to the equation does not exist globally in time and blows up in a finite time for the scaling critical space. Besides there exists a concentration point such that the solution exhibits the concentration in the critical norm. This type of blow up was observed in the scaling critical two dimensions. The proof is based on the profile decomposition and the Shannon inequality in the weighted space.

Mathematics Subject Classification Primary 35K15 · 35K55 · 35Q60 secondary 78A35

1 Drift-diffusion system in higher dimensions

We consider the finite time blow-up for the solution to the drift-diffusion system in spatially higher dimensions. Let u be a solution to the Cauchy problem of the drift-diffusion system:

$$\begin{cases} \partial_t u - \Delta u + \nabla \cdot (u \nabla \psi) = 0, & t > 0, x \in \mathbb{R}^n, \\ -\Delta \psi = u, & t > 0, x \in \mathbb{R}^n, \\ u(0, x) = u_0(x), & x \in \mathbb{R}^n, \end{cases} \quad (1.1)$$

where $u = u(t, x)$ denotes particle density and ψ denotes a potential of the particle field. The equation (1.1) is relevant to a model of self-interacting particles (see Biler–Nadzieja [7], see also Biler [2]), the semi-conductor device simulations (see [15, 21, 32]) and a model of the aggregation of mold known as the Keller–Segel system (see [22]). In some models, the equation can be derived from a singular limit problem from a fluid mechanical approximation of

Communicated by A. Mondino.

✉ Takeshi Suguro
suguro@kurims.kyoto-u.ac.jp

¹ Mathematical Institute, Tohoku University, Sendai 980-8578, Japan

² Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan

³ Faculty of Science Division I, Tokyo University of Science, Tokyo 162-8601, Japan

gravitation gaseous stars (cf. [14, 24]). If the solution has enough regularity and integrability, then the problem exhibits an instability of the solution, namely, under certain conditions on the initial data, the solution blows up in finite time. The blow up results can be found in [2, 6, 13, 18, 27, 28, 33, 34, 37, 43, 44]. On the other hand, the global existence and stability are known for two dimensional case ([20, 25, 26, 35–38]). This property is naturally inherited to the system (1.1).

Since u denotes the density of particles, it is natural to consider a non-negative solution, and in this case, the L^1 norm of the solution $u(t)$ is preserved in time and this corresponds to the mass conservation law. Under this setting the question of whether the solution exists globally in time or blows up in a finite time is a basic problem. If the initial data is large and decays fast at spatial infinity, the solution blows up in a finite time. In this paper, we discuss such instability of solution under weaker assumptions on *the initial data that decays slower at space infinity*.

The system (1.1) involves the Poisson equation and the solution u is influenced by a non-local effect from the Green’s function of the Poisson equation. Hence, the large time behavior of the solution is largely depending on the behavior of the solution at spatial infinity. Hence, the weight condition on the data may give a subtle effect on the large time behavior of the solution. Therefore, to eliminate the weight condition or reduce the condition is an interesting problem to (1.1).

The local existence of the solution in both the semi-group approach and the energy method is now well established (cf. [27]). To state results, we define some function spaces: For $s > 0$ and $1 \leq p \leq \infty$,

$$L_s^p(\mathbb{R}^n) \equiv \{f \in L_{\text{loc}}^p(\mathbb{R}^n); \langle x \rangle^s f(x) \in L^p(\mathbb{R}^n)\},$$

where $\langle \cdot \rangle = (1 + |\cdot|^2)^{1/2}$. Noting that $L_s^p(\mathbb{R}^n) \subset L^1(\mathbb{R}^n)$ if $s > n/p$, we recall the existence and uniqueness of the solution for the n dimensional drift-diffusion equation in a critical space $L^{\frac{n}{2}}(\mathbb{R}^n)$.

Definition. Let $n \geq 3$ and $1 \leq p < \infty$. For $u_0 \in L^p(\mathbb{R}^n)$, we call u a mild solution to the system (1.1) if $u(t)$ solves the integral equation

$$u(t) = e^{t\Delta}u_0 - \int_0^t e^{(t-s)\Delta}\nabla \cdot (u(s)\nabla\psi(s)) ds$$

in $C([0, T]; L^p(\mathbb{R}^n))$, where

$$\psi = (-\Delta)^{-1}u \equiv \frac{1}{(n-2)\omega_{n-1}}|x|^{-(n-2)} * u,$$

with $\omega_{n-1} \equiv 2\pi^{\frac{n}{2}}/\Gamma(\frac{n}{2})$ as the surface volume of a unit sphere and $\Gamma(\cdot)$ denotes the gamma function.

Proposition 1.1 (Local well-posedness and conservation laws) *Let $n \geq 3$ and $\frac{n}{2} \leq p < n$. For any $u_0 \in L^p(\mathbb{R}^n)$, there exists $T > 0$ and a unique mild solution (u, ψ) to (1.1) with the initial data u_0 such that $u \in C([0, T]; L^p(\mathbb{R}^n)) \cap L^\theta(0, T; L^q(\mathbb{R}^n))$ with $2/\theta + n/q = 2$ and $q > \frac{n}{2}$. Moreover, the solution has higher regularity $u \in C([0, T]; W^{2,p}(\mathbb{R}^n)) \cap C^1((0, T); L^p(\mathbb{R}^n))$ and it is a strong solution for (1.1). Besides there exists a maximal existence time $T = T_* \leq \infty$ such that if $T_* < \infty$, then for any $\frac{n}{2} < p \leq \infty$,*

$$\overline{\lim}_{t \rightarrow T_*} \|u(t)\|_p = \infty.$$

Furthermore, the solution satisfies the following properties:

(1) If the initial data $u_0 \in L_b^1(\mathbb{R}^n)$ for $b > 0$, then the solution satisfies

$$u \in C([0, T]; L^p(\mathbb{R}^n) \cap L_b^1(\mathbb{R}^n)).$$

(2) If $u_0(x) \geq 0$, then $u(t, x) \geq 0$ for any $(t, x) \in (0, T) \times \mathbb{R}^n$.

(3) If $u_0 \in L^1(\mathbb{R}^n)$, then

$$\|u(t)\|_1 = \|u_0\|_1. \tag{1.2}$$

(4) If in addition $u_0 \in L_b^1(\mathbb{R}^n)$, where $b > 0$ and $p \geq \frac{n}{2}$, then the solution (u, ψ) satisfies

$$H[u(t)] + \int_0^t \int_{\mathbb{R}^n} u(\tau) |\nabla(\log u(\tau) - \psi(\tau))|^2 dx d\tau = H[u_0], \tag{1.3}$$

where

$$H[u(t)] \equiv \int_{\mathbb{R}^n} u(t) \log u(t) dx - \frac{1}{2} \int_{\mathbb{R}^n} u(t) \psi(t) dx. \tag{1.4}$$

(5) If $u_0 \in L^1_2(\mathbb{R}^n)$, then the second moment of the solution satisfies the following identity:

$$\begin{aligned} \int_{\mathbb{R}^n} |x - x'|^2 u(t) dx &= \int_{\mathbb{R}^n} |x - x'|^2 u_0(x) dx + 2(n-2) \int_0^t H[u(s)] ds + 2nt \|u_0\|_1 \\ &\quad - 2(n-2) \int_0^t \int_{\mathbb{R}^n} u(s) \log u(s) dx ds, \end{aligned} \tag{1.5}$$

where $x' \in \mathbb{R}^n$ is an arbitrary point.

The local well-posedness in L^p space is essentially due to the earlier works by Weissler [50] and Giga [15], where they consider the nonlinear heat equations and incompressible Navier–Stokes equations. Since the scaling structure to (1.1) is similar to the Navier–Stokes equation, we may apply those theories and obtain the existence and uniqueness of the mild solution by Kurokiba–Ogawa [28] (see also for the critical case [25] and in the weighted space [27]). If the initial data is non-negative and integrable, then the weak maximum principle and the conservation law of the average assure that the weak solution preserves the total mass (1.2). This is natural consequence from the equation originally appears from the conservation laws (cf. [14, 24]).

On the other hand, if we consider the invariant scaling property, namely, the equation has a scaling invariant property that for $\lambda > 0$,

$$\begin{cases} u_\lambda(t, x) \equiv \lambda^2 u(\lambda^2 t, \lambda x), \\ \psi_\lambda(t, x) \equiv \psi(\lambda^2 t, \lambda x) \end{cases}$$

is invariant scaling for the system. The critical space coincide with the invariant scale is

$$L^\infty(\mathbb{R}_+; L^{\frac{n}{2}}(\mathbb{R}^n)) \times L^\infty(\mathbb{R}_+; L^\infty(\mathbb{R}^n))$$

and for two dimensional case it is $L^\infty(\mathbb{R}_+; L^1(\mathbb{R}^2)) \times L^\infty(\mathbb{R}_+; L^\infty(\mathbb{R}^2))$. The global existence in the scaling critical spaces is a direct consequence of Fujita–Kato’s principle.

Proposition 1.2 (Global existence) *Let $n \geq 3$, $p \geq \frac{n}{2}$, and $b > 0$. Assume that the initial data u_0 is non-negative in $L^p(\mathbb{R}^n) \cap L_b^1(\mathbb{R}^n)$, and for some constant $B_n > 0$ depending only on n such that $\|u_0\|_{\frac{n}{2}} < B_n$.*

(1) Then the corresponding solution u obtained by Proposition 1.1 exists globally in time.

(2) *The solution decays in time:*

$$\|u(t)\|_p \leq C(1+t)^{-\frac{n}{2}\left(1-\frac{1}{p}\right)}.$$

One can find the constant B_n can be chosen as

$$B_n = \frac{8}{nS_n^2} = 8\pi(n-2) \left(\frac{\Gamma(\frac{n}{2})}{\Gamma(n)}\right)^{\frac{2}{n}}$$

(see for instance [13], see also [11] for an improved constant), where S_n denotes the best possible constant of Sobolev’s inequality (cf. Talenti [47]). Under this assumption we see that for some $T > 0$, it holds

$$\|u(t)\|_{\frac{n}{2}} \leq \|u_0\|_{\frac{n}{2}}$$

for $t \in [0, T)$. Then $\|u(t)\|_{\frac{n}{2}}$ is uniformly bounded by B_n . This implies a uniform bound for the solution.

The main subject of this paper is to show an instability result of the mild solution to (1.1). When $n = 2$, the solution to (1.1) exists globally in time or non-negative initial data $u_0 \in L^1(\mathbb{R}^2) \cap L^2_s(\mathbb{R}^2)$ satisfying

$$\int_{\mathbb{R}^2} u_0(x) dx \leq 8\pi.$$

On the other hand, if

$$\int_{\mathbb{R}^2} u_0(x) dx > 8\pi,$$

then the solution blows up in a finite time. Namely, there exists $T_m < \infty$ such that

$$\overline{\lim}_{t \rightarrow T_m} \|u(t)\|_p = \infty$$

for all $1 < p \leq \infty$ (see one dimensional modification [9]).

In Biler [2] and Nagai [33, 34], the finite time blow up of the positive solutions are shown (cf. Kurokiba–Ogawa [27]), under certain conditions for two dimensional case.

While if we consider the higher dimensional case, the invariant space is shifted in $L^{\frac{n}{2}}$ and the L^1 conservation law does not work very well. In this sense, the problem is a “super critical” case. On the other hand, the usage of the entropy functional yields new difficulty to show the finite time blow-up. Biler [3] obtained a finite time blow-up result for the case of bounded domain with a boundary condition. Corrias–Perthame–Zaag [13] obtained the finite time blowing up result for the higher dimensional cases who did not use the entropy functional. Namely, there exists a large constant $M = M(n) > 0$ such that if the initial data satisfies

$$M \leq \frac{\|u_0\|_1^{\frac{n}{n-2}}}{\int_{\mathbb{R}^n} |x - \bar{x}|^2 u_0(x) dx},$$

where \bar{x} is the second mass center u_0 , that is,

$$\int_{\mathbb{R}^n} |x - \bar{x}|^b u_0(x) dx \equiv \inf_{x \in \mathbb{R}^n} \int_{\mathbb{R}^n} |y - x|^b u_0(y) dy.$$

Then the solution blows up in a finite time. In both results, the initial data and solutions are both assumed to be in $L^2_1(\mathbb{R}^n)$, namely, the second moment of the solution remains finite for

all time up to the maximal existence time. This is indeed a natural condition of the system if we regard that the $u(t, x) dx$ as a probability measure assuming $\|u_0\|_1 = 1$ with positivity.

The following statement is essentially due to Biler [3] in a bounded domain and refined by Calvez–Corrias–Ebde [11] for the whole space case and Ogawa–Wakui [41].

Proposition 1.3 (Blow-up criterion) *Let $n \geq 3$ and $b > 0$. For $u_0 \in L^{\frac{n}{2}}(\mathbb{R}^n) \cap L^1_b(\mathbb{R}^n)$, $u_0 \geq 0$, assume further that for some constant $C_{n,b} > 0$,*

$$H[u_0] < -\frac{n}{b} \|u_0\|_1 \log \left(\frac{C_{n,b}}{\|u_0\|_1^{1+\frac{2}{n}}} \int_{\mathbb{R}^n} |x - \bar{x}|^b u_0(x) dx \right), \tag{1.6}$$

where \bar{x} is the b -th mass center of u_0 .

(1) When $b \geq 2$, then

$$C_{n,b} \geq \frac{2\pi e^{\frac{n}{n-2}}}{n} \tag{1.7}$$

and the solution u to (1.1) blows up in a finite time. Namely, for any $\frac{n}{2} < p \leq \infty$, there exists some $T_* < \infty$ such that

$$\overline{\lim}_{t \rightarrow T_*} \|u(t)\|_p = \infty. \tag{1.8}$$

(2) If $u_0 \geq 0$ is radially symmetric and $b > 0$, then

$$C_{n,b} \geq \frac{bc_{n,b} e^{1+\frac{b(1+\delta/n)}{n-2}}}{n} \tag{1.9}$$

and the solution u to (1.1) blows up in a finite time in the sense of (1.8), where $c_{n,b}$ is the constant defined in Proposition 2.1.

(3) When $0 < b < 2$, then the solution does not remain uniformly bounded in $L^{\frac{n}{2}}(\mathbb{R}^n)$.

Our main result is the concentration phenomena for the blowing up solution:

Theorem 1.4 *Let $n \geq 3$ and $b \geq 2$. For $u_0 \in L^{\frac{n}{2}}(\mathbb{R}^n) \cap L^1_b(\mathbb{R}^n)$, $u_0 \geq 0$, assume that (1.6) holds with $C_{n,b} = 2\pi e^{\frac{n}{n-2}}/n$.*

(1) Then the blowing up solution u to (1.1) concentrates the following sense: Let $T_* > 0$ be the blow up time and $\{t_k\}_{k \in \mathbb{N}}$ satisfy $t_k \rightarrow T_*$ and $\|u(t_k)\|_{\frac{n}{2}} \rightarrow \infty$ as $k \rightarrow \infty$. Then for any $\varepsilon > 0$, there exist a subsequence of $\{t_k\}_{k \in \mathbb{N}}$ and $x_* \in \mathbb{R}^n$ such that

$$\left(\frac{2n}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}} \leq \liminf_{k \rightarrow \infty} \int_{B_\varepsilon(x_*)} u(t_k, x)^{\frac{n}{2}} dx, \tag{1.10}$$

where C_{HLS} is the best possible constant of the Hardy–Littlewood–Sobolev inequality, that is,

$$\int_{\mathbb{R}^n} f(-\Delta)^{-1} f dx \leq C_{\text{HLS}} \|f\|_1 \|f\|_{\frac{n}{2}}. \tag{1.11}$$

(2) Furthermore, if $T_* < \infty$, then for any $\varepsilon > 0$, there exist a subsequence of $\{t_k\}_{k \in \mathbb{N}}$ and $\{x_k\}_{k \in \mathbb{N}} \subset \mathbb{R}^n$ such that

$$\left(\frac{2n}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}} \leq \liminf_{k \rightarrow \infty} \int_{B_\varepsilon(\sqrt{T_k - t_k}(x_k))} u(t_k, x)^{\frac{n}{2}} dx, \tag{1.12}$$

where C_{HLS} is defined by the above.

By the assumption (1.6) with $C_{n,b} = 2\pi e^{\frac{n}{n-2}}/n$, we see that

$$-\frac{1}{2} \int_{\mathbb{R}^n} u_0(x)(-\Delta)^{-1} u_0(x) dx < -\frac{n}{2} \|u_0\|_1 \log \left(\frac{2\pi e^{\frac{n}{n-2}}}{n \|u_0\|_1^{1+\frac{2}{n}}} \int_{\mathbb{R}^n} |x - \bar{x}|^2 u_0(x) dx \right) - \int_{\mathbb{R}^n} u_0(x) \log u_0(x) dx.$$

It follows from the Hardy–Littlewood–Sobolev inequality (1.11) and Shannon inequality (2.1) (see Proposition 2.1 below) that

$$-\frac{C_{\text{HLS}}}{2} \|u_0\|_1 \|u_0\|_{\frac{n}{2}} < -\frac{n}{2} \|u_0\|_1 \log \left(e^{\frac{2}{n-2}} \right),$$

which implies that the initial data $u_0 \geq 0$ satisfies

$$\int_{\mathbb{R}^n} u_0(x)^{\frac{n}{2}} dx > \left(\frac{2n}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}}. \tag{1.13}$$

The constant of the right hand side coincides with one appearing in (1.10). We also see that

$$B_n = \frac{8}{nS_n^2} < \frac{2n}{(n-2)C_{\text{HLS}}}.$$

for $n \geq 3$ (see Proposition 2.4 below).

For a radially symmetric solution to (1.1), one can extend the condition of the weighted Lebesgue space. Since the assumption on the initial data is different from the case of $b \geq 2$, the constant appearing in the concentration phenomena changes as follows:

Theorem 1.5 *Let $n \geq 3$ and $b > 0$. For a radially symmetric function $u_0 \in L^{\frac{n}{2}}(\mathbb{R}^n) \cap L_b^1(\mathbb{R}^n)$, $u_0 \geq 0$.*

(1) *If $b \geq 2$ and u_0 satisfies (1.6) with $C_{n,b} = 2\pi e^{\frac{n}{n-2}}/n$, then the blowing up solution u to (1.1) concentrates the following sense: Let $0 < T_* < \infty$ be the blow up time and $\{t_k\}_{k \in \mathbb{N}}$ satisfy $t_k \rightarrow T_*$ and $\|u(t_k)\|_{\frac{n}{2}} \rightarrow \infty$ as $k \rightarrow \infty$. Then for any $\varepsilon > 0$, there exists a subsequence of $\{t_k\}_{k \in \mathbb{N}}$ such that*

$$\left(\frac{2n}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}} \leq \liminf_{k \rightarrow \infty} \int_{B_{\varepsilon\sqrt{T_*-t_k}}(0)} u(t_k, x)^{\frac{n}{2}} dx. \tag{1.14}$$

(2) *If $b > 0$ and u_0 satisfies (1.6) with (1.9) for any $\delta > 0$, then the blowing up solution u to (1.1) concentrates the following sense: Let $0 < T_* < \infty$ be the blow up time and $\{t_k\}_{k \in \mathbb{N}}$ satisfy $t_k \rightarrow T_*$ and $\|u(t_k)\|_{\frac{n}{2}} \rightarrow \infty$ as $k \rightarrow \infty$. Then for any $\varepsilon > 0$, there exist a subsequence of $\{t_k\}_{k \in \mathbb{N}}$ such that*

$$\left(\frac{2(n+\delta)}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}} \leq \liminf_{k \rightarrow \infty} \int_{B_{\varepsilon\sqrt{T_*-t_k}}(0)} u(t_k, x)^{\frac{n}{2}} dx. \tag{1.15}$$

Theorem 1.5 gives the $L^{\frac{n}{2}}$ -concentration rate of the radially symmetric blow-up solution and is the analogous result of a nonlinear Schrödinger equation (see Merle–Tsutsumi [31] and Tsutsumi [48]).

Remark In the derivation of (1.13), if we assume that u_0 satisfies (1.6) with (1.9) instead of $C_{n,b} = 2\pi e^{\frac{n}{n-2}}/n$, then we see that

$$\int_{\mathbb{R}^n} u_0(x)^{\frac{n}{2}} dx > \left(\frac{2(n+\delta)}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}}.$$

While we relax the weight condition from $b \geq 2$ to $b > 0$ in Theorem 1.5, the assumption of the initial data needs to be tightened when $b > 0$. The result of Theorem 1.5 (2) follows from the analogous argument of Theorem 1.4 naturally.

It is worth comparing the above result and the case of the degenerate drift-diffusion system:

$$\begin{cases} \partial_t u - \Delta u^\alpha + \nabla \cdot (u \nabla \psi) = 0, & t > 0, x \in \mathbb{R}^n, \\ -\Delta \psi = u, & t > 0, x \in \mathbb{R}^n, \\ u(0, x) = u_0(x), & x \in \mathbb{R}^n, \end{cases} \quad (1.16)$$

where $\alpha > 1$ denotes the adiabatic constant originated from the pressure term $P(\rho) = c\rho^\alpha$ in the barotropic damped compressible Navier–Stokes–Poisson system (cf. [14, 24]). It is known that there are two critical exponents $\alpha_* = 2 - 4/(n+2)$ and $\alpha^* = 2 - 2/n$ (see [8, 12, 40, 42, 45, 46, 49]). In the case $\alpha_* \leq \alpha \leq \alpha^*$ (see [23]), the threshold for the global existence of the weak solution to (1.16) is identified as

$$\|u_0\|_1^\beta \|u_0\|_\alpha^\gamma < \|V\|_1^\beta \|V\|_\alpha^\gamma,$$

where

$$\beta \equiv \frac{2}{2-\alpha} - \frac{n}{\alpha}, \quad \gamma \equiv n - \frac{2}{2-\alpha}$$

and V is the optimizer for the Hardy–Littlewood–Sobolev inequality

$$\int_{\mathbb{R}^n} f(x)(-\Delta)^{-1} f(x) dx \leq C_{\text{HLS},\alpha} \|f\|_1^{1-\sigma} \|f\|_\alpha^{1+\sigma}$$

with

$$\sigma \equiv \frac{\alpha}{\alpha-1} \frac{n-2}{n} - 1.$$

In particular case of $\beta = 0$ when $\alpha = \alpha_*$, the threshold is

$$\int_{\mathbb{R}^n} u_0(x)^{\alpha_*} dx < \int_{\mathbb{R}^n} V(x)^{\alpha_*} dx = \left(\frac{2n}{(n-2)C_{\text{HLS},\alpha_*}} \right)^{\frac{n}{2}}.$$

From this estimate, the above lower bound for the concentration coincides the threshold for the global existence of the weak solution to the degenerate drift-diffusion equation in higher dimensions.

The proof of Theorem 1.4 is based on the profile decomposition in $L^1(\mathbb{R}^n)$. We take $\{t_k\}_{k \in \mathbb{N}}$ such that $t_k \rightarrow T$ as $k \rightarrow \infty$ and introduce the rescaled solution sequence

$$u_k(x) \equiv \lambda_k^{-n} u(t_k, \lambda_k^{-1} x) \quad \text{with } \lambda_k \equiv \|u(t_k)\|_{\frac{n}{2}}^{-\frac{1}{n-2}}$$

for a blowing up solution u to (1.1). Then $\|u_k\|_1 = \|u_0\|_1$ and $\|u_k\|_{\frac{n}{2}} = 1$. In Bedrossian–Kim [1], they showed the profile decomposition in $L^1(\mathbb{R}^n)$, but their technique allows that the profile depends on k . In this paper, we improve the extraction of the profile independent of k from $\{u_k\}_{k \in \mathbb{N}}$. In order to deny the possibility that $\{u_k\}$ is vanishing, we use the bump

function method for the rescaled equation (see Biler [2], Biler–Cieślak–Karch–Zienkiewicz [4], Biler–Karch–Zienkiewicz [5]).

2 Preliminaries

The following inequality originally due to Shannon is useful to estimate the logarithmic functional (see [29, 41]).

Proposition 2.1 (Generalized Shannon’s inequality) *Let $n \geq 2$ and $b > 0$. There exists a constant $C_{n,b} > 0$ such that for any non-negative function $f \in L^1_b(\mathbb{R}^n)$,*

$$-\int_{\mathbb{R}^n} f(x) \log f(x) \, dx \leq \frac{n}{b} \|f\|_1 \log \left(\frac{C_{n,b}}{\|f\|_1^{1+\frac{b}{n}}} \int_{\mathbb{R}^n} |x - \bar{x}|^b f(x) \, dx \right), \quad (2.1)$$

where \bar{x} is the b -th mass center of f and

$$C_{n,b} = \frac{bec_{n,b}}{n}, \quad c_{n,b} = \left(\frac{2\pi^{\frac{n}{2}} \Gamma(\frac{n}{b})}{b \Gamma(\frac{n}{2})} \right)^{\frac{b}{n}}$$

is the best possible. In particular, if $b = 2$, then

$$C_{n,2} = \frac{2\pi e}{n}.$$

Lemma 2.2 *Let $n \geq 2$ and $b > 0$. There exists a constant $C > 0$ such that for all $f \in L^{\frac{n}{2}}(\mathbb{R}^n) \cap L^1_b(\mathbb{R}^n)$,*

$$\|f\|_1 \leq C \|f\|_{\frac{n}{2}}^{\frac{b}{n-2+b}} \left(\int_{\mathbb{R}^n} |x|^b f(x) \, dx \right)^{\frac{n-2}{n-2+b}}.$$

From this lemma, we see that

$$\|u_0\|_1^{\frac{n-2+b}{n-2}} \left(\int_{\mathbb{R}^n} |x|^b u(t) \, dx \right)^{-1} \leq C^{\frac{n-2+b}{n-2}} \|u(t)\|_{\frac{n}{2}}^{\frac{b}{n-2}}$$

and there is a limitation of the blow-up speed of the $L^{\frac{n}{2}}$ -norm of the solution u to (1.1) and the speed of the b -th moment.

The following inequality is well-known and will be used in later often:

Proposition 2.3 (Hardy–Littlewood–Sobolev inequality) *Let $n \geq 3$. Then there exists a constant $C_{\text{HLS}} > 0$ which is depending only on n such that for any function $f \in L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$,*

$$\| |\nabla|^{-1} f \|_2^2 \leq C_{\text{HLS}} \|f\|_1 \|f\|_{\frac{n}{2}}. \quad (2.2)$$

Moreover, there exists the extremal function $V \in L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$ such that V is radially symmetric and decreasing function satisfying the Euler–Lagrange equation

$$\begin{cases} (-\Delta)^{-1} V = \frac{n}{4} C_{\text{HLS}} V^{\frac{n}{2}-1} + \frac{1}{2} C_{\text{HLS}} \chi_{\text{supp } V} & \text{in } B_R, \\ V > 0 & \text{in } B_R, \\ V = 0 & \text{in } \mathbb{R}^n \setminus B_R \end{cases} \quad (2.3)$$

for some $0 < R < \infty$.

Proof of Proposition 2.3 The Hardy–Littlewood–Sobolev inequality implies that it follows that

$$\iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|f(x)||f(y)|}{|x-y|^{n-2}} dx dy \leq C_{n,\alpha} \|f\|_\alpha \|f\|_r,$$

where $1 < \alpha, r < \infty$ satisfy

$$\frac{1}{\alpha} + \frac{n-2}{n} + \frac{1}{r} = 2, \quad \text{i.e.,} \quad \frac{1}{\alpha} - \frac{2}{n} + \frac{1}{r} = 1.$$

By Hölder’s inequality, we have

$$\iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|f(x)||f(y)|}{|x-y|^{n-2}} dx dy \leq C_{n,\alpha} \|f\|_1^{1-\theta} \|f\|_\alpha^{1+\theta},$$

where $0 \leq \theta \leq 1$ satisfies

$$1 + \theta = \frac{\alpha}{\alpha-1} \frac{n-2}{n}.$$

In particular, let $\alpha = \frac{n}{2}$, then $\theta = 0$, so that the inequality

$$\iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|f(x)||f(y)|}{|x-y|^{n-2}} dx dy \leq C_n \|f\|_1 \|f\|_{\frac{n}{2}}$$

holds for any function $f \in L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$. By the definition of the Poisson kernel, we can rewrite as

$$\left| \int_{\mathbb{R}^n} f(x) (-\Delta)^{-1} f(x) dx \right| \leq C_{\text{HLS}} \|f\|_1 \|f\|_{\frac{n}{2}},$$

where C_{HLS} denotes the best possible constant of the Hardy-Littlewood-Sobolev inequality and $C_{\text{HLS}} = C_n c_n$. This implies the inequality (2.2). The attainability and Euler–Lagrange equation (2.3) of the inequality (2.2) is proved by Kimijima–Nakagawa–Ogawa [23] (see also [10]). □

Proposition 2.4 (Comparison of constants) *The best constant C_{HLS} is estimated from the above by*

$$C_{\text{HLS}} < S_n^2, \tag{2.4}$$

where S_n is the best constant of Sobolev’s inequality

$$\|f\|_{\frac{2n}{n-2}} \leq S_n \|\nabla f\|_2,$$

given by

$$S_n^2 = \frac{1}{\pi n(n-2)} \left(\frac{\Gamma(\frac{n}{2})}{\Gamma(n)} \right)^{\frac{2}{n}}$$

for all $f \in \dot{H}^1(\mathbb{R}^n)$. In particular, for $n \geq 3$,

$$\frac{8}{n S_n^2} < \frac{2n}{(n-2) C_{\text{HLS}}}. \tag{2.5}$$

Remark The precise constant for S_n is known by Talenti [47]. In the view of (1.10), it is interesting to observe that

$$\frac{2n}{(n-2)C_{\text{HLS}}}\Big|_{n=2} = \frac{2n}{C_n} \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})}\Big|_{n=2} = 8\pi.$$

Indeed, the best constant C_{HLS} can be decomposed into the explicit constant and implicit one as

$$C_{\text{HLS}} = \frac{1}{(n-2)\omega_{n-1}} C_n,$$

where C_n is the best constant defined by

$$C_n \equiv \sup_{\substack{f \in L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n), \\ f \geq 0}} \frac{\iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{f(x)f(y)}{|x-y|^{n-2}} dx dy}{\|f\|_1 \|f\|_{\frac{n}{2}}}.$$

In particular, the best constant C_n becomes 1 when $n = 2$. In this case, the role of the constant coincides with the threshold in the case of $n = 2$.

Proof of Proposition 2.4 It is easy to see that the best constant S_n^2 also gives the best constant for the inequality

$$\| |\nabla|^{-1} f \|_2^2 \leq S_n^2 \|f\|_{\frac{2n}{n+2}}$$

for any $f \in L^{2n/(n+2)}(\mathbb{R}^n)$ by the duality argument (see for instance [40]). Hence, by Hölder’s inequality, one can observe that

$$\int_{\mathbb{R}^n} f(x)(-\Delta)^{-1} f(x) dx \leq S_n^2 \|f\|_1 \|f\|_{\frac{n}{2}}$$

holds for any $f \in L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$. This shows $C_{\text{HLS}} \leq S_n^2$. To see that $C_{\text{HLS}} < S_n^2$, we assume on the contrary that $C_{\text{HLS}} = S_n^2$ for $n \geq 3$. Then we may show that there exists an extremal function $V \in L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$ such that it attains the best constant, that is,

$$\| |\nabla|^{-1} V \|_2^2 = S_n^2 \|V\|_1 \|V\|_{\frac{n}{2}}.$$

Then by the Hölder and Sobolev inequalities,

$$\| |\nabla|^{-1} V \|_2^2 = S_n^2 \|V\|_1 \|V\|_{\frac{n}{2}} \geq S_n^2 \|V\|_{\frac{2n}{n+2}}^2 \geq \| |\nabla|^{-1} V \|_2^2.$$

This shows that V is also the extremal function that attains the best possible constant of Sobolev’s inequality, that is,

$$\| |\nabla|^{-1} V \|_2^2 = S_n^2 \|V\|_{\frac{2n}{n+2}}^2.$$

Since the extremal function of this inequality is uniquely identified up to translation and dilation, namely,

$$V(x) = a_n \left(1 + \frac{1}{n(n-2)} |x|^2 \right)^{-\frac{n+2}{2}}, \text{ where } a_n \equiv \left(\frac{2n}{n-2} \right)^{\frac{n+2}{2}}. \tag{2.6}$$

On the other hand, the extremal function of the Hardy–Littlewood–Sobolev inequality (2.2) must satisfy the Euler–Lagrange equation (2.3). Clearly, the Talenti function (2.6) does not satisfy (2.3). This is a contradiction, and hence, we obtain (2.4). Moreover, it follows from (2.4) that

$$\frac{8}{nS_n^2} < \frac{8}{nC_{HLS}} < \frac{2n}{(n-2)C_{HLS}}$$

for any $n \geq 3$. Thus, we conclude that (2.5) holds.

As an alternative proof of (2.4), since the extremal function of this inequality is uniquely identified up to translation and dilation, one can compute $\|V\|_1 \|V\|_{\frac{n}{2}}$ and $\|V\|_{2n/(n+2)}$ explicitly. Since it holds that

$$\|V\|_q^q = \frac{1}{2} a_n^q (n(n-2))^{\frac{n}{2}} \omega_{n-1} B\left(\frac{n}{2}, \frac{q(n+2)}{2} - \frac{n}{2}\right),$$

where $B(\cdot, \cdot)$ is the beta function, the problem can be reduced as the comparison between

$$B\left(\frac{n}{2}, 1\right) B\left(\frac{n}{2}, \frac{n^2}{4}\right)^{\frac{2}{n}} = \frac{2}{n} \left(\frac{\Gamma(\frac{n}{2})\Gamma(\frac{n^2}{4})}{\Gamma(\frac{n}{2} + \frac{n^2}{4})}\right)^{\frac{2}{n}} \geq \Gamma\left(\frac{n}{2}\right)^{\frac{2}{n}} \left(\frac{(2/n)^{\frac{n}{2}} \Gamma(\frac{n^2}{4})}{\Gamma(\frac{(n+1)^2}{4})}\right)^{\frac{2}{n}}$$

$$\text{and } B\left(\frac{n}{2}, \frac{n}{2}\right)^{\frac{n+2}{n}} = \left(\frac{\Gamma(\frac{n}{2})^2}{\Gamma(n)}\right)^{1+\frac{2}{n}} = \Gamma\left(\frac{n}{2}\right)^{1+\frac{2}{n}} \left(\frac{\Gamma(\frac{n}{2})}{\Gamma(n)}\right)^{1+\frac{2}{n}}.$$

One can check those values are different from each other by numerical computation¹. For the higher dimensions, one can show it by applying the Stirling formula

$$\Gamma(z) \simeq (2\pi)^{\frac{1}{2}} e^{-z} z^{z-\frac{1}{2}} \quad \text{for } z \gg 1.$$

The ratio of norms of V is estimated by

$$\begin{aligned} \frac{\|V\|_1 \|V\|_{\frac{n}{2}}}{\|V\|_{\frac{2n}{n+2}}^2} &= \frac{2}{n} \frac{\Gamma(n)}{\Gamma(\frac{n}{2})^2} \left(\frac{\Gamma(n)\Gamma(\frac{n^2}{4})}{\Gamma(\frac{n}{2})\Gamma(\frac{n}{2} + \frac{n^2}{4})}\right)^{\frac{2}{n}} \\ &\simeq (2\pi)^{-\frac{1}{2}} 2^{\frac{n}{2}+2} n^{-\frac{3}{2}} \left(\frac{1}{2} + \frac{1}{n}\right)^{-\frac{n}{2}-1+\frac{1}{n}} > 1 \quad \text{for } n \gg 1. \end{aligned}$$

For example, in the case of $n = 200$, we compute

$$(2\pi)^{-\frac{1}{2}} 2^{\frac{n}{2}+2} n^{-\frac{3}{2}} \left(\frac{1}{2} + \frac{1}{n}\right)^{-\frac{n}{2}-1+\frac{1}{n}} = (2\pi)^{-\frac{1}{2}} 2^{126-\frac{1}{2}} 5^{-\frac{9}{2}} \left(\frac{125}{63}\right)^{126-\frac{1}{250}} > 1.$$

□

3 Profile decomposition in L^1

The following decomposition is originally due to Gerard [16] (cf. Nawa [39]) and extended by many authors. Here we show a slightly modified version of the result due to Bedrossian–Kim [1] (see also Hmidi–Keraani [19]).

¹ The authors checked this for $n = 4$ rigorously and up to $n = 300$ numerically.

Lemma 3.1 (Profile decomposition in L^1) *Let $\{f_k\}_{k \in \mathbb{N}} \subset L^1(\mathbb{R}^n) \cap L \log L(\mathbb{R}^n)$ satisfy*

$$f_k \geq 0, \quad \|f_k\|_1 = M, \quad \int_{\mathbb{R}^n} f_k(x) \log f_k(x) dx \leq L \tag{3.1}$$

for some constant $M, L > 0$. Then for all $\varepsilon > 0$, there exists a subsequence of $\{f_k\}$ (not relabeled), $J \in \mathbb{N} \cup \{0\}$, $\{x_k^{(j)}\}_{k \in \mathbb{N}, j=1, \dots, J} \subset \mathbb{R}^n$, $\{R^{(j)}\}_{j=1}^J \subset \mathbb{R}_+$, and function sequences $\{F^{(j)}\}_{j=1}^J$, $\{w_k\}_{k \in \mathbb{N}}$, $\{e_k\}_{k \in \mathbb{N}} \subset L^1(\mathbb{R}^n)$ which satisfy

$$f_k(x) = \sum_{j=1}^J F^{(j)}(x - x_k^{(j)}) + w_k(x) + e_k(x) \quad \text{a.a. } x \in \mathbb{R}^n$$

with the following properties:

- (1) *The profile $\{F^{(j)}\}$ is a nonnegative function sequence satisfying that for each $k \in \mathbb{N}$, $\text{supp } F^{(j)} \subset B_{R_j}(\mathbf{0})$ and for $j \neq j'$, $|x_k^{(j)} - x_k^{(j')}| \rightarrow \infty$ as $k \rightarrow \infty$. Moreover, for each $k \in \mathbb{N}$, $B_{R^{(j)}}(x_k^{(j)})$ and $\text{supp } w_k$ are all disjoint for any $j = 1, \dots, J$.*
- (2) *$\{w_k\}$ is the vanishing part, that is, for all $R > 0$,*

$$\lim_{k \rightarrow \infty} \sup_{x \in \mathbb{R}^n} \int_{B_R(x)} w_k(x) dx = 0,$$

and $0 \leq w_k \leq f_k$ almost everywhere $x \in \mathbb{R}^n$.

- (3) *$\{e_k\}$ is the error term, that is,*

$$\overline{\lim}_{k \rightarrow \infty} \|e_k\|_1 < \varepsilon$$

and $e_k \leq f_k$ almost everywhere $x \in \mathbb{R}^n$.

In particular, passing $k \rightarrow \infty$, we have the almost orthogonality

$$\|f_k\|_1 = \sum_{j=1}^J \|F^{(j)}\|_1 + \|w_k\|_1 + \varepsilon.$$

Proof of Lemma 3.1 By the argument of the concentration compactness lemma in [30], there exists a subsequence $\{f_k\}_{k \in \mathbb{N}}$ (not relabeled) such that one of the following situation occurs for $\{f_k\}$: (1) the compactness, (2) the vanishing, or (3) the dichotomy. We fix $\varepsilon > 0$ arbitrarily. *Case 1. Compactness:* If the compactness occurs, then there exists a sequence $\{x_k^{(1)}\}_{k \in \mathbb{N}} \subset \mathbb{R}^n$ such that

$$\int_{B_{R^{(1)}}(x_k^{(1)})} f_k(y) dy \geq M - \frac{\varepsilon}{2}.$$

for some radius $R^{(1)} > 0$ independent of k . We set

$$\tilde{f}_k(x) \equiv f_k(x + x_k^{(1)}) \chi_{B_{R^{(1)}}(\mathbf{0})}(x)$$

for $x \in \mathbb{R}^n$. By the assumption (3.1), we see that

$$\int_{\mathbb{R}^n} \tilde{f}_k(x) \log \tilde{f}_k(x) dx \leq L.$$

On the other hand, we have

$$\int_{\mathbb{R}^n} \tilde{f}_k(x) \log \tilde{f}_k(x) dx = \int_{B_{R^{(1)}(x_k^{(1)})}} f_k(x) \log f_k(x) dx \geq -e^{-1} |B_{R^{(1)}}|.$$

Thus, it holds that

$$\left| \int_{\mathbb{R}^n} \tilde{f}_k(x) \log \tilde{f}_k(x) dx \right| \leq \max\{L, e^{-1} |B_{R^{(1)}}|\},$$

which implies that $\{\tilde{f}_k\}$ is uniformly integrable. Then the Dunford–Pettis theorem implies that there exist a subsequence $\{\tilde{f}_{k_l}\}_{l \in \mathbb{N}} \subset \{\tilde{f}_k\}$ and a nonnegative function $\tilde{F}^{(1)} \in L^1(B_{R^{(1)}}(0))$ such that \tilde{f}_{k_l} converges weakly to $\tilde{F}^{(1)}$ in $L^1(B_{R^{(1)}}(0))$. Thus, we set a profile $F^{(1)}$ as the zero expansion of $\tilde{F}^{(1)}$, that is,

$$F^{(1)}(x) \equiv \begin{cases} \tilde{F}^{(1)}(x - x_k^{(1)}), & x \in B_{R^{(1)}}(x_k^{(1)}), \\ 0, & \text{otherwise.} \end{cases}$$

Moreover, by the weak lower continuity of norm, we have

$$M - \frac{\varepsilon}{2} \leq \|F^{(1)}\|_1 \leq M.$$

On the other hand, we set

$$e_{k_l}(x) \equiv f_{k_l}(x) - F^{(1)}(x),$$

then e_{k_l} converges weakly to 0 in L^1 . In this case, we define $w_k \equiv 0$. Then the claim in Lemma 3.1 holds in $J = 1$.

Case 2. Vanishing: If the vanishing occurs, it follows that

$$\lim_{k \rightarrow \infty} \sup_{x \in \mathbb{R}^n} \int_{B_R(x)} f_k(y) dy = 0$$

for arbitrary $R > 0$. Define $w_k \equiv f_k$ and $e_k \equiv 0$, then we have the claim of Lemma 3.1 with $J = 0$.

Case 3. Dichotomy: If the dichotomy occurs, there exists $\mu \in (0, M)$ such that there exist the non-negative function sequences $\{f_{k,1}\}_{k \in \mathbb{N}}, \{f_{k,2}\}_{k \in \mathbb{N}} \subset L^1(\mathbb{R}^n)$ which satisfy

$$f_k = f_{k,1} + f_{k,2} + h_k,$$

where we define $h_k \equiv f_k - (f_{k,1} + f_{k,2})$. More precisely, there exist $\{x_k^{(1)}\}_{k \in \mathbb{N}} \subset \mathbb{R}^n$, $R^{(1)} > 0$, and $\{R_k^{(1)}\}_{k \in \mathbb{N}} \subset \mathbb{R}_+$ such that

$$f_{k,1} = f_k|_{B_{R^{(1)}}(x_k^{(1)})}, \quad f_{k,2} = f_k|_{\mathbb{R}^n \setminus B_{R_k^{(1)}}(x_k^{(1)})} \quad \text{with} \quad \lim_{k \rightarrow \infty} R_k^{(1)} = \infty$$

with the following estimates:

$$\begin{cases} \overline{\lim}_{k \rightarrow \infty} \|\|f_{k,1}\|_1 - \mu\| < \frac{\varepsilon}{2}, \\ \overline{\lim}_{k \rightarrow \infty} \|\|f_{k,2}\|_1 - (M - \mu)\| < \frac{\varepsilon}{2}, \\ \overline{\lim}_{k \rightarrow \infty} \|h_k\|_1 < \frac{\varepsilon}{2}, \end{cases}$$

and

$$\lim_{k \rightarrow \infty} \text{dist}(\text{supp } f_{k,1}, \text{supp } f_{k,2}) = \infty. \tag{3.2}$$

These properties imply that $f_{k,1}$, $f_{k,2}$, and h_k are the compact, escape, and error term, respectively. Firstly, similarly to Case 1, we set

$$\tilde{f}_{k,1}(x) \equiv f_{k,1}(x + x_k^{(1)}) = f_k(x + x_k^{(1)})\chi_{B_{R(1)}(0)}(x),$$

and use the Dunford–Pettis theorem, we can get a weak convergence subsequence of $\tilde{f}_{k,1}$ so that we can set a profile $F^{(1)}$, which has a compact support and satisfies

$$\mu - \varepsilon \leq \|F^{(1)}\|_1 \leq \mu.$$

On the other hand, the escape term $f_{k,2}$ satisfies

$$\overline{\lim}_{l \rightarrow \infty} \|f_{k,2}\|_1 = M - \mu.$$

Secondary, we divide some cases whether the mass of the escape term is small or not. If we can take $\mu \in (0, M)$ such that

$$\overline{\lim}_{k \rightarrow \infty} \|f_{k,2}\|_1 < \frac{\varepsilon}{2},$$

then we define $e_k \equiv f_k - F^{(1)}(x - x_k^{(1)})$, and hence, the claim in Lemma 3.1 holds with $J = 1$. In the case of

$$\overline{\lim}_{k \rightarrow \infty} \|f_{k,2}\|_1 \geq \frac{\varepsilon}{2},$$

we reset

$$\bar{f}_{k,2} \equiv \frac{1}{\|f_{k,2}\|_1} f_{k,2}.$$

Then $\|\bar{f}_{k,2}\|_1 = 1$. We apply the concentration compactness lemma to $\{\bar{f}_{k,2}\}_{k \in \mathbb{N}}$. Then there exists a subsequence $\{\bar{f}_{k,2}\}$ (not relabeled) such that one of the following situation occur for $\{\bar{f}_{k,2}\}$: (1) the compactness, (2) the vanishing, or (3) the dichotomy.

Case 3.1. Compactness: If the compactness occurs, then there exists $\{x_k^{(2)}\}_{k \in \mathbb{N}} \subset \mathbb{R}^n$ such that

$$\int_{B_{R(2)}(x_k^{(2)})} \bar{f}_{k,2}(y) dy \geq 1 - \frac{\varepsilon}{2(M - \mu)}$$

for some radius $R^{(2)} > 0$. We note that passing $k \rightarrow \infty$, then $|x_k^{(1)} - x_k^{(2)}| \rightarrow \infty$ by (3.2). Then we set

$$\tilde{f}_{k,2}(x) \equiv f_{k,2}(x + x_k^{(2)})\chi_{B_{R(2)}(0)}(x)$$

Similarly to Case 1, we get a profile $F^{(2)}$ as a weak convergence limit of $\tilde{f}_{k,2}$ and define

$$e_k(x) \equiv f_k(x) - (F^{(1)}(x) + F^{(2)}(x)).$$

Then the claim in Lemma 3.1 holds in $J = 2$.

Case 3.2. Vanishing: If the vanishing occurs, similarly to Case 2, we define

$$w_k \equiv f_{k,2}, \quad e_k \equiv f_k - F^{(1)} - f_{k,2}.$$

Then we have the claim of Lemma 3.1 with $J = 1$.

Case 3.3. Dichotomy: If the dichotomy occurs, there exists $\nu \in (0, 1)$ such that there exists the non-negative function sequences $\{\bar{f}_{k,2,1}\}, \{\bar{f}_{k,2,2}\} \subset L^1(\mathbb{R}^n)$ which satisfy

$$\bar{f}_{k,2} = \bar{f}_{k,2,1} + \bar{f}_{k,2,2} + \bar{h}_k,$$

where we define $\bar{h}_k \equiv \bar{f}_{k,2} - (\bar{f}_{k,2,1} + \bar{f}_{k,2,2})$. For each sequences, there exist $\{x_k^{(2)}\}_{k \in \mathbb{N}} \subset \mathbb{R}^n$, $R^{(2)} > 0$, and $\{R_k^{(2)}\}_{k \in \mathbb{N}} \subset \mathbb{R}_+$ such that

$$\bar{f}_{k,2,1} = \bar{f}_{k,2}|_{B_{R^{(2)}}(x_k^{(2)})}, \quad \bar{f}_{k,2,2} = \bar{f}_{k,2}|_{\mathbb{R}^n \setminus B_{R_k^{(2)}}(x_k^{(2)})} \quad \text{with} \quad \lim_{k \rightarrow \infty} R_k^{(2)} = \infty$$

with the following estimates:

$$\begin{cases} \overline{\lim}_{l \rightarrow \infty} \|\bar{f}_{k,2,1}\|_1 - \nu < \frac{\varepsilon}{4(M - \mu)}, \\ \overline{\lim}_{l \rightarrow \infty} \|\bar{f}_{k,2,2}\|_1 - (1 - \nu) < \frac{\varepsilon}{4(M - \mu)}, \\ \overline{\lim}_{l \rightarrow \infty} \|\bar{h}_k\|_1 < \frac{\varepsilon}{4(M - \mu)}. \end{cases}$$

By the definition of $\bar{f}_{k,2}$, we can define $h_{k,2} \equiv \|f_{k,2}\|_1 \bar{h}_k$ so that we have

$$\overline{\lim}_{k \rightarrow \infty} \|h_{k,2}\|_1 < \frac{\varepsilon}{2^3}.$$

If we can take $\nu \in (0, 1)$ such that

$$\overline{\lim}_{l \rightarrow \infty} \|f_{k,2,2}\|_1 < \frac{\varepsilon}{2}, \quad f_{k,2,2} \equiv \|f_{k,2}\|_1 \bar{f}_{k,2,2}$$

then we take a subsequence $\{k_l\}$ which satisfies the above estimates and define

$$e_k(x) \equiv f_k(x) - F^{(1)}(x) - F^{(2)}(x).$$

Then the claim in Lemma 3.1 holds with $J = 2$. In the case of

$$\overline{\lim}_{k \rightarrow \infty} \|f_{k,2,2}\|_1 \geq \frac{\varepsilon}{2},$$

we apply the same argument above many times.

The proof of Lemma 3.1 relies upon the induction argument. We omit to prove in detail. Note that the argument must be terminated in a finite step. Indeed, we assume on the contrary, these steps continue infinitely. Then it holds that

$$\overline{\lim}_{k \rightarrow \infty} \|f_{k,l,2}\|_1 \geq \frac{\varepsilon}{2}$$

for any $l \in \mathbb{N}$. Take the sum with respect to l , then the total mass $\|f_k\|_1$ diverges, that is,

$$\infty > M = \|f_k\|_1 \geq \sum_{l=1}^{\infty} \|f_{k,l,2}\|_1 \geq \sum_{l=0}^{\infty} \frac{\varepsilon}{2} = \infty.$$

This contradicts the assumption of Lemma 3.1. □

Remark In [1], they also showed the profile decomposition on $L^1(\mathbb{R}^n)$. We emphasize that Lemma 3.1 assures the profile independent of k while the profile in [1] depends on k . The independence of the profile on k is valid in the proof of Theorem 1.4.

4 Concentration

Lemma 4.1 *Let the initial data u_0 satisfy the assumption (1.6) with (1.7), then there exists the corresponding solution $u(t)$ to (1.1) blows up in a finite time T_* . Then it holds that for any $0 < t < T_*$,*

$$\frac{2n}{n-2} \|u_0\|_1 \leq \| |\nabla|^{-1} u(t) \|_2^2. \tag{4.1}$$

Proof of Lemma 4.1 For simplicity, we consider the case of $b = 2$. For the solution $u(t)$ to the equation (1.1), by the entropy dissipation (1.3) and definition of the entropy (1.4), we have

$$\int_{\mathbb{R}^n} u(t) \log u(t) \, dx - H[u_0] \leq \frac{1}{2} \| |\nabla|^{-1} u(t) \|_2^2.$$

By Shannon’s inequality (2.1), it follows that

$$-\frac{n}{2} \|u(t)\|_1 \log \left(\frac{2\pi e}{n \|u(t)\|_1^{1+\frac{2}{n}}} \int_{\mathbb{R}^n} |x - \bar{x}|^2 u(t) \, dx \right) - H[u_0] \leq \frac{1}{2} \| |\nabla|^{-1} u(t) \|_2^2.$$

Since the mass conservation (1.2) holds, then we can rewrite as

$$-\frac{n}{2} \|u_0\|_1 \log \left(\frac{2\pi e}{n \|u_0\|_1^{1+\frac{2}{n}}} \int_{\mathbb{R}^n} |x - \bar{x}|^2 u(t) \, dx \right) - H[u_0] \leq \frac{1}{2} \| |\nabla|^{-1} u(t) \|_2^2. \tag{4.2}$$

While by the virial law (1.5) for (1.1), we have

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^n} |x - \bar{x}|^2 u(t) \, dx \\ &= 2n \|u_0\|_1 + 2(n-2) H[u(t)] - 2(n-2) \int_{\mathbb{R}^n} u(t) \log u(t) \, dx \\ &\leq 2(n-2) \left[H[u_0] + \frac{n}{n-2} \|u_0\|_1 + \frac{n}{2} \|u_0\|_1 \log \left(\frac{2\pi e \int_{\mathbb{R}^n} |x - \bar{x}|^2 u(t) \, dx}{n \|u_0\|_1^{1+\frac{2}{n}}} \right) \right], \end{aligned} \tag{4.3}$$

and hence, under the assumption

$$\begin{aligned} H[u_0] &< \frac{n}{2} \|u_0\|_1 \log \left(\frac{n \|u_0\|_1^{1+\frac{2}{n}}}{2\pi e \int_{\mathbb{R}^n} |x - \bar{x}|^2 u_0 \, dx} \right) \\ &= \frac{n}{2} \|u_0\|_1 \log \left(\frac{n \|u_0\|_1^{1+\frac{2}{n}}}{2\pi e \int_{\mathbb{R}^n} |x - \bar{x}|^2 u_0 \, dx} \right) - \frac{n}{n-2} \|u_0\|_1, \end{aligned} \tag{4.4}$$

we have from (4.3) and (4.4)

$$\frac{d}{dt} \int_{\mathbb{R}^n} |x - \bar{x}|^2 u(t) \, dx \Big|_{t=0} \leq 0.$$

This implies that

$$\int_{\mathbb{R}^n} |x - \bar{x}|^2 u(t) \, dx \leq \int_{\mathbb{R}^n} |x - \bar{x}|^2 u_0 \, dx. \tag{4.5}$$

Thus, it follows from (4.2) and (4.5) that

$$-\frac{n}{2} \|u_0\|_1 \log \left(\frac{2\pi e}{n \|u_0\|_1^{1+\frac{2}{n}}} \int_{\mathbb{R}^n} |x - \bar{x}|^2 u_0 \, dx \right) - H[u_0] \leq \frac{1}{2} \|\nabla^{-1} u(t)\|_2^2.$$

From the assumption (4.4), we obtain

$$\frac{n}{n-2} \|u_0\|_1 \leq \frac{1}{2} \|\nabla^{-1} u(t)\|_2^2$$

for any $0 < t < T_*$. □

In what follows, we show Theorem 1.4 (1). We assume that the initial data $u_0 \in L^{\frac{n}{2}}(\mathbb{R}^n) \cap L^1_b(\mathbb{R}^n)$ satisfies the blow-up condition (1.6) with (1.7) and the solution $u(t)$ to (1.1) blows up at $T_* > 0$. Namely,

$$\limsup_{t \rightarrow T_*} \|u(t)\|_{\frac{n}{2}} = \infty.$$

Let $\{t_k\}_{k \in \mathbb{N}}$ be a sequence that gives the supremum of $\|u(t)\|_{\frac{n}{2}}$. We introduce the scaling transform by $\lambda > 0$ that

$$S_\lambda u(t, x) \equiv \lambda^{-n} u(t, \lambda^{-1} x).$$

This is the L^1 -invariant scaling $\|S_\lambda u(t)\|_1 = \|u(t)\|_1 = \|u_0\|_1$ and it also holds that

$$\|S_\lambda u(t)\|_{\frac{n}{2}} = \lambda^{2-n} \|u(t)\|_{\frac{n}{2}}.$$

For a blow-up solution $u(t)$ and blow-up time sequence $\{t_k\}$, we set

$$\lambda_k \equiv \|u(t_k)\|_{\frac{n}{2}}^{\frac{1}{n-2}} \quad \text{and} \quad \lambda_0 \equiv \|u_0\|_{\frac{n}{2}}^{\frac{1}{n-2}}. \tag{4.6}$$

Then we find that the rescale solution

$$v(t, x) \equiv \lambda_k^{-n} u(t, \lambda_k^{-1} x) \quad \text{for } t_{k-1} < t \leq t_k, \quad x \in \mathbb{R}^n$$

is bounded in $L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$. Note that the scaling is L^1 -invariant and hence the L^1 -norm of v is invariant. The rescaled solution $v(t, x)$ now solves the Cauchy problem of a semistationary equation

$$\begin{cases} \partial_t v - \lambda_k^2 \Delta v + \lambda_k^n \nabla \cdot (v \nabla \phi) = 0, & t_{k-1} < t \leq t_k, \quad x \in \mathbb{R}^n, \\ -\Delta \phi = v, & t > 0, \quad x \in \mathbb{R}^n, \\ v(0, x) = v_0(x) \equiv \lambda_0^{-n} u_0(\lambda^{-1} x), & x \in \mathbb{R}^n, \end{cases} \tag{4.7}$$

where $t_0 = 0$. The solution preserves its total mass $\|v(t)\|_1 = \|u(t)\|_1 = \|u_0\|_1$ for any time $t \geq 0$ and $\|v(t_k)\|_{\frac{n}{2}} = 1$ at $t = t_k$.

We set the rescaled solution sequence

$$u_k(x) \equiv v(t_k) = S_{\lambda_k} u(t_k, x) = \lambda_k^{-n} u(t_k, \lambda_k^{-1} x) \tag{4.8}$$

satisfies

$$\|u_k\|_1 = \|u_0\|_1, \quad \|u_k\|_{\frac{n}{2}} = 1. \tag{4.9}$$

In this case, one can apply Proposition 3.1 with $\{u_k\}_{k \in \mathbb{N}}$:

Proposition 4.2 (Profile decomposition in L^p) *Let $\{u_k\}_{k \in \mathbb{N}}$ be a non-negative sequence in $L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$ with*

$$\|u_k\|_1 = \|u_0\|_1 \quad \text{and} \quad \|u_k\|_{\frac{n}{2}} = 1$$

defined by the above. Then for all $\varepsilon > 0$, there exists a subsequence of $\{u_k\}$ (not re-labeled), $J \in \mathbb{N}$, $\{x_k^{(j)}\}_{k \in \mathbb{N}, j=1, \dots, J} \subset \mathbb{R}^n$, $\{R^{(j)}\}_{j=1}^J \subset \mathbb{R}_+$, and function sequences $\{U^{(j)}\}_{j=1}^J$, $\{w_k\}_{k \in \mathbb{N}}$, $\{e_k\}_{k \in \mathbb{N}} \subset L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$ which satisfy the following properties:

(1) u_k is decomposed as

$$u_k(x) = \sum_{j=1}^J U^{(j)}(x - x_k^{(j)}) + w_k(x) + e_k(x) \quad \text{a.a. } x \in \mathbb{R}^n, \quad (4.10)$$

where $\{U^{(j)}\}_{j=1}^J$ is a nonnegative function sequence with $\text{supp } U^{(j)} \subset B_{R^{(j)}}(0)$. Moreover, for each $k \in \mathbb{N}$, $B_{R^{(j)}}(x_k^{(j)})$ and $\text{supp } w_k$ are all disjoint for any $j = 1, 2, \dots, J$. For any $R > 0$, it holds that

$$\lim_{k \rightarrow \infty} \sup_{x \in \mathbb{R}^n} \int_{B_R(x)} w_k(x) dx = 0 \quad (4.11)$$

and the error term satisfies

$$\overline{\lim}_{k \rightarrow \infty} \|e_k\|_1 < \varepsilon \quad (4.12)$$

(2) *Almost orthogonality:*

$$\|u_k\|_1 = \sum_{j=1}^J \|U^{(j)}\|_1 + \|w_k\|_1 + \varepsilon, \quad (4.13)$$

and for any $j \in \{1, 2, \dots, J\}$, it follows that

$$\|U^{(j)}\|_{\frac{n}{2}} \leq (1 + \varepsilon) \|u_k\|_{L^{\frac{n}{2}}(B_{R^{(j)}}(x_k^{(j)}))}. \quad (4.14)$$

(3) *Drift term estimate:*

$$\overline{\lim}_{k \rightarrow \infty} \|\nabla^{-1} w_k\|_2^2 = \overline{\lim}_{k \rightarrow \infty} \|\nabla^{-1} e_k\|_2^2 = 0. \quad (4.15)$$

We note that the vanishing in the sense of Lions' concentration compactness lemma implies $J = 0$. We emphasize that the sequence $\{u_k\}_{k \in \mathbb{N}}$ is not vanishing and one can extract at least the profile of $\{u_k\}$.

Proof of Proposition 4.2 We apply Lemma 3.1 with $\{u_k\}_{k \in \mathbb{N}} \subset L^1(\mathbb{R}^n)$. We note that $L^{\frac{n}{2}}$ -boundedness implies that $\{u_k\}_{k \in \mathbb{N}}$ is uniformly integrable. The decomposition (4.10), (4.11), (4.12), and (4.13) are shown directly in Lemma 3.1. By the construction of the profile, we see that

$$\tilde{u}_k \equiv u_k(\cdot + x_k^{(1)}) \chi_{B_{R^{(1)}}(0)}(\cdot) \rightharpoonup U^{(1)} \quad \text{in } L^1(B_{R^{(1)}}(0)).$$

Since $\{\tilde{u}_k\}$ is uniformly bounded in $L^{\frac{n}{2}}(B_{R^{(1)}}(0))$, $\{\tilde{u}_k\}_{k \in \mathbb{N}}$ also converges to $U^{(1)}$ weakly in $L^{\frac{n}{2}}(B_{R^{(1)}}(0))$. This argument can be applied for any $j = 1, 2, \dots, J$. Thus, we obtain

(4.14). Moreover, we have

$$\|e_k\|_{\frac{n}{2}} \leq \|u_k\|_{\frac{n}{2}} + \sum_{j=1}^J \|U^{(j)}\|_{\frac{n}{2}} + \|w_k\|_{\frac{n}{2}} \leq J + 2,$$

which implies that $\{e_k\}_{k \in \mathbb{N}}$ is uniformly bounded in $L^{\frac{n}{2}}$.

In what follows, we show (4.15). By the construction of the error term e_k , we have

$$\overline{\lim}_{k \rightarrow \infty} \|e_k\|_1 = 0.$$

We note that $\{e_k\}_{k \in \mathbb{N}}$ is uniformly bounded in $L^{\frac{n}{2}}(\mathbb{R}^n)$. Hence, the Hardy–Littlewood–Sobolev inequality (2.2) shows that

$$\| |\nabla|^{-1} e_k \|_2^2 \leq C_{\text{HLS}} \|e_k\|_1 \|e_k\|_{\frac{n}{2}} \leq C \|e_k\|_1 \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Next, we shall estimate the vanishing term w_k . We fix $\varepsilon > 0$ arbitrarily. For $R > 0$, we separate the non-local term into three parts as follows:

$$\begin{aligned} \| |\nabla|^{-1} w_k \|_2^2 &= c_n \left(\iint_{\{|x-y| < R^{-1}\}} + \iint_{\{R^{-1} < |x-y| < R\}} + \iint_{\{|x-y| > R\}} \right) \frac{w_k(x)w_k(y)}{|x-y|^{n-2}} dx dy \\ &\equiv I_1 + I_2 + I_3, \end{aligned}$$

where $c_n = 1/((n-2)\omega_{n-1})$. For the integral I_1 , by the Hausdorff–Young inequality, we have

$$I_1 = c_n \int_{\mathbb{R}^n} w_k(y) \int_{|x-y| < R^{-1}} \frac{w_k(x)}{|x-y|^{n-2}} dx dy \leq C \|w_k\|_p^2 \|\cdot\|^{2-n} \chi_{|\cdot| < R^{-1}}(\cdot) \|q\|_q,$$

where $\chi_A(\cdot)$ denotes the characteristic function of a set A , $1 \leq p \leq \frac{n}{2}$ and $1 \leq q < \frac{n}{n-2}$ satisfy

$$2 = \frac{2}{p} + \frac{1}{q}.$$

Then the integral of the kernel can be estimated as

$$\int_{|x| < R^{-1}} |x|^{-q(n-2)} dx = \omega_{n-1} \int_0^{R^{-1}} r^{-q(n-2)+n-1} dr = \frac{\omega_{n-1}}{n-q(n-2)} R^{-n+q(n-2)}.$$

By the uniform boundedness of L^1 and $L^{\frac{n}{2}}$ for $\{u_k\}$, there exists $R_0 > 0$ independent of k such that for any $R > R_0$,

$$I_1 \leq C R_0^{-\frac{n}{q}+n-2} < \frac{\varepsilon}{3}.$$

For the integral I_3 , it follows that

$$I_3 = c_n \int_{\mathbb{R}^n} w_k(y) \int_{|x-y| > R} \frac{w_k(x)}{|x-y|^{n-2}} dx dy \leq c_n R^{-(n-2)} \|w_k\|_1^2.$$

Thus, there exists $R_1 > 0$ independent of k such that for any $R > R_1$,

$$I_3 \leq C R_1^{-(n-2)} < \frac{\varepsilon}{3}.$$

We set $\tilde{R} \equiv \max\{R_0, R_1\}$. Lastly, the integral I_2 can be estimated as

$$I_2 = c_n \int_{\mathbb{R}^n} w_k(y) \int_{\tilde{R}^{-1} < |x-y| < \tilde{R}} \frac{w_k(x)}{|x-y|^{n-2}} dx dy \leq c_n \tilde{R}^{n-2} \|w_k\|_1 \sup_{x \in \mathbb{R}^n} \int_{B_{\tilde{R}}(x)} w_k(y) dy.$$

Since w_k is the vanishing term, we obtain

$$\overline{\lim}_{k \rightarrow \infty} I_2 \leq c_n \tilde{R}^{n-2} \lim_{k \rightarrow \infty} \sup_{x \in \mathbb{R}^n} \int_{B_{\tilde{R}}(x)} w_k(y) dy = 0.$$

Thus, we conclude

$$\overline{\lim}_{k \rightarrow \infty} \|\lvert \nabla \rvert^{-1} w_k\|_2 = 0.$$

In order to extract at least one profile, we shall show that $\{v(t_k)\}_{k \in \mathbb{N} \cup \{0\}}$ is not a vanishing sequence by using the bump function method (see [2, 4, 5]). We set the bump function as

$$\eta(x) \equiv (1 - |x|^2)_+^2.$$

For $\varepsilon \in (0, 1/\sqrt{3})$, the Hessian of η satisfies

$$H\eta \leq -c_\varepsilon I \quad \text{for } |x| \leq \varepsilon, \tag{4.16}$$

where $c_\varepsilon \equiv 4(1 - 3\varepsilon^2)$. For $R > 0$ and $a \in \mathbb{R}^n$, we define

$$\eta_R(x) \equiv R^{-4}(R^2 - |x - a|^2)_+^2,$$

which satisfies

$$\Delta \eta_R(x) = 4R^{-4}(n + 2)|x - a|^2 - 4nR^{-2} \geq -4nR^{-2}, \tag{4.17}$$

in particular, η_R is concave on $|x - a| < R\sqrt{n/(n + 2)}$ as (4.16). Multiplying the equation (4.7) by η_R and integrating over $x \in \mathbb{R}^n$, then we have

$$\frac{d}{dt} \int_{\mathbb{R}^n} v(t)\eta_R dx = \lambda_k^2 \int_{\mathbb{R}^n} v(t)\Delta \eta_R dx + \lambda_k^n \int_{\mathbb{R}^n} v(t)\nabla \phi(t) \cdot \nabla \eta_R dx.$$

By the property (4.17), we see that

$$\frac{d}{dt} \int_{\mathbb{R}^n} v(t)\eta_R dx \geq -4nR^{-2}\lambda_k^2 \int_{B_R(a)} v(t) dx + \lambda_k^n \int_{\mathbb{R}^n} v(t)\nabla \phi(t) \cdot \nabla \eta_R dx.$$

On the second term in the right hand side, using Poisson representation and symmetry, we decompose

$$\begin{aligned} & \int_{\mathbb{R}^n} v(t)\nabla \phi(t) \cdot \nabla \eta_R dx \\ &= \iint_{\mathbb{R}^n \times \mathbb{R}^n} v(t, x)(\nabla_x \mathcal{N}(x - y)v(t, y)) \cdot \nabla \eta_R(x) dx dy \\ &= \frac{1}{2} \iint_{\mathbb{R}^n \times \mathbb{R}^n} v(t, x)v(t, y)\nabla_x \mathcal{N}(x - y) \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)) dx dy \equiv J_1 + J_2, \end{aligned}$$

where we set

$$J_1 \equiv \frac{1}{2} \iint_{B_{\varepsilon R}(a) \times B_{\varepsilon R}(a)} v(t, x)v(t, y)\nabla_x \mathcal{N}(x - y) \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)) dx dy,$$

$$J_2 \equiv \frac{1}{2} \iint_{\mathbb{R}^n \times \mathbb{R}^n \setminus B_{\varepsilon R}(a) \times B_{\varepsilon R}(a)} v(t, x)v(t, y)\nabla_x \mathcal{N}(x - y) \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)) dx dy.$$

For J_1 , the integrand is rewritten by

$$\begin{aligned} & \nabla_x \mathcal{N}(x - y) \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)) \\ &= -\frac{1}{\omega_{n-1}} \frac{x - y}{|x - y|^{n-2}} \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)). \end{aligned}$$

We fix any $\varepsilon \in (0, \sqrt{n/(n+2)})$. By the concavity of η_R as (4.16), we have

$$\begin{aligned} J_1 &= -\frac{1}{2\omega_{n-1}} \iint_{B_{\varepsilon R}(a) \times B_{\varepsilon R}(a)} v(t, x)v(t, y) \frac{x - y}{|x - y|^{n-2}} \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)) \, dx \, dy \\ &\geq \frac{c_\varepsilon R^{-2}}{2\omega_{n-1}} \iint_{B_{\varepsilon R}(a) \times B_{\varepsilon R}(a)} \frac{v(t, x)v(t, y)}{|x - y|^{n-2}} \, dx \, dy. \end{aligned}$$

Since we know that

$$\frac{1}{|x - y|^{n-2}} \geq \left(\frac{1}{2\varepsilon R}\right)^{n-2} \quad \text{for } |x - a|, |y - a| \leq \varepsilon R,$$

it follows that

$$\begin{aligned} & \iint_{B_{\varepsilon R}(a) \times B_{\varepsilon R}(a)} \frac{v(t, x)v(t, y)}{|x - y|^{n-2}} \, dx \, dy \\ &\geq \left(\frac{1}{2\varepsilon R}\right)^{n-2} \left(\int_{B_R(a)} v(t, x) \, dx - \int_{\varepsilon R \leq |x-a| \leq R} v(t, x) \, dx \right)^2 \\ &\geq \left(\frac{1}{2\varepsilon R}\right)^{n-2} \left(\int_{B_R(a)} v(t, x) \, dx \right)^2 - 2 \left(\frac{1}{2\varepsilon R}\right)^{n-2} \\ &\quad \times \int_{B_R(a)} v(t, x) \, dx \int_{\varepsilon R \leq |x-a| \leq R} v(t, x) \, dx. \end{aligned}$$

If we set a constant as

$$C_\varepsilon \equiv \inf_{|x-a| \geq \varepsilon R} (1 - \eta_R(x)) = 1 - (1 - \varepsilon^2)^2 > 0,$$

then we have

$$\begin{aligned} - \int_{\varepsilon R \leq |x-a| \leq R} v(t, x) \, dx &\geq - \int_{\varepsilon R \leq |x-a| \leq R} v(t, x) \frac{1 - \eta_R(x)}{C_\varepsilon} \, dx \\ &\geq -\frac{1}{C_\varepsilon} \left(\int_{B_R(a)} v(t, x) \, dx - \int_{\mathbb{R}^n} v(t, x) \eta_R(x) \, dx \right). \end{aligned}$$

Thus, we obtain

$$\begin{aligned} J_1 &= \frac{1}{2} \iint_{B_{\varepsilon R}(a) \times B_{\varepsilon R}(a)} v(t, x)v(t, y) \nabla_x \mathcal{N}(x - y) \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)) \, dx \, dy \\ &\geq \frac{c_\varepsilon R^{-n}}{2(2\varepsilon)^{n-2} \omega_{n-1}} \int_{B_R(a)} v(t, x) \, dx \left[\int_{B_R(a)} v(t, x) \, dx \right. \\ &\quad \left. - \frac{2}{C_\varepsilon} \left(\int_{B_R(a)} v(t, x) \, dx - \int_{\mathbb{R}^n} v(t, x) \eta_R(x) \, dx \right) \right]. \end{aligned}$$

On the other hand, we recall that the integration over other regions is defined as

$$J_2 \equiv \frac{1}{2} \iint_{\mathbb{R}^n \times \mathbb{R}^n \setminus B_{\varepsilon R}(a) \times B_{\varepsilon R}(a)} v(t, x)v(t, y) \nabla_x \mathcal{N}(x - y) \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)) \, dx \, dy.$$

In order to estimate J_2 , it suffices to consider the integral over

$$\{(x, y) \in \mathbb{R}^n \times \mathbb{R}^n; |x - a| < R, |y - a| \geq \varepsilon R\}$$

by the symmetry property of J_2 and the fact that $\nabla \eta_R(x)$ vanishes on the set $\{x \in \mathbb{R}^n; |x - a| > R\}$. For $|x - a| < R$ and $|y - a| \geq \varepsilon R$, we have

$$\nabla_x \mathcal{N}(x - y) \cdot (\nabla_x \eta_R(x) - \nabla_y \eta_R(y)) \leq \frac{2R^{-2}}{\omega_{n-1}} \frac{1}{|x - y|^{n-2}}.$$

Thus, we have

$$|J_2| \leq \frac{2R^{-2}}{\omega_{n-1}} \iint_{\{|x-a|<R\} \times \{|y-a|\geq \varepsilon R\}} \frac{v(t, x)v(t, y)}{|x - y|^{n-2}} dx dy.$$

By the Hardy–Littlewood–Sobolev inequality and $\|v(t_k)\|_{\frac{n}{2}} = 1$, we have

$$J_2 \geq -\frac{2(n-2)C_{\text{HLS}}}{\omega_{n-1}} R^{-2} \int_{B_R(a)} v(t_k, x) dx$$

at $t = t_k$. Hence, we have

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^n} v(t) \eta_R dx \Big|_{t=t_k} \\ & \geq -4nR^{-2} \lambda_k^2 \int_{B_R(a)} v(t_k) dx + J_1 + J_2 \\ & \geq -4nR^{-2} \lambda_k^2 \int_{B_R(a)} v(t_k) dx - \frac{2(n-2)C_{\text{HLS}}}{\omega_{n-1}} R^{-2} \lambda_k^n \int_{B_R(a)} v(t_k) dx \\ & \quad + \frac{c_\varepsilon R^{-n}}{2(2\varepsilon)^{n-2} \omega_{n-1}} \lambda_k^n \int_{B_R(a)} v(t_k) dx \left[\int_{B_R(a)} v(t_k) dx \right. \\ & \quad \left. - \frac{2}{C_\varepsilon} \left(\int_{B_R(a)} v(t_k) dx - \int_{\mathbb{R}^n} v(t_k) \eta_R dx \right) \right] \\ & = R^{-2} \lambda_k^n \int_{B_R(a)} v(t_k) dx \left[-4n \lambda_k^{-(n-2)} + \frac{c_\varepsilon R^{-(n-2)}}{2(2\varepsilon)^{n-2} \omega_{n-1}} \int_{B_R(a)} v(t_k) dx \right. \\ & \quad \left. - \frac{2(n-2)C_{\text{HLS}}}{\omega_{n-1}} - \frac{c_\varepsilon R^{-n}}{C_\varepsilon (2\varepsilon)^{n-2} \omega_{n-1}} \left(\int_{B_R(a)} v(t_k) dx - \int_{\mathbb{R}^n} v(t_k) \eta_R dx \right) \right]. \end{aligned}$$

There exists $k_0 \in \mathbb{N}$ and $\varepsilon_0 > 0$ such that for any $k \geq k_0$,

$$-4n \lambda_k^{-(n-2)} + \frac{c_{\varepsilon_0} R^{-(n-2)}}{2(2\varepsilon_0)^{n-2} \omega_{n-1}} \int_{B_R(a)} v(t_k) dx - \frac{2(n-2)C_{\text{HLS}}}{\omega_{n-1}} > C_0$$

for some constant $C_0 > 0$. Thus, there exists $R_0 = R(\varepsilon_0) > 0$ such that for any $k \geq k_0$,

$$C_0 - \frac{c_{\varepsilon_0} R_0^{-n}}{C_{\varepsilon_0} (2\varepsilon_0)^{n-2} \omega_{n-1}} \left(\int_{B_{R_0}(a)} v(t_k) dx - \int_{\mathbb{R}^n} v(t_k) \eta_{R_0}(x) dx \right) > 0$$

since $\|v(t_k)\|_1 = 1$. Therefore, it follows that for any $k \geq k_0$,

$$\int_{\mathbb{R}^n} v(t_k) \eta_{R_0} dx \geq C$$

for some constant $C > 0$, which implies that $\{v(t_k)\}_{k \in \mathbb{N}}$ is not vanishing sequence. □

The error estimate (4.15) gives the following decomposition:

Proposition 4.3 (Profile decomposition) *There exist an integer $J \in \mathbb{N}$ and a non-negative function sequence $\{U^{(j)}\}_{j=1}^J \subset L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$ with $\text{supp } U^{(j)} \subset B_{R^{(j)}}(x_k^{(j)})$ for some sequences $\{x_k^{(j)}\}_{k \in \mathbb{N}, j=1,2,\dots,J} \subset \mathbb{R}^n$ and $\{R^{(j)}\}_{j=1}^J \subset \mathbb{R}_+$ such that for any $\varepsilon > 0$, there exists $k_* \geq 1$ such that for all $k \geq k_*$,*

$$(1 - \varepsilon) \|\nabla|^{-1}u_k\|_2^2 \leq \sum_{j=1}^J \|\nabla|^{-1}U^{(j)}\|_2^2, \tag{4.18}$$

where $\text{supp } U^{(j)}$ are disjoint and $|x_k^{(j)} - x_k^{(j')}| \rightarrow \infty$ as $k \rightarrow \infty$ if $j \neq j'$.

Proof of Theorem 1.4(1) On the right hand side of the inequality (4.1), putting $t = t_k$ and scaling with respect to λ_k , then the right hand side is rewritten by

$$\begin{aligned} \|\nabla|^{-1}u(t_k)\|_2^2 &= \int_{\mathbb{R}^n} u(t_k, x)(-\Delta)^{-1}u(t_k, x) dx \\ &= \lambda_k^{2n} \int_{\mathbb{R}^n} u_k(\lambda_k x)(-\Delta)^{-1}u_k(\lambda_k x) dx = \lambda_k^{n-2} \|\nabla|^{-1}u_k\|_2^2. \end{aligned}$$

By the auxiliary inequality (4.1), it holds that for all $k = 1, 2, \dots$,

$$\frac{2n}{n-2} \|u_0\|_1 \leq \lambda_k^{n-2} \|\nabla|^{-1}u_k\|_2^2. \tag{4.19}$$

By the above inequality (4.19) and the profile decomposition (4.18), for any $\varepsilon > 0$, there exists $k_* \in \mathbb{N}$ such that for any $k \geq k_*$, we see that

$$\frac{2n}{n-2} \|u_0\|_1 \leq \frac{1}{1-\varepsilon} \lambda_k^{n-2} \sum_{j=1}^J \|\nabla|^{-1}U_k^{(j)}\|_2^2.$$

By the Hardy–Littlewood–Sobolev inequality (2.2) and (4.13),

$$\begin{aligned} \frac{2n}{n-2} \|u_0\|_1 &\leq \frac{C_{\text{HLS}}}{1-\varepsilon} \lambda_k^{n-2} \sum_{j=1}^J \|U_k^{(j)}\|_1 \|U_k^{(j)}\|_{\frac{n}{2}} \\ &\leq \frac{C_{\text{HLS}}}{1-\varepsilon} \lambda_k^{n-2} \max_{j=1,2,\dots,J} \|U_k^{(j)}\|_{\frac{n}{2}} \sum_{j=1}^J \|U_k^{(j)}\|_1 \\ &\leq \frac{1+\varepsilon}{1-\varepsilon} C_{\text{HLS}} \lambda_k^{n-2} \|u_0\|_1 \max_{j=1,2,\dots,J} \|U_k^{(j)}\|_{\frac{n}{2}}. \end{aligned}$$

Thus, there exists $j_0 \in \{1, 2, \dots, J\}$ such that

$$\left(\frac{2n}{(n-2)C_{\text{HLS}}}\right)^{\frac{n}{2}} \leq \left(\frac{(1+\varepsilon)\lambda_k^{n-2}}{1-\varepsilon}\right)^{\frac{n}{2}} \int_{\mathbb{R}^n} U^{(j_0)}(y)^{\frac{n}{2}} dy. \tag{4.20}$$

Moreover, by (4.14), there exist $x_k^{(j_0)} \in \mathbb{R}^n$ and $R_0 > 0$ such that

$$\int_{\mathbb{R}^n} U^{(j_0)}(y)^{\frac{n}{2}} dy \leq (1+\varepsilon)^{\frac{n}{2}} \int_{|y-x_k^{(j_0)}| < R_0} u_k(y)^{\frac{n}{2}} dy$$

$$= (1 + \varepsilon)^{\frac{n}{2}} \int_{|y-x_k^{(j_0)}| < R_0} \lambda_k^{-\frac{n}{2}} u(t_k, \lambda_k^{-1} y)^{\frac{n}{2}} dy.$$

Substituting this to the inequality (4.20), then by scaling, we conclude that

$$\left(\frac{2n}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}} \leq \left(\frac{(1+\varepsilon)^2}{1-\varepsilon} \right)^{\frac{n}{2}} \int_{|x-\lambda_k^{-1}x_k^{(j_0)}| < \lambda_k^{-1}R_0} u(t_k, x)^{\frac{n}{2}} dx.$$

By passing a subsequence if necessary, there exists a point $x_* \in \mathbb{R}^n$ such that $y_k \equiv \lambda_k^{-1}x_k^{(j_0)} \rightarrow x_*$. Indeed, if we assume that $\{\lambda_k^{-1}x_k^{(j_0)}\}_{k \in \mathbb{N}}$ is an unbounded sequence, then $\lambda_k^{-1}|x_k^{(j_0)}| \rightarrow \infty$ ($k \rightarrow \infty$). For $k \in \mathbb{N}$ sufficiently large, we see that

$$M \equiv \int_{B_{R_0}(x_k^{(j_0)})} U^{(j_0)}(y) dy \leq \int_{B_{\lambda_k^{-1}R_0}(\lambda_k^{-1}x_k^{(j_0)})} u(t_k, x) dx.$$

Then, we have

$$\begin{aligned} & \int_{B_{\lambda_k^{-1}R_0}(y_k)} |x-x_0|^2 u(t_k, x) dx \\ & \geq 2 \int_{B_{\lambda_k^{-1}R_0}(y_k)} |y_k-x_0|^2 u(t_k, x) dx - 2 \int_{B_{\lambda_k^{-1}R_0}(y_k)} |x-y_k|^2 u(t_k, x) dx \\ & \geq 2M|y_k-x_0|^2 - 2\|u_0\|_1 \end{aligned}$$

for any $k \geq k_0$. While from the assumption of the initial condition, there exists a constant $C_0 > 0$ such that for any $k \in \mathbb{N}$,

$$\int_{\mathbb{R}^n} |x-\bar{x}|^2 u(t_k, x) dx \leq C_0. \tag{4.21}$$

Since $\{\lambda_k^{-1}x_k^{(j_0)}\}_{k \in \mathbb{N}}$ is not bounded, there exists $k_1 \in \mathbb{N}$ such that for any $k \geq k_1$,

$$2M|y_k-x_0|^2 - 2\|u_0\|_1 > C_0,$$

which contradicts (4.21). Therefore, there exists a point $x_* \in \mathbb{R}^n$ such that $\lambda_k^{-1}x_k^{(j_0)} \rightarrow x_*$, and hence, we obtain (1.10). \square

5 Proof of Theorem 1.4 (2)

In what follows, we show Theorem 1.4 (2). We assume that the initial data $u_0 \in L^{\frac{n}{2}}(\mathbb{R}^n) \cap L^1_b(\mathbb{R}^n)$ satisfies the blow-up condition (1.6) and the solution $u(t)$ to (1.1) blows up in a finite time. Namely, for some $T_* < \infty$,

$$\limsup_{t \rightarrow T_*} \|u(t)\|_{\frac{n}{2}} = \infty.$$

Let $\{t_k\}_{k \in \mathbb{N}}$ be a sequence that gives the supremum of $\|u(t)\|_{\frac{n}{2}}$. We introduce the backward self-similar transform

$$\tilde{u}(s, y) \equiv (T_* - t)u(t, \sqrt{T_* - t}y) \quad \text{with } s = -\log\left(1 - \frac{t}{T_*}\right).$$

Then \tilde{u} satisfies

$$\begin{cases} \partial_s \tilde{u} - \Delta \tilde{u} + \frac{1}{2} \nabla \cdot (y \tilde{u}) - \frac{n-2}{2} \tilde{u} + \nabla \cdot (\tilde{u} \nabla \tilde{\psi}) = 0, & s > 0, y \in \mathbb{R}^n, \\ -\Delta \tilde{\psi} = \tilde{u}, & s > 0, y \in \mathbb{R}^n, \\ \tilde{u}(0, y) = T_* u_0(\sqrt{T_*} y), & y \in \mathbb{R}^n. \end{cases}$$

We note that

$$\|\tilde{u}(s)\|_1 = e^{-\frac{n-2}{2}s} M \quad \text{with } M \equiv \|\tilde{u}(0)\|_1 = T_*^{-\frac{n-2}{2}} \|u_0\|_1.$$

We set $\{s_k\}_{k \in \mathbb{N}}$ as

$$s_k \equiv -\log \left(1 - \frac{t_k}{T_*} \right).$$

We consider the scaling transform by $\lambda > 0$ that

$$S_\lambda u(t, x) \equiv \lambda^{-n} u(t, \lambda^{-1} x).$$

This is the L^1 -invariant scaling $\|S_\lambda u(t)\|_1 = \|u(t)\|_1 = \|u_0\|_1$ and it also holds that

$$\|S_\lambda u(t)\|_{\frac{n}{2}} = \lambda^{2-n} \|u(t)\|_{\frac{n}{2}}.$$

For a blow-up solution $u(t)$ and blow-up time sequence $\{t_k\}$, we set

$$\lambda_k \equiv \|u(t_k)\|_{\frac{n}{2}}^{\frac{1}{n-2}} = \|\tilde{u}(s_k)\|_{\frac{n}{2}}^{\frac{1}{n-2}} \quad \text{and} \quad \lambda_0 \equiv \|u_0\|_{\frac{n}{2}}^{\frac{1}{n-2}}. \tag{5.1}$$

Then we find that the rescale solution

$$v(s, x) \equiv \lambda_k^{-n} \tilde{u}(s, \lambda_k^{-1} x) \quad \text{for } s_{k-1} < s \leq s_k, x \in \mathbb{R}^n$$

is bounded in $L^{\frac{n}{2}}(\mathbb{R}^n)$ and $\{e^{-\frac{n-2}{2}s_k} v(s_k)\}_{k \in \mathbb{N}}$ is bounded in $L^1(\mathbb{R}^n)$. The rescaled solution $v(t, x)$ now solves the Cauchy problem of a semi-stationary equation

$$\begin{cases} \partial_s v - \lambda_k^2 \Delta v + \frac{\lambda_k^2}{2} \nabla \cdot (y v) - \frac{n-2}{2} \lambda_k^2 v + \lambda_k^n \nabla \cdot (v \nabla \phi) = 0, & s_{k-1} < s \leq s_k, y \in \mathbb{R}^n, \\ -\Delta \phi = v, & t > 0, x \in \mathbb{R}^n, \\ v(0, y) = v_0(y) \equiv \lambda_0^{-n} u_0(\lambda^{-1} y), & y \in \mathbb{R}^n, \end{cases} \tag{5.2}$$

where $s_0 = 0$. The solution preserves its total mass $\|v(s)\|_1 = \|\tilde{u}(s)\|_1$ for any time $s \geq 0$ and $\|v(s_k)\|_{\frac{n}{2}} = 1$.

We set the rescaled solution sequence

$$v_k(y) \equiv v(s_k) = S_{\lambda_k} \tilde{u}(s_k, y) = \lambda_k^{-n} \tilde{u}(s_k, \lambda_k^{-1} y) \tag{5.3}$$

satisfies

$$e^{-\frac{n-2}{2}s_k} \|v_k\|_1 = M \quad \text{and} \quad \|v_k\|_{\frac{n}{2}} = 1. \tag{5.4}$$

In this case, one can apply Proposition 3.1 with $\{v_k\}_{k \in \mathbb{N}}$:

Proposition 5.1 (Profile decomposition in L^p) *Let $\{v_k\}_{k \in \mathbb{N}}$ be a non-negative sequence in $L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$ with (5.4) defined by the above. Then for all $\varepsilon > 0$, there exists a subsequence of $\{v_k\}$ (not relabeled), $J \in \mathbb{N}$, $\{y_k^{(j)}\}_{k \in \mathbb{N}, j=1, \dots, J} \subset \mathbb{R}^n$, $\{R^{(j)}\}_{j=1}^J \subset \mathbb{R}_+$, and nonnegative function sequences $\{V_k^{(j)}\}_{k \in \mathbb{N}, j=1, \dots, J}$, $\{w_k\}_{k \in \mathbb{N}}$, $\{e_k\}_{k \in \mathbb{N}} \subset L^1(\mathbb{R}^n) \cap L^{\frac{n}{2}}(\mathbb{R}^n)$ which satisfy the following properties:*

(1) v_k is decomposed as

$$v_k(x) = \sum_{j=1}^J V_k^{(j)}(x - y_k^{(j)}) + w_k(x) + e_k(x) \quad \text{a.a. } x \in \mathbb{R}^n, \quad (5.5)$$

where $\{V_k^{(j)}\}_{k \in \mathbb{N}}^{j=1, \dots, J}$ is a nonnegative function sequence with $\text{supp } V_k^{(j)} \subset B_{R^{(j)}}(0)$. Moreover, for each $k \in \mathbb{N}$, $B_{R^{(j)}}(y_k^{(j)})$ and $\text{supp } w_k$ are all disjoint for any $j = 1, 2, \dots, J$. For any $R > 0$, it holds that

$$\lim_{k \rightarrow \infty} \sup_{x \in \mathbb{R}^n} \int_{B_R(x)} w_k(x)^p dx = 0 \quad \text{for any } 1 \leq p \leq \frac{n}{2} \quad (5.6)$$

and the error term satisfies

$$\overline{\lim}_{k \rightarrow \infty} \|e_k\|_{\frac{n}{2}} < \varepsilon \quad (5.7)$$

(2) Almost orthogonality:

$$\|v_k\|_1 = \sum_{j=1}^J \|V_k^{(j)}\|_1 + \|w_k\|_1 + \|e_k\|_1, \quad (5.8)$$

and for any $j \in \{1, 2, \dots, J\}$, it holds that

$$\|V_k^{(j)}\|_{\frac{n}{2}} = \|v_k\|_{L^{\frac{n}{2}}(B_{R^{(j)}}(y_k^{(j)}))}. \quad (5.9)$$

(3) Drift term estimate:

$$\overline{\lim}_{k \rightarrow \infty} e^{-\frac{n-2}{2}s_k} \|\ |\nabla|^{-1} w_k \|_2^2 = \overline{\lim}_{k \rightarrow \infty} e^{-\frac{n-2}{2}s_k} \|\ |\nabla|^{-1} e_k \|_2^2 = 0. \quad (5.10)$$

In addition, if $\{v_k\}_{k \in \mathbb{N}}$ is radially symmetric, then $J = 1$ and $\{y_k^{(1)}\}_{k \in \mathbb{N}}$ is bounded.

For the proof of Proposition 5.1, see Appendix.

We set

$$u_k(x) \equiv \lambda_k^{-1} u(t_k, \lambda_k^{-1} x)$$

and

$$\begin{cases} \tilde{U}_k^{(j)}(x) \equiv (T_* - t_k)^{-1} V_k^{(j)}\left(\frac{x}{\sqrt{T_* - t_k}}\right), \\ \tilde{w}_k(x) \equiv (T_* - t_k)^{-1} w_k\left(\frac{x}{\sqrt{T_* - t_k}}\right), \\ \tilde{e}_k(x) \equiv (T_* - t_k)^{-1} e_k\left(\frac{x}{\sqrt{T_* - t_k}}\right). \end{cases}$$

Then we see that

$$u_k(x) = \sum_{j=1}^J \tilde{U}_k^{(j)}(x - y_k^{(j)}) + \tilde{w}_k(x) + \tilde{e}_k(x) \quad \text{a.a. } x \in \mathbb{R}^n.$$

The estimate (5.8) implies that

$$\sum_{j=1}^J \|\tilde{U}_k^{(j)}\|_1 \leq (1 + \varepsilon) \|u_k\|_1 = (1 + \varepsilon) \|u_0\|_1.$$

By (5.10) and changing variables, we have

$$\overline{\lim}_{k \rightarrow \infty} \left\| |\nabla|^{-1} \tilde{u}_k \right\|_2^2 = \overline{\lim}_{k \rightarrow \infty} \left\| |\nabla|^{-1} \tilde{e}_k \right\|_2^2 = 0,$$

in particular, there exists $k_* \in \mathbb{N}$ such that for all $k \geq k_*$,

$$\left\| |\nabla|^{-1} u_k \right\|_2^2 \leq \frac{1}{1 - \varepsilon} \sum_{j=1}^J \left\| |\nabla|^{-1} \tilde{U}_k^{(j)} \right\|_2^2.$$

Proof of Theorem 1.4 (2) By the similar argument of the proof of Theorem 1.4 (1), for any $\varepsilon > 0$, there exists $k_* \in \mathbb{N}$ such that for any $k \geq k_*$, we see that

$$\frac{2n}{n-2} \|u_0\|_1 \leq \lambda_k^{n-2} \left\| |\nabla|^{-1} u_k \right\|_2^2 \leq \frac{1 + \varepsilon}{1 - \varepsilon} C_{\text{HLS}} \lambda_k^{n-2} \|u_0\|_1 \max_{j=1,2,\dots,J} \|V_k^{(j)}\|_{\frac{n}{2}}.$$

Thus, there exists $j_0 \in \{1, 2, \dots, J\}$ such that

$$\left(\frac{2n}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}} \leq \left(\frac{1 + \varepsilon}{1 - \varepsilon} \right)^{\frac{n}{2}} \lambda_k^{\frac{n(n-2)}{2}} \int_{\mathbb{R}^n} V_k^{(j_0)}(y)^{\frac{n}{2}} dy. \tag{5.11}$$

Moreover, by (5.9), there exist $x_k^{(j_0)} \in \mathbb{R}^n$ and $R_0 > 0$ such that

$$\int_{\mathbb{R}^n} V_k^{(j_0)}(y)^{\frac{n}{2}} dy = \int_{|y-y_k^{(j_0)}| < R_0} \tilde{u}_k(y)^{\frac{n}{2}} dy = \int_{|y-y_k^{(j_0)}| < R_0} \lambda_k^{-\frac{n^2}{2}} \tilde{u}(t_k, \lambda_k^{-1}y)^{\frac{n}{2}} dy.$$

Substituting this to the inequality (5.11), then by scaling, we conclude that

$$\begin{aligned} \left(\frac{2n}{(n-2)C_{\text{HLS}}} \right)^{\frac{n}{2}} &\leq \left(\frac{1 + \varepsilon}{1 - \varepsilon} \right)^{\frac{n}{2}} \lambda_k^{\frac{n(n-2)}{2}} \lambda_k^{-\frac{n^2}{2}} \int_{|y-y_k^{(j_0)}| < R_0} \tilde{u}(s_k, \lambda_k^{-1}y)^{\frac{n}{2}} dy \\ &\leq \left(\frac{1 + \varepsilon}{1 - \varepsilon} \right)^{\frac{n}{2}} \int_{|x-\lambda_k^{-1}y_k^{(j_0)}| < \lambda_k^{-1}R_0} \tilde{u}(s_k, x)^{\frac{n}{2}} dx. \end{aligned}$$

Therefore, if we set $x_k \equiv \lambda_k^{-1} \sqrt{T - t_k} y_k^{(j_0)}$, then we obtain (1.12). □

6 Radially symmetric case

In this section, we consider the case that $u_0 \in L^{\frac{n}{2}}(\mathbb{R}^n) \cap L^1_b(\mathbb{R}^n)$ is radially symmetric and nonnegative for $b > 0$. We assume that the initial data u_0 satisfies the assumption (1.6) with (1.9) for $\delta > 0$. Let $u(t, x)$ be a blowing up solution to (1.1). We set the scaling parameter (5.1) and consider the rescaled solution (5.3).

Since the assumption on the data is different from the case of $b \geq 2$, the lower estimate has to be arranged.

Lemma 6.1 *Let the initial data u_0 be radially symmetric and satisfy (1.6) with (1.9) for $\delta > 0$, then there exists the corresponding radially symmetric solution $u(t)$ to (1.1) blows up in a finite time T . Then it holds that for any $0 < t < T$,*

$$\frac{2(n + \delta)}{n - 2} \|u_0\|_1 \leq \left\| |\nabla|^{-1} u(t) \right\|_2^2. \tag{6.1}$$

Proof of Theorem 1.5 By applying the similar argument in the proof of Theorem 1.4 (2), we obtain (1.14). By Proposition 5.1, we note that the center $\{y_k^{(1)}\}_{k \in \mathbb{N}}$ of the profile $V^{(1)}$ is bounded. Thus, we have

$$\lambda_k^{-1} \sqrt{T - t_k} y_k^{(1)} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

After admitting Lemma 6.1, we conclude the analogous result to (1.12): For any $\varepsilon > 0$,

$$\left(\frac{2(n + \delta)}{(n - 2)C_{\text{HLS}}} \right)^{\frac{n}{2}} \leq \liminf_{k \rightarrow \infty} \int_{B_{\varepsilon \sqrt{T - t_k}}(y_k^{(1)})} u(t_k, x)^{\frac{n}{2}} dx,$$

which implies that (1.15).

To see Lemma 6.1, we need to introduce the modified moment and derive its dynamics from the *modified virial law*. Let $\phi(x)$ be a radially symmetric smooth function such that

$$\phi(x) = \begin{cases} |x|^2, & 0 \leq |x| < 1, \\ \text{smooth}, & 1 \leq |x| < 2, \\ |x|^b, & 2 \leq |x| \end{cases}$$

and set $r = |x|$. We define

$$\Phi_R(r) = R^2 \phi\left(\frac{r}{R}\right)$$

for any $R \geq 1$.

Proposition 6.2 (Modified virial law) For $\frac{n}{2} \leq p < n$, $b > 0$ let $u \in C([0, T]; L_b^1(\mathbb{R}^n) \cap L^p(\mathbb{R}^n))$ be a solution to (1.1) with initial data $u_0 \in L_b^1(\mathbb{R}^n)$ with positive initial data $u_0(x) \geq 0$. Then if $n \geq 4$ or $n = 3$ and $b < 1$, then it holds for any $t \in (0, T)$ that

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^n} \Phi_R(x) u(t) dx \\ & \leq 2n \|u(s)\|_1 - (n - 2) \int_{\mathbb{R}^n} u(s) \psi(s) dx + (n - 2) \int_{B_R^c(0)} u(t) (-\Delta)^{-1} u(t) dx \quad (6.2) \\ & + \int_{B_R^c(0)} \Psi_R(r) \partial_r^2 \psi(t) \psi(t) dx + \frac{1}{4} \int_{B_R^c(0)} \Delta^2 \Phi_R |\psi(t)|^2 dx, \end{aligned}$$

where

$$\Psi_R(r) = \frac{\Phi'_R(r)}{r} - \Phi''_R(r) = \begin{cases} 0, & |x| \leq R, \\ \text{smooth}, & R < |x| \leq 2R, \\ -b(b - 2) \left(\frac{r}{R}\right)^{b-2}, & 2R < |x| \end{cases}$$

is supported in $B_R^c(0)$.

Lemma 6.3 Let $\Psi_R(x) \in L^1(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ be a radially symmetric function supported in $B_R^c(0)$ and $u(t)$ is radially symmetric.

(1) Then there exists a constant $C = C(\phi) > 0$ depending on Φ_R and η such that

$$\int_{\mathbb{R}^n} \Psi_R(r) \partial_r^2 (-\Delta)^{-1} u(t) (-\Delta)^{-1} u(t) dx \leq CR^{-(n-2)} \|u_0\|_1^2.$$

(2) For any $R > 0$, it holds that

$$\frac{1}{4} \int_{\mathbb{R}^n} \Delta^2 \Phi_R |\psi(t)|^2 dx \leq C(\phi) R^{-(n-2)} \|u_0\|_1^2.$$

Proof for Proposition 6.2 and Lemma 6.3, see [41].

Proposition 6.4 Let the initial data be radially symmetric in $L^{\frac{n}{2}}(\mathbb{R}^n) \cap L^1_b(\mathbb{R}^n)$ and satisfy the condition (1.6) with (1.9), then

$$\int_{\mathbb{R}^n} |x|^b u(t) dx \leq \int_{\mathbb{R}^n} |x|^b u_0 dx. \tag{6.3}$$

Proof of Proposition 6.4 By Lemma 6.3, choosing $R > 0$ sufficiently large such that

$$\int_{B_R^c(0)} \Psi_R(r) \partial_r^2 \psi(t) \psi(t) dx + \frac{1}{4} \int_{B_R^c(0)} \Delta^2 \Phi_R |\psi(t)|^2 dx \leq 2(n-2) C R^{-(n-2)} M^2 \leq 2\delta M. \tag{6.4}$$

By the modified virial law (6.2), (6.4), and Shannon’s inequality (2.1), we have for $M \equiv \|u_0\|_1$,

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^n} \Phi_R(x) u(t) dx \\ & \leq 2(n-2) \left[H[u(t)] + \frac{n}{n-2} M + \frac{\delta}{n-2} M - \int_{\mathbb{R}^n} u(t) \log u(t) dx \right] \\ & \leq 2(n-2) \left[H[u(0)] + \frac{n+\delta}{n-2} M + \frac{n}{b} M \log \left(\frac{bc_{n,b} e}{nM^{1+\frac{b}{n}}} \int_{\mathbb{R}^n} |x|^b u(t) dx \right) \right] \\ & = 2(n-2) \left[H[u_0] + \frac{n}{b} M \log \left(\frac{bc_{n,b} e^{1+\frac{b(n+\delta)}{n(n-2)}}}{nM^{1+\frac{b}{n}}} \int_{\mathbb{R}^n} |x|^b u(t) dx \right) \right]. \end{aligned} \tag{6.5}$$

Hence under the condition (1.6), the right hand side of (6.5) is negative if $t = 0$. Then

$$X(t) \equiv \int_{\mathbb{R}^n} \Phi_R(x) u(t) dx$$

is decreasing function of t in $[0, \eta)$. Since it holds that

$$\int_{\mathbb{R}^n} |x|^b u(t) dx \leq R^{b-2} X(t) + R^b M,$$

the b -th moment of u does not increase if the initial data satisfies

$$H[u_0] < -\frac{n}{b} M \log \left(\frac{bc_{n,b} e^{1+\frac{b(n+\delta)}{n(n-2)}}}{nM^{1+\frac{b}{n}}} \int_{\mathbb{R}^n} |x|^b u_0 dx \right),$$

where $c_{n,b}$ is the constant appearing in Proposition 2.1. Thus, we obtain the inequality (6.3).□

Proof of Lemma 6.1 From (1.3) we have

$$\int_{\mathbb{R}^n} u(t) \log u(t) dx - H[u(t)] \leq \frac{1}{2} \|\nabla^{-1} u(t)\|_2^2.$$

By the Shannon inequality (Proposition 2.1), it follows that

$$\frac{n}{b} \|u_0\|_1 \log \left(\frac{n \|u_0\|_1^{1+\frac{b}{n}}}{bc_{n,b} e \int_{\mathbb{R}^n} |x - \bar{x}|^b u(t) dx} \right) - H[u(t)] \leq \frac{1}{2} \|\nabla^{-1} u(t)\|_2^2. \quad (6.6)$$

We obtain from (6.3) and (6.6) that

$$\frac{n}{b} \|u_0\|_1 \log \left(\frac{n \|u_0\|_1^{1+\frac{b}{n}}}{bc_{n,b} e \int_{\mathbb{R}^n} |x - \bar{x}|^b u_0(x) dx} \right) - H[u_0] \leq \frac{1}{2} \|\nabla^{-1} u(t)\|_2^2.$$

From the assumption (1.6) with (1.9), we have

$$\frac{2(n + \delta)}{n - 2} \|u_0\|_1 \leq \|\nabla^{-1} u(t)\|_2^2,$$

and hence, we obtain the inequality (6.1). □

Acknowledgements The first author would like to thank Professor Taub Hmidi for his sending his result on the profile decomposition. He is also grateful to Professor Kazuhiro Kuwae for information on the b -th moment center. The work of T. Ogawa is partially supported by JSPS Grant-in-aid for Scientific Research S #19H05597. The work of T. Suguro is partially supported by JSPS Grant-in-aid for JSPS Fellows #19J20763 and Research Activity Start-up #22K20336. The work of H. Wakui is supported by JSPS Grant-in-aid for JSPS Fellows #20J00940.

Data Availability Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix A. Proof of Proposition 5.1

Proof of Proposition 5.1 We apply the similar argument in the proof of Lemma 3.1 with $\{v_k^{\frac{n}{2}}\}_{k \in \mathbb{N}} \subset L^1(\mathbb{R}^n)$. We notice that the profile given by the convergence of $\{v_k\}$ in $L^{\frac{n}{2}}$ is not always non-zero function. From this reason, we define the profile, which depends on k , as

$$V_k^{(j)}(x) \equiv v_k(x + y_k^{(j)}) \chi_{B_{R^{(j)}}(0)}(x).$$

The decomposition (5.5), (5.8), and (5.9) are shown directly in Lemma 3.1. In what follows, we show that the error estimates (5.10).

We note that $\{e^{-\frac{n-2}{2}s_k} e_k\}_{k \in \mathbb{N}}$ is uniformly bounded in $L^1(\mathbb{R}^n)$. Hence, the Hardy–Littlewood–Sobolev inequality (2.2) and (5.7) show that

$$e^{-\frac{n-2}{2}s_k} \|\nabla^{-1} e_k\|_2^2 \leq C_{\text{HLS}} e^{-\frac{n-2}{2}s_k} \|e_k\|_1 \|e_k\|_{\frac{n}{2}} \leq C \|e_k\|_{\frac{n}{2}} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Next, we shall estimate the vanishing term w_k . We fix $\varepsilon > 0$ arbitrarily. For $R > 0$, we separate the non-local term into three parts as follows:

$$\begin{aligned} \|\lvert \nabla \rvert^{-1} w_k\|_2^2 &= c_n \left(\iint_{\{|x-y|<R^{-1}\}} + \iint_{\{R^{-1}<|x-y|<R\}} + \iint_{\{|x-y|>R\}} \right) \frac{w_k(x) \overline{w_k(y)}}{|x-y|^{n-2}} dx dy \\ &\equiv I_1 + I_2 + I_3, \end{aligned}$$

where $c_n = 1/((n-2)\omega_{n-1})$. For the integral I_1 , by the Hausdorff–Young inequality, we have

$$I_1 = c_n \int_{\mathbb{R}^n} w_k(y) \int_{\{|x-y|<R^{-1}\}} \frac{w_k(x)}{|x-y|^{n-2}} dx dy \leq C \|w_k\|_{\frac{n}{2}}^2 \left\| \lvert \cdot \rvert^{-(n-2)} \chi_{\{| \cdot | < R^{-1}\}}(\cdot) \right\|_q,$$

where $\chi_A(\cdot)$ denotes the characteristic function of a set A , and q satisfies $q = \frac{n}{2(n-2)}$. By the uniform boundedness of $L^{\frac{n}{2}}$ for $\{w_k\}$, there exists $R_0 > 0$ independent of k such that for any $R > R_0$,

$$I_1 \leq C R_0^{-\frac{n}{2}} < \frac{\varepsilon}{3}.$$

For the integral I_3 , we decompose \mathbb{R}^n into countable cubes whose centers are lattice points, that is, $\mathbb{R}^n = \bigcup_{j \in \mathbb{Z}^n} Q(j)$, where $Q(j) \equiv \{y \in \mathbb{R}^n; \max_{1 \leq i \leq n} |y_i - j_i| \leq 1/2\}$. It follows from the cubic decomposition and the Hölder and weak Hausdorff–Young inequalities that

$$\begin{aligned} I_3 &= c_n \int_{\mathbb{R}^n} w_k(y) \int_{\{|x-y|>R\}} \frac{w_k(x)}{|x-y|^{n-2}} dx dy \\ &\leq C \int_{\mathbb{R}^n} w_k(y) \sum_{\substack{j, j' \in \mathbb{Z}^n, \\ |j-j'| \geq R}} (\lvert \cdot \rvert^{-(n-2)} \chi_{Q(j')}) * (w_k \chi_{Q(j)})(y) dy \\ &\leq C \|w_k\|_{\frac{2n}{n+2}} \sum_{|j-j'| \geq R} \left\| (\lvert \cdot \rvert^{-(n-2)} \chi_{Q(j')}) * (w_k \chi_{Q(j)}) \right\|_{\frac{2n}{n-2}} \\ &\leq C \|w_k\|_{\frac{2n}{n+2}} \sum_{|j-j'| \geq R} \|w_k \chi_{Q(j)}\|_{\frac{2n}{n+2}} \left\| \lvert \cdot \rvert^{-(n-2)} \chi_{Q(j')} \right\|_{\frac{n}{n-2}, w} \\ &\leq C 3^n \|w_k\|_{\frac{2n}{n+2}}^2 \left(\sum_{|j'| \geq R} \left\| \lvert \cdot \rvert^{-(n-2)} \chi_{Q(j')} \right\|_{\frac{n}{n-2}, w} \right)^{\frac{n-2}{2n}}, \end{aligned}$$

where the last estimate is derived by Hölder’s inequality and the covering. By the radially decreasing property, $|x|^{-(n-2)}$ attains the maximum in $Q(j')$ at the closest point $y_{j'}$ to the origin. For $k \in \mathbb{Z}^n$ satisfying $\max |k_i| \geq R$, we see that

$$\left\| \lvert \cdot \rvert^{-(n-2)} \chi_{Q(j')} \right\|_{\frac{n}{n-2}, w}^{\frac{2n}{n-2}} \leq \left(\int_{Q(j')} |x|^{-n} dx \right)^2 \leq C |y_{j'}|^{-2n}.$$

We draw a line from the origin to $y_{j'}$ and order cubes intersecting with this line. The first cube is $Q(0)$ and the second cube is the cube which the line meets when it goes out of $Q(0)$, and so on. We denote $I(Q(j'))$ the second-to-last cube in this order. Then we have

$$|y_{j'}|^{-2n} \leq \int_{I(Q(j', \frac{1}{2}))} |x|^{-2n} dx.$$

Thus, we see that

$$\sum_{|j'| \geq R} \left\| |\cdot|^{-(n-2)} \chi_{Q(j')} \right\|_{\frac{2n}{n-2}, w} \leq C \sum_{|j'| \geq R} \int_{Q(j')} |y|^{-2n} dy \leq C \int_R^\infty r^{-n-1} dr \leq CR^{-n}.$$

Therefore, by the uniform boundedness of L^1 for $\{e^{-\frac{n-2}{2}s_k} v_k\}$ and Hölder's inequality, there exists $R_1 > 0$ independent of k such that for any $R > R_1$,

$$e^{-\frac{n-2}{2}s_k} I_3 \leq CR_1^{-n} < \frac{\varepsilon}{3}.$$

We set $\tilde{R} \equiv \max\{R_0, R_1\}$. Lastly, the integral I_2 can be estimated as

$$I_2 = c_n \int_{\mathbb{R}^n} w_k(y) \int_{\tilde{R}^{-1} < |x-y| < \tilde{R}} \frac{w_k(x)}{|x-y|^{n-2}} dx dy \leq c_n \tilde{R}^{n-2} \|w_k\|_1 \sup_{x \in \mathbb{R}^n} \int_{B_{\tilde{R}}(x)} w_k(y) dy.$$

Since w_k is the vanishing term, (5.6) gives

$$\overline{\lim}_{k \rightarrow \infty} e^{-\frac{n-2}{2}s_k} I_2 \leq c_n \tilde{R}^{n-2} \limsup_{k \rightarrow \infty} \sup_{x \in \mathbb{R}^n} \int_{B_{\tilde{R}}(x)} w_k(y) dy = 0.$$

Thus, we conclude

$$\overline{\lim}_{k \rightarrow \infty} e^{-\frac{n-2}{2}s_k} \|\ |\nabla|^{-1} w_k \|_2 = 0.$$

By the similar argument in the proof of Proposition 4.2, we see that $\{v_k\}_{k \in \mathbb{N}}$ is also not a vanishing sequence, and hence, we have $J \geq 1$.

In the case that $\{v_k\}_{k \in \mathbb{N}}$ is radially symmetric, we see that $\{y_k^{(j)}\}_{k \in \mathbb{N}}$ is bounded for any $j = 1, 2, \dots, J$. Indeed, if not, it follows from the radially symmetrically and weak $L^{\frac{n}{2}}$ -convergence of $\{v_k\}$ that

$$\|v_k\|_{\frac{n}{2}} \geq C |y_k^{(j)}|^n \int_{B_{R(j)}(y_k^{(j)})} v_k(y)^{\frac{n}{2}} dy \geq C |y_k^{(j)}|^n \int_{B_{R(j)}(0)} V^{(j)}(y)^{\frac{n}{2}} dy$$

for $k \in \mathbb{N}$ sufficiently large. The right hand side diverges to infinity as $k \rightarrow \infty$, which contradicts with $\|v_k\|_{\frac{n}{2}} = 1$. Moreover, if we assume that $J \geq 2$, then for $j \neq j'$, we see that

$$|y_k^{(j)} - y_k^{(j')}| \rightarrow \infty \text{ as } k \rightarrow \infty.$$

For the same reason, this is a contradiction. □

References

1. Bedrossian, J., Kim, I.C.: Global existence and finite time blow-up for critical Patlak-Keller-Segel models with inhomogeneous diffusion. *SIAM J. Math. Anal.* **45**, 934–964 (2013)
2. Biler, P.: Existence and nonexistence of solutions for a model of gravitational interaction of particles, III. *Colloq. Math.* **68**, 229–239 (1995)
3. Biler, P.: Local and global solvability of some parabolic systems modeling chemotaxis. *Adv. Math. Sci. Appl.* **8**, 715–743 (1998)
4. Biler, P., Cieślak, T., Karch, G., Zienkiewicz, J.: Local criteria for blowup in two-dimensional chemotaxis models. *Discrete Contin. Dyn. Syst.* **37**, 1841–1856 (2017)
5. Biler, P., Karch, G., Zienkiewicz, J.: Large global-in-time solutions to a nonlocal model of chemotaxis. *Adv. Math.* **330**, 834–875 (2018)

6. Biler, P., Nadzieja, T.: Existence and nonexistence of solutions for a model of gravitational interactions of particles I. *Colloq. Math.* **66**, 319–334 (1994)
7. Biler, P., Nadzieja, T.: A nonlocal singular parabolic problem modeling gravitational interaction of particles. *Adv. Diff. Equ.* **3**, 177–197 (1998)
8. Biler, P., Nadzieja, T., Stanczy, R.: Nonisothermal systems of self-attracting Fermi-Dirac particles. *Banach Center Publ.* **66**, 61–78 (2004)
9. Bournaveas, N., Calvez, V.: The one-dimensional Keller-Segel model with fractional diffusion of cells. *Nonlinearity* **23**, 923–935 (2010)
10. Blanchet, A., Carrillo, J.A., Laurençot, P.: Critical mass for a Patlak-Keller-Segel model with degenerate diffusion in higher dimensions. *Calc. Var. P.D.E.* **35**, 133–168 (2009)
11. Calvez, V., Corrias, L., Ebde, M.: Blow-up, concentration phenomenon and global existence for the Keller-Segel model in high dimension. *Comm. Partial Differ. Equ.* **37**, 561–584 (2012)
12. Chen, L., Liu, J.G., Wang, J.: Multidimensional degenerate Keller-Segel system with critical diffusion exponent $2n/(n+2)^*$. *SIAM J. Math. Anal.* **44**(2), 1077–1102 (2012)
13. Corrias, L., Perthame, B., Zaag, H.: Global solutions of some chemotaxis and angiogenesis system in high space dimensions. *Milan J. Math.* **72**, 1–28 (2004)
14. Feireisl, E., Laurençot, P.: Non-isothermal Smoluchowski-Poisson equations as a singular limit of the Navier-Stokes-Fourier-Poisson system. *J. Math. Pures Appl.* **88**, 325–349 (2007)
15. Gajewski, H., Gröger, K.: On the basic equations for carrier transport in semiconductors. *J. Math. Anal. Appl.* **113**, 12–35 (1986)
16. Gérard, P.: Description du défaut de compacité de l’injection de Sobolev. *ESAIM Control Optim. Calc. Var.* **3**, 213–233 (1998)
17. Giga, Y.: Solutions for semilinear Parabolic equations in L^p and regularity of weak solutions of the Navier-Stokes system. *J. Differ. Equ.* **61**, 186–212 (1986)
18. Herrero, M.A., Velázquez, J.J.L.: Singularity patterns in a chemotaxis model. *Math. Ann.* **306**, 583–623 (1996)
19. Hmidi, T., Keraani, S.: Remarks on the blowup for the L^2 critical nonlinear Schrödinger equations. *SIAM J. Math. Anal.* **38**, 1035–1047 (2006)
20. Iwabuchi, T.: Global well-posedness for Keller-Segel system in Besov type spaces. *J. Math. Anal. Appl.* **379**, 930–948 (2011)
21. Jüngel, A.: Qualitative behavior of solutions of a degenerate nonlinear drift-diffusion model for semiconductors. *Math. Model. Meth. Appl. Sci.* **5**, 497–518 (1995)
22. Keller, E.F., Segel, L.A.: Initiation of slime mold aggregation viewed as an instability. *J. Theor. Biol.* **26**, 399–415 (1970)
23. Kimijima, A., Nakagawa, K., Ogawa, T.: Threshold of global behavior of solutions to a degenerate drift-diffusion system in between two critical exponents. *Calc. Var. Partial Differ. Equ.* **53**, 441–472 (2015)
24. Kobayashi, T., Ogawa, T.: Fluid mechanical approximation to the degenerated drift-diffusion and chemotaxis equations in barotropic model. *Indiana Univ. Math. J.* **62**(3), 1021–1054 (2013)
25. Kozono, H., Sugiyama, Y., Yahagi, Y.: Existence and uniqueness theorem on weak solutions to the parabolic-elliptic Keller-Segel system. *J. Differ. Equ.* **253**, 2295–2313 (2012)
26. Kurokiba, M., Nagai, T., Ogawa, T.: The uniform boundedness of the radial solution for drift-diffusion system. *Comm. Pure Appl. Anal.* **5**, 97–106 (2006)
27. Kurokiba, M., Ogawa, T.: Finite time blow-up of the solution for a nonlinear parabolic equation of drift-diffusion type. *Differ. Integral Equ.* **16**, 427–452 (2003)
28. Kurokiba, M., Ogawa, T.: L^p well-posedness of the for the drift-diffusion system arising from the semiconductor device simulation. *J. Math. Anal. Appl.* **342**, 1052–1067 (2008)
29. Kurokiba, M., Ogawa, T.: Two dimensional drift-diffusion system in a critical weighted space. *Differ. Integral Equ.* **28**(7–8), 753–776 (2015)
30. Lions, P.-L.: The concentration-compactness principle in the calculus of variations. The locally compact case I. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1**, 109–145 (1984)
31. Merle, F., Tsutsumi, Y.: L^2 concentration of blowup solutions for the nonlinear Schrödinger equation with critical power nonlinearity. *J. Differ. Equ.* **84**, 1035–1047 (1990)
32. Mock, M.S.: An initial value problem from semiconductor device theory, *SIAM. J. Math.* **5**, 597–612 (1974)
33. Nagai, T.: Blow-up of radially symmetric solutions to a chemotaxis system. *Adv. Math. Sci. Appl.* **5**, 581–601 (1995)
34. Nagai, T.: Blowup of non-radial solutions to parabolic-elliptic systems modeling chemotaxis in two-dimensional domains. *J. Inequal. Appl.* **6**, 37–55 (2001)
35. Nagai, T., Ogawa, T.: Brezis-Merle inequalities and application to the global existence of the Keller-Segel equations. *Comm. Contemporary Math.* **13**(5), 795–812 (2011)

36. Nagai, T., Ogawa, T.: Global existence of solutions to a parabolic-elliptic system of drift-diffusion type in \mathbb{R}^2 , *Funkcial. Ekvac.* **59**, No. 2 (2016),
37. Nagai, T., Senba, T., Suzuki, T.: Chemotactic collapse in a parabolic system of mathematical biology. *Hiroshima J. Math.* **30**, 463–497 (2000)
38. Nagai, T., Senba, T., Yoshida, K.: Application of the Trudinger-Moser inequality to a parabolic system of chemotaxis. *Funkcial. Ekvac.* **40**(3), 411–433 (1997)
39. Nawa, H.: Asymptotic and limiting profiles of blowup solutions of the nonlinear Schrödinger equation with critical power. *Comm. Pure Appl. Math.* **52**, 193–270 (1999)
40. Ogawa, T.: The degenerate drift-diffusion system with the Sobolev critical exponent. *Disc. Contin. Dyn. Syst. Ser S* **4**, 875–886 (2011)
41. Ogawa, T., Wakui, H.: Non-uniform bound and finite time blow up for solutions to a drift-diffusion equation in higher dimensions. *Anal. Appl.* **14**, 145–183 (2016)
42. Ogawa, T., Wakui, H.: Finite time blow up and non-uniform bound for solutions to a degenerate drift-diffusion equation with the mass critical exponent under non-weight condition. *Manuscr. Math.* **159**, 475–509 (2019)
43. Senba, T.: Blowup behavior of radial solutions to Jäger-Luckhaus system in high dimensional domain. *Funkcialaj Ekvac.* **48**, 247–271 (2005)
44. Senba, T.: Blowup in infinite time of radial solutions to parabolic-elliptic system in high-dimensional Euclidean spaces. *Nonlinear Anal.* **70**, 2549–2562 (2009)
45. Sugiyama, Y.: Application of the best constant of the Sobolev inequality to degenerate Keller-Segel models. *Adv. Differ. Equ.* **12**, 121–144 (2007)
46. Suzuki, T., Takahashi, R.: Degenerate parabolic equations with critical exponent derived from the kinetic theory II, Blowup threshold. *Differ. Integral Equ* **22**, 1153–1172 (2009)
47. Talenti, G.: Best constant in Sobolev inequality. *Ann. Mat. Pura Appl.* **110**, 353–372 (1976)
48. Tsutsumi, Y.: Rate of L^2 concentration of blow-up solutions for the nonlinear Schrödinger equation with critical power. *Nonlinear Anal.* **15**, 719–724 (1990)
49. Wakui, H.: The rate of concentration for the radially symmetric solution to a degenerate drift-diffusion equation with the mass critical exponent. *Arch. Math. (Basel)* **111**, 535–548 (2018)
50. Weissler, F.: Existence and non-existence of global solutions for a semilinear heat equation. *Israel J. Math.* **38**, 29–40 (1981)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.