

Adherence to a Multidisciplinary Lifestyle Program for Patients With Atrial Fibrillation and Obesity

Citation for published version (APA):

Tenbult, N., Kraal, J., Brouwers, R., Spee, R., Eijssbouts, S., & Kemps, H. (2022). Adherence to a Multidisciplinary Lifestyle Program for Patients With Atrial Fibrillation and Obesity: Feasibility Study. *JMIR Formative Research*, 6(4), Article e32625. <https://doi.org/10.2196/32625>

Document license:

CC BY

DOI:

[10.2196/32625](https://doi.org/10.2196/32625)

Document status and date:

Published: 01/04/2022

Document Version:

Publisher's PDF, also known as Version of Record (includes final page, issue and volume numbers)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.tue.nl/taverne

Take down policy

If you believe that this document breaches copyright please contact us at:

openaccess@tue.nl

providing details and we will investigate your claim.

Original Paper

Adherence to a Multidisciplinary Lifestyle Program for Patients With Atrial Fibrillation and Obesity: Feasibility Study

Nicole Tenbult¹, MANP; Jos Kraal^{1,2}, PhD; Rutger Brouwers¹, MD, PhD; Ruud Spee^{1,3}, MD, PhD; Sabine Eijssbouts³, MD, PhD; Hareld Kemps^{1,3,4}, MD, PhD

¹Telemedicine and Rehabilitation in Chronic Disease, Flow, Center for Prevention, Máxima MC, Veldhoven/Eindhoven, Netherlands

²Faculty Industrial Design Engineering, Delft, Netherlands

³Department of Cardiology, Máxima MC, Veldhoven/Eindhoven, Netherlands

⁴Department of Industrial Design, University of Technology Eindhoven, Eindhoven, Netherlands

Corresponding Author:

Nicole Tenbult, MANP

Telemedicine and Rehabilitation in Chronic Disease

Flow, Center for Prevention

Máxima MC

Dominee Theodor Fliednerstraat 1

Veldhoven/Eindhoven, 5631 BM

Netherlands

Phone: 31 040 8888220

Email: nicole.van.limpt@mmc.nl

Abstract

Background: Atrial fibrillation is commonly associated with obesity. Observational studies have shown that weight loss is associated with improved prognosis and a decrease in atrial fibrillation frequency and severity. However, despite these benefits, nonadherence to lifestyle programs is common.

Objective: In this study, we evaluated adherence to and feasibility of a multidisciplinary lifestyle program focusing on behavior change in patients with atrial fibrillation and obesity.

Methods: Patients with atrial fibrillation and obesity participated in a 1-year goal-oriented cardiac rehabilitation program. After baseline assessment, the first 3 months included a cardiac rehabilitation intervention with 4 fixed modules: lifestyle counseling (with an advanced nurse practitioner), exercise training, dietary consultation, and psychosocial therapy; relaxation sessions were an additional optional treatment module. An advanced nurse practitioner monitored the personal lifestyle of each individual patient, with assessments and consultations at 3 months (ie, immediately after the intervention) and at the end of the year (ie, 9 months after the intervention). At each timepoint, level of physical activity, personal goals and progress, atrial fibrillation symptoms and frequency (Atrial Fibrillation Severity Scale), psychosocial stress (Generalized Anxiety Disorder-7), and depression (Patient Health Questionnaire-9) were assessed. The primary endpoints were adherence (defined as the number of visits attended as percentage of the number of planned visits) and completion rates of the cardiac rehabilitation intervention (defined as performing at least of 80% of the prescribed sessions). In addition, we performed an exploratory analysis of effects of the cardiac rehabilitation program on weight and atrial fibrillation symptom frequency and severity.

Results: Patients with atrial fibrillation and obesity (male: n=8; female: n=2; age: mean 57.2 years, SD 9.0; baseline weight: mean 107.2 kg, SD 11.8; baseline BMI: mean 32.4 kg/m², SD 3.5) were recruited. Of the 10 participants, 8 participants completed the 3-month cardiac rehabilitation intervention, and 2 participants did not complete the cardiac rehabilitation intervention (both because of personal issues). Adherence to the fixed treatment modules was 95% (mean 3.8 sessions attended out of mean 4 planned) for lifestyle counseling, 86% (mean 15.2 sessions attended out of mean 17.6 planned) for physiotherapy sessions, 88% (mean 3.7 sessions attended out of mean 4.1 planned) for dietician consultations, and 60% (mean 0.6 sessions attended out of mean 1.0 planned) for psychosocial therapy; 70% of participants (7/10) were referred to the optional relaxation sessions, for which adherence was 86% (mean 2 sessions attended out of mean 2.4 planned). The frequency of atrial fibrillation symptoms was reduced immediately after the intervention (before: mean 35.6, SD 3.8; after: mean 31.2, SD 3.3), and this was sustained at 12 months (mean 24.8, SD 3.2). The severity of atrial fibrillation complaints immediately after the intervention (mean 20.0, SD 3.7) and at 12 months (mean 9.3, SD 3.6) were comparable to that at baseline (mean 16.6, SD 3.3).

Conclusions: A 1-year multidisciplinary lifestyle program for obese patients with atrial fibrillation was found to be feasible, with high adherence and completion rates. Exploratory analysis revealed a sustained reduction in atrial fibrillation symptoms; however, these results remain to be confirmed in large-scale studies.

(*JMIR Form Res* 2022;6(4):e32625) doi: [10.2196/32625](https://doi.org/10.2196/32625)

KEYWORDS

cardiac rehabilitation; atrial fibrillation; obesity; participation; completion; adherence; lifestyle

Introduction

Worldwide, atrial fibrillation is the most common sustained cardiac arrhythmia in adults [1]. It has been estimated that 17.9 million people will suffer this condition in Europe by 2060 [2]. The prevalence of atrial fibrillation also increases with the presence of traditional cardiovascular disease risk factors including diabetes, obesity, and hypertension [3]. Importantly, several longitudinal studies [4,5] have demonstrated the association between obesity and cardiac dysrhythmias with increased risk of atrial fibrillation. Moreover, patients with obesity often have a higher burden and faster progression of atrial fibrillation symptoms compared with patients having a normal weight. Not surprisingly, sustainable weight loss is, therefore, associated with a decrease in atrial fibrillation symptoms [6]. Lifestyle modification, including education, psychosocial support, dietary counseling, and physical training, are effective strategies to achieve weight loss in order to reduce atrial fibrillation symptoms and improve quality of life [7]. European Society of Cardiology guidelines [8] for the management of atrial fibrillation recommend lifestyle modification to achieve weight reduction for obese patients with atrial fibrillation; however, in practice, the effects of traditional lifestyle modification programs are often not maintained in the long-term because regular cardiac rehabilitation programs, which focus on restoring the physical condition, improving lifestyle, and providing psychological support throughout, do not sufficiently focus on behavior change [9]. In addition, only a small proportion of patients with atrial fibrillation are included in cardiac rehabilitation programs, despite proven clinical benefits [10,11]. Low participation rates are caused by suboptimal referral by medical professionals. In addition, participation in cardiac rehabilitation programs for patients with atrial fibrillation is not reimbursed by health insurance in the Netherlands. Although at the start of cardiac rehabilitation, more than 80% of patients are overweight and more than 50% have metabolic syndrome [12], most cardiac rehabilitation programs do not contain a program specifically aimed at weight loss [12,13]. In addition, most efforts to change health behaviors have had limited success due to dropout and declines in adherence [14]. Therefore, addressing barriers that influence adherence, such as a lack of personalization or tailoring of the program to a specific disease is key for improving the effectiveness of lifestyle modification programs [15,16].

In this study, we aimed to investigate the adherence and feasibility of a lifestyle program for overweight atrial fibrillation patients specifically designed to improve long-term adherence by focusing on improving self-management and sustainable behavior change.

Methods

Study Design

We conducted a feasibility study; the results of this study will be used to optimize the intervention, specifically targeted to long-term behavior change, in an associated parallel multicenter open-label randomized controlled trial (Optimal Cardiac Rehabilitation XL; Netherlands Trial Register NL5589) that will evaluate the cost-effectiveness of a multidisciplinary cardiac rehabilitation program in obese patients with atrial fibrillation and coronary artery disease.

Ethics

The study protocol was reviewed and approved by the medical research ethics committee of Máxima Medical Center. All participants were informed about the goals of the study, and all participants provided written informed consent.

Participants

Patients with atrial fibrillation who presented at the cardiac rehabilitation unit (Centre Flow, Máxima Medical Center in Veldhoven/Eindhoven) in 2016 and 2017 were eligible for participation. Patients with atrial fibrillation and obesity were referred by the cardiologist, and subsequently, invited for an intake assessment with an advanced nurse practitioner. All participants were screened for trainability and practical possibilities. To personalize the lifestyle intervention, participants were counseled to formulate personal lifestyle goals based on SMART (Specific, Measurable, Achievable, Realistic and relevant, and Timed [17]), for a 1-year lifestyle program consisting of physical training, nutrition guidance, lifestyle counseling, and psychosocial therapy, with relaxation therapy as optional additional treatment module.

For this study, we chose a sample of 10 participants, rather than using a calculation-based sample size because the goal of this study was not to evaluate clinical effectiveness but to evaluate the feasibility of the intervention in this specific patient population.

Inclusion and Exclusion Criteria

Patients with symptomatic atrial fibrillation and a BMI higher or equal to 29 kg/m² were asked to participate in the study. Exclusion criteria were progressive symptoms of heart failure, myocardial ischemia during exercise testing (ST depression \geq 2 mm), sinus tachycardia at rest with a frequency greater than 110 bpm or atrial fibrillation greater than 100 bpm at rest, and severe cognitive impairment (memory, attention, concentration and psychiatric disorders).

Intervention

An individual program was created with the patient, based on their exercise capacity and personal goals. To improve exercise capacity, aerobic training was included, as well as strength training. Training frequency, training duration, training intensity, length of the work and rest intervals, and dosage training were used as variables to determine prescribed aerobic and strength training. The first 6 weeks of the intervention consisted of 2 training sessions per week, and the 6 weeks thereafter consisted of training sessions once per week. Blood pressure and heart rate were checked before, during, and after sessions. Training sessions (total duration: 60 minutes) consisted of aerobic training (treadmill or bicycle ergometer; a build-up time of 2 weeks at 40% to 50% of maximum rate of oxygen consumption was recommended) and strength training. For each strength training exercise (seated leg press, chest press, biceps curls, triceps curls, core muscle exercises, low-dose hip abductor-adductor exercise, and low row back exercise), 2 sets were performed with 1 to 2 minute breaks in-between, and each set consisted of 12 repetitions. In addition to the exercise program, all participants received an individual intake assessment with a dietician and a psychologist. If indicated (ie, depending on care need), treatments were planned with the dietician or psychologist. Finally, participants were offered an optional relaxation module, which consisted of breathing exercise and body awareness relaxation techniques aimed at teaching participants to create more moments for rest and to cope better with stress.

At the end of the 3-month intervention and at the end of the year, participant progress was evaluated with the advanced nurse practitioner, and personal goals were adjusted (participants were coached on active lifestyle and personal goals). Participants were instructed to continue to pursue these goals in their home environment. We used behavior change techniques such as goal setting, action planning, motivational interviewing, information about healthy lifestyle and reviewing outcome goals.

Main Outcomes and Measures

The primary endpoints of the study were percentage of participants who completed the intervention and adherence to the treatment modules (percentage of visits attended out of those planned). Secondary endpoints included body weight, atrial fibrillation frequency and severity, exercise capacity, anxiety, and depression. The frequency and symptom severity of atrial fibrillation were assessed with the Atrial Fibrillation Severity Scale, which is a validated questionnaire [18] that combines measures of frequency and patient-perceived severity. Exercise capacity was measured using the 6-minute walking test. The 6-minute walking test was performed in a 25-meter-long corridor at a speed of the participants' preference with the instruction to cover the greatest possible distance during 6 minutes without running. The 1-repetition maximum, a measure of muscle strength, was defined as the maximum weight that could be pushed or lifted once. Anxiety and depression complaints were assessed using validated questionnaires (Patient Health Questionnaire-9 [19] and Generalized Anxiety Disorder-7 [20]). Weight and height were measured at all follow-up visits, and BMI was calculated. Secondary outcome measures were

assessed at baseline, at 3 months (ie, immediately after the intervention), and at 12 months.

Statistical Analysis

Program completion and adherence to the treatment modules and changes in secondary endpoints (body weight and Atrial Fibrillation Severity Scale score) were analyzed by calculating descriptive statistics.

Results

A total of 10 participants participated in the feasibility study. The majority of participants were male (male: $n=8$; female: $n=2$; age: mean 57.2 years, SD 9.0; baseline weight: mean 107.2 kg, SD 11.8; baseline BMI: mean 32.4 kg/m², SD 3.5). Two participants did not complete their cardiac rehabilitation programs due to personal problems.

Of the 10 participants, 8 participants completed all 4 appointments with the advanced nurse practitioner to monitor their lifestyle (mean 3.8, SD 0.42 appointments attended per participant); 2 participants canceled their last appointment, resulting in an adherence of 95% (mean 3.8 sessions attended out of mean 4 planned). All participants were referred to the exercise training program sessions, for which adherence was 86% (mean 15.2 sessions attended out of mean 17.6 planned); 9 participants were referred to a dietician (1 patient had already received dietetic treatment in a referring hospital), for which adherence was 88% (mean 3.7 sessions attended out of mean 4.1 planned); and all participants were referred to a medical psychologist, for which adherence was 60% (mean 0.6 sessions attended out of mean 1.0 planned). Of the 10 participants, 9 participants were referred to, of whom 7 participants participated in, relaxation sessions for which adherence to the relaxation sessions was 86% (mean 2 sessions attended out of mean 2.4 planned).

After the initial 3-month cardiac rehabilitation period, weight decreased from mean 107.2 kg (SD 11.8) to mean 102.5 kg (SD 13.7), and there was weight loss at 12 months compared with baseline (Table 1). The frequency of atrial fibrillation symptoms was reduced immediately after the intervention (before: mean 35.6, SD 3.8; after: mean 31.2, SD 3.3), and this was sustained at 12 months (mean 24.8, SD 3.2). Severity of atrial fibrillation complaints immediately after the intervention (mean 20.0, SD 3.7) and at 12 months (mean 9.3, SD 3.6) were comparable with that at baseline (mean 16.6, SD 3.3). Depression and anxiety scores immediately after the intervention were comparable with those at baseline. The distance walked on the 6-minute walking test was higher after the intervention (before: mean 538.9, SD 70.6 meters; after: mean 595.3, SD 72.8 meters) and at 12 months (mean 600.8, SD 94.0). Quadriceps strength (1-repetition maximum) showed an increase immediately after rehabilitation (before: mean 179.5, SD 49.3 kg; after: mean 245.3, SD 63.7 kg) and at 12 months (mean 259.7, SD 52.7 kg). The 1-repetition maximums for biceps (before: mean 22.1, SD 8.9 kg; after: mean 25.6, SD 6.0 kg) and triceps (before: mean 27.1, SD 9.7 kg; after: mean 35.6, SD 6.8 kg) also showed increases in strength immediately after cardiac rehabilitation, and at 1 year,

remained higher (biceps: mean 29.8, SD 3.4, triceps: mean 39.6, SD 2.5) than those at baseline.

Table 1. Outcome measures at baseline, at 3 months (immediately after the cardiac rehabilitation intervention), and at 12 months (follow-up).

	Baseline, mean (SD)	Postintervention (at 3 months), mean (SD)	Follow-up (at 12 months), mean (SD)
Weight (kg)	107.2 (11.8)	102.5 (13.7)	105.1 (16.1)
Atrial Fibrillation Severity Scale			
Atrial fibrillation frequency	35.6 (3.8)	31.2 (3.3)	24.8 (3.2)
Atrial fibrillation severity	20.0 (3.7)	16.6 (3.3)	9.3 (3.6)
6-Minute walking test (m)	538.9 (70.6)	595.3 (72.8)	600.8 (94.0)
Patient Health Questionnaire–9	6.8 (4.3)	4.6 (4.3)	— ^a
Generalized Anxiety Disorder–7	4.4 (5.3)	2.7 (2.9)	—
1-Repetition maximum			
Quadriceps (kg)	179.5 (49.3)	245.3 (63.7)	259.7 (52.7)
Biceps (kg)	22.1 (8.9)	25.6 (6.0)	29.8 (3.4)
Triceps (kg)	27.1 (9.7)	35.6 (6.8)	39.6 (2.5)

^aNo data.

Discussion

General

This study shows that a multidisciplinary lifestyle program focused on sustainable behavior change in individuals who are overweight with atrial fibrillation is feasible with high completion rates (8/10 participants, 80%) and high adherence to the treatment modules (60% to 95%). Although exploratory analysis revealed sustained improvements in both the frequency and the severity of the atrial fibrillation symptoms and modest weight reduction, these results need to be confirmed in large-scale randomized controlled trials.

To the best of our knowledge, this is the first study to evaluate cardiac rehabilitation adherence specifically in obese patients with symptomatic atrial fibrillation. One study [21] showed that 81.3% of general patients referred for cardiac rehabilitation attended at least one-half of the sessions; however, nearly half of the patients did not follow any dietary recommendations or did not increase their physical activity levels, and in another, the dropout rate (24% [22]) was comparable to that in our study (20%). However, whereas participation in exercise training was high (90% [22]), participation in other modules (25% for dietary counseling, 9% for relaxation therapy, 9% for psychological therapy [22]) was substantially lower than those in our study. Data on adherence to the planned sessions were lacking [22].

The relatively high completion and adherence rates in our study may be explained by the design of the intervention. First, the intervention was designed to be a nurse-coordinated multidisciplinary intervention, rather than one that focused on exercise training or dietary counseling only, which is in line with the notion that holistic care models targeting are most effective for behavior change [23]. Second, counseling on expectations and individualized goal setting constituted an important part of the intervention. In fact, a previous study [24] showed that unrealistic expectations form an important barrier

to lifestyle behavior change [24]. Also, it is well recognized that goal setting is an effective way of achieving behavior change [24,17]. Third, the high adherence rates may be explained by the fact that we used motivational interviewing, which is a communication skill set to help facilitate change and adherence to health behaviors. In fact, Pack et al [25] showed that continual, purposeful, and planned quality improvement efforts significantly increased patient participation in cardiac rehabilitation—motivational interviewing was an important strategy in achieving this.

Our exploratory analysis revealed that cardiac rehabilitation resulted in sustained reductions in the frequency and severity of atrial fibrillation symptoms. However, in contrast with other studies [6,26,27] that have shown that weight loss has beneficial effects on atrial fibrillation symptoms, the reduction in symptoms in our study was not associated with weight loss, suggesting that other physiological mechanisms, such as improvement of traditional cardiovascular risk factors and an increase in physical fitness, play a role. In line with this assumption, Pathak et al [6] observed a significant dose-response relationship between improvements in physical fitness and reduction in the risk of atrial fibrillation recurrence. Other studies [26,28] have shown positive effects of comprehensive management of concomitant cardiovascular risk factors on atrial fibrillation recurrence. Although these studies [26,28,29] underline that targeting modifiable risk factors, obesity and physical fitness should be an essential pillar of atrial fibrillation care, the optimal approach to achieve sustained long-term effects in obese patients with atrial fibrillation remains to be established.

Limitations

An important limitation of this study was the small number of study participants. We, therefore, described conclusions only about feasibility and not about the outcomes. Furthermore, a control group was not included because the primary goal of this study was to evaluate feasibility and adherence; therefore, the

reported effects of the intervention on weight and atrial fibrillation symptoms should be regarded as exploratory. The results of this feasibility study will be applied to an ongoing study.

Conclusions

This study showed that a nurse-coordinated multidisciplinary lifestyle program for obese patients with atrial fibrillation is feasible, with a high level of completion and adherence. Effects on atrial fibrillation symptoms and weight reduction remain to be determined in a randomized controlled trial.

Acknowledgments

This study was not funded. The authors would like to thank all patients and professionals for contributing to this research.

Conflicts of Interest

None declared.

References

1. Steinberg B, Turner J, Lyons A, Biber J, Chelu MG, Fang JC, et al. Systematic collection of patient-reported outcomes in atrial fibrillation: feasibility and initial results of the Utah mEVAL AF programme. *Europace* 2020 Mar 01;22(3):368-374 [FREE Full text] [doi: [10.1093/europace/euz293](https://doi.org/10.1093/europace/euz293)] [Medline: [31702780](https://pubmed.ncbi.nlm.nih.gov/31702780/)]
2. Lippi G, Sanchis-Gomar F, Cervellin G. Global epidemiology of atrial fibrillation: an increasing epidemic and public health challenge. *Int J Stroke* 2021 Feb 19;16(2):217-221. [doi: [10.1177/1747493019897870](https://doi.org/10.1177/1747493019897870)] [Medline: [31955707](https://pubmed.ncbi.nlm.nih.gov/31955707/)]
3. Kamil-Rosenberg S, Kokkinos P, Grune de Souza E Silva C, Yee WLS, Abella J, Chan K, et al. Association between cardiorespiratory fitness, obesity, and incidence of atrial fibrillation. *Int J Cardiol Heart Vasc* 2020 Dec;31:100663 [FREE Full text] [doi: [10.1016/j.ijcha.2020.100663](https://doi.org/10.1016/j.ijcha.2020.100663)] [Medline: [33145394](https://pubmed.ncbi.nlm.nih.gov/33145394/)]
4. Wang TJ, Parise H, Levy D, D'Agostino RB, Wolf PA, Vasani RS, et al. Obesity and the risk of new-onset atrial fibrillation. *JAMA* 2004 Nov 24;292(20):2471-2477. [doi: [10.1001/jama.292.20.2471](https://doi.org/10.1001/jama.292.20.2471)] [Medline: [15562125](https://pubmed.ncbi.nlm.nih.gov/15562125/)]
5. Mahajan R, Lau DH, Brooks AG, Shipp NJ, Wood JPM, Manavis J, et al. Atrial fibrillation and obesity: reverse remodeling of atrial substrate with weight reduction. *JACC Clin Electrophysiol* 2021 May;7(5):630-641. [doi: [10.1016/j.jacep.2020.11.015](https://doi.org/10.1016/j.jacep.2020.11.015)] [Medline: [33640353](https://pubmed.ncbi.nlm.nih.gov/33640353/)]
6. Pathak RK, Middeldorp ME, Meredith M, Mehta AB, Mahajan R, Wong CX, et al. Long-term effect of goal-directed weight management in an atrial fibrillation cohort: a long-term follow-up study (LEGACY). *J Am Coll Cardiol* 2015 May 26;65(20):2159-2169 [FREE Full text] [doi: [10.1016/j.jacc.2015.03.002](https://doi.org/10.1016/j.jacc.2015.03.002)] [Medline: [25792361](https://pubmed.ncbi.nlm.nih.gov/25792361/)]
7. Lowres N, Neubeck L, Freedman SB, Briffa T, Bauman A, Redfern J. Lifestyle risk reduction interventions in atrial fibrillation: a systematic review. *Eur J Prev Cardiol* 2012 Oct 07;19(5):1091-1100. [doi: [10.1177/1741826711422505](https://doi.org/10.1177/1741826711422505)] [Medline: [23126003](https://pubmed.ncbi.nlm.nih.gov/23126003/)]
8. Hindricks G, Potpara T, Dagres N, Arbelo E, Bax JJ, Blomström-Lundqvist C, et al. Corrigendum to: 2020 ESC guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *Eur Heart J* 2021 Oct 21;42(40):4194. [doi: [10.1093/eurheartj/ehab648](https://doi.org/10.1093/eurheartj/ehab648)] [Medline: [34520521](https://pubmed.ncbi.nlm.nih.gov/34520521/)]
9. Middleton KR, Anton SD, Perri MG. Long-term adherence to health behavior change. *Am J Lifestyle Med* 2013 Jun 14;7(6):395-404 [FREE Full text] [doi: [10.1177/1559827613488867](https://doi.org/10.1177/1559827613488867)] [Medline: [27547170](https://pubmed.ncbi.nlm.nih.gov/27547170/)]
10. de Vries H, Kemps HM, van Engen-Verheul MM, Kraaijenhagen RA, Peek N. Cardiac rehabilitation and survival in a large representative community cohort of Dutch patients. *Eur Heart J* 2015 Jul 21;36(24):1519-1528. [doi: [10.1093/eurheartj/ehv111](https://doi.org/10.1093/eurheartj/ehv111)] [Medline: [25888007](https://pubmed.ncbi.nlm.nih.gov/25888007/)]
11. van Engen-Verheul M, de Vries H, Kemps H, Kraaijenhagen R, de Keizer N, Peek N. Cardiac rehabilitation uptake and its determinants in the Netherlands. *Eur J Prev Cardiol* 2013 May 08;20(2):349-356. [doi: [10.1177/2047487312439497](https://doi.org/10.1177/2047487312439497)] [Medline: [22345699](https://pubmed.ncbi.nlm.nih.gov/22345699/)]
12. Ades P, Savage P, Harvey-Berino J. The treatment of obesity in cardiac rehabilitation. *J Cardiopulm Rehabil Prev* 2010;30(5):289-298 [FREE Full text] [doi: [10.1097/HCR.0b013e3181d6f9a8](https://doi.org/10.1097/HCR.0b013e3181d6f9a8)] [Medline: [20436355](https://pubmed.ncbi.nlm.nih.gov/20436355/)]
13. Menezes AR, Lavie CJ, De Schutter A, Milani RV, O'Keefe J, DiNicolantonio JJ, et al. Lifestyle modification in the prevention and treatment of atrial fibrillation. *Prog Cardiovasc Dis* 2015 Sep;58(2):117-125. [doi: [10.1016/j.pcad.2015.07.001](https://doi.org/10.1016/j.pcad.2015.07.001)] [Medline: [26184674](https://pubmed.ncbi.nlm.nih.gov/26184674/)]
14. Kelly MP, Barker M. Why is changing health-related behaviour so difficult? *Public Health* 2016 Jul;136:109-116 [FREE Full text] [doi: [10.1016/j.puhe.2016.03.030](https://doi.org/10.1016/j.puhe.2016.03.030)] [Medline: [27184821](https://pubmed.ncbi.nlm.nih.gov/27184821/)]
15. Burgess E, Hassmén P, Welvaert M, Pumpa K. Behavioural treatment strategies improve adherence to lifestyle intervention programmes in adults with obesity: a systematic review and meta-analysis. *Clin Obes* 2017 May 15;7(2):105-114. [doi: [10.1111/cob.12180](https://doi.org/10.1111/cob.12180)] [Medline: [28199047](https://pubmed.ncbi.nlm.nih.gov/28199047/)]

16. Leung AWY, Chan RSM, Sea MMM, Woo J. An overview of factors associated with adherence to lifestyle modification programs for weight management in adults. *Int J Environ Res Public Health* 2017 Aug 16;14(8):922 [FREE Full text] [doi: [10.3390/ijerph14080922](https://doi.org/10.3390/ijerph14080922)] [Medline: [28813030](https://pubmed.ncbi.nlm.nih.gov/28813030/)]
17. Bovend'Eerd TJH, Botell RE, Wade DT. Writing SMART rehabilitation goals and achieving goal attainment scaling: a practical guide. *Clin Rehabil* 2009 May;23(4):352-361. [doi: [10.1177/0269215508101741](https://doi.org/10.1177/0269215508101741)] [Medline: [19237435](https://pubmed.ncbi.nlm.nih.gov/19237435/)]
18. Dorian P, Guerra PG, Kerr CR, O'Donnell SS, Crystal E, Gillis AM, et al. Validation of a new simple scale to measure symptoms in atrial fibrillation: the Canadian Cardiovascular Society Severity in Atrial Fibrillation scale. *Circ Arrhythm Electrophysiol* 2009 Jul;2(3):218-224. [doi: [10.1161/CIRCEP.108.812347](https://doi.org/10.1161/CIRCEP.108.812347)] [Medline: [19808471](https://pubmed.ncbi.nlm.nih.gov/19808471/)]
19. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001 Oct;16(9):606-613 [FREE Full text] [doi: [10.1046/j.1525-1497.2001.016009606.x](https://doi.org/10.1046/j.1525-1497.2001.016009606.x)] [Medline: [11556941](https://pubmed.ncbi.nlm.nih.gov/11556941/)]
20. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006 May 22;166(10):1092-1097. [doi: [10.1001/archinte.166.10.1092](https://doi.org/10.1001/archinte.166.10.1092)] [Medline: [16717171](https://pubmed.ncbi.nlm.nih.gov/16717171/)]
21. Kotseva K, Wood D, De Bacquer D, De Backer G, Rydén L, Jennings C, EUROASPIRE Investigators. EUROASPIRE IV: A European Society of Cardiology survey on the lifestyle, risk factor and therapeutic management of coronary patients from 24 European countries. *Eur J Prev Cardiol* 2016 May 16;23(6):636-648. [doi: [10.1177/2047487315569401](https://doi.org/10.1177/2047487315569401)] [Medline: [25687109](https://pubmed.ncbi.nlm.nih.gov/25687109/)]
22. Brouwers RWM, Houben VJG, Kraal JJ, Spee RF, Kemps HMC. Predictors of cardiac rehabilitation referral, enrolment and completion after acute myocardial infarction: an exploratory study. *Neth Heart J* 2021 Mar 08;29(3):151-157 [FREE Full text] [doi: [10.1007/s12471-020-01492-0](https://doi.org/10.1007/s12471-020-01492-0)] [Medline: [33030659](https://pubmed.ncbi.nlm.nih.gov/33030659/)]
23. Jennings C, Astin F. A multidisciplinary approach to prevention. *Eur J Prev Cardiol* 2017 Jun 16;24(3_suppl):77-87. [doi: [10.1177/2047487317709118](https://doi.org/10.1177/2047487317709118)] [Medline: [28618913](https://pubmed.ncbi.nlm.nih.gov/28618913/)]
24. Burgess E, Hassmén P, Pumpa KL. Determinants of adherence to lifestyle intervention in adults with obesity: a systematic review. *Clin Obes* 2017 Jun;7(3):123-135 [FREE Full text] [doi: [10.1111/cob.12183](https://doi.org/10.1111/cob.12183)] [Medline: [28296261](https://pubmed.ncbi.nlm.nih.gov/28296261/)]
25. Pack Q, Johnson L, Barr L, Daniels S, Wolter A, Squires R, et al. Improving cardiac rehabilitation attendance and completion through quality improvement activities and a motivational program. *J Cardiopulm Rehabil Prev* 2013;33(3):153-159 [FREE Full text] [doi: [10.1097/HCR.0b013e31828db386](https://doi.org/10.1097/HCR.0b013e31828db386)] [Medline: [23595006](https://pubmed.ncbi.nlm.nih.gov/23595006/)]
26. Abed HS, Wittert GA, Leong DP, Shirazi MG, Bahrami B, Middeldorp ME, et al. Effect of weight reduction and cardiometabolic risk factor management on symptom burden and severity in patients with atrial fibrillation: a randomized clinical trial. *JAMA* 2013 Dec 20;310(19):2050-2060. [doi: [10.1001/jama.2013.280521](https://doi.org/10.1001/jama.2013.280521)] [Medline: [24240932](https://pubmed.ncbi.nlm.nih.gov/24240932/)]
27. Pathak RK, Middeldorp ME, Lau DH, Mehta AB, Mahajan R, Twomey D, et al. Aggressive risk factor reduction study for atrial fibrillation and implications for the outcome of ablation: the ARREST-AF cohort study. *J Am Coll Cardiol* 2014 Dec 02;64(21):2222-2231 [FREE Full text] [doi: [10.1016/j.jacc.2014.09.028](https://doi.org/10.1016/j.jacc.2014.09.028)] [Medline: [25456757](https://pubmed.ncbi.nlm.nih.gov/25456757/)]
28. Hong K, Glover B. The impact of lifestyle intervention on atrial fibrillation. *Curr Opin Cardiol* 2018 Jan;33(1):14-19. [doi: [10.1097/HCO.0000000000000470](https://doi.org/10.1097/HCO.0000000000000470)] [Medline: [29049041](https://pubmed.ncbi.nlm.nih.gov/29049041/)]
29. Chamberlain AM, Agarwal SK, Ambrose M, Folsom AR, Soliman EZ, Alonso A. Metabolic syndrome and incidence of atrial fibrillation among blacks and whites in the Atherosclerosis Risk in Communities (ARIC) Study. *Am Heart J* 2010 May;159(5):850-856 [FREE Full text] [doi: [10.1016/j.ahj.2010.02.005](https://doi.org/10.1016/j.ahj.2010.02.005)] [Medline: [20435195](https://pubmed.ncbi.nlm.nih.gov/20435195/)]

Abbreviations

BMI: body mass index

Edited by A Mavragani; submitted 30.08.21; peer-reviewed by D Filos, J Edwards; comments to author 12.11.21; revised version received 23.12.21; accepted 13.01.22; published 29.04.22

Please cite as:

Tenbult N, Kraal J, Brouwers R, Spee R, Eijbsbouts S, Kemps H

Adherence to a Multidisciplinary Lifestyle Program for Patients With Atrial Fibrillation and Obesity: Feasibility Study

JMIR Form Res 2022;6(4):e32625

URL: <https://formative.jmir.org/2022/4/e32625>

doi: [10.2196/32625](https://doi.org/10.2196/32625)

PMID:

©Nicole Tenbult, Jos Kraal, Rutger Brouwers, Ruud Spee, Sabine Eijbsbouts, Hareld Kemps. Originally published in JMIR Formative Research (<https://formative.jmir.org>), 29.04.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Formative Research, is properly cited. The

complete bibliographic information, a link to the original publication on <https://formative.jmir.org>, as well as this copyright and license information must be included.