

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/162696/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Yim, Ka Man and Nanda, Vidit 2023. Topological inference of the Conley index. *Journal of Dynamics and Differential Equations* 10.1007/s10884-023-10310-1

Publishers page: <http://dx.doi.org/10.1007/s10884-023-10310-1>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.





Topological Inference of the Conley Index

Ka Man Yim¹ · Vidit Nanda²

Received: 1 July 2022 / Revised: 13 August 2023 / Accepted: 28 August 2023
© The Author(s) 2023

Abstract

The Conley index of an isolated invariant set is a fundamental object in the study of dynamical systems. Here we consider smooth functions on closed submanifolds of Euclidean space and describe a framework for inferring the Conley index of any compact, connected isolated critical set of such a function with high confidence from a sufficiently large finite point sample. The main construction of this paper is a specific index pair which is local to the critical set in question. We establish that these index pairs have positive reach and hence admit a sampling theory for robust homology inference. This allows us to estimate the Conley index, and as a direct consequence, we are also able to estimate the Morse index of any critical point of a Morse function using finitely many local evaluations.

Keywords Conley index theory · Homology inference · Manifold learning

1 Introduction

There is a pair of spaces at the heart of every topological quest to study gradient-like dynamics. Such space-pairs appear, for instance, whenever one encounters a closed m -dimensional Riemannian manifold M endowed with a Morse function $f : M \rightarrow \mathbb{R}$. The fundamental result of Morse theory [19] asserts that if f admits a single critical value in an interval $[a, b] \subset \mathbb{R}$, and if this value corresponds to a unique critical point $p \in M$ of Morse index μ , then the sublevel set

$$M_{f \leq b} := \{x \in M \mid f(x) \leq b\}$$

✉ Vidit Nanda
nanda@maths.ox.ac.uk

Ka Man Yim
YimKM@cardiff.ac.uk

¹ School of Mathematics, Cardiff University, Cardiff, UK

² Mathematical Institute, University of Oxford, Oxford, UK

is obtained from $M_{f \leq a}$ by gluing a closed m -dimensional disk \mathbb{D} along a $(\mu - 1)$ -dimensional boundary sphere \mathbb{S} . The relevant pair,¹ in this case is (\mathbb{D}, \mathbb{S}) , and it follows by excision that there are isomorphisms

$$\mathbf{H}_\bullet(M_{f \leq b}, M_{f \leq a}) \simeq \mathbf{H}_\bullet(\mathbb{D}, \mathbb{S})$$

of (integral) relative homology groups. Thus, when working over field coefficients, the homology groups of $M_{f \leq b}$ are obtained by altering those of $M_{f \leq a}$ in precisely one of two ways—either the μ -th Betti number is incremented by one, or the $(\mu - 1)$ -st Betti number is decremented by one.

Attempts to extend this story beyond the class of Morse functions run head-first into two significant complications—first, since the critical points need not be isolated, one must confront arbitrary critical subsets; and second, there is no single number analogous to the Morse index which completely characterises the change in topology from $M_{f \leq a}$ to $M_{f \leq b}$. The first complication is encountered in Morse-Bott theory [1], where the class of admissible functions is constrained to ones whose critical sets are normally nondegenerate submanifolds of M . The second complication is ubiquitous in Goresky-MacPherson's stratified Morse theory [11], where the class of admissible functions is constrained to those which only admit isolated critical points. In both cases, there are satisfactory analogues of (\mathbb{D}, \mathbb{S}) obtained by separately considering tangential and normal Morse data; however, the constraints imposed on functions $M \rightarrow \mathbb{R}$ in these extensions of Morse theory are far too severe from the perspective of dynamical systems.

1.1 The Conley Index

Conley index theory [5, 22] provides a powerful generalisation of Morse theory which has been adapted to topological investigations of dynamics.

Consider an arbitrary smooth function $f : M \rightarrow \mathbb{R}$ and the concomitant gradient flow $\sigma : \mathbb{R} \times M \rightarrow M$. A subset $S \subset M$ is *invariant* under f if $\sigma(t, x)$ lies in S whenever (t, x) lies in $\mathbb{R} \times S$; and such an S is *isolated* if there exists a compact subset $N \subset M$ containing S in its interior, so that S is the largest invariant subset of f inside N . Assuming that S is isolated in this sense, let $N_- \subset N$ be any compact subset disjoint from S satisfying the following two conditions: first, any flow line in N entering N_- cannot re-enter $N \setminus N_-$; and second, the flow lines that leave N_- are precisely the flow lines that leave N entirely. Pairs of the form (N, N_-) are called *index pairs* for S , and the relative homology $\mathbf{H}_\bullet(N, N_-)$ does not depend on the choice of index pair; this relative homology is called the **homological Conley index** of S . The notion of index pairs subsumes not only the pair (\mathbb{D}, \mathbb{S}) from Morse theory, but also the analogous local Morse data for Morse-Bott functions and stratified Morse functions.

The Conley index enjoys three remarkable properties as an algebraic-topological measurement of isolated invariant sets:

- (1) being relative homology classes, Conley indices are efficiently computable [14, 15];
- (2) if $\mathbf{H}_\bullet(N, N_-)$ is nontrivial for an index pair, then one is guaranteed the existence of a nonempty invariant set in the interior of N ; and finally,
- (3) the Conley index of an isolated invariant set S remains constant across sufficiently small perturbations of $f : M \rightarrow \mathbb{R}$ even though S itself might fluctuate wildly.

As a result of these attributes, the Conley index has found widespread applications to several interesting dynamical problems across pure and applied mathematics. We have no hope

¹ Several other pairs of spaces would satisfy the same homological properties, e.g., $(M_{f \leq b}, M_{f \leq a})$ itself, or $(M_{a \leq f \leq b}, M_{f=a})$. The goal is usually to find the simplest and most explicit space-pair.

of providing an exhaustive list of these success stories here, but we can at least point the interested reader to its application in the disproof of the triangulation conjecture [18], the study of fixed points of Hamiltonian diffeomorphisms [6], travelling waves in predator–prey systems [9], heteroclinic orbits in fast-slow systems [10], chaos in the Lorenz equations [21], and local Lefschetz trace formulas for weakly hyperbolic maps [12].

1.2 Topological Inference

An enduring theme within applied algebraic topology involves recovering the homology of an unknown subset X of Euclidean space \mathbb{R}^d with high confidence from a finite point cloud $P \subset \mathbb{R}^d$ that lies on, or more realistically, near X . This task is impossible unless one assumes some form of regularity on X —no amount of finite sampling will unveil the homology groups of Cantor sets and other fractals.

The authors of [23] consider the case where X is a compact Riemannian manifold and P is drawn uniformly and independently from either X or from a small tubular neighbourhood of X in \mathbb{R}^d . Their main result furnishes, for sufficiently small radii $\epsilon > 0$ and probabilities $\delta \in (0, 1)$, an explicit lower bound $B = B_X(\epsilon, \delta)$. If the cardinality of P exceeds B , then it holds with probability exceeding $(1 - \delta)$ that the homology of X is isomorphic to that of the union P^ϵ of ϵ -balls around points of P . Similar results have subsequently appeared for inferring homology of manifolds with boundary [24], of a large class of Euclidean compacta [4], and of induced maps on homology [8].

A crucial regularity assumption underlying all of these results is that the map induced on homology by the inclusion $X \hookrightarrow X^\epsilon$ is an isomorphism for all suitably small $\epsilon > 0$. When X is smooth, this can be arranged by requiring the radius ϵ to be controlled by the injectivity radius of the embedding $X \hookrightarrow \mathbb{R}^d$, often called the *reach* of X —see [7].

1.3 This Paper

Here we consider a compact, connected and isolated critical set S of a smooth function $f : M \rightarrow \mathbb{R}$ defined on a closed submanifold $M \subset \mathbb{R}^d$. Our contributions are threefold:

- (1) we construct a specific index pair $(\mathcal{N}, \mathcal{N}_-)$ for S in terms of auxiliary data pertaining to some isolating neighbourhood of S in M ; moreover,
- (2) we establish that both \mathcal{N} and \mathcal{N}_- have positive reach when viewed as subsets of \mathbb{R}^d ; and finally,
- (3) we provide a sampling theorem for inferring the Conley index $\mathbf{H}_\bullet(\mathcal{N}, \mathcal{N}_-)$ from finite point samples of \mathcal{N} and \mathcal{N}_- .

The auxiliary data required in our construction of $(\mathcal{N}, \mathcal{N}_-)$ is a smooth real-valued function g , which is defined on an isolating neighbourhood and whose vanishing locus equals S . These are not difficult to find—one perfectly acceptable choice of g is the norm-squared of the gradient $\|\nabla f\|^2$. Using any such g along with a smoothed step function, we construct a perturbation $h : M \rightarrow \mathbb{R}$ of f which agrees with f outside the isolating neighbourhood. The set \mathcal{N} is then obtained by intersecting a sublevel set of f with a superlevel set of h ; and similarly, \mathcal{N}_- is obtained by intersecting the same sublevel set of f with an interlevel set of h . The endpoints of all intervals considered in these (sub, super and inter) level sets are regular values of f and h , i.e., ∇f and ∇h . Here is a simplified version of our main result, summarising Proposition 5.2 and Theorem 5.5.

Theorem (A) Let $(\mathcal{N}, \mathcal{N}_-)$ be our constructed index pair for S . Assume that $\mathbb{X} \subset \mathcal{N}$ is a (uniform, independent) finite point sample, and set $\mathbb{X}_- := \mathbb{X} \cap \mathcal{N}_-$. Then:

- (1) If the density of $(\mathbb{X}, \mathbb{X}_-)$ in $(\mathcal{N}, \mathcal{N}_-)$ exceeds an explicit threshold t_1 , then $\mathbf{H}_\bullet(\mathbb{X}^\epsilon, \mathbb{X}_-^\epsilon)$ is isomorphic to the Conley index of S over an open interval of choices of ϵ ;
- (2) A point sample with sufficient density can be realised with high probability $1 - \kappa$ from a uniform, i.i.d. sample of $(\mathcal{N}, \mathcal{N}_-)$, if the number of points exceeds a threshold t_2 ; and
- (3) The thresholds t_1 and t_2 depend only on the reach of the manifold, C^1 data of f and g on the isolating neighbourhood, and bounds on the norm of the second derivatives of f and g .

An essential step in our proof involves showing that \mathcal{N} and \mathcal{N}_- have positive reach. Our strategy for establishing this fact is to prove a considerably more general result, which we hope will be of independent interest. We call $E \subset \mathbb{R}^d$ a **regular intersection** if it can be written as

$$E = \bigcap_{i=1}^{\ell} f_i^{-1}(I_i)$$

for some integer $\ell > 0$; here each $f_i : M \rightarrow \mathbb{R}$ is a smooth function with 0 a regular value,² and each I_i is either the point $\{0\}$ or the interval $(-\infty, 0]$. The geometry of such intersections is coarsely governed by two positive real numbers μ and Λ —here μ bounds from below the singular values of all Jacobian minors of $(f_1, \dots, f_\ell) : M \rightarrow \mathbb{R}^\ell$ evaluated on specific strata of E specified in Definition 4.1, while Λ bounds from above the operator norm of all the Hessians Hf_i .

Lemma (B) Every (μ, Λ) -regular intersection of ℓ smooth functions $M \rightarrow \mathbb{R}$ has reach τ bounded from below by

$$\frac{1}{\tau} \leq \frac{1}{\tau_M} + \sqrt{\ell} \cdot \frac{\Lambda}{\mu},$$

where τ_M is the reach of M .

See Lemma 4.4 for the full statement and proof of this result.

1.4 Related Work

Our construction of the index pair $(\mathcal{N}, \mathcal{N}_-)$ for an isolated critical set S is inspired by Milnor’s construction for the case where S is a critical point of a Morse function [19, Section I.3]. Index pairs for isolated critical points of smooth functions have been thoroughly explored by Gromoll and Meyer [13]; the work of Chang and Ghoussoub [3] provides a convenient dictionary between Conley’s index pairs and a generalised version of these Gromoll-Meyer pairs. Also close in spirit and generality to our $(\mathcal{N}, \mathcal{N}_-)$ are the *systems of Morse neighbourhoods* around arbitrary isolated critical sets in the recent work of Kirwan and Penington [16].

1.5 Outline

In Sect. 2 we briefly introduce index pairs and the Conley index. In Sect. 3, we give an explicit construction of $(\mathcal{N}, \mathcal{N}_-)$. Section 4 is devoted to proving Lemma (B). In Sect. 5,

² That is, the gradient $\nabla_x f_i$ is nonzero $\forall x \in f_i^{-1}(0)$

we specialise the above results for regular intersections to our index pairs $(\mathcal{N}, \mathcal{N}_-)$ —in particular, we derive a sufficient sampling density for the recovery of the Conley index and give a bound on the number of uniform independent point samples required to attain this density with high confidence.

2 Conley Index Preliminaries

The definitions and results quoted in this section are sourced from Sects. III.4 and III.5 of Conley’s monograph [5]; see also Mischaikow’s survey [22] for a gentler introduction to this material.³

Let $m \leq d$ be a pair of positive integers, and consider a closed m -dimensional Riemannian submanifold of d -dimensional Euclidean space \mathbb{R}^d . Throughout this paper, we fix a smooth function $f : M \rightarrow \mathbb{R}$ and denote by $\nabla_x f$ its gradient evaluated at a point $x \in M$. The *gradient flow* of f is the solution $\sigma : \mathbb{R} \times M \rightarrow M$ to the initial value problem:

$$\frac{\partial \sigma(t, x)}{\partial t} = -\nabla_x f \text{ and } \sigma(0, x) = x. \tag{1}$$

We call $x \in M$ a *critical point* of f if $\nabla_x f = 0$, whence $\sigma(\mathbb{R}, x) = x$. Let **Crit**(f) denote the set of critical points of f , and **Crit** _{c} (f) the set of compact connected components of **Crit**(f). More generally, a subset $S \subset M$ is *invariant* under σ whenever $\sigma(\mathbb{R}, S) \subset S$. We say that S is *isolated* if there exists a compact set $K \subset M$ such that S is in the interior of K , and is precisely the set of points that cannot be sent outside of K by the flow σ ; explicitly, we must have

$$S = \{x \in K \mid \sigma(\mathbb{R}, x) \subset K\} \subset \text{int}(K). \tag{2}$$

Any such K is called an *isolating neighbourhood* of S .

Definition 2.1 Let $N_- \subset N$ be pair of compact subsets of M . We call (N, N_-) an **index pair** for the isolated invariant set S if the following axioms are satisfied:

- (IP1) the closure $\text{cl}(N \setminus N_-)$ is an isolating neighbourhood of S ;
- (IP2) the set N_- is *positively invariant* in N : that is, for any $x \in N_-$ with $\sigma([0, t], x) \subset N$ for $t > 0$, we have $\sigma([0, t], x) \subset N_-$;
- (IP3) the set N_- is an *exit set*; namely, if for some $x \in N$ and $t > 0$, we have $\sigma(t, x) \notin N$, then there exists some $s \in [0, t]$ with $\sigma([0, s], x) \subset N$ and $\sigma(s, x) \in N_-$.

Every isolated invariant set S of σ admits an index pair (N, N_-) —see [5, Sect. III.4] for a proof. The content of [5, Sect. III.5] is that if (L, L_-) is any other index pair for S , then the pointed homotopy types of N/N_- and L/L_- coincide. As a result, the relative integral homology groups $\mathbf{H}_\bullet(N, N_-)$ and $\mathbf{H}_\bullet(L, L_-)$ are isomorphic and the following notion is well-defined.

Definition 2.2 The (homological) **Conley index** of an isolated invariant set S , denoted $\mathbf{Con}_\bullet(S)$, is the relative homology $\mathbf{H}_\bullet(N, N_-)$ of any index pair for S .

It follows immediately from the additivity of homology that if S decomposes as a finite disjoint union $\bigsqcup_i S_i$ of isolated invariant subsets, then $\mathbf{Con}_\bullet(S)$ is isomorphic to the direct

³ While the Conley index is defined for isolated invariant sets of any flow $\mathbb{R} \times M \rightarrow M$ or map $M \rightarrow M$, we have confined our presentation here to gradient flows as those are directly relevant to this paper.

sum $\bigoplus_i \mathbf{Con}_\bullet(S_i)$. Therefore, it suffices to restrict attention to the case where S is connected. In this paper we consider only the special case $S \in \mathbf{Crit}_c(f)$, i.e., isolated invariant sets which are compact, connected and critical. It follows that the restriction of f to S is constant, and we will assume henceforth (without loss of generality) that $f(S) = 0$.

3 Constructing Index Pairs

Let $M \subset \mathbb{R}^d$ be a closed Riemannian submanifold and $f : M \rightarrow \mathbb{R}$ a smooth function. We consider a (connected, compact) critical set $S \in \mathbf{Crit}_c(f)$ with $f(S) = 0$ and isolating neighbourhood K , as described in Eq. (2). For each x in M , we write $H_x f$ to denote the Hessian matrix of second partial derivatives of f evaluated at x . Our goal in this section is to explicitly construct an index pair for S in the sense of Definition 2.1. We describe the ingredients of our construction below, and give an illustration of our construction in Fig. 1.

Definition 3.1 A smooth⁴ map $g : K \rightarrow \mathbb{R}$ is called a **bounding function** for S if there exists a pair of real numbers $r_0 < r_1$ for which the following properties hold:

- (G1) $S \subset g^{-1}(-\infty, r_0)$;
- (G2) $S = g^{-1}(-\infty, r_1] \cap \mathbf{Crit}(f)$;
- (G3) $g^{-1}[r_0, r_1] \cap \mathbf{Crit}(g) = \emptyset$.

We call $[r_0, r_1]$ a *regularity interval* for the bounding function g .

We note that as K is itself bounded, $g^{-1}(-\infty, r_1]$ is compact. Bounding functions always exist for isolated sets in $\mathbf{Crit}_c(f)$ —one convenient choice is furnished by the normsquare of the gradient ∇f of f ,

Lemma 3.2 *The function $g : K \rightarrow \mathbb{R}$ given by $g(x) = \|\nabla_x f\|^2$ is a bounding function for S .*

Proof Writing ∂K for the boundary of K in M , set $s = \sup_{x \in \partial K} \|\nabla_x f\|^2$ and note that $s > 0$ because K is an isolating neighbourhood of S . Since critical sets of smooth functions are closed, and K – by virtue of being an isolating neighbourhood – is compact, we have that $\mathbf{Crit}(g) \cap K$ is compact. As g is continuous, $g(\mathbf{Crit}(g))$ is compact in \mathbb{R} , and thus the regular values of g are open in \mathbb{R} . Applying Sard's theorem, regular values of g then form an open dense subset of $[0, s]$. Consequently, there exists an interval of regular values $[r_0, r_1] \subset (0, s]$ of g . Since $r_0 > 0$, and S is the only set of critical points in K , Items (G1) to (G3) of Definition 3.1 are trivially satisfied. \square

We further assume knowledge of the following numerical data.

Assumption 3.3 For a given bounding function $g : K \rightarrow \mathbb{R}$ for S , we assume:

- (G4) There is a constant $q_0 > 0$ so that the inequality

$$\frac{\|\nabla f\|}{\|\nabla g\|} \frac{(r_1 - r_0)}{2} \geq q_0$$

holds on $g^{-1}[r_0, r_1]$;

⁴ Since K has a boundary, we mean that g is smooth on an open subset of M containing K .

(G5) There are regular values (α, r) and (α, s) of $(f, g) : K \rightarrow \mathbb{R}^2$ satisfying

$$0 < \alpha \leq \frac{q_0}{2} \text{ and } \frac{r_0 + r_1}{2} < s \leq r < r_1.$$

Remark 3.4 When using $g = \|\nabla f\|^2$, we have $\nabla g = 2 \cdot Hf \cdot \nabla f$, where Hf is the Hessian of f . As long as this Hessian remains nonsingular on $g^{-1}[r_0, r_1]$, the two assumptions above are readily satisfied. In particular, we can rephrase Item (G4) to the statement that

$$\frac{\|Hf \cdot \nabla f\|}{\|\nabla f\|} \leq \frac{(r_1 - r_0)}{q_0};$$

Since the left side is bounded by the operator norm of Hf , if we set

$$b = \inf_{g(x) \in [r_0, r_1]} \|H_x f\|$$

then any $q_0 \leq \frac{(r_1 - r_0)}{b}$ suffices. Similarly, the function $(f, \|\nabla f\|^2) : K \rightarrow \mathbb{R}^2$ is singular at $x \in K$ if and only if $\nabla_x f$ is an eigenvector of $H_x f$, so points of K chosen at random will generically be regular.

A regularity interval $[r_0, r_1]$ for g may be used to construct a smooth step function which decreases from q_0 to 0. In turn, the function q facilitates the construction of a local perturbation of f near S , which we call h . This perturbation h is the last piece of information required to construct an index pair for S (Fig. 1).




Fig. 1 Left: the sublevel set $f^{-1}(-\infty, \alpha]$; Centre left: orange contour lines correspond to those of f , and grey contour lines correspond to $g = r_0$ and $g = r_1$. Note how outside $g^{-1}(-\infty, r_1]$, the functions f and h coincide, while within $g^{-1}(-\infty, r_1]$, the local addition of a non-zero $q(g)$ perturbation causes the contours of h to deform and deviate away from f . Centre right: an example of $(\mathcal{N}, \mathcal{N}_-)$. The green region corresponds to \mathcal{N}_- and the union of the green and blue regions constitute \mathcal{N} . The orange lines are contour lines of $h = \beta$ and $h = \gamma$; the black lines are the contour lines of $f = \alpha$; and the grey lines are contour lines of $g = r$ and $g = s$. Right: $(\mathcal{N}, \mathcal{N}_-)$ depicted with a stream plot of the flow along $-\nabla_x f$ superimposed (Color figure online)

Definition 3.5 The **step function** $q : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ is defined as

$$q(t) = \begin{cases} q_0 & t < r_0 \\ q_0 \left(1 + \exp\left(\frac{r_1-r_0}{r_1-t} + \frac{r_1-r_0}{r_0-t}\right) \right)^{-1} & t \in [r_0, r_1], \\ 0 & t > r_1 \end{cases} \tag{3}$$

and the q -**perturbation** of f is the smooth function $h : M \rightarrow \mathbb{R}$ given by

$$h(x) = \begin{cases} f(x) + q(g(x)) & \text{if } x \in K \\ f(x) & \text{otherwise} \end{cases} \tag{4}$$

Item (H3) of Lemma 3.6 shows that h is smooth despite its piecewise definition: f only disagrees with h on a strict subset of the interior of K .

Define the constants

$$\beta = \alpha + q(r) \text{ and } \gamma = \alpha + q(s), \tag{5}$$

where α has been chosen in Item (G4) while r and s are chosen in Item (G5). Let $(\mathcal{N}, \mathcal{N}_-)$ be the pair of spaces given by

$$\mathcal{N} = f^{-1}(-\infty, \alpha] \cap h^{-1}[\beta, \infty) \text{ and } \mathcal{N}_- = f^{-1}(-\infty, \alpha] \cap h^{-1}[\beta, \gamma]. \tag{6}$$

(Note that $\mathcal{N}_- \subset \mathcal{N} \subset K$ holds by construction). Here is the main result of this section.

We now show that $(\mathcal{N}, \mathcal{N}_-)$ from Eq. (6) satisfy (IP1) to (IP3), and thus form an index pair for S . Lemma 3.7 establishes how $(\mathcal{N}, \mathcal{N}_-)$ satisfy (IP1), and Lemma 3.8 establishes how $(\mathcal{N}, \mathcal{N}_-)$ satisfy (IP2) and (IP3). Before proceeding to prove these lemmas, we outline some relevant features of the function h from Eq. (4).

Lemma 3.6 *The function $h : M \rightarrow \mathbb{R}$ satisfies the following properties:*

- (H1) $f(x) \leq h(x)$;
- (H2) $g(x) \leq r_0 \iff h(x) = f(x) + q(0)$;
- (H3) $g(x) \geq r_1 \iff h(x) = f(x)$;
- (H4) $\langle \nabla h, \nabla f \rangle \geq 0$ with equality only attained on **Crit** (f); and,
- (H5) **Crit** (h) = **Crit** (f).

Proof The only properties here which don't follow directly from Definition 3.5 are Item (H4) and Item (H5). For Item (H4), note by Eq. (4) that $\nabla h = \nabla f$ holds outside $g^{-1}[r_0, r_1]$ since the derivative of q vanishes in this region. So we consider $x \in g^{-1}[r_0, r_1]$, and calculate

$$\nabla_x f = \nabla_x f + q'(g(x)) \cdot \nabla_x g.$$

It is readily checked that $|q'(t)|$ is maximised at $t = \frac{r_0+r_1}{2}$, where its value is $\frac{2q_0}{r_1-r_0}$. Therefore,

$$\begin{aligned} \langle \nabla_x h, \nabla_x f \rangle &= \langle \nabla_x f, \nabla_x f \rangle + q'(g(x)) \langle \nabla_x g, \nabla_x f \rangle \\ &\geq \|\nabla_x f\| \|\nabla_x g\| \left(\frac{\|\nabla_x f\|}{\|\nabla_x g\|} - |q'(g(x))| \right) && \text{Cauchy-Schwarz, item (G2)} \\ &\geq \|\nabla_x f\| \|\nabla_x g\| \left(\frac{\|\nabla_x f\|}{\|\nabla_x g\|} - \frac{2q_0}{r_1-r_0} \right) && \text{Bound on } |q'| \\ &> 0 && \text{item(G4)} \end{aligned}$$

As a consequence of item (H4), we know that $\nabla h \neq 0$ on the set $g^{-1}[r_0, r_1]$. Since $\nabla h = \nabla f$ whenever $g \leq r_0$ or $g \geq r_1$, have **Crit** (f) = **Crit** (h), as required by Item (H5). \square

The next result forms the first step in our proof of Theorem 3.9.

Lemma 3.7 $(\mathcal{N}, \mathcal{N}_-)$ satisfy (IP1) : the closure $\text{cl}(\mathcal{N} \setminus \mathcal{N}_-)$ is an isolating neighbourhood of S .

Proof Before verifying Eq. (2) with $K := \text{cl}(\mathcal{N} \setminus \mathcal{N}_-)$, we first check that the closed set K is compact by confirming that the ambient set \mathcal{N} is bounded. To this end, note that for any $x \in \mathcal{N}$, we have:

$$\begin{aligned} h(x) &= f(x) + q(g(x)) && \text{by Eq. 4} \\ &\leq \alpha + q(g(x)) && \text{by Eq. 6} \end{aligned}$$

A second appeal to Eq. (6) gives $h(x) \geq \beta$ for $x \in \mathcal{N}$, whence $\beta - \alpha \leq q(g(x))$. But $\beta - \alpha = q(r)$ by Eq. (5) and q is strictly decreasing on $[r_0, r_1]$, which forces $g(x) \leq r \leq r_1$. Thus \mathcal{N} lies within $g^{-1}(-\infty, r_1]$, which is compact as it is a closed subset of an isolating neighbourhood which is compact itself.

Next we establish that S lies in the interior of K by showing that $S \subset f^{-1}(-\infty, \alpha) \cap h^{-1}(\gamma, \infty)$. For this purpose, note that $f(S) = 0$ by assumption and $\alpha > 0$ by Item (G4), so $S \subset f^{-1}(-\infty, \alpha]$ is immediate. And since $S \subset g^{-1}(-\infty, r_0]$ by Item (G1), we have from Item (H2) that $h(S) = q_0$. Now,

$$\begin{aligned} \gamma &= \alpha + q(s) && \text{by Eq. 5} \\ &\leq \alpha + \frac{q_0}{2} && \text{since } s > \frac{r_0 + r_1}{2} \text{ by Item (G5)} \\ &< q_0 && \text{since } \alpha < \frac{q_0}{2} \text{ by Item (G5)} \end{aligned}$$

Thus, $h(S) = q_0$ exceeds γ , and so $S \subset h^{-1}[\gamma, \infty)$ as desired.

Finally, to see that S is the maximal invariant subset of K , begin with the facts $S \subset \mathcal{N}$ and $\mathcal{N} \subset g^{-1}(-\infty, r]$ established above, so we have

$$S \subset \mathcal{N} \cap \mathbf{Crit}(f) \subset g^{-1}(-\infty, r] \cap \mathbf{Crit}(f).$$

Item (G2) guarantees $g^{-1}(-\infty, r] \cap \mathbf{Crit}(f) = S$, so we conclude that $S = \mathcal{N} \cap \mathbf{Crit}(f)$. Since S lies on a single level set $f = 0$, there are no connecting orbits between points of S , and so S is the maximal invariant subset of K . □

Lemma 3.8 \mathcal{N}_- is positively invariant in \mathcal{N} , satisfying (IP2); and furthermore, it is an exit set of \mathcal{N} , thus satisfying (IP3).

Proof We note from Eq. (1) that σ flows along the gradient $-\nabla f$, so f is non-increasing along the flow. Thus if $x \in \mathcal{N}$, then $f(\sigma(t, x)) \leq \alpha$; and by Item (H4), h is also non-increasing along the flow. Thus if $x \in \mathcal{N}_-$, then $\gamma \geq h(\sigma(t, x))$. Since $\mathcal{N}_- = f^{-1}(-\infty, \alpha] \cap h^{-1}[\beta, \gamma]$, if $x \in \mathcal{N}_-$, then any $\sigma(t, x) \in \mathcal{N}$ is in \mathcal{N}_- if $t > 0$, therefore it is positively invariant in \mathcal{N} and satisfies Item (IP2).

We now show that \mathcal{N}_- satisfies the exit set condition Item (IP3). Consider any $x \in \mathcal{N}$, such that $\sigma(t, x) \notin \mathcal{N}$ for some $t > 0$. Then either $f(\sigma(t, x)) > \alpha$ or $h(\sigma(t, x)) < \beta$. Since f cannot increase along $\sigma(t, x)$ and $f(x) \leq \alpha$, we have $f(\sigma(s, x)) \leq \alpha$. Similarly, because h cannot increase along $\sigma(t, x)$, then we must have $h(\sigma(t, x)) < \beta$. As $h(x) \geq \beta$, there must be some $s \in [0, t)$ where $h(\sigma(s, x)) = \beta$ by continuity. Therefore, there is some $s \in [0, t]$ such that $\sigma(s, x) \in \mathcal{N}_-$ for any $x \in \mathcal{N}$ that flows outside \mathcal{N} at some $t > 0$. □

We can now state the main result of this section, which follows from $(\mathcal{N}, \mathcal{N}_-)$ satisfying (IP1) to (IP3), which we have shown in Lemmas 3.7 and 3.8.

Theorem 3.9 *The pair $(\mathcal{N}, \mathcal{N}_-)$ from Eq. (6) is an index pair for S .*

4 The Geometry of Regular Intersections

Given our definition of $(\mathcal{N}, \mathcal{N}_-)$ in Eq. (6), the problem of inferring Conley indices is subsumed by the more general task of inferring the homology of subsets generated by taking finite intersections of level and sublevel sets of smooth functions at regular values. We parametrise this class of subsets as follows.

Definition 4.1 A non-empty subset E of a compact Riemannian manifold M is called a (μ, Λ) -**regular intersection** for real numbers $\mu > 0$ and $\Lambda \geq 0$ if there exist (finitely many) smooth functions $f_1, \dots, f_\ell : M \rightarrow \mathbb{R}$, such that E can be written as a finite intersection

$$E = \bigcap_{i=1}^{\ell} f_i^{-1}(I_i) \tag{7}$$

where each $f_i^{-1}(I_i)$ is either a level set with $I_i = \{0\}$ or a sublevel set with $I_i = (-\infty, 0]$, with 0 being a regular value of each f_i ; moreover, these f_i satisfy the following criteria:

- (R1) For any $1 \leq k \leq \ell$ and set of indices $1 \leq i_1 < i_2 < \dots < i_k \leq \ell$ with $f_{(i_1, \dots, i_k)} = (f_{i_1}, \dots, f_{i_k}) : M \rightarrow \mathbb{R}^k$, the Jacobian $d_p f_{(i_1, \dots, i_k)}$ is surjective at all points p in the intersection $f_{(i_1, \dots, i_k)}^{-1}(0) \cap E$, and the smallest non-zero singular value of this Jacobian is greater or equal to μ .
- (R2) The supremum $\sup_{p \in M} \|H_p f_i\|$ of the norm of each f_i 's Hessian on M is bounded above by Λ ; here

$$\|H_p f_i\| := \sup_{\|X\|_{\mathbb{R}^m} = 1} \|H_p f_i(X)\|_{\mathbb{R}^m} . \tag{8}$$

Next we show that regular intersections are topologically well-behaved; in the statement below, we write $\text{int}(A)$ to indicate the interior of a subset A of M , and $\text{cl}(A)$ to denote its closure in M .

Lemma 4.2 *Every regular intersection of the form*

$$E = \bigcap_{i=1}^{\ell} f_i^{-1}(-\infty, 0]$$

is a regular closed subset of M , i.e. $E = \text{cl}(\text{int}(E))$. In particular,

$$\text{int}(E) = \bigcap_{i=1}^{\ell} f_i^{-1}(-\infty, 0).$$

Proof We first check that $\text{int}(E)$ has the desired form; to this end, note that for each sublevel set $f_i^{-1}(-\infty, 0]$ taken at a regular value, the open sublevel set $f_i^{-1}(-\infty, 0)$ is the interior of $f_i^{-1}(-\infty, 0]$ (see [17, Proposition 5.46]). Thus, as $f_i^{-1}(-\infty, 0)$ is the largest open set in

$f_i^{-1}(-\infty, 0]$, the intersection $\bigcap_{i=1}^{\ell} f_i^{-1}(-\infty, 0)$ must contain any open set of E , including $\text{int}(E)$. However, as $\bigcap_{i=1}^{\ell} f_i^{-1}(-\infty, 0) \subset E$, it follows that

$$\text{int}(E) \subseteq \bigcap_{i=1}^{\ell} f_i^{-1}(-\infty, 0) \subset E \implies \text{int}(E) = \bigcap_{i=1}^{\ell} f_i^{-1}(-\infty, 0).$$

Since it follows from the definition of closure that $E \supseteq \text{cl}(\text{int}(E))$, it suffices to show that $E \subseteq \text{cl}(\text{int}(E))$. Consider $p \in E \setminus \text{int}(E)$, where without loss of generality we assume that $f_1(p) = \dots = f_k(p) = 0$ and $f_i(p) < 0$ for $i > k$. Let $u_i = \frac{\nabla_p f_i}{\|\nabla_p f_i\|}$. Let \tilde{u}_i be the component of u_i orthogonal to all u_j where $j \neq i$ and $1 \leq j \leq k$. Since E is a regular intersection (Definition 4.1), we have $\tilde{u}_i \neq 0$ and $\langle \tilde{u}_i, u_j \rangle$ is positive and non-zero if and only if $i = j$. Define

$$v = - \sum_{i=1}^k \tilde{u}_i, \tag{9}$$

so $\langle v, u_i \rangle < 0$ for all $i = 1, \dots, k$ (thus patently $v \neq 0$). Consider a continuous curve $\gamma(t)$ on M where $\gamma(0) = p$ and $\dot{\gamma}(0) = v$. As f_i are continuous, $f_i(p) = 0$ and $\langle \nabla_p f_i, \dot{\gamma}(0) \rangle < 0$ for $i \leq k$; and $f_i(p) < 0$ for $i > k$, there is some sufficiently small $\epsilon > 0$ such that for all $t \in (0, \epsilon)$, we have $f_i(\gamma(t)) < 0$ for all $i \in \{1, \dots, n\}$. We thus have for any $p \in E \setminus \text{int}(E)$ a sequence of points $\gamma(-t)$ for $t \in (0, \epsilon)$ in E , whose limit is p . Therefore, $E \subseteq \text{cl}(\text{int}(E))$ as desired. \square

We now proceed to analyse the geometry of regular intersections through the perspective of [7]. For any closed subset $A \subset \mathbb{R}^d$, let $d_A(x) := d_{\mathbb{R}^d}(x, A)$ denote the distance of any point $x \in \mathbb{R}^d$ to A , and let $\mathbf{NN}_A(x) \subseteq A$ be the set of nearest neighbours of x in A . As A is closed, $\mathbf{NN}_A(x)$ is a non-empty closed subset of \mathbb{R}^d . We let $\mathbf{UP}(A)$ be the set of points $x \in \mathbb{R}^d$ for which admit a unique nearest neighbour in A :

$$\mathbf{UP}(A) := \left(x \in \mathbb{R}^d : \#\mathbf{NN}_A(x) = 1 \right), \tag{10}$$

There is a projection map $\xi_A : \mathbf{UP}(A) \rightarrow A$ that sends each x to its unique nearest neighbour in A . For p in A , we define the subset $\mathbf{UP}(A, p) = \{x \in \mathbf{UP}(A) : \xi_A(x) = p\}$.

We also consider the complement of $\mathbf{UP}(A)$ in \mathbb{R}^d , which is the *medial axis* of A :

$$\mathbf{Med}(A) := \left(x \in \mathbb{R}^d : \#\mathbf{NN}_A(x) > 1 \right) = \mathbb{R}^d \setminus \mathbf{UP}(A). \tag{11}$$

The *local feature size* of $p \in A$ is

$$\tau_A(p) := d_{\mathbb{R}^d}(p, \mathbf{Med}(A)). \tag{12}$$

We say A has positive local feature size if $\tau_A(p) > 0$ for all $p \in A$.

Definition 4.3 The **reach** of A is the infimum of the local feature size over A

$$\tau_A := \inf_{p \in A} \tau_A(p). \tag{13}$$

We say that A has **positive reach** if $\tau_A > 0$.

One can also show that

$$\tau_A = \inf_{x \in \mathbf{Med}(A)} d_A(x). \tag{14}$$

Closed submanifolds of Euclidean space have positive reach [17], but in general the class of positive-reach subsets of \mathbb{R}^d includes many non-manifold spaces. Our goal in this Section is to prove the following result, which is Lemma (B) from the Introduction.

Lemma 4.4 *Every (μ, Λ) -regular intersection $E = \bigcap_{i=1}^{\ell} f^{-1}(I_i)$ has its reach bounded from below by $\rho_k > 0$, which is given by*

$$\frac{1}{\rho_k} = \frac{1}{\tau_M} + \sqrt{k} \cdot \frac{\Lambda}{\mu} \tag{Eq. 28}$$

where for any $p \in E$, the number of functions f_i which is zero on p is at most $k \leq l$.

In order to arrive at this result, we first recall some fundamental facts about the reach.

4.1 Geometric Consequences of the Reach

If $A \subset \mathbb{R}^d$ has positive reach, then every $x \in \mathbb{R}^d$ with $d(x, A) < \tau_A$ has a unique projection $\xi_A(x)$ in A . This is expressed in Federer’s tubular neighbourhood theorem [7], recalled below.

Theorem 4.5 *Let A be a subset of \mathbb{R}^d with $\tau_A > 0$. Then for $r \leq \tau_A$, the set*

$$A^r := \bigcup_{p \in A} B_r(p)$$

is entirely contained within $\mathbf{UP}(A)$ from Eq. (10).

The reach also places constraints on the length of shortest paths between two points on a shape; here is the content of [2, Theorem 1 & Corollary 1]. In the statement below, $B_r(\cdot)$ denotes a closed Euclidean ball around a point x whereas $B_r(x)$ denotes the corresponding open ball.

Theorem 4.6 *Assume that $A \subset \mathbb{R}^d$ has positive reach. Consider points $p, q \in A$ contained $B_r(\cdot)$ with $r < \tau_A$. Then:*

- (i) *There is a geodesic path connecting p and q in A which lies entirely within $B_r(x) \cap A$.*
- (ii) *The length of this geodesic path is bounded above by*

$$d_A(p, q) \leq 2\tau_A \operatorname{asin} \left(\frac{\|p - q\|}{2\tau_A} \right). \tag{15}$$

Let $\mathbf{Med}(A, p) = \{x \in \mathbf{Med}(A) : p \in \mathbf{NN}_A(x)\}$. For $A \subset \mathbb{R}^d$ closed, consider the function $\tau_A^+ : A \rightarrow [0, \infty]$ defined by


$$\tau_A^+(p) = \begin{cases} d_{\mathbb{R}^d}(p, \mathbf{Med}(A, p)) & \text{if } \mathbf{Med}(A, p) \neq \emptyset \\ \infty & \text{if } \mathbf{Med}(A, p) = \emptyset. \end{cases} \tag{16}$$

This function furnishes lower bounds for the reach in the following sense.

Lemma 4.7 *Let $A \subset \mathbb{R}^d$ be a closed subset. Then:*

- (i) *For any $p \in A$, we have $\tau_A^+(p) \geq \tau_A(p)$; and*
- (ii) *$\inf_{p \in A} \tau_A^+(p) = \tau_A$.*

Fig. 2 An illustration of the constraint placed on points $q \in A$ in relation to point $p \in A$ due to Lemma 4.8: implies any point $q \in A$ must lie outside the ball of radius $\tau_A^+(p)$ (boundary indicated by a dashed line), if the radial vector from p to the centre of the ball points along $x - p$ for some $x \in \mathbf{UP}(A, p)$



Proof If $\mathbf{Med}(A, p) = \emptyset$, then it follows from Eq. (16) that $\tau_A^+(p) = \infty \geq \tau_A(p)$. Otherwise, $\mathbf{Med}(A)$ is non-empty since it contains $\mathbf{Med}(A, p)$. Therefore,

$$\tau_A^+(p) = d_{\mathbb{R}^d}(p, \mathbf{Med}(A, p)) \geq d_{\mathbb{R}^d}(p, \mathbf{Med}(A)) = \tau_A(p).$$

We turn now to the second assertion. For $x \in \mathbf{Med}(A)$, choose some point $p_x \in A$ such that $p_x \in \mathbf{NN}_A(x)$. From the definition of τ_A^+ in Eq. 16, we have $d(x, A) \geq \tau_A^+(p_x)$. Thus

$$\tau_A = \inf_{x \in \mathbf{Med}(A)} d(x, A) \geq \inf_{x \in \mathbf{Med}(A)} \tau_A^+(p_x) \geq \inf_{p \in A} \tau_A^+(p).$$

As we have shown above that $\tau_A^+(p) \geq \tau_A(p)$, we also have an inequality in the opposite direction: $\inf_{p \in A} \tau_A^+(p) \geq \tau_A$. Combining these two inequalities, we obtain $\tau_A = \inf_{p \in A} \tau_A^+(p)$. □

We have considered $\tau_A^+(p)$ rather than the local feature size due to a convenient geometric property [7, Theorem 4.8(7)].

Lemma 4.8 *Let A be a closed subset of \mathbb{R}^d and consider $x \in \mathbf{UP}(A, p)$. Then for any $q \in A$,*

$$\frac{\|q - p\|}{2\tau_A^+(p)} > \left\langle \frac{q - p}{\|q - p\|}, \frac{x - p}{\|x - p\|} \right\rangle. \tag{17}$$

The geometric implications of this inequality are illustrated in Fig. 2.

4.2 The Reach of Manifolds

Here we collect some relevant facts about the reach of closed submanifolds of Euclidean space.

Fix a closed m -dimensional submanifold $M \subset \mathbb{R}^d$, and let $T_p M \subset T_p \mathbb{R}^d$ be the plane tangent to M at p in the ambient Euclidean space. By $\zeta_p : M \rightarrow T_p M$ we denote the restriction to M of the orthogonal projection $\mathbb{R}^d \rightarrow T_p M$. The normal space $N_p M$ to M at p is the kernel of this projection, i.e., the $(d - m)$ -dimensional orthogonal complement to $T_p M$ in \mathbb{R}^d . For any non-zero vectors $u \in T_p M$ and $v \in T_q M$ (where p is not necessarily the same point as q), we let $\angle(u, v)$ be the angle between u and v parallel transported in the ambient Euclidean space to $T_0 \mathbb{R}^d$ where 0 is the (arbitrarily chosen) origin. Here is [7, Theorem 4.8(12)].

Theorem 4.9 For any $p \in M$ and $r < \tau_M(p)$, we have

$$\mathbf{UP}(M, p) \cap B_r(p) = (p + N_p M) \cap B_r(p). \tag{18}$$

Below we have reproduced [2, Lemma 5 & Corollary 3].

Lemma 4.10 For any p and q in M with $\|p - q\| \leq 2\tau_M$, consider a geodesic $\gamma : [0, s] \rightarrow M$ (given by Theorem 4.6) with $\gamma(0) = p$ and $\gamma(s) = q$. Let $v(t) \in T_{\gamma(t)}M$ be the parallel transport of a unit vector $v \in T_{\gamma(0)}M$ along γ to $T_{\gamma(t)}M$. Then

$$\angle(v(0), v(t)) \leq \frac{t}{\tau_M} \text{ and} \tag{19}$$

$$\sin\left(\frac{\angle(v(0), v(t))}{2}\right) \leq \frac{\|\gamma(t) - \gamma(0)\|}{2\tau_M}. \tag{20}$$

And finally, we recall the following result from [23, Proposition 6.1] relating the curvature of manifolds to the reach.

Lemma 4.11 If $M \subset \mathbb{R}^d$ is a compact Riemannian submanifold, then for any $p \in M$ and unit vectors u, v in $T_p M$, the operator norm of the second fundamental form $\Pi : T_p M \times T_p M \rightarrow N_p M$ is bounded above by

$$\|\Pi u v\| \leq \frac{1}{\tau_M}. \tag{21}$$

4.3 Projection onto Tangent Planes

Let M be a smooth closed submanifold of \mathbb{R}^d . For each point $p \in M$, we write $B_r(p)$ to indicate the Euclidean ball of radius $r > 0$ around p , and $\zeta_p : M \rightarrow T_p M$ to indicate the restriction of the orthogonal projection from \mathbb{R}^d onto $T_p M$. The following result is a variant of [23, Lemma 5.4].

Proposition 4.12 For each $p \in M$ and $r < \sqrt{2}\tau_M$, the restriction of ζ_p to $B_r(p) \cap M$ is a local diffeomorphism.

Proof It suffices to show that the Jacobian $d_q \zeta_p$ is injective at all $q \in B_r(p) \cap M$, as the dimensions of the domain and codomain are the same (see [17, Proposition 4.8]).

Suppose $d_q \zeta_p$ is singular at some $q \in B_r(p) \cap M$. Then there is some unit vector $u \in T_q M$, when parallel transported in the ambient Euclidean space to $T_p \mathbb{R}^d$ along the line segment \overline{qp} , is orthogonal to $T_p M \subset T_p \mathbb{R}^d$. As $\|p - q\| < 2\tau_M$, we can apply Theorem 4.6 and infer that there is a geodesic $\gamma : [0, s] \rightarrow M$ where $\gamma(0) = q$ and $\gamma(s) = p$. Let $v \in T_p M$ be the parallel transport of $u \in T_q M$ along this geodesic.

As $u \in N_p M$, it is orthogonal to $v \in T_p M$ and we have $\angle(u, v) = \frac{\pi}{2}$. Applying the bound for $\angle(u, v)$ in Lemma 4.10, we have

$$\frac{t}{\tau_M} \geq \frac{\pi}{2}.$$

Substituting this into Theorem 4.6, we obtain

$$r \geq 2\tau_M \sin\left(\frac{t}{2\tau_M}\right) \geq 2\tau_M \sin\left(\frac{\pi}{4}\right) = \sqrt{2}\tau_M,$$

which contradicts our assumption that $r < \sqrt{2}\tau_M$; thus, $d_q \zeta_p$ is injective at q as desired. \square




Fig. 3 Geometric illustrations for the proof of Proposition 4.13. **A** Illustration of a geometric argument in deriving Eq. (23) in the proof of Proposition 4.13. N.B. for ambient dimension $d \geq 2$, $\zeta_p(q)$ need not lie on the line joining q and (q') . **B** Illustration of a geometric argument in deriving Eq. (24) in the proof of Proposition 4.13

Next we prove a slight extension of this result.

Proposition 4.13 *For all $p \in M$, the map ζ_p is a smooth embedding of $B_{\tau_M}(p) \cap M$ into $T_p M$.*

Proof Consider $r < \sqrt{2}\tau_M$. It suffices to show that ζ_p restricted to $B_r(p) \cap M$ is a smooth immersion (i.e. $d\zeta_p$ is injective), that ζ_p is an open map, and that ζ_p is injective (see [17, Proposition 4.22]). As ζ_p is a local diffeomorphism on $B_r(p)$ for $r < \sqrt{2}\tau_M$ (see Proposition 4.12), it is an open map ([17, Proposition 4.6]). Thus all that remains is to show that ζ_p is injective for r sufficiently small.

Suppose ζ_p is not injective on $B_r(p) \cap M$ and consider the illustration in Fig. 3a. Suppose there are two distinct points q and q' in $B_r(p) \cap M$ that project onto the same point in $\zeta_p(q) = \zeta_p(q') \in T_p M$. Since Theorem 4.9 implies $q - \zeta_p(q)$ and $q' - \zeta_p(q')$ are vectors in $N_p M$, the vector $w = q - q' = (q - \zeta_p(q)) - (q' - \zeta_p(q'))$ is also a vector in $N_p M$.

As a shorthand, let $v = q - p$ and $v' = q' - p$. Consider then $\theta = \angle(w, -v)$ and $\theta' = \angle(w, v')$; applying Lemma 4.8,

$$\cos(\theta) = \left\langle \frac{-v}{\|v\|}, \frac{w}{\|w\|} \right\rangle \leq \frac{\|v\|}{2\tau_M} < \frac{r}{2\tau_M}, \tag{22}$$

and similarly, $\cos(\theta') < \frac{r}{2\tau_M}$. As $r < \tau_M$, angles θ and θ' must be strictly greater than $\frac{\pi}{3}$. Consider then the triangle formed by p, q and q' . Applying the cosine rule, we have

$$\begin{aligned} \|w\|^2 &= \|v\|^2 + \|v'\|^2 - 2\|v\|\|v'\|\cos(\pi - (\theta + \theta')) \\ &< 2r^2(1 + \cos(\theta + \theta')) \\ &= 4r^2 \cos[2] \left(\frac{\theta + \theta'}{2} \right). \end{aligned}$$

The inequality is due to the bound q and q' being in $B_r(p)$, whence $\|v\| = \|q - p\|$ and $\|v'\| = \|q' - p\|$ are both strictly less than r . Maximising the right hand side by choosing θ and θ' to be as small as possible subject to the constraints of Eq. (22), we obtain

$$\|w\| < \frac{r^2}{\tau_M}. \tag{23}$$

Suppose now $w \in N_q M$ and consider the point $q_{\text{mid}} = q + \frac{w}{2} = \frac{q+q'}{2}$. Since we have assumed $r < \sqrt{2}\tau_M$, Eq. (23) implies $\|w\| < 2\tau_M$, and thus $\|q_{\text{mid}} - q\| < \tau_M$. Since we have assumed $(q_{\text{mid}} - q) \propto w \in N_q M$, Theorem 4.9 implies q_{mid} has a unique projection onto M , which is q . However, q_{mid} is equidistant to both q and q' which lie in M ; hence we reach a contradiction and w must thus have a non-zero component in $T_q M$ (as a vector subspace of $T_q \mathbb{R}^d$).

Consider the illustration in Fig. 3b. Let u be the non-zero projection of w onto $T_q M$ and let $\phi = \angle(u, w)$. If $w = u$, then $\phi = 0$; else, if $w - u \neq 0$, w has a component $w - u$ in $N_q M$. Thus we can apply Lemma 4.8 once again and obtain a

$$\sin(\phi) = \cos\left(\frac{\pi}{2} - \phi\right) = \left\langle \frac{w}{\|w\|}, \frac{w-u}{\|w-u\|} \right\rangle \leq \frac{\|w\|}{2\tau_M} < \frac{r^2}{2\tau_M^2} \tag{24}$$

where we substituted Eq. (23) in the final line.

As $\|p - q\| < \sqrt{2}\tau_M$, there is a geodesic γ on M such that $\gamma(0) = q$ and $\gamma(s) = p$ due to Theorem 4.6. Let us parallel transport $u \in T_q M$ along γ to $T_p M$. Let u' be the transported vector in $T_p M$. Applying Lemma 4.10,

$$\sin\left(\frac{\angle(u, u')}{2}\right) \leq \frac{r}{2\tau_M} \implies \cos(\angle(u, u')) \geq 1 - \frac{1}{2}\left(\frac{r}{\tau_M}\right)^2.$$

As $r < \sqrt{2}\tau_M$, the right hand side is positive and hence $\angle(u, u') < \frac{\pi}{2}$.

The triangle inequality on S^{d-1} implies

$$\phi = \angle(u, w) \geq \angle(u', w) - \angle(u, u') = \frac{\pi}{2} - \angle(u, u') \tag{25}$$

where the last equality is due to $u' \in T_p M$ and $w \in N_p M$. As u is a non-zero projection of w onto $T_q M$, the angle $\phi = \angle(u, w)$ between u and w is at most $\phi < \frac{\pi}{2}$. In addition, as we have shown that $\angle(u, u') < \frac{\pi}{2}$, the sine function is monotonic on both sides of the inequality. Hence, we can combine the inequalities of Eqs. 24 and 25 and obtain the following:

$$\begin{aligned} \frac{r^2}{2\tau_M^2} > \sin(\phi) &\geq \sin\left(\frac{\pi}{2} - \angle(u, u')\right) = \cos(\angle(u, u')) \geq 1 - \frac{1}{2}\left(\frac{r}{\tau_M}\right)^2 \\ \implies r > \tau_M. \end{aligned}$$

We have thus shown that any two points q and q' that project onto the same point in $T_p M$ must be at least τ_M away from p . Thus the projection is injective on $B_r(p) \cap M$. \square

4.4 Bounding the Reach of Regular Intersections

It will be helpful to first focus on the cases where the regular intersection at hand is a level set or sublevel set of a single smooth function $f : M \rightarrow \mathbb{R}$ on a compact submanifold M of \mathbb{R}^d . We suppose without loss of generality that 0 is a regular value of f . Consequently, the level set $f^{-1}(0)$ is a codimension-1 submanifold of M , and $f^{-1}(-\infty, 0]$ is a codimension-0 submanifold of M with $f^{-1}(0)$ as its boundary (see [20]). Since f is continuous, both sets $f^{-1}(0)$ and $f^{-1}(-\infty, 0]$ are closed in M . As M is compact in \mathbb{R}^d , it follows that these sets are also compact. Thus $f^{-1}(0)$ is a compact submanifold of \mathbb{R}^d , and therefore $f^{-1}(0)$ has positive reach.

Proposition 4.14 *Suppose $f : M \rightarrow \mathbb{R}$ is smooth on a positive reach closed submanifold $M \subset \mathbb{R}^d$. Consider $x \in \mathbb{R}^d$ and $p \in f^{-1}(0)$. Assume $\|\nabla_p f\| = \mu > 0$. Then:*

- (i) *If $p \in \mathbf{NN}_{f^{-1}(0)}(x)$, then $x - p = n + \lambda \nabla_p f$ where $n \in N_p M$.*
- (ii) *Let Λ be an upper bound on the norm of the Hessian of f on $B_{\tau_M}(\{x\}) \cap M$ and*

$$\frac{1}{\rho} = \frac{\Lambda}{\mu} + \frac{1}{\tau_M}. \tag{26}$$

If $x - p = n + \lambda \nabla_p f$ and $\|x - p\| < \rho$, then $\xi_{f^{-1}(0)}(x) = p$.

- (iii) *Consequently, we have*

$$\mathbf{UP}(f^{-1}(0), p) \cap B_\rho(p) = \{x \in B_\rho(p) : x - p = n + \lambda \nabla_p f \text{ and } n \in N_p M\}, \tag{27}$$

along with $\tau_{f^{-1}(0)}^+(p) \geq \rho$.

- (iv) *Consider some $x \in \mathbb{R}^d$ such that $B_r(x) \cap M \neq \emptyset$, and $r < \tau_M$. If $B_r(\{x\}) \cap f^{-1}(0) = p$ and $B_r(x) \cap f^{-1}(0) = \emptyset$, then f is either non-negative or non-positive on $B_r(x) \cap M$.*

Proof Let $g(y) = \frac{\|x-y\|^2}{2}$ and $\tilde{g} : M \rightarrow [0, \infty)$ be the restriction of g to M . We note that $\nabla \tilde{g}$ is the projection of $x - p$ onto $T_p M$.

- (i) If $\nabla_p \tilde{g} = 0$, then $x - p \perp T_p M$; else, $\tilde{g}^{-1}(r^2/2)$ is tangent to $f^{-1}(0)$ at p , and $\nabla_p \tilde{g}$ is parallel with $\nabla_p f$. As $\nabla \tilde{g}$ is the projection of $x - p$ onto $T_p M$, we can thus write $x - p = n + \lambda \nabla_p f$ where $n \in N_p M$.
- (ii) As $f^{-1}(0)$ is closed, x must have a nearest neighbour in $f^{-1}(0)$. Suppose $p \notin \mathbf{NN}_{f^{-1}(0)}(x)$ and consider $q \in \mathbf{NN}_{f^{-1}(0)}(x)$. Then $\|x - q\| \leq \|x - p\| = r$. Suppose $r < \tau_M$. Then, $B_r(\{x\}) \cap M$ is connected by Theorem 4.6. Thus there is a geodesic $\gamma : [0, s] \rightarrow M$ between $\gamma(0) = p$ and $\gamma(s) = q$ parametrised by arc length. Consider $\hat{f} = f \circ \gamma : [0, s] \rightarrow \mathbb{R}$ and $\hat{g} = g \circ \gamma : [0, s] \rightarrow \mathbb{R}$. Then by definition $\hat{f}(0) = \hat{f}(s)$ and $\hat{g}(0) \geq \hat{g}(s)$. If we Taylor expand \hat{f} and \hat{g} , we have for some s_1 and s_2 in $(0, s)$

$$\begin{aligned} \hat{f}(s) &= \hat{f}(0) + \left. \frac{d\hat{f}}{dt} \right|_{t=0} s + \left. \frac{1}{2} \frac{d^2 \hat{f}}{dt^2} \right|_{t=s_1} s^2 \implies \left. \frac{d\hat{f}}{dt} \right|_{t=0} + \left. \frac{1}{2} \frac{d^2 \hat{f}}{dt^2} \right|_{t=s_1} s = 0 \\ \hat{g}(s) &= \hat{g}(0) + \left. \frac{d\hat{g}}{dt} \right|_{t=0} s + \left. \frac{1}{2} \frac{d^2 \hat{g}}{dt^2} \right|_{t=s_2} s^2 \implies \left. \frac{d\hat{g}}{dt} \right|_{t=0} + \left. \frac{1}{2} \frac{d^2 \hat{g}}{dt^2} \right|_{t=s_2} s \leq 0. \end{aligned}$$

Evaluating the derivatives, and substituting $x - p = n + \lambda \nabla_p f$, we obtain

$$\begin{aligned} \langle \dot{\gamma}(0), \nabla_p f \rangle + \frac{1}{2} \langle \dot{\gamma}(s_1), \nabla_{\dot{\gamma}(s_1)} \nabla f(\gamma(s_1)) \rangle s &= 0 \text{ and} \\ \lambda \langle \dot{\gamma}(0), \nabla_p f \rangle + \frac{1}{2} (1 + \langle \gamma(s_2) - x, \ddot{\gamma}(s_2) \rangle) s &\leq 0, \\ \implies 1 - \lambda \langle \dot{\gamma}(s_1), \nabla_{\dot{\gamma}(s_1)} \nabla f(\gamma(s_1)) \rangle + \langle \gamma(s_2) - x, \ddot{\gamma}(s_2) \rangle &\leq 0. \end{aligned}$$

As $B_r(x) \cap M$ is geodesically convex for $t < \tau_M$ and $\|x - q\| \leq \|x - p\| < \tau_M$, we thus have $\gamma([0, s]) \subset B_r(\{x\})$. Hence $\|\gamma(s_2) - x\| \leq r$. Applying the bound on $\|\ddot{\gamma}\|$ Lemma 4.11 and the Hessian of f , we have

$$|\lambda| \Lambda + \frac{r}{\tau_M} \geq 1.$$

In addition, since $x - p = n + \lambda \nabla_p f$ and $n \perp \nabla_p f$, we have $|\lambda| \leq \frac{r}{\mu}$. Thus the existence of $q \in \mathbf{NN}_{f^{-1}(0)}(x)$ not equal to p , and $\|x - q\| \leq \|x - p\| = r$ implies

$$r \geq \left(\frac{\Lambda}{\mu} + \frac{1}{\tau_M} \right)^{-1} =: \rho$$

Hence, if $r = \|x - p\| < \rho$, the point x must project onto $f^{-1}(0)$ at p .

- (iii) As a consequence of Items Proposition 3.14 (i) and Proposition 3.14 (ii), any $x \in B_\rho(p)$ that projects onto $f^{-1}(0)$ at p is a linear combination of some $n \in N_p M$ and $\nabla_p f$, and vice versa any $x \in B_\rho(p)$ that is a linear combination of some $n \in N_p M$ and $\nabla_p f$ projects onto $f^{-1}(0)$ at p . For $x \in \mathbf{Med}(f^{-1}(0), p)$, Item Proposition 3.14 (i) implies $x - p = n + \lambda \nabla_p f$. As $x \notin \mathbf{UP}(f^{-1}(0), p)$, $\|x - p\| \geq \rho$. Thus $\tau_{f^{-1}(0)}^+(p) \geq \rho$.
- (iv) Since $r < \tau_M$, then Theorem 4.6 implies $B_r(\lfloor x \rfloor) \cap M$ is connected. We claim that $B_r(x) \cap f^{-1}(0) = \emptyset$ implies the sign of f must be non-positive or non-negative on $B_r(\lfloor x \rfloor) \cap M$. Suppose that is not the case; then there are two points of opposite sign in $B_r(\lfloor x \rfloor) \cap M$. Consider a path connecting those two points contained in $B_r(x) \cap M$. As f is continuous, there must be some point along the path where f is zero. However this contradicts $B_r(x) \cap f^{-1}(0) = \emptyset$.

□

The above observations about sublevel and level sets give us a stepping stone towards deriving the bound for the reach of regular intersections of multiple sublevel and level sets.

Proposition 4.15 Consider a (μ, Λ) -regular intersection as in Definition 4.1:

$$E = \bigcap_{i=1}^{\ell} f_i^{-1}(I_i) \tag{Equation 7}$$

For $0 \leq k \leq \ell$, define

$$\frac{1}{\rho_k} = \frac{1}{\tau_M} + \sqrt{k} \frac{\Lambda}{\mu}. \tag{28}$$

Consider $x \in \mathbb{R}^d$ and $p \in E$. Without loss of generality, assume $f_i(p) = 0$ for $i \leq k$, and $f_i(p) < 0$ otherwise. Then:

- (i) If $p \in \mathbf{NN}_E(x)$, then there are coefficients λ_i and $n \in N_p M$, such that

$$x - p = n + \sum_{i=1}^k \lambda_i \nabla_p f_i, \tag{29}$$

where $\lambda_i \geq 0$ if $I_i = (-\infty, 0]$.

- (ii) Conversely, if $x - p$ can be written in the form Eq. (29), and if $\|x - p\| < \rho_k$, then we have $\xi_E(x) = p$.
- (iii) Consequently,

$$\mathbf{UP}(E, p) \cap B_{\rho_k}(p) = \{x \in B_{\rho_k}(p) : x - p \text{ satisfies Eq. (29)}\}. \tag{30}$$

and $\tau_E^+(p) \geq \rho$.

Proof Let $r = \|x - p\|$ and consider the open Euclidean ball $B_r(x)$.

- (i) If $B_r(x) \cap M = \emptyset$, then $p \in \mathbf{NN}_M(x)$ and $x - p \in N_pM$. Else, consider the case $B_r(x) \cap M \neq \emptyset$ and write $x - p = n + t + \sum_{i=1}^k \lambda_i \nabla_p f_i$, where $t \in T_pM$ is orthogonal to $\nabla_p f_1, \dots, \nabla_p f_k$. Let $\gamma : [0, 1] \rightarrow M$ be a smooth curve wholly contained in E , with $\gamma(0) = p$ and $\dot{\gamma}(0) = v$. As $f_i(p) = 0$ for $i \leq k$, for $I_i = (-\infty, 0]$

$$\exists \epsilon > 0 \quad \text{s.t. } f_i(\gamma(t)) < 0 \quad \forall t \in (0, \epsilon) \iff \langle v, \nabla_p f_i \rangle < 0$$

and for all i ,

$$f_i(\gamma(t)) = 0 \iff \langle \dot{\gamma}(t), \nabla_{\gamma(t)} f_i \rangle = 0$$

As $p \in \mathbf{NN}_E(x)$, the ball $B_r(x)$ cannot intersect any point in E ; as such, we also have

$$\langle v, x - p \rangle \leq 0 \implies \langle v, t \rangle + \sum_{i=1}^k \lambda_i \langle v, \nabla_p f_i \rangle \leq 0 \tag{*}$$

where we substituted our expression for $x - p$ and noted that $n \in N_pM$ and $v \in T_pM$. Choosing γ so that $f_i(\gamma(t)) = 0$ for $i \leq k$ and $v = t$, we substitute into (*) and deduce

$$\langle t, t \rangle \leq 0 \implies t = 0.$$

Then, as $\nabla_p f_1, \dots, \nabla_p f_k$ are linearly independent, we can choose v such that for one $j \leq k$ where $I_j = (-\infty, 0]$, we have $\langle v, \nabla_p f_j \rangle < 0$ and $\langle v, \nabla_p f_i \rangle = 0$ for $i \neq j$. Substituting this choice of v into (*),

$$\lambda_j \langle v, \nabla_p f_j \rangle \leq 0 \text{ and } \langle v, \nabla_p f_j \rangle < 0 \implies \lambda_j \geq 0.$$

Thus $x - p = n + \sum_{i=1}^k \lambda_i \nabla_p f_i$ where $\lambda_i \geq 0$ if $I_i = (-\infty, 0]$.

- (ii) First, let us consider the case where $\lambda_i = 0$. Then $x - p \in N_pM$. As $r = \|x - p\| < \tau_M(p)$, we can apply Theorem 4.9 and deduce that $\xi_M(x) = p$. As $E \subset M$, therefore $\xi_E(x) = p$. Then let us consider the case where $\lambda_i \neq 0$ for some i . Let $\tilde{f} = \sum_{i=1}^k \alpha_i f_i$ where $\alpha_i = \lambda_i / \|\lambda\|$, where $\|\lambda\| = \sqrt{\sum_{i=1}^k \lambda_i^2}$. Since $f_i(q) \leq 0$ for $q \in E$ we note that $E \subset \tilde{f}^{-1}(-\infty, 0]$. In particular, $p \in \tilde{f}^{-1}(0)$. Since E is a regular intersection, we can check that $\|\nabla \tilde{f}\| \geq \mu > 0$ due to Item (R1). Furthermore, due to Item (R2), for any $q \in M$ and unit vector $X \in T_qM$,

$$\begin{aligned} |\langle X, \nabla_X \nabla \tilde{f} \rangle| &= \left| \sum_{i=1}^k \alpha_i \langle X, \nabla_X \nabla f_i \rangle \right| \\ &\leq \Lambda \sup_{\|\alpha\|=1} \sum_{i=1}^k \alpha_i = \sqrt{k} \Lambda. \end{aligned}$$

Thus $\tilde{\Lambda} = \sqrt{k} \Lambda$ is an upper bound on the Hessian of \tilde{f} . Because $(x - p) \propto (n + \nabla \tilde{f}(p))$ and

$$r = \|x - p\| < \rho_k = \left(\frac{1}{\tau_M} + \frac{\tilde{\Lambda}}{\mu} \right)^{-1},$$

we can apply Proposition 4.14 to p in $\tilde{f}^{-1}(0)$ and deduce that $\xi_{\tilde{f}^{-1}(0)}(x) = p$. Consider $B_r([x]) \cap M$. Since $\xi_{\tilde{f}^{-1}(0)}(x) = p$, we have $B_r([x]) \cap \tilde{f}^{-1}(0) = p$ and $B_r(x) \cap$

$\tilde{f}^{-1}(0) = \emptyset$. Furthermore, since $r < \tau_M$, the sign of \tilde{f} is either non-negative or non-positive on $B_r(\lceil x \rceil) \cap M$ according to Item Proposition 3.14 (iv). We now determine the sign of \tilde{f} on $B_r(\lceil x \rceil) \cap M$. Since there is some i for which $\lambda_i \neq 0$, let us consider a smooth curve $\sigma(t)$ on M where $\sigma(0) = p$ and $\dot{\sigma}(t) = \nabla_{\sigma(t)} \tilde{f}$. Since $\langle \dot{\sigma}(0), x - p \rangle \propto \left\| \nabla_p \tilde{f} \right\|^2 > 0$, the curve enters $B_r(\lceil x \rceil)$ at p . As \tilde{f} increases away from 0 along the curve and for sufficiently small $\epsilon > 0$, $\sigma(\epsilon) \in B_r(x) \cap M$, we deduce that $\tilde{f} > 0$ on $B_r(x) \cap M$ as we have shown that \tilde{f} is of constant sign on $B_r(x) \cap M$. In other words, $\tilde{f}^{-1}(-\infty, 0] \cap B_r(x) = \emptyset$. As we have shown that $B_r(\lceil x \rceil) \cap \tilde{f}^{-1}(0) = p$, the continuity of \tilde{f} implies $\tilde{f}^{-1}(-\infty, 0] \cap B_r(\lceil x \rceil) = p$. Because $E \subset \tilde{f}^{-1}(-\infty, 0]$ and $\tilde{f}^{-1}(-\infty, 0] \cap B_r(\lceil x \rceil) = p$, we must therefore have $E \cap B_r(\lceil x \rceil) = p$ i.e. $\xi_E(x) = p$.

(iii) Follows the reasoning of the proof of Item Proposition 3.14 (iii). □

We have now arrived at the proof of Lemma 4.4.

Proof of Lemma 4.4 As we have produced a bound on $\tau_E^+(p)$ for any $p \in E$, Item Proposition 3.14 (iii), this follows from applying Item 3.7 (ii) that $\tau_E = \inf_{p \in E} \tau_E^+(p) \geq \rho_k$ where ρ_k is as defined in Eq. (28).

In the next subsection, we establish theoretical guarantees for recovering the homology of regular intersections via the more general homology inference results for subsets with positive reach.

4.5 Homological Inference of Subsets with Positive Reach

It was shown in [23, 24] that a sufficiently dense point sample of compact manifolds with and without boundary can be used to recover their homotopy type. These arguments are readily generalised to the following inference result for Euclidean subsets with positive reach.

Proposition 4.16 *Let A be a subset of \mathbb{R}^d with positive reach τ_A . Suppose we have a finite point sample $\mathbb{X} \subset A$ that is δ -dense in A where $\delta < \frac{\bar{\tau}}{4}$, where $\bar{\tau} \in (0, \tau_A]$ is some positive lower bound on the reach of A . Then \mathbb{X}^ϵ deformation retracts to A if*

$$\frac{\epsilon}{\bar{\tau}} \in \left(\frac{1}{2} - \sqrt{\frac{1}{4} - \frac{\delta}{\bar{\tau}}}, \frac{1}{2} + \sqrt{\frac{1}{4} - \frac{\delta}{\bar{\tau}}} \right). \tag{31}$$

Given the explicit lower bound for the reach of regular intersections in Lemma 4.4, the following homology inference result for regular intersections is a direct corollary of Proposition 4.16.

Corollary 4.17 *Suppose we have a finite point sample $\mathbb{X} \subset E$ is δ -dense in a regular intersection $E = \bigcap_{i=1}^k f_i^{-1}(I_i)$ (as given in Eq. (7)) where I_i is either 0 or $(-\infty, 0]$. Let k be the maximum number of level sets $f_i^{-1}(0)$ that intersect in E . If $\delta < \frac{\rho_k}{4}$, where ρ_k is given by Eq. (28), then \mathbb{X}^ϵ deformation retracts to A if*

$$\frac{\epsilon}{\rho_k} \in \left(\frac{1}{2} - \sqrt{\frac{1}{4} - \frac{\delta}{\rho_k}}, \frac{1}{2} + \sqrt{\frac{1}{4} - \frac{\delta}{\rho_k}} \right). \tag{32}$$

We devote the remainder of this subsection to deriving Proposition 4.16, closely following the original argument for manifolds in [23, Proposition 4.1].

Lemma 4.18 *Let A be a subset of \mathbb{R}^d with positive reach. Suppose for some $B \subset \mathbb{R}^d$, we have $A \subset B \subset \mathbf{UP}(A)$. If B is star-shaped relative to A – i.e. if for every $p \in B$, the line segment $\overline{p\xi_A(p)}$ is also contained in B – then B deformation retracts to A .*

Proof Let us consider the function $F : [0, 1] \times B \rightarrow \mathbb{R}^d$

$$F(t, p) = (1 - t) \cdot p + t \cdot \xi_A(p).$$

Since projection maps ξ_A are continuous [7, Theorem 4.8(2)], the map F is continuous. Furthermore, if $\overline{p\xi_A(p)}$ is contained in B , then $F(t, p) \subset B$. We can easily check that $F(0, p) = p$, $F(1, p) = \xi_A(p) \in A$, and $F(t, p) = \xi_A(p) = p$. Therefore, F is a strong deformation retraction $F : [0, 1] \times B \rightarrow B$ of B onto A . \square

Lemma 4.19 *Assume the conditions of Proposition 4.16. Consider $x \in \mathbb{X}$ and $p \in B_\epsilon(x) \cap \mathbf{UP}(A, q)$ where $\epsilon \in (0, \tau_A)$. If $q \notin B_\epsilon(x)$, then there is a unique point $y = \partial B_\epsilon(x) \cap \overline{qp}$, such that*

$$\|y - q\| \leq \frac{\epsilon^2}{\tau_A}.$$

Proof Since $\epsilon < \tau_A$, and $p \in B_\epsilon(x)$, we know p has a unique projection $\xi_A(p) = q$. By assumption $q \notin B_\epsilon(x)$ and therefore $q \neq x$. As q is the nearest neighbour of p in A , we have $r = \|p - q\| < \|p - x\| < \epsilon < \tau_A$.

Let $q(t) = (1 - t)q + tp$ for $t \in [0, 1]$ parametrise the line segment \overline{qp} . Since $q \in B_{tr}(\{q(t)\}) \cap A \subset B_r(\{q\}) \cap A = q$ for all $t \in [0, 1]$, the entire line segment \overline{qp} lies in $\mathbf{UP}(A, q)$ Since.

We consider the continuous function $c(t) = \|q(t) - x\|$. Since $g(0) > \epsilon$ and $g(1) < \epsilon$, by continuity there must be some $t_* \in (0, 1)$ such that $c(t_*) = \|q(t_*) - x\| = \epsilon$. Since p is in the interior of $B_\epsilon(x)$ and $B_\epsilon(x)$ is convex, t_* must be unique. Thus for $y = q(t_*)$, we have $\overline{yp} \subset B_\epsilon(\{x\})$.

Since $\pi(y) = q \in A$ and $x \in A$, we can now apply Lemmas 4.8 and 4.7, and deduce that

$$\left\langle \frac{y - q}{\|y - q\|}, \frac{x - q}{\|x - q\|} \right\rangle \leq \frac{\|x - q\|}{2\tau_A}.$$

As such,

$$\begin{aligned} \epsilon^2 &= \|y - x\|^2 = \|(y - q) + (x - q)\|^2 \\ &= \|y - q\|^2 + \|x - q\|^2 - 2 \langle y - q, x - q \rangle \\ &\geq \|y - q\|^2 + \|x - q\|^2 \left(1 - \frac{\|y - q\|}{\tau_A}\right) \\ &\geq \|y - q\|^2 + \epsilon^2 \left(1 - \frac{\|y - q\|}{\tau_A}\right) \\ \implies \|y - q\| &\leq \frac{\epsilon^2}{\tau_A}. \end{aligned}$$

where in the fourth line we applied our assumption that $q \notin B_\epsilon(x)$ and the fact that $\|y - q\| = \|c(t_*) - q\| < \tau_A$. \square

Here is the promised proof of Proposition 4.16.

Proof of Proposition 4.16 Consider any point $p \in \mathbb{X}^\epsilon$ and let $\xi_A(p) = q$. Since $p \in \mathbb{X}^\epsilon$, the point p is contained in at least one open Euclidean ball $B_\epsilon(x)$ for some $x \in \mathbb{X}$. If $q \in B_\epsilon(x)$ for one such x , then \overline{pq} is also contained within $B_\epsilon(x)$ as Euclidean balls are convex. Therefore, \overline{pq} is contained in \mathbb{X}^ϵ .

Let us consider the case where q is not contained in any of the Euclidean balls containing p . From Lemma 4.19, there exists some $y \in \overline{pq}$ such that $\overline{pq} \subset B_\epsilon([x])$, and $\|y - q\| \leq \epsilon^2/\tau_A$. We can subdivide the line segment \overline{pq} into two segments \overline{qy} and \overline{py} , the latter being contained in $B_\epsilon(x)$. If there is some $x' \in \mathbb{X}$ such that the closed line segment \overline{qy} is contained in $B_\epsilon(x')$, then the entirety of \overline{pq} is contained in \mathbb{X}^ϵ . This can be achieved if both q and y are contained in $B_\epsilon(x')$. By the δ -density assumption, we can pick a point $x' \in \mathbb{X}$ such that $q \in B_\delta(x')$. If we assume $\delta < \epsilon$, then $q \in B_\epsilon(x')$.

If y is to be contained in $B_\epsilon(x')$, we require $\|x' - y\| < \epsilon$. If we choose $\delta < \epsilon - \epsilon^2/\bar{\tau}$, then by the triangle inequality,

$$\begin{aligned} \|x' - y\| &\leq \|x' - q\| + \|q - y\| \\ &< \delta + \frac{\epsilon^2}{\tau_A} \\ &< \epsilon - \frac{\epsilon^2}{\bar{\tau}} + \frac{\epsilon^2}{\tau_A} \leq \epsilon. \end{aligned}$$

Therefore for choices of ϵ that satisfy $\delta < \epsilon - \epsilon^2/\bar{\tau}$, the line segment \overline{pq} is contained in \mathbb{X}^ϵ . Since this holds for any choice of p , by the Lemma 4.18, this implies \mathbb{X}^ϵ deformation retracts to A .

Finally, we have $\delta < \epsilon - \epsilon^2/\bar{\tau}$ if and only if

$$\frac{\epsilon}{\bar{\tau}} \in \left(\frac{1}{2} - \sqrt{\frac{1}{4} - \frac{\delta}{\bar{\tau}}}, \frac{1}{2} + \sqrt{\frac{1}{4} - \frac{\delta}{\bar{\tau}}} \right).$$

5 Homological Inference of Index Pairs from Point Samples

Having constructed an index pair $(\mathcal{N}, \mathcal{N}_-)$ of S in Sect. 3, we will show in this Section that the Conley index $\mathbf{Con}_\bullet(S) \cong \mathbf{H}_\bullet(\mathcal{N}, \mathcal{N}_-)$ can be inferred from a sufficiently large finite point sample $\mathbb{X} \subset \mathcal{N}$ as described in Theorem (A) of the Introduction. We proceed by showing that \mathcal{N} and \mathcal{N}_- are both regular intersections, and thus have positive reach.

Lemma 5.1 *If the Hessians of f and g are bounded, then \mathcal{N} and \mathcal{N}_- as constructed for $S \in \mathbf{Crit}_c(f)$ as defined in Eq. (6) are regular intersections.*

Proof As the Hessians of f and g is bounded and q has bounded second derivative, the Hessian of h (Eq. (4)) is also bounded. Thus item (R2) is satisfied.

Since $\mathbf{Crit}(f) = \mathbf{Crit}(h)$ ((Item (H5)) and $S = \mathbf{Crit}(f) \cap \mathcal{N} \subset h^{-1}(\gamma, \infty) \cap f^{-1}(-\infty, \alpha)$ ((Item 3.7), ∇f and ∇h are non-zero on level sets $f^{-1}(\alpha)$, $h^{-1}(\beta)$, and $h^{-1}(\gamma)$). Because we have assumed (Item (G5), the Jacobian of $(f, h) : M \rightarrow \mathbb{R}^2$ is surjective on $f^{-1}(\alpha) \cap h^{-1}(\beta)$ and $f^{-1}(\alpha) \cap h^{-1}(\gamma)$ respectively. Since these are bounded and thus compact, the infimum of the second largest singular value of the Jacobian of (f, h) on these sets is positive. We thus satisfy Item (R1). \square

Given this characterisation of $(\mathcal{N}, \mathcal{N}_-)$ in terms of regular intersections, the following homology inference guarantee follows from Corollary 4.17.

Proposition 5.2 *Let \mathcal{N} and \mathcal{N}_- be as constructed in Eq. (6); these are (μ, Λ) -regular intersections for parameters $\mu > 0$ and $\Lambda > 0$ by Lemma 5.1. Fix $\delta \in (0, \frac{\rho_2}{4})$, where*

$$\frac{1}{\rho_2} = \frac{1}{\tau_M} + \sqrt{2} \frac{\mu}{\Lambda},$$

Let $\mathbb{X} \subset \mathcal{N}$ be a finite point sample that is δ -dense in \mathcal{N} so that $\mathbb{X}_- := \mathbb{X} \cap \mathcal{N}_-$ is δ -dense in \mathcal{N}_- . Then $\mathbf{H}_\bullet(\mathcal{N}, \mathcal{N}_-) \cong \mathbf{H}_\bullet(\mathbb{X}^\epsilon, \mathbb{X}_-^\epsilon)$ for ϵ in the open interval

$$\frac{\epsilon}{\rho_2} \in \left(\frac{1}{2} - \sqrt{\frac{1}{4} - \frac{\delta}{\rho_2}}, \frac{1}{2} + \sqrt{\frac{1}{4} - \frac{\delta}{\rho_2}} \right). \tag{33}$$

Proof A positive lower bound on the reaches of \mathcal{N} and \mathcal{N}_- is prescribed by Lemma 4.4. In particular, since each point in either \mathcal{N} or \mathcal{N}_- lies at the intersection of at most two level sets of f and h , the reaches of \mathcal{N} and \mathcal{N}_- are bounded below by $\rho_2 > 0$ (where ρ_2 is given by Eq. (28) for the case $k = 2$). We can then apply Proposition 4.16 to recover the homotopy type of \mathcal{N} and \mathcal{N}_- from \mathbb{X}^ϵ and \mathbb{X}_-^ϵ . As we have isomorphisms $\mathbf{H}_\bullet(\mathcal{N}) \xrightarrow{\cong} \mathbf{H}_\bullet(\mathbb{X}^\epsilon)$ and $\mathbf{H}_\bullet(\mathcal{N}_-) \xrightarrow{\cong} \mathbf{H}_\bullet(\mathbb{X}_-^\epsilon)$, applying the five lemma to the two long exact sequences of the pairs $(\mathbb{X}^\epsilon, \mathbb{X}_-^\epsilon)$ and $(\mathcal{N}, \mathcal{N}_-)$ gives the isomorphism $\mathbf{H}_\bullet(\mathcal{N}, \mathcal{N}_-) \cong \mathbf{H}_\bullet(\mathbb{X}^\epsilon, \mathbb{X}_-^\epsilon)$. \square

5.1 Homology Inference via Uniform Sampling

We now consider random i.i.d. samples from a compactly supported measure on a Riemannian manifold M . A Riemannian manifold is endowed with a unique Riemannian measure μ_g ; for $p \in M$ and coordinate neighbourhood $(U, \{x_i\})$ of p such that $d_p x_i$ are orthonormal in the cotangent space T_p^*M ,

$$\mu_g(p) = |d_p x_1 \wedge \dots \wedge d_p x_m|. \tag{34}$$

The authors of [23] studied how many random i.i.d. draws from a probability measure ν would suffice to obtain an ϵ -dense point sample of some measurable subset of M . We consider ν that is continuous with respect to μ_g : that is, for any measurable subset $A \subset M$, $\mu_g(A) = 0$ implies $\nu(A) = 0$. Here we will follow the argument from [23, Proposition 7.2], summarised below.

Lemma 5.3 *Let A be a compact, measurable subset of a properly embedded Riemannian submanifold of \mathbb{R}^d . Suppose ν is a probability measure supported on A that is continuous with respect to μ_g , and let \mathbb{X} be a finite set of i.i.d. draws according to ν . If*

$$\#\mathbb{X} \geq \frac{1}{K \left(\frac{\epsilon}{2}\right)} \left(\log \left(\frac{\nu(A)}{K \left(\frac{\epsilon}{4}\right)} \right) + \log \left(\frac{1}{\kappa} \right) \right). \tag{35}$$

where

$$K(r) = \inf_{p \in A} \nu(A \cap B_r(p)), \tag{36}$$

then \mathbb{X} is ϵ -dense in A with probability $> 1 - \kappa$.

The key parameter that controls the bound on the right hand side of Eq. (35) is $K(r)$; in particular, we require $K(r) > 0$ for $r > 0$ so that the bound is finite. That is the case if A is a compact, regular closed subset of M .

Lemma 5.4 *Let $A \subset M$ be a compact regular closed subset. For any $r > 0$, and measure ν supported on A that is continuous with respect to μ_g on M ,*

$$K(r) = \inf_{p \in A} \nu(B_r(p) \cap A) > 0.$$

Proof We first show that $\nu(B_r(p) \cap A) > 0$ for $p \in E$. Since ν is supported on A , and continuous with respect to μ_g , it suffices to show that $B_r(p) \cap A$ contains some open subset of M for all $p \in A$. Since A is a regular closed subset of M , $p \in \text{cl}(\text{int}(A)) = A$, and thus we also have $B_r(p) \cap \text{int}(A) \neq \emptyset$. Since $B_r(p) \cap A$ contains a non-empty open set $B_r(p) \cap \text{int}(A)$ for any $p \in A$, we therefore surmise that $\nu(B_r(p) \cap A) > 0$. As $\nu(B_r(p) \cap A)$ is continuous with respect to p and positive (Lemma 6.1), and A is compact, its infimum $K(r)$ over $p \in A$ is positive. \square

As regular intersections are regular closed sets (Lemma 4.2), Lemma 5.4 implies that a finite sampling bound on the type in Eq. (35) can be obtained if $K(r)$ is bounded below away from zero. We focus on the case where the regular intersection can be written as an intersection of two function sublevel sets for simplicity of analysis. Furthermore, our index pairs constructed above

$$\mathcal{N} = h^{-1}[\beta, \infty) \cap f^{-1}(-\infty, \alpha] \text{ and } \mathcal{N}_- = f^{-1}(-\infty, \alpha] \cap h^{-1}[\beta, \gamma], \quad (\text{Eq. (6)})$$

fall under this category: \mathcal{N}_- is the intersection of $f^{-1}(-\infty, \alpha]$ with the sublevel set $\tilde{h}^{-1}(-\infty, 0]$ of $\tilde{h} = (h - \beta)(h - \gamma)$. The remainder of this section is devoted to proving the following result.

Theorem 5.5 *Let $E = f_1^{-1}(-\infty, 0] \cap f_2^{-1}(-\infty, 0]$ be a regular intersection on M and $f_1^{-1}(0) \cap f_2^{-1}(0) \neq \emptyset$. Let \mathbb{X} be a finite set of i.i.d. draws according to the Riemannian density μ_g on M , restricted to E . Then for $\epsilon < \rho_2$, if*

$$\#\mathbb{X} \geq \frac{1}{K(\frac{\epsilon}{2})} \left(\log \left(\frac{\nu(E)}{K(\frac{\epsilon}{4})} \right) + \log \left(\frac{1}{\kappa} \right) \right). \quad (37)$$

then \mathbb{X} is ϵ -dense in A with probability $> 1 - \kappa$, where

$$K(r) = \inf_{p \in A} \mu_g(A \cap B_r(p)) \geq \mathbf{V}_{\text{HS2}}^{(m)} \left(\frac{\bar{\eta}_E(r)}{2} \cos(\theta \left(\frac{\bar{\eta}_E(r)}{2} \right)), \rho, \phi_{12} \right) > 0, \quad (38)$$

and ϕ_{12} , $\mathbf{V}_{\text{HS2}}^{(m)}(\cdot, \cdot, \cdot)$, and $\bar{\eta}_E$ are as defined in Lemma 5.14, Notation 5.6, and Proposition 5.12 respectively.

This bound captures two aspects of how the volume of a Euclidean ball intersected with M can be diminished when we restrict to E . One aspect is the local geometry near the ‘cusps’ of the intersection $f_1^{-1}(0) \cap f_2^{-1}(0)$, and this is parametrised by an angle parameter ϕ_{12} . This angle, which we formally define in Lemma 5.14, is the smallest angle subtended by ∇f_1 and ∇f_2 in $T_p M$ for $p \in f_1^{-1}(0) \cap f_2^{-1}(0)$, and $\mathbf{V}_{\text{HS2}}^{(m)}(\cdot, \cdot, \phi_{12})$ is a strictly decreasing function of ϕ_{12} which is non-zero for $\phi_{12} < \pi$. The closer ∇f_1 and ∇f_2 are to aligning oppositely, the thinner the intersection is near the cusps, and the smaller the volume.

The other pertinent aspect of the geometry of E is the thickness of E away from the cusps. This is parametrised by $\bar{\eta}_E(r)$, which takes into account the curvature effects of the individual level sets $f_i^{-1}(0) \cap \partial E$ in the boundary, as well as the thickness at which the two individual level sets approach each other. The interaction of the two level sets is parametrised by what we call a bottleneck thickness, which we define in Definition 5.11, and illustrate in

Fig. 5. We provide a lower bound on $\bar{\eta}_E(r)$ in terms of properties of the constituent functions f_1 and f_2 in Proposition 5.12.

Notation 5.6 *Our volume bounds will be phrased in terms of the volumes of the following geometric quantities and objects:*

- (1) For a manifold with reach τ_M , let $\theta(r) = \arcsin(\frac{r}{2\tau_M})$;
- (2) \mathcal{B}_r denotes a ball of radius r in T_pM , centered at the origin of T_pM which is identified with p in the ambient Euclidean space. We denote the volume of \mathcal{B}_r as

$$V_B^{(m)}(r) = \text{vol}(\mathcal{B}_r). \tag{39}$$

- (3) For $v \in T_pM$, we let $\mathfrak{B}(v)$ denote the ball in T_pM centred at v with radius $\|v\|$ (so that v is a radial vector of the ball). We let

$$V_{HS}^{(m)}(r, s, \cdot) = \text{vol}(\mathfrak{B}(su) \cap \mathcal{B}_r) \tag{40}$$

be the volume of the hyperspherical cap that arises from the intersection on the right and side, for any unit vector $u \in \mathbb{S}_pM$. If t is another unit vector in \mathbb{S}_pM , and $\angle(u, t) = \varphi > 0$, then we let

$$V_{HS2}^{(m)}(r, s, \varphi) := \text{vol}(\mathfrak{B}(su) \cap \mathfrak{B}(st) \cap \mathcal{B}_r). \tag{41}$$

We give an illustration of $\mathfrak{B}(su) \cap \mathfrak{B}(st) \cap \mathcal{B}_r$ in the two dimensional case in Fig. 4.

5.2 Review: the Case of Manifolds and Regular Domains

In [23], they bounded the volumes and probability measure of $A \cap B_r(p)$ (where in their case $A = M$) by considering the volume of its image under the orthogonal projection $\zeta_p : M \rightarrow T_pM$ onto the hyperplane T_pM . We proceed with the same argument and begin by stating the following two lemmas that [23] implicitly relies on. These lemmas in turn rely on the result in Proposition 4.13 which gives a lower bound on the maximum radius r such that ζ_p is a diffeomorphism onto its image when restricted to $B_r(p) \cap M$.

We first recall if we have a smooth map $F : M \rightarrow N$ between two Riemannian manifolds of the same dimensions, then we can pull back a density ν on N to obtain a density $F^*\nu$ on M . Locally, if $\nu = u|dy_1 \wedge \dots \wedge dy_m|$ on some local coordinate patch $(V, (y_i))$ of $F(p)$, then we can locally express the pullback $F^*\nu$ as

$$F^*\nu(p) = F^*(u|dy_1 \wedge \dots \wedge dy_m|)(p) = |\det(d_p F)| \cdot u(F(p)) \cdot |dx_1 \wedge \dots \wedge dx_m|. \tag{42}$$

where $(U, (x_i))$ is a local coordinate patch of p in M , and dF is the Jacobian matrix of F with respect to coordinates (x_i) and (y_i) on U and N respectively. The pullback satisfies an important property: if F is a diffeomorphism, then one can show that

$$\int_M F^*\nu = \int_N \nu. \tag{43}$$

Lemma 5.7 *Suppose $U \subseteq B_r(p) \cap M$ where $r \leq \tau_M$. Then*

$$\text{vol}(U) \geq \text{vol}(\zeta_p(U)). \tag{44}$$

where $\text{vol}(U) = \mu_g(U)$, and $\text{vol}(\zeta_p(U))$ is the m -dimensional volume of $\zeta_p(U)$ in the tangent plane.




Fig. 4 An illustration of the volume $\mathfrak{B}(su) \cap \mathfrak{B}(st) \cap \mathfrak{B}_r$ (as defined in Notation 5.6) in the plane containing vectors s and t in $T_p M$. The shaded region represents the intersection of the three balls, where the ball with solid boundary represents \mathfrak{B}_r , and the balls with dashed boundaries represent $\mathfrak{B}(su)$ and $\mathfrak{B}(st)$ respectively

Proof Let ν be the Riemannian density of the tangent plane. Recall ζ_p restricted to U is a diffeomorphism onto its image and let $F = \zeta_p^{-1}$ be its inverse. Then

$$\text{vol}(U) = \int_U \mu_g = \int_{\zeta_p(U)} F^* \mu_g.$$

Let y_1, \dots, y_m be a set of coordinates on the tangent plane that are orthonormal with respect to the ambient Euclidean metric restricted to the tangent plane, and consider the density at $z = \zeta_p(q) \in \zeta_p(U)$. If x_1, \dots, x_m is an orthonormal set of coordinates at $q \in T_q M$, then we can explicitly write the density $F^* \mu_g$ as

$$\begin{aligned} F^* \mu_g(z) &= F^* |dx_1 \wedge \dots \wedge dx_m|(z) \\ &= |\det dF| |dy_1 \wedge \dots \wedge dy_m| \\ &= \frac{1}{|\det d\zeta_p|} |dy_1 \wedge \dots \wedge dy_m| \\ &\geq |dy_1 \wedge \dots \wedge dy_m|. \end{aligned}$$

where the inequality in the final line is due to $|\det d\zeta_p| \leq 1$ (Lemma 6.2). Since $|dy_1 \wedge \dots \wedge dy_m|$ is the Riemannian density on the tangent plane at $z = \zeta_p(q)$,

$$\text{vol}(U) = \int_{\zeta_p(U)} F^* \mu_g \geq \text{vol}(\zeta_p(U)).$$

□

We now derive a lower bound on $\text{vol}(B_\epsilon(p))$ by describing the image of such balls in T_pM under the projection ζ_p . We first recall such descriptions for manifolds with boundary, where [24] differentiates into two cases where $p \in \partial M$ or $p \in \text{int}(M)$.

Lemma 5.8 [[23, Lemma 5.3] and [24, Lemma 4.7]] *Let M be a properly embedded manifold with positive reach. Recall Notation 5.6, we have for any $p \in M$ and $r < \tau_M$,*

$$\zeta_p(B_r(p) \cap M) \supset \mathcal{B}_{r \cos(\theta(r))} \tag{45}$$

Consequently, $\text{vol}(B_r(p) \cap M) \geq \mathbf{V}_B^{(m)}(r \cos(\theta(r)))$.

Lemma 5.9 [[24, Lemma 4.6]] *Let $M \subset \mathbb{R}^d$ be a properly embedded submanifold with boundary. For $p \in \partial M$, let $n \in T_pM \cap N_p\partial M$ be the inward pointing unit normal at p . If ζ_p is a diffeomorphism when restricted to $B_\delta(p) \cap M$, where $\delta < \min(\tau_M, \tau_{\partial M})$, then for $\epsilon \in (0, \delta)$,*

$$\zeta_p(B_\epsilon(p) \cap M) \supset \mathcal{B}_{\epsilon \cos(\theta(\epsilon))} \cap \mathfrak{B}(\epsilon n)$$

where we recall notation from Notation 5.6. Consequently, $\text{vol}(B_\epsilon(p) \cap M) \geq \mathbf{V}_{HS}^{(m)}(\epsilon \cos(\theta(\epsilon)), \epsilon)$.

Applying Proposition 4.13 to deduce the minimal radius δ in Lemma 5.9, we obtain the following result for regular domains.

Lemma 5.10 *Consider $f^{-1}(-\infty, 0]$, where $f : M \rightarrow \mathbb{R}$ is a smooth function on a submanifold M of \mathbb{R}^d with positive reach τ_M and f satisfies Items (R1) and (R2). Suppose for any $p \in f^{-1}(0)$, we have $\|\nabla_p f\| \geq \mu > 0$. Let*

$$\frac{1}{\rho} = \frac{1}{\tau_M} + \frac{\Lambda}{\mu}. \tag{46}$$

Recall Notation 5.6. If $\epsilon < \rho$, then

(i) For $p \in f^{-1}(0)$, and $n = -\frac{\nabla_p f}{\|\nabla_p f\|}$,

$$\zeta_p(B_\epsilon(p) \cap f^{-1}(-\infty, 0]) \supset \mathcal{B}_{\epsilon \cos(\theta(\epsilon))} \cap \mathfrak{B}(\rho n),$$

(ii) Consequently, for any $q \in f^{-1}(-\infty, 0]$,

$$\text{vol}(B_\epsilon(q) \cap E) \geq \mathbf{V}_{HS}^{(m)}\left(\frac{\epsilon}{2} \cos(\theta\left(\frac{\epsilon}{2}\right)), \rho\right) \tag{47}$$

is a function of ϵ, ρ, τ_M , and the dimension of the manifold m .

Proof (i) Because $\rho < \tau_M$, Proposition 4.13 implies ζ_p is a diffeomorphism on $B_\rho(p) \cap M$; thus, combined with the fact that $\rho < \min(\tau_{f_i^{-1}(0)}, \tau_{f_i^{-1}(-\infty, 0]})$ (Lemma 4.4), we can apply Lemma 5.9 and deduce the inclusion as stated.

(ii) The proof adopts the proof of [24, Lemma 4.6] for regular domains being a special case of submanifolds with boundary, which breaks down the analysis into two cases, whether $d(p, \partial E) > \frac{\epsilon}{2}$ or otherwise. In the first case, we can trivially bound $\text{vol}(B_\epsilon(p) \cap E)$ from below by $\text{vol}(B_{\epsilon/2}(p) \cap E)$. Since $B_{\epsilon/2}(p) \cap \partial E = \emptyset$, we have $B_{\epsilon/2}(p) \cap E = B_{\epsilon/2}(p) \cap M$. The volume of the latter is bounded below by $V_B^{(m)}(\frac{\epsilon}{2} \cos(\frac{\epsilon}{2}))$ (by Lemma 5.8), which is in turn bounded below by $V_{HS}^{(m)}(\frac{\epsilon}{2} \cos(\theta(\frac{\epsilon}{2})), \rho)$ by definition (see Notation 5.6). For the case where $d(p, \partial E) \leq \frac{\epsilon}{2}$, choose a nearest neighbor x of p in ∂E . As $B_{\epsilon/2}(x) \subset B_\epsilon(p)$, we can bound the volume of the former by the latter. The latter's volume is bounded by that of its projection (Lemma 5.7), and we arrive at the stated bound. □

5.3 Regular Intersections of Two Functions

We now consider a regular intersections of two function $E = f_1^{-1}(-\infty, 0] \cap f_2^{-1}(-\infty, 0]$. We bound the volume of $B_\epsilon(p) \cap E$ from below by dividing the set of points $p \in E$ into two cases. Let E_r denote the set of points

$$E_r = \{x \in E : d(x, f_1^{-1}(0) \cap f_2^{-1}(0)) \geq r\}. \tag{48}$$

Then either:

- (Case 1) $p \in E_\epsilon$, i.e., $B_\epsilon(p)$ may intersect both $f_1^{-1}(0)$ and $f_2^{-1}(0)$, but not $f_1^{-1}(0) \cap f_2^{-1}(0)$; else
- (Case 2) $p \notin E_\epsilon$, i.e., $B_\epsilon(p)$ intersects $f_1^{-1}(0) \cap f_2^{-1}(0)$.

In Item (Case 1), we require an additional lengthscale to control the geometry of E in the ϵ -ball about p , which we illustrate in Fig. 5.

Definition 5.11 Let E be a regular intersection where $E = f_1^{-1}(-\infty, 0] \cap f_2^{-1}(-\infty, 0]$. The r -bottleneck thickness $\eta_E(r)$ of E is

$$\eta_E(r) = \sup\{\epsilon \in (0, r) : \forall x \in E_r, B_\epsilon(x) \cap f_i^{-1}(0) \neq \emptyset \text{ for at most one } i \in (1, 2)\} \tag{49}$$




Fig. 5 An illustration of the bottleneck distance $\eta_E(r)$ of a regular intersection of two sublevel sets

Note that in the case where $f_1^{-1}(0) \cap f_2^{-1}(0) = \emptyset$ (i.e. $E = E_r$ for all r), we can write $\partial E = f_1^{-1}(0) \sqcup f_2^{-1}(0)$. The bottleneck thickness η_E is then the largest radius for which we can thicken ∂E such that the thickenings of $f_1^{-1}(0)$ and $f_2^{-1}(0)$ do not intersect. Thus η_E is a homological critical value in the thickening of ∂E , implying the bottleneck thickness η_E is an upper bound on the reach of the manifold ∂E in \mathbb{R}^d . Furthermore, writing $\partial E = F^{-1}(0)$ for $F = f_1 f_2$, one can check that dF is nowhere zero on ∂E and we can derive an explicit bound on the reach of ∂E using Proposition 4.14, thus in turn providing a lower bound on η_E .

In the more general case, we can also interpret η_E as thus. For any point x on E that are at least r away from $f_1^{-1}(0) \cap f_2^{-1}(0)$ on E , the bottleneck thickness $\eta_E(r)$ prescribes the largest radius such that for $t < \eta_E(r)$,

$$B_t(x) \cap E = B_t(x) \cap f_i^{-1}(-\infty, 0]$$

for i either 1 or 2. Thus, for such points, we can apply results such as Lemma 5.10 to bound the volume of $B_r(x) \cap M$. We now derive a positive lower bound on $\eta_E(r)$.

Proposition 5.12 *Let E be a regular intersection where $E = f_1^{-1}(-\infty, 0] \cap f_2^{-1}(-\infty, 0]$. Let τ_i denote the reach of the submanifold $f_i^{-1}(0)$ for $i \in (1, 2)$. Let $F = f_1 f_2 : M \rightarrow \mathbb{R}$. For $0 < r/2 < \min(\tau_1, \tau_2)$, let $\mu_F(r) = \inf_{x \in E_{r/2} \cap F^{-1}(0)} \|\nabla F\|$, and Λ_F be the supremum of the norm of the Hessian of F on M . Let*

$$\frac{1}{\rho_F(r)} = \frac{1}{\tau_M} + \frac{\Lambda_F}{\mu_F(r)}. \tag{50}$$

Then $\eta_E(r) \geq \bar{\eta}_E(r)$ where

$$\bar{\eta}_E(r) = \min\left(\frac{r}{2}, \rho_F(r)\right) > 0. \tag{51}$$

Proof We show that $\rho_F(r) > 0$ by showing $\mu_F(r) > 0$. For $x \in E_{r/2} \cap F^{-1}(0)$, since $E_{r/2}$ excludes $f_1^{-1}(0) \cap f_2^{-1}(0)$, we observe that $f_1(x)$ and $f_2(x)$ cannot both be zero at x . Moreover, since E is a regular intersection, by Definition 4.1 we have $df_i \neq 0$. Thus, for such a point x , either

$$dF(x) = f_1 df_2 \neq 0 \quad \text{ord} F(x) = f_2 df_1 \neq 0.$$

Furthermore, one can check that $E_{r/2} \cap F^{-1}(0)$ is compact and thus the quantity

$$\mu_F(r) = \inf_{x \in E_{r/2} \cap F^{-1}(0)} \|\nabla F\|$$

is positive.

Consider then the Euclidean ball $B_\epsilon(p)$ where $p \in E_r$ and $\epsilon < \frac{r}{2}$. Since we have restricted $p \in E_r$, the ball $B_\epsilon(p)$ cannot intersect $f_1^{-1}(0) \cap f_2^{-1}(0)$. Assume

$$B_\epsilon(p) \cap f_1^{-1}(0) \neq \emptyset, \text{ and } B_\epsilon(p) \cap f_2^{-1}(0) \neq \emptyset.$$

Since $\epsilon < r/2 < \tau_i$ by assumption, p has unique projections onto $f_1^{-1}(0)$ and $f_2^{-1}(0)$ respectively (Theorem 4.5); let these projections be p_i and note that $\|p - p_i\| < r/2$. Note that $f_1(p_1) = f_2(p_2) = 0$, but $f_1(p_2), f_2(p_1) \neq 0$. As $p \in E_r$, and $\|p - p_i\| < r/2$, we see that $p_i \in E_{r/2}$.

Since $B_\epsilon(p)$ cannot intersect $f_1^{-1}(0) \cap f_2^{-1}(0)$, we can suppose without loss of generality that $p \neq p_1$ and $\|p - p_1\| \geq \|p - p_2\|$. Since p_1 is a nearest neighbour of p in

$f_1^{-1}(0)$, Item Proposition 3.14 (i) implies

$$p - p_1 = n + \lambda \nabla f_1(p_1) = n + \frac{\lambda}{f_2(p_1)} \nabla F(p_1)$$

for some $n \in N_{p_1}M$. Because $p_1 \in B_\epsilon(p)$ and $B_\epsilon(p) \cap f_1^{-1}(0) \cap f_2^{-1}(0) = \emptyset$, we observe that $f_2(p_1) \neq 0$, and $\nabla F(p_1) = f_2(p_1) \nabla f_1(p_1)$. Thus, we can write

$$p - p_1 = n + \frac{\lambda}{f_2(p_1)} \nabla F(p_1).$$

In other words, the unit vector of $p - p_1$ lies in the normal space of $F^{-1}(0)$ at p . If $\epsilon \leq \rho_F(r)$, then we have $\|p - p_1\| < \rho_F(r)$. Item Proposition 3.14 (ii) then implies p_1 is the *unique* nearest neighbour of p in $F^{-1}(0)$. However, this contradicts our assumption that $\|p - p_1\| \geq \|p - p_2\|$, as $p_2 \in F^{-1}(0)$ too. Therefore, $\epsilon > \rho_F(r)$.

Put in other words, either $\eta_E(r) \geq \frac{r}{2}$; else, $\eta_E(r) \geq \rho_F(r)$. We conclude that

$$\eta_E(r) \geq \min\left(\frac{r}{2}, \rho_F(r)\right).$$

□

Having obtained a lower bound for $\eta_E(r)$, we can bound the volume $B_r(p) \cap E$ for $p \in E_r$.

Lemma 5.13 Consider $E = f_1^{-1}(-\infty, 0] \cap f_2^{-1}(-\infty, 0]$. Suppose f_1 and f_2 satisfy the conditions placed on f in Lemma 5.10, and ρ be as defined in Lemma 5.10. Let $\bar{\eta}_E(\epsilon)$ is as defined in Proposition 5.12. Then for $p \in E_\epsilon$,

$$\text{vol}(B_\epsilon(p) \cap E) \geq \mathbf{V}_{\text{HS}}^{(m)}\left(\frac{\bar{\eta}_E(\epsilon)}{2} \cos\left(\theta\left(\frac{\bar{\eta}_E(\epsilon)}{2}\right)\right), \rho\right).$$

Proof By definition, $\bar{\eta}_E(\epsilon) \leq \epsilon$ (Proposition 5.12). As $\epsilon < \rho$, and ρ is a lower bound on the reaches τ_i of $f_i^{-1}(0)$ by Lemma 4.4, we have $\bar{\eta}_E(\epsilon) < \tau_i$. And the conditions of Proposition 5.12 are satisfied so that $\bar{\eta}_E(\epsilon)$ is a positive lower bound on $\eta_E(\epsilon)$. Thus, for $p \in E_\epsilon$, the Euclidean ball $B_{\bar{\eta}_E(\epsilon)}(p)$ can only intersect at most one level set $f_i^{-1}(0)$. In other words, we can write without loss of generality that

$$B_\epsilon(p) \cap E \supset B_{\bar{\eta}_E(\epsilon)}(p) \cap E = B_{\bar{\eta}_E(\epsilon)}(p) \cap f_1^{-1}(-\infty, 0]$$

Since f_1 satisfies the conditions placed on f in Lemma 5.10, we apply the volume bound in Lemma 5.10 to deduce

$$\begin{aligned} \text{vol}(B_\epsilon(p) \cap E) &\geq \text{vol}(B_{\bar{\eta}_E(\epsilon)}(p) \cap f_1^{-1}(-\infty, 0]) \\ &\geq \mathbf{V}_{\text{HS}}^{(m)}\left(\frac{\bar{\eta}_E(\epsilon)}{2} \cos\left(\theta\left(\frac{\bar{\eta}_E(\epsilon)}{2}\right)\right), \rho\right). \end{aligned}$$

□

Lemma 5.14 Consider $E = f_1^{-1}(-\infty, 0] \cap f_2^{-1}(-\infty, 0]$, and suppose f_1 and f_2 satisfy the conditions placed on f in Lemma 5.10, and ρ be as defined in Lemma 5.10. Then, if $\epsilon < \rho$, and $p \in f_1^{-1}(0) \cap f_2^{-1}(0)$, then

$$\begin{aligned} \text{vol}(B_\epsilon(p) \cap E) &\geq \text{vol}(B_{\epsilon \cos(\theta(\epsilon))}(p) \cap \mathfrak{B}(\rho n_1) \cap \mathfrak{B}(\rho n_2)) \\ &=: \mathbf{V}_{\text{HS}^2}^{(m)}(\epsilon \cos(\theta(\epsilon)), \rho, \phi_{12}) > 0 \end{aligned}$$

where $\cos(\phi_{12}) = \inf_{x \in f_1^{-1}(0) \cap f_2^{-1}(0)} |\langle n_1(x), n_2(x) \rangle| > -1$ for $n_i(x) = \frac{\nabla_x f_i}{\|\nabla_x f_i\|}$.

Proof Given f_i satisfy the conditions placed on f in Lemma 5.10, and ρ be as defined in Lemma 5.10,

$$\zeta_p (B_\epsilon (p) \cap f_i^{-1}(-\infty, 0]) \supset \mathcal{B}_{\epsilon \cos \theta(\epsilon)} \cap \mathfrak{B}(\rho n_i)$$

Because $\epsilon < \rho$, the projection ζ_p is a diffeomorphism on $B_\epsilon (p) \cap M$, and thus

$$\begin{aligned} \zeta_p (B_\epsilon (p) \cap E) &= \zeta_p (B_\epsilon (p) \cap f_1^{-1}(-\infty, 0]) \cap \zeta_p (B_\epsilon (p) \cap f_2^{-1}(-\infty, 0]) \\ &\supset \mathcal{B}_{\epsilon \cos \theta(r)} \cap \mathfrak{B}(\rho n_1) \cap \mathfrak{B}(\rho n_2). \end{aligned}$$

As E is a regular intersection, n_1 and n_2 are linearly independent (Item (R1)); and $\varphi_{12} > 0$ as the supremum is taken over a compact set by assumption that E is a regular intersection (Definition 4.1). We now show that this implies $\mathfrak{B}(\rho n_1) \cap \mathfrak{B}(\rho n_2) \neq \emptyset$. We make two observations, first, that $\mathfrak{B}(\rho n_1)$ and $\mathfrak{B}(\rho n_2)$ are tangent to the origin; and second, the centres ρn_1 and ρn_2 of $\mathfrak{B}(\rho n_1)$ and $\mathfrak{B}(\rho n_2)$ respectively are not collinear with the origin as n_1 and n_2 are linearly independent. Thus $\mathfrak{B}(\rho n_1) \cap \mathfrak{B}(\rho n_2) \neq \emptyset$.

Since p is in the closure of $\mathfrak{B}(\rho n_1) \cap \mathfrak{B}(\rho n_2)$, for $\epsilon > 0$,

$$B_{\epsilon \cos(\theta(\epsilon))} (p) \cap \mathfrak{B}(\rho n_1) \cap \mathfrak{B}(\rho n_2) \neq \emptyset.$$

Finally, since this subset is a non-empty intersection of open sets, it is also open, and it has positive volume. □

Lemma 5.15 Consider $E = f_1^{-1}(-\infty, 0] \cap f_2^{-1}(-\infty, 0]$, and suppose f_1 and f_2 satisfy the conditions placed on f in Lemma 5.10, and ρ be as defined in Lemma 5.10. Then, if $\epsilon < \rho$, and $p \in E$, then

$$\text{vol} (B_\epsilon (p) \cap E) \geq \mathbf{V}_{\text{HS2}}^{(m)} \left(\frac{\bar{\eta}_E(\epsilon)}{2} \cos(\theta \left(\frac{\bar{\eta}_E(\epsilon)}{2} \right)), \rho, \phi_{12} \right) > 0$$

where ϕ_{12} , $\mathbf{V}_{\text{HS2}}^{(m)}(\cdot, \cdot, \cdot)$, and $\bar{\eta}_E$ are as defined in Lemma 5.14, Notation 5.6, and Proposition 5.12 respectively.

Proof Combining Lemma 5.13 and Lemma 5.14, we have, for $p \in E$ and $\epsilon < \rho$,

$$\text{vol} (B_\epsilon (p) \cap E) \geq \begin{cases} \mathbf{V}_{\text{HS2}}^{(m)} (\epsilon \cos(\epsilon), \rho, \phi_{12}) & \text{if } p \notin E_\epsilon \\ \mathbf{V}_{\text{HS}}^{(m)} \left(\frac{\bar{\eta}_E(\epsilon)}{2} \cos(\theta \left(\frac{\bar{\eta}_E(\epsilon)}{2} \right)), \rho \right) & \text{otherwise.} \end{cases}$$

Since $\epsilon > \bar{\eta}_E(\epsilon)$ (Proposition 5.12), and for any $t > 0$,

$$\begin{aligned} \mathbf{V}_{\text{HS2}}^{(m)} (t, \rho, \phi_{12}) &= \text{vol} (\mathcal{B}_{t \cos(\theta(t))} \cap \mathfrak{B}(\rho n_1) \cap \mathfrak{B}(\rho n_2)) \\ &\leq \text{vol} (\mathcal{B}_{t \cos(\theta(t))} \cap \mathfrak{B}(\rho n_1)) \\ &= \mathbf{V}_{\text{HS}}^{(m)} (t \cos(\theta(t)), \rho), \end{aligned}$$

our bound holds for either $p \in E_\epsilon$ or otherwise. □

6 Technical Lemmas

Lemma 6.1 *Let A be measurable subset of a Riemannian manifold $M \subset \mathbb{R}^d$ with $\mu_g(A) > 0$, where μ_g is the Riemannian density on M . If ν is a measure supported on M that is continuous with respect to μ_g , then for any radius $r > 0$ the function $p \mapsto \nu(B_r(p)) \cap A$ is continuous on A .*

Proof Consider c such that $\|c - p\| = \delta < r$. Then

$$\begin{aligned} B_{r-\delta}(p) &\subset B_r(c) \subset B_{r+\delta}(p) \\ \implies \nu(B_{r-\delta}(p) \cap A) &\leq \nu(B_r(c) \cap A) \leq \nu(B_{r+\delta}(p) \cap A). \end{aligned}$$

As $\nu(B_{r+\delta}(p) \cap A)$ monotonically increases with δ , as we decrease δ , the monotonicity and continuity of the measure with respect to δ ensures that $\nu(B_{r\pm\delta}(p) \cap A) \xrightarrow{\delta \rightarrow 0} \nu(B_r(p) \cap A)$. Thus, by the sandwich theorem,

$$\lim_{c \rightarrow p} \nu(B_r(c) \cap A) = \nu(B_r(p) \cap A).$$

Thus, $\nu(B_r(p) \cap A)$ is continuous with respect to p . \square

For $M \subset \mathbb{R}^d$ Recall that $\zeta_p : M \rightarrow T_p M$ is the orthogonal projection onto the m -dimensional hyperplane tangent to M at p .

Lemma 6.2 *Let $d_q \zeta_p$ be the Jacobian of ζ_p at $q \in M$ with respect to orthonormal coordinates in $T_q M$ and $T_p M$. Then $|\det d_p \zeta_p| \leq 1$.*

Proof Let $P : \mathbb{R}^d \rightarrow \mathbb{R}^m$ be the orthogonal projection onto an m -dimensional hyperplane in \mathbb{R}^d . As ζ_p is the restriction of P to M , therefore $d\zeta_p : \mathcal{T}(M) \rightarrow \mathbb{R}^m$ is the restriction of $P : \mathcal{T}(\mathbb{R}^d) \rightarrow \mathbb{R}^m$ to $\mathcal{T}(M)$. Choosing an orthonormal set of coordinates (x_i) in \mathbb{R}^d such that $P(x_1, \dots, x_d) = (x_1, \dots, x_m)$, we can write $dP = 1_m \oplus 0_{m,d-m}$ where 1_m is the identity matrix corresponding to the first m coordinates, and $0_{m,d-m}$ is the $m \times (d-m)$ matrix of zeros. Since $d\zeta_p$ is a restriction of dP to an m -dimensional subspace $T_p M \subset T_p \mathbb{R}^d$, the absolute value of the determinant of $d_p \zeta_p$ is at most 1. \square

Acknowledgements We are grateful to Frances Kirwan and Konstantin Mischaikow for insightful discussions. KMY would like to thank Pete Grindrod CBE for his guidance and mentorship, as well as Primoz Skraba and Gesine Reinert for their thoughtful feedback on this work. VN is supported by the EPSRC Grant EP/R018472/1. KMY is supported by a UKRI Future Leaders Fellowship (MR/W01176X/1; PI: J Harvey), and the EPSRC Centre For Doctoral Training in Industrially Focused Mathematical Modelling (EP/L015803/1). We thank the two reviewers for their helpful comments and suggestions which improved this paper.

Author Contributions This paper contains results of the first author's PhD research, which was conducted under the guidance of the second author.

Declarations

Conflict of interest The authors declare no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Banyaga, A., Hurtubise, D.: Morse-Bott homology. *Trans. AMS* **362**, 3997–4043 (2010)
2. Boissonnat, J.-D., Lieutier, A., Winttraecken, M.: The reach, metric distortion, geodesic convexity and the variation of tangent spaces. *J. Appl. Comput. Topol.* **3**(1–2), 29–58 (2019)
3. Chang, K.C., Ghoussoub, N.: The Conley index and the critical groups via an extension of Gromoll-Meyer theory. *Topol. Methods Nonlinear Anal.* **7**(1), 77 (1996)
4. Chazal, F., Cohen-Steiner, D., Lieutier, A.: A sampling theory for compact sets in Euclidean space. *Discr. Comput. Geome.* **41**(3), 461–479 (2009)
5. Conley, C.: Isolated invariant sets and the Morse index: expository lectures. In: Number 38 in Regional conference series in mathematics. American Mathematical Society, Providence, RI, (1978)
6. Conley, C.C., Zehnder, E.: The Birkhoff-Lewis fixed point theorem and a conjecture of V I Arnold. *Invent. Math.* **73**(1), 33–49 (1983)
7. Federer, H.: Curvature measures. *Trans. Am. Math. Soc.* **93**(3), 418 (1959)
8. Ferry, S., Mischaikow, K., Nanda, V.: Reconstructing functions from random samples. *J. Comput. Dyn.* **1**, 2 (2014)
9. Gardner, R., Smoller, J.: The existence of periodic travelling waves for singularly perturbed predator-prey equations via the Conley index. *J. Diff. Equ.* **47**, 133–161 (1983)
10. Gedeon, T., Kokubu, H., Oka, H., Reineck, J.F.: The Conley index for fast-slow systems. I One-dimensional slow variable. *J. Dyn. Diff. Equ.* **11**, 427–470 (1999)
11. Goresky, M., MacPherson, R.: *Stratified Morse Theory*. Springer-Verlag, (1988)
12. Goresky, M., MacPherson, R.: Local contribution to the Lefschetz fixed point formula. *Invent. Math.* **111**, 1–33 (1993)
13. Gromoll, D., Meyer, W.: On differentiable functions with isolated critical points. *Topology* **8**(4), 361–369 (1969)
14. Harker, S., Mischaikow, K., Mrozek, M., Nanda, V.: Discrete Morse theoretic algorithms for computing homology of complexes and maps. *Found. Comput. Math.* **14**, 151–184 (2014)
15. Kaczynski, T., Mischaikow, K., Mrozek, M.: *Computational homology*. In: *Applied Mathematical Sciences*, vol. 157. Springer, Berlin (2004)
16. Kirwan, F., Penington, G.: Morse theory without nondegeneracy. *Quart. J. Math.* **72**(1–2), 455–514 (2021)
17. Lee, J.M.: *Introduction to smooth manifolds*. In: *Graduate Texts in Mathematics*, vol. 218. Springer, New York, (2012)
18. Manolescu, C.: The Conley index, gauge theory, and triangulations. *J. Fixed Pt Theory Appl.* **13**(2), 431–457 (2013)
19. Milnor, J.: *Morse theory*. Princeton University Press, (1973)
20. Milnor, J.: *Topology from the differentiable viewpoint*. Princeton University Press, (1997)
21. Mischaikow, K., Mrozek, M.: Chaos in the Lorenz equations: a computer-assisted proof. *Bullet. Am. Math. Soc.* **32**, 66–72 (1995)
22. Mischaikow, K., Mrozek, M.: Conley index. *Handb. Dyn. Syst.* **2**, 393–460 (2002)
23. Niyogi, P., Smale, S., Weinberger, S.: Finding the homology of submanifolds with high confidence from random samples. *Discr. Comput. Geome.* **39**(1), 419–441 (2008)
24. Wang, Y., Wang, B.: Topological inference of manifolds with boundary. *Comput. Geome.* **88**, 101606 (2020)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.