



<b>Title</b>	Flavanoid Epoxides - XIII: Acid and Base Catalysed Reactions of 2'-Tostyloxychalcone epoxides. Mechanism of the Algar-Flynn-Oyamada Reaction
<b>Authors(s)</b>	Gormley, T. R. (Thomas Ronan), O'Sullivan, W.I.
<b>Publication date</b>	1973
<b>Publication information</b>	Gormley, T. R. (Thomas Ronan), and W.I. O'Sullivan. "Flavanoid Epoxides - XIII: Acid and Base Catalysed Reactions of 2'-Tostyloxychalcone Epoxides. Mechanism of the Algar-Flynn-Oyamada Reaction" 29, no. 2 (1973).
<b>Publisher</b>	Elsevier
<b>Item record/more information</b>	<a href="http://hdl.handle.net/10197/6996">http://hdl.handle.net/10197/6996</a>
<b>Publisher's version (DOI)</b>	10.1016/S0040-4020(01)93304-6

Downloaded 2023-10-05T14:16:07Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd\_oa)



© Some rights reserved. For more information

## FLAVANOID EPOXIDES—XIII<sup>1</sup>

### ACID AND BASE CATALYSED REACTIONS OF 2'-TOSYLOXYCHALCONE EPOXIDES. MECHANISM OF THE ALGAR-FLYNN-OYAMADA REACTION

T. R. GORMLEY and W. I. O'SULLIVAN\*

Department of Chemistry, University College, Dublin 4, Ireland

(Received in the UK 28 August 1972; Accepted for publication 29 September 1972)

**Abstract**—2'-Tosyloxychalcone epoxide (6a) on reaction with alkali gave flavonol (4a) while similar treatment of 6'-methoxy-2'-tosyloxychalcone epoxide (6b), both at room temperature and at the boiling point of the reaction medium, afforded 5-methoxyaurone (5b). The latter result indicates that an epoxide is not an intermediate in the production of flavanols from 2'-hydroxy-6'-methoxychalcone epoxides on treatment with alkaline hydrogen peroxide (AFO Reaction) at the higher temperature. Epoxide 6a on treatment with boron trifluoride etherate gave a mixture of flavanone and flavonol while epoxide 6b gave formyldeoxybenzoin (9) under similar conditions.

The Algar-Flynn-Oyamada<sup>2</sup> (AFO) reaction involves the alkaline hydrogen peroxide oxidation of 2'-hydroxychalcones firstly into flavanonols<sup>3</sup> (3) and then into flavonols (4) (Scheme I, route 1). If an OMe<sup>4</sup> or Me<sup>5</sup> substituent is present in the 6'-position of the chalcone, aurones (5) (Scheme I, route 2) rather than flavonols are obtained, provided the 2-<sup>6</sup> or 4-<sup>7</sup> positions do not carry an hydroxyl group, and that the reaction is effected at room temperature. When the reaction is carried out in MeOH or EtOH 6'-methoxychalcones give flavonols as the main products.<sup>8</sup> Dean and Podimuang<sup>9</sup> proposed routes 3 or 4 (Scheme I) for the formation of flavanonols and flavonols from 2'-hydroxychalcones in the AFO reaction rather than the previously accepted route which presumed the intermediate formation of a chalcone epoxide (route 1). The accepted routes (1 and 2) to 5-methoxyflavonols (at temperatures above 20°) and to 4-methoxyaurones from 2'-hydroxy-6'-methoxychalcones on AFO oxidation were retained by these authors.

The results we now report on the reactions of 6'-methoxy-2'-tosyloxychalcone epoxide are consistent with, but do not prove, route 2 to aurone but they are not in accord with route 1 to flavonols from 2'-hydroxy-6'-methoxychalcones in the AFO reaction carried out at the boiling point of the reaction medium.

2'-Tosyloxychalcone epoxide (6a) and 6'-methoxy-2'-tosyloxychalcone epoxide (6b) were


prepared by the action of alkaline H<sub>2</sub>O<sub>2</sub> on the parent 2'-tosyloxychalcones. Treatment of epoxide 6a with alkali at room temperature gave flavonol 4a<sup>11</sup> while epoxide 6b on similar treatment gave 5-methoxyaurone (5b) (along with, in one instance, a small amount of 6-methoxyflavonol; 3b). When the latter reaction was carried out in EtOH at the b.p. aurone 5b was the only product.

Various suggestions have been presented as to why cyclisation of the putative AFO 6-methoxychalcone epoxide intermediate takes place by attack of the 2'-O<sup>-</sup> group at the  $\alpha$ -rather than at the  $\beta$ -position. One such suggestion is that displacement of the carbonyl group out of the plane of the phenolic ring by steric interaction with the 6'-methoxyl group increases the distance of the phenolic oxygen from the  $\beta$ -position more than from the  $\alpha$ -position. Another is that the said steric interaction favours the product with the smaller heterocyclic ring.<sup>12</sup> Dean and Podimuang<sup>9</sup> suggested that such effects are not large and that at temperatures above 20° they diminish rapidly and pyrone derivatives are again the main products. However, the results of the present work indicate that once an intermediate chalcone epoxide is formed in the AFO reaction the presence of a 6'-methoxyl substituent plays a practically decisive role, over a range of temperatures, in determining the position of attack by the 2'-O<sup>-</sup> group, i.e.  $\alpha$ -position (route 2), on cyclisation to form products. A more plausible explanation for flavonol formation on the oxidation of 2'-hydroxy-6'-methoxychalcones at temperatures greater than 20° is that the inhibition to resonance between the 2'-O<sup>-</sup> ion and the carbonyl group† caused by steric interaction between the 6'-methoxyl group and the ketone group, which facilitates epoxide formation, is overcome at higher temperatures and that the reaction then pro-

\*To whom all correspondence should be addressed.

†The epoxidation of  $\alpha,\beta$ -unsaturated ketones by the hydroperoxide ion is hindered because of the coulombic repulsion of the reagent as well as internal electronic inactivation as indicated in structure 7.<sup>9</sup>


SCHEME 1



ceeds by routes 3 or 4 without the formation of an intermediate epoxide.

Two further experiments gave results which are in agreement with these conclusions. When 2'-tosyloxy-2,4,4',6'-tetramethoxychalcone (8) in MeOH was allowed to react with alkaline  $H_2O_2$  at room temperature for 24 h the corresponding flavanone was obtained in low yield. This result suggests that hydrolysis of the tosylate group had taken place and that the partial deactivation of the  $\alpha,\beta$ -unsaturated ketone moiety by the 2'-O<sup>-</sup> ion

(which is not fully conjugated with the carbonyl group) and by the 2'- and 4'-methoxyl groups disfavoured epoxide formation and the reaction proceeded by the alternative routes 3 or 4. Had epoxidation occurred the corresponding aurone would have been the expected product. In another experiment 2-chloro-4',6'-dimethoxy-2'-tosyl-oxyacetophenone on condensation with anisaldehyde under alkaline conditions afforded 4,4',6-trimethoxyaurone.<sup>12</sup> It is probable that this reaction proceeded by a Darzens type condensation



to give an intermediate epoxide<sup>13</sup> which then cyclised to the aurone (Scheme 2). Attempts to isolate tosyloxychalcone epoxides in the latter two reactions were without success.

produce the  $\alpha$ -formyldeoxybenzoin **9**. The reaction of 4'-methoxyl substituted 2'-benzyloxychalcone epoxides with  $\text{BF}_3$  was reported<sup>15</sup> to give analogous results.

#### Reactions of 2'-tosyloxychalcone epoxides with boron trifluoride


The reaction of epoxide **6a** with boron trifluoride etherate under  $\text{N}_2$  afforded a mixture of flavanonol (**3a**; 70%) and flavonol (**4a**; 10%). When air was not excluded from the reaction the yields were 33% of **3a** and 37% of **4a**. The action of  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  on epoxide **6b** produced  $\alpha$ -formyldeoxybenzoin **9**, which on heating with alkali gave desoxybenzoin **10**.

Flavanonols are produced by the reaction of 2'-benzyloxychalcone epoxides with acids and their formation was formulated as follows:<sup>14</sup>

#### EXPERIMENTAL

**2'-Tosyloxychalcone.** A mixture of 2'-hydroxychalcone (2.5 g), *p*-TsCl (1.5 g),  $\text{K}_2\text{CO}_3$  (1.5 g) and dry acetone (15 ml) was heated under reflux for 2 h and added to water. The oil which separated was dissolved in ether and the resulting solution washed successively with  $\text{NaHCO}_3$  (5%), NaOH (10%) and water and then dried ( $\text{Na}_2\text{SO}_4$ ). The solid obtained on removal of solvent was recrystallised (MeOH) to give 2'-tosyloxychalcone (1.8 g), m.p. and m.m.p.  $84^\circ$  (IR spectrum identical with that of an authentic<sup>16</sup> sample).

**2'-Tosyloxy-6'-methoxychalcone** was prepared in the same way from 2'-hydroxy-6'-methoxychalcone in 61%




A similar mechanism may be operative in the cyclisation of epoxide **6a** by means of  $\text{BF}_3 \cdot \text{Et}_2\text{O}$ . The steric factors, suggested earlier, which hindered attack of the 2'-O<sup>-</sup> moiety at the  $\beta$ -position of 6'-methoxyl substituted chalcone epoxide in alkali would also, presumably, be operative in impeding attack of the 2'-tosyloxy group at the  $\beta$ -position in 6'-methoxy-2'-tosyloxychalcone epoxide on reaction with  $\text{BF}_3 \cdot \text{Et}_2\text{O}$ . An alternative pathway (Scheme 3) is followed with this compound involving migration of the aroyl group to

yield; white plates from MeOH, m.p.  $82^\circ$ ;  $\nu_{\text{max}}$  (KBr) 1639, 1592  $\text{cm}^{-1}$  (Found: C, 67.1; H, 4.9; S, 7.7; MeO, 7.8.  $\text{C}_{23}\text{H}_{20}\text{O}_5\text{S}$  requires: C, 67.6; H, 4.9; S, 7.6; MeO, 7.6%).

**2'-Tosyloxy-2,4,4',6'-tetramethoxychalcone (8)** was similarly prepared from 2'-hydroxy-2,4,4',6'-tetramethoxychalcone in 57% yield; white needles from MeOH, m.p.  $137\text{--}138^\circ$  (Found: C, 62.2; H, 5.3.  $\text{C}_{26}\text{H}_{26}\text{O}_8\text{S}$  requires: C, 62.3; H, 5.2%).

**Formation of 2'-tosyloxychalcone oxide 6a.** To a solution of 2'-tosyloxychalcone (4.5 g) in MeOH (60 ml) was added  $\text{H}_2\text{O}_2$  (12 ml; 30%) and NaOH (5 ml; 8%) at  $0^\circ$ . After 1 h the mixture was acidified with HCl (10%) and



separated crystals recrystallised (EtOH) to give needles of 2'-tosyloxychalcone oxide (3.6 g), m.p. 132°. (Found: C, 67.3; H, 4.7; S, 8.2.  $C_{22}H_{18}O_5S$  requires: C, 67.0; H, 4.6; S, 8.1%).  $\nu_{max}$  (KBr) 1690–1700  $cm^{-1}$ .  $\tau(CDCl_3; TMS)$ , 7.65 (s,  $-CH_3$ ), 5.88, 5.83 (d's,  $J$  1.7 Hz, epoxidic protons) 2.0–2.95 (m, aromatic protons). KI test positive.

**Formation of 2'-tosyloxy-6'-methoxychalcone oxide 6b.** To a suspension of 2'-tosyloxy-6'-methoxychalcone (1.8 g) in MeOH (50 ml) was added  $H_2O_2$  (6 ml; 30%) and NaOH (2.5 ml; 8%) and the mixture stirred at room temp for 2 h. A white solid precipitated which was collected and recrystallised from EtOH to give plates of 2'-tosyloxy-6'-methoxychalcone (1.8 g) in MeOH (50 ml) was added  $H_2O_2$  (6 ml; 30%) and NaOH (2.5 ml; 8%) and the mixture stirred at room temp for 2 h. A white solid precipitated which was collected and recrystallised from EtOH to give plates of 2'-tosyloxy-6'-methoxychalcone oxide (1.1 g), m.p. 120°.  $\nu_{max}$  (KBr) 1704  $cm^{-1}$ . KI test positive. (Found: C, 64.9; H, 4.9; S, 7.3; OMe, 6.7.  $C_{22}H_{20}O_6S$  requires: C, 65.1; H, 4.7; S, 7.5; OMe, 7.3%).

**Reaction of 2'-tosyloxy-2,4,4',6'-tetramethoxychalcone 8 with alkaline hydrogen peroxide.** To a suspension of 8 (2.3 g) in MeOH (35 ml) was added  $H_2O_2$  (6 ml; 30%) and NaOH (2.5 ml; 8%). After 24 h a white solid separated which recrystallised (MeOH) to give needles of 2',4',5,7-tetramethoxydihydroflavonol (0.7 g), m.p. 178–180°. (Found: C, 63.5; H, 5.6.  $C_{18}H_{20}O_7$  requires: C, 63.3; H, 5.66%).

**Reaction of 2'-tosyloxychalcone oxide (6a) with alkali.** To a solution of 6a (0.5 g) in MeOH (60 ml) was added NaOH aq. (8 ml; 8%) and the mixture stirred for 15 h at 17° and then acidified with HCl (10%). The oil which separated was dissolved in ether and the solution extracted successively with  $NaHCO_3$  (5%) and NaOH (10%). On removal of solvent a white compound was obtained which on recrystallisation (EtOH) gave unreacted 6a, m.p. and m.m.p. 131°. Acidification of the NaOH extract with HCl (10%) gave flavonol (4a) (0.1 g; 34%) m.p. 169°, identical (m.p. and IR) with an authentic sample.

Similar experiments were run with 6a except that reaction times, temperatures and concentrations of NaOH were altered. In each case, only flavonol 4a was obtained. The following conditions were used and the yields of 4a (%) obtained: 6a (0.7 g), NaOH (14 ml, 8%), 70°, 2 h, 4a (52%); 6a (1 g), NaOH (20 ml; 16%), 70°, 2 h, 4a (98%); 6a (1 g), NaOH (50 ml; 8%), 0°, 18 h, 4a (65%).

**Reaction of 2'-tosyloxy-6'-methoxychalcone oxide (6b) with alkali.** To a suspension of 6b (0.5 g) in MeOH (200 ml) was added NaOH (16 ml; 16%) and the mixture stirred for 24 h at room temp. The mixture was acidified with HCl (10%) and the MeOH removed. On chromatography (silica gel) the residual solid gave two compounds. The first recrystallised (EtOH aq) to give needles of 4-methoxyaurone (5b) (0.3 g), m.p. and m.m.p. 150°. The second on recrystallisation from EtOAc gave needles of 5-methoxydihydroflavonol (3b) (10 mg), m.p. 200° (decomp.); no depression on admixture with an authentic sample.<sup>18</sup>

Similar experiments were run with 6b except that reaction times, temperatures and concentrations of NaOH were altered. In each case only aurone 5b was obtained. The following conditions were used and the yield of aurone 5b (%) is given: 6b (0.35 g), NaOH (5 ml; 12%), 17°, 24 h, 5b (85%); 6b (0.5 g), NaOH (8 ml; 16%), 70°, 2 h, 5b (98%).

**Reaction of 2'-tosyloxychalcone oxide 6a with  $BF_3$ .**

To a solution of oxide 6a (0.5 g) in dry ether (25 ml) in an atmosphere of  $N_2$  was added  $BF_3 \cdot Et_2O$  (6 ml; 45%) and the mixture allowed react at 15° for 0.5 h and then added to ice. The resulting oil was dissolved in ether and the ethereal solution extracted successively with  $NaHCO_3$  (5%) and NaOH (5%). Evaporation of the dried ( $Na_2SO_4$ ) ethereal solution gave dihydroflavonol (3a) (0.24 g), m.p. and m.m.p. 181°. No material was isolated on acidification and ether extraction of the alkaline extracts.

The reaction was repeated at 35° in air. The neutral ether fraction afforded dihydroflavonol (3a) (0.12 g) and the NaOH extract on neutralisation with HCl (10%) gave flavanol (4a) (0.12 g), m.p. 169°.

**Reaction of 2'-tosyloxy-6'-methoxychalcone oxide (6b) with  $BF_3 \cdot Et_2O$ .** To a suspension of oxide 6b in dry ether (25 ml) was added  $BF_3 \cdot Et_2O$  (6 ml; 45%) and the mixture left at room temp for 0.5 h then added to water. After 24 h the white solid which separated was recrystallised (EtOH) to give plates of 3-(6-methoxy-2-tosyloxyphenyl)-3-oxo-2-phenylpropanal (9) (0.4 g), m.p. 137–138°. (Found: C, 65.2; H, 4.8; S, 6.7; OMe, 7.9.  $C_{22}H_{20}O_6S$  requires: C, 65.1; H, 4.7; S, 7.5; OMe, 7.3%).  $\nu_{max}$  (KBr) 1623  $cm^{-1}$ . 9 gave a yellow colour with  $FeCl_3$  (EtOH) and formed a copper chelate.

**Deformylation of desoxybenzoin 9.** A solution of desoxybenzoin (9) (0.2 g) in ethanolic KOH (2N; 25 ml) was heated under reflux for 2 h. The cooled mixture was acidified with dil. HCl and the EtOH removed under reduced pressure. The residual liquid was taken up in ether. The solid obtained on removal of the solvent from the dried ethereal solution crystallised from aq. EtOH as white plates of 2'-hydroxy-6'-methoxy-2-phenylacetophenone (0.1 g), m.p. 71°. (Found: C, 74.8; H, 6.0; OMe, 11.5;  $C_{18}H_{14}O_3$  requires: C, 74.4; H, 5.8; OMe, 12.8%).  $\nu_{max}$  (KBr) 1615–1620  $cm^{-1}$ . It gave a positive ethanolic  $FeCl_3$  test.

**Acknowledgement**—The authors thank Professor E. M. Philbin for her interest in this work.

#### REFERENCES

- Communication XII, B. A. Brady, W. I. O'Sullivan and A. M. Duffield, *Org. Mass Spectrom.* 5, 199 (1972)
- J. Algar and J. P. Flynn, *Proc. Roy. Irish Acad.* 1, 42B, (1934); T. Oyamada, *J. Chem. Soc. Japan* 55, 1256 (1934)
- M. Murakami and T. Irie, *Proc. Imp. Acad. Tokyo* 11, 229 (1935)
- T. A. Geissman and D. K. Fukushima, *J. Am. Chem. Soc.* 70, 1686 (1948)
- N. Narasimhachari and T. R. Seshadri, *Proc. Indian Acad. Sci.* 30A, 216 (1949)
- T. H. Simpson and W. B. Whalley, *J. Chem. Soc.* 166 (1955)
- N. Anand, R. N. Iyer and K. Venkataraman, *Proc. Indian Acad. Sci.* 29A, 203 (1949)
- B. Cummins, D. M. X. Donnelly, J. F. Eades, H. Fletcher, F. Ó Cinnéide, E. M. Philbin, J. Swirski, T. S. Wheeler and R. K. Wilson, *Tetrahedron* 19, 499 (1963)
- F. M. Dean and V. Podimung, *J. Chem. Soc.* 3978 (1965)
- T. R. Seshadri, *The Chemistry of Flavonoid Compounds* p. 156; ed. T. A. Geissman, Pergamon Press, Oxford (1962) F. M. Dean, *Naturally Occurring Oxygen Ring Compounds* p. 337. Butterworths, London, (1963)
- K. A. Barr, F. M. Dean and H. D. Locksley, *J. Chem. Soc.* 2425 (1959)

- <sup>12</sup>A. F. Lennox, *Ph. D. Thesis*, The National University of Ireland (1967)
- <sup>13</sup>J. E. Gowan, P. M. Hayden and F. S. Wheeler, *J. Chem. Soc.* 862 (1955)
- <sup>14</sup>R. Bognar and J. Stefanovsky, *Tetrahedron* **18**, 143 (1962); R. Bognar and G. Litkei, *Acta Chim. (Budapest)* **67**, 83 (1971)
- <sup>15</sup>S. K. Grover, A. C. Jain, T. R. Seshadri, *Indian J. Chem.* **1**, 517 (1963)
- <sup>16</sup>R. Lawless, *Ph.D. Thesis*, The National University of Ireland (1962)
- <sup>17</sup>J. F. K. Eades, *Ph.D. Thesis*, The National University of Ireland (1961)