

Research



Cite this article: Oras E *et al.* 2023 Parallel worlds and mixed economies: multi-proxy analysis reveals complex subsistence systems at the dawn of early farming in the northeast Baltic. *R. Soc. Open Sci.* **10**: 230880. <https://doi.org/10.1098/rsos.230880>

Received: 22 June 2023

Accepted: 8 September 2023

Subject Category:

Earth and environmental science

Subject Areas:

evolution/environmental science/behaviour

Keywords:

early agriculture, neolithic transition, ancient diet, biomolecular archaeology, organic residue analysis, stable isotope analysis

Author for correspondence:

Ester Oras

e-mail: ester.oras@ut.ee

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6845558>.

Parallel worlds and mixed economies: multi-proxy analysis reveals complex subsistence systems at the dawn of early farming in the northeast Baltic

Ester Oras^{1,2,3}, Mari Tõrv^{1,2}, Kristiina Johanson², Eve Rannamäe², Anneli Poska⁴, Lembi Lõugas⁵, Alexandre Lucquin⁶, Jasmine Lundy⁶, Samantha Brown⁷, Shidong Chen¹, Liivi Varul⁸, Vanda Haferberga⁹, Dardega Legzdiņa⁹, Gunita Zariņa⁹, Lucy Cramp¹⁰, Volker Heyd¹¹, Michaela Reay¹², Łukasz Pospieszny^{10,13}, Harry K. Robson⁶, Kerkko Nordqvist¹⁴, Carl Heron¹⁵, Oliver E. Craig⁶ and Aivar Kriiska²

¹Institute of Chemistry, University of Tartu, Ravila 14 a, 50411 Tartu, Estonia

²Institute of History and Archaeology, University of Tartu, Jakobi 2, 51005 Tartu, Estonia

³Swedish Collegium for Advanced Study (SCAS), Linneum, Thunbergsvägen 2, 75238 Uppsala, Sweden

⁴Department of Geology, Tallinn University of Technology, Ehitajate Tee 5, 19086 Tallinn, Estonia

⁵Archaeological Research Collection, Tallinn University, Rüütli 10, 10130 Tallinn, Estonia

⁶BioArCh, Department of Archaeology, University of York, Environment Building, Wentworth Way, YO10 5DD York, UK

⁷Institute for Archaeological Sciences, Department of Geosciences, University of Tübingen, Rümelinstrasse 23, 72070 Tübingen, Germany

⁸School of Humanities, Division of History, Tallinn University, Narva rd 25, 10120 Tallinn, Estonia

⁹Institute of Latvian History, University of Latvia, Kalpaka blvd 4, LV-1050 Riga, Latvia

¹⁰Department of Anthropology and Archaeology, University of Bristol, 43 Woodland Road, BS8 1UU Bristol, UK

¹¹Department of Cultures, University of Helsinki, Unioninkatu 38, 00014 Helsinki, Finland

¹²Organic Geochemistry Unit, School of Chemistry, University of Bristol, BS8 1TS Bristol, UK

¹³Institute of Archaeology, University of Gdańsk, ul. Bielańska 5, 80-851 Gdańsk, Poland

¹⁴Helsinki Collegium for Advanced Studies, University of Helsinki, Fabianinkatu 24, 00014 Helsinki, Finland

¹⁵Department of Scientific Research, The British Museum, WC1B 3DG London, UK

EO, 0000-0002-7212-629X; MT, 0000-0001-9473-2568; ER, 0000-0003-1186-5512; AP, 0000-0002-8778-1430; LL, 0000-0003-2011-2141; AL, 0000-0003-4892-6323;

JL, 0000-0001-8250-7697; SB, 0000-0001-5001-525X; SC, 0000-0002-3495-2443; LV, 0000-0002-8498-3352; VH, 0000-0002-9241-1389; GZ, 0000-0002-7769-4341; LC, 0000-0003-2012-2753; VH, 0000-0002-0889-6513; MR, 0000-0001-6335-1690; ŁP, 0000-0003-4676-5802; HKR, 0000-0002-4850-692X; KN, 0000-0001-8696-1707; CH, 0000-0002-5206-7464; OEC, 0000-0002-4296-8402; AK, 0000-0002-0900-7626

The transition from foraging to farming was a key turning point in ancient socio-economies. Yet, the complexities and regional variations of this transformation are still poorly understood. This multi-proxy study provides a new understanding of the introduction and spread of early farming, challenging the notions of hierarchical economies. The most extensive biological and biomolecular dietary overview, combining zooarchaeological, archaeobotanical, dietary stable isotope and pottery lipid residue analyses is presented, to unravel the nature and extent of early farming in the 3rd millennium cal BCE in the northeast Baltic. Farming was introduced by incoming Corded Ware cultural groups (CWC), but some dietary segregation existed within these communities, with some having more access to domesticates, others incorporating more wild resources into their diet. The CWC groups coexisted in parallel with local hunter–fisher–gatherers (HFG) without any indication of the adoption of domesticates. There was no transition from foraging to farming in the 3rd millennium cal BCE in the NE Baltic. Instead, we see a complex system of parallel worlds with local HFGs continuing forager lifeways, and incoming farmers practising mixed economies, with the continuation of these subsistence strategies for at least a millennium after the first encounter with domesticated animals.

1. Introduction

The introduction of farming has been considered one of the key elements of ancient subsistence changes and human–animal interactions. Recent research has disputed the notion of universal and linear economic change [1–6], highlighting the importance of deeply contextualized regional studies for revealing the diversity by which people ‘transitioned’ or adopted their subsistence strategies.

The roots of European agriculture go back to the Levant some 11 000 years ago, where it emerged from the local adoption of some native plants and animals, a process lasting more than a millennium [7,8]. The spread of early farming to other parts of Europe has been explained through the exchange of ideas (cultural diffusion) or population replacement [9]. The arrival of farming in the east Baltic (nowadays Lithuania, Latvia, Estonia, Finland) has been directly related to the latter, i.e. arrival of Corded Ware cultural groups (CWC) in the 3rd millennium cal BCE [10–16], with their migratory steppe-related ancestry [17–19] and burial customs, as well as material culture differing from the local hunter–fisher–gatherer (HFG) populations [20–23]. Yet, the biological legacy and the aftermath of the arrival of early farming in the east Baltic, set in the context of local forager communities, northern climatic conditions with dense woodland and aquatic landscapes, abundant wild resources and missing indigenous species suitable for domestication, has remained ambiguous. Recent biomolecular findings have confirmed that the introduction of new, domesticated species took place from the 3rd millennium cal BCE onwards [6,24–26], but we lack clarity concerning the extent and character that the subsistence and gross diet actually changed and for whom.

The goal of this study is to disentangle the complexity of early farming livelihood among CWC groups and assess its impact on local HFG populations in northern latitudes beyond the 56th parallel north, focusing on Estonian and Latvian (labelled as northeast Baltic from here onwards) 3rd millennium cal BCE material. To do so, we bring together an exceptional combination of a new bioarchaeological dataset covering human bone collagen stable isotope and pottery lipid residue analyses, zooarchaeological records together with recent direct accelerator mass spectrometry (AMS) radiocarbon dates, as well as archaeobotanical information from previous studies. Our findings show multifaceted regional adaptive strategies revealing both segregated dietary practices and mixed economy during the introductory phase of early farming, clearly deviating from the grand-scale narratives of linear and hierarchical models of ancient subsistence change.

We use the more general term ‘farming’, rather than ‘agriculture’, referring mostly to animal husbandry, not necessarily including the element of crop cultivation, although in the context of the northeast Baltic early farming seems to mostly rely on stockbreeding. The term ‘domesticates’ indicates the species (both plant and animal) that rely on human input in their growth cycle and survival. The 4th–3rd millennium cal BCE is traditionally classified as Neolithic in the northeast Baltic, whereas the majority of the period is characterized by pottery-using HFG lifeways. By HFG




Figure 1. (a) Map of the analysed sites in this study, and (b) an example of a CWC burial Sope B I, Estonia. (c) Examples of CWC pottery: a beaker from the Narva-Jõesuu IIB burial ground (Estonia) and two fragments of Estonian-type Corded Ware pots from the Lemmetsa I settlement (Estonia) and (d) early domesticates' bones found as grave goods from the Sope B I burial. Note that the Abora I settlement also includes other Porous and Lubāna pottery types besides the CWC vessels.

populations, we refer to local forager groups with different regional and/or cultural affiliations: mostly Typical/Late Comb Ware groups (3900–1800 cal BCE, the two ceramic types are often chronologically difficult to separate without direct dating; hence here collated as Comb Ware culture (CMC)), but also the users of regional Porous (3300–1800 cal BCE, a specific pottery group defined in Latvian archaeology from the Lake Lubāns region) and Lubāna (2600–1500 cal BCE) pottery types in Latvia. By CWC (2800–2000 cal BCE), we refer to archaeological complexes which differ from both earlier and contemporaneous ones: individuals interred in burials with a distinctive (mostly) crouched body position and associated grave goods, such as Corded Ware pottery, various stone axes and adzes, flint knives, both domesticated and wild animal bones and tools made from them; but also the users of Corded Ware pottery, which is a clearly distinguishable 3rd millennium cal BCE ceramic type [27] (figure 1).

2. Results

2.1. Early domesticated animal bone and plant records

Faunal remains from CWC settlements are scarce and highly fragmented, often mixed with preceding and later materials. In Estonia, early domesticated livestock has been mostly identified in CWC burial contexts [12], where they are found either as unprocessed animal remains or worked bones (i.e. tools or their preceding blanks). In Latvia, besides at least one chisel made from an undated sheep/goat bone from Zvejnieki CWC burial no. 137 [28,29], the earliest domesticates have been suggested based on their find context from the 4th and 3rd millennia cal BCE settlements of Zvidze, Kreiči, Sārnate, Abora I, Lagaža and Eiņi [10,11,30]. However, since all are multi-period sites without direct AMS radiocarbon dates, this evidence remains inconclusive.

A range of new radiocarbon AMS dates from a selection of domesticated animal bones from several CWC contexts shows that the introduction of early domesticates in the northeast Baltic took place between 2730 and 2490 cal BCE (modelled starting time, 95.4%; figure 2 and table 1; see also electronic supplementary material, table S1). Although the aurochs (*Bos primigenius*; the ancestor of the domestic cattle) were commonly found across Europe [33], their population in Estonia receded by the 3rd millennium cal BCE, and there are no traces of local domestication of the aurochs or interbreeding with imported domestic cattle [34,35]. Hence the dated specimen must be the domesticated congener.




Figure 2. The AMS dates of early domesticates from the northeast Baltic. Calibrated with OxCal 4.4.4. using IntCal20 atmospheric curve [31], all the dates are rounded by 10. For data, table 1.

The two sheep/goat samples from the Sope B I burial were subjected to zooarchaeology by mass spectrometry (ZooMS) analysis [36] to clarify the species, but the results were unsuccessful (electronic supplementary material, table S2). These two samples are from different skeletal elements, and their dates represent a single event statistically (2840–2490 cal BCE; χ^2 -test d.f. = 1; $T = 0.113$; $T(5\%) = 3.841$, with an overall agreement index (A_{comb} 130%), hence they could belong to the same individual. As the taxonomic differentiation between the domestic pig (*Sus domesticus*) and wild boar (*Sus scrofa*) is impeded by bone morphology [12,37], all the suids dating around 2880–2350 cal BCE are cautiously identified as *Sus* sp., although the possibility of domesticated suids has been suggested for the Loona settlement specimen [34].

In terms of domesticated plants, cereal-type grains have not been identified from any CWC context in the northeast Baltic. Single grains of barley (*Hordeum vulgare*) and possibly einkorn (*Triticum monococcum*) have been reported from the Kreiči settlement in Latvia [38]. However, the site was excavated in the 1950s, its contexts are partially mixed with finds from the 5th to 2nd millennium cal BCE, perhaps even later, while the grains are no longer available for direct AMS dating [39]. A barley grain and an impression of another cereal grain have previously been documented in relation to a CWC sherd from the Iru settlement [11,40], but the sherd and the grain are unfortunately lost, making any further analysis impossible. A potential cereal grain imprint, most likely *Hordeum vulgare*, was noted in a Lubāna type sherd from the Abora I settlement during sampling for organic residue analysis reported in this article, but the foodcrust has not been directly dated.

The scarcity of macrobotanical remains has instigated the extensive use of pollen data for reconstructing early cereal cultivation in the northeast Baltic [10,11,41]. Cereal-type pollen grains identified as unspecified cereal (*Cerealia*-type), barley (*Hordeum*-type), wheat (*Triticum*-type) and oat (*Avena*-type) have been previously noted from a few sites from the 7th millennium cal BCE (electronic

Table 1. An overview of the AMS radiocarbon dates of the earliest domestic livestock in Estonia. All finds, besides Loona, are associated with CWC contexts. Calibrated with OxCal 4.4.4 using IntCal20 atmospheric curve [31], all the dates are rounded by 10.

species	arch. site	site type (burial)	sample ID	skeletal element (U, unworked; P, pretreated tool blank)	¹⁴ C lab no.	¹⁴ C dating BP	cal BCE (95.4%)	reference for ¹⁴ C date
<i>Ovis aries/Capra hircus</i>	Sope	burial (B I)	AI 2607	Os coxae (U)	UBA-40305	4057 ± 27	2840–2490	this study
<i>Ovis aries/Capra hircus</i>	Sope	burial (B I)	AI 2607: 2	Radius (P)	UBA-45549	4073 ± 36	2840–2490	this study
<i>Bos taurus</i>	Ardlu	burial (II)	AI 3499: 66	Os coxae (U)	UBA-38464	4034 ± 41	2850–2460	this study
<i>Sus</i> sp.	Sope	burial(B I)	AI 2607: 3	Tibia (P)	UBA-40304	3964 ± 28	2580–2340	this study
<i>Sus</i> sp.	Tallinn Müller's Field (Pärnu Rd. 37, 41)	settlement	AI 8244	Scapula (U)	UBA-45532	3970 ± 35	2580–2340	this study
<i>Sus</i> sp.	Kunila ^a	burial	AI 3989/AZ-3	Ilium (U)	UBA-45542	3830 ± 29	2460–2140	this study
<i>Sus</i> sp.	Kunila ^a	burial	AI 3898/AZ-6	Ilium + ischium (U)	UBA-45547	3889 ± 36	2470–2200	this study
<i>Sus</i> sp.	Kunila ^a	burial	AI 3989/AZ-4	Ischium (U)	UBA-45543	3883 ± 32	2470–2210	this study
<i>Sus</i> sp. ^b	Loona	settlement	AI 4129	Unknown	Ua-4825	4050 ± 80	2880–2350	Lõugas <i>et al.</i> , 1996 [32]

^aExact burial affiliation unknown.

^bSpecimen absent from the collections.

supplementary material, table S3). Yet, the overall number of samples (and sites) reporting finds of cereal-type pollen remains low up until the beginning of the 4th millennium BCE. An abrupt increase in both sample and site frequency and overall number of cereal-type pollen grains is visible in both Estonia and Latvia from *ca* 3500 cal BCE onwards, whereas in the following millennium the pollen evidence stabilizes or even shows some regression [42]. Gradual increase of open landscapes together with slightly growing evidence of cereal-type pollen finds has been recorded from the beginning of the 2nd millennium cal BCE onwards [41,43].

However, several methodological peculiarities like the uncertainties of dating, pollen identification and estimating pollen source area [39,44,45] must be considered. Most importantly, pollen taxonomy is not directly compatible with conventional botanical taxonomic precision, and one pollen type can include pollen from plants belonging to different species, families or sometimes even genera [46]. Most cereals commonly grown in Europe produce pollen identifiable to the level of pollen type—hence the *Hordeum*-type, *Triticum*-type, *Avena*-type and *Cerealia*-type identifications—and all these types include some species of wild grasses [42,47,48]. Among such wild grass species are several members of the grass family (e.g. *Elymus arenarius*, *Glyceria fluitans*, *Bromus inermis*) common in the northeast Baltic, which produce pollen grains with similar size and/or surface structure to those of the cereals. Hence, arguing for early crop cultivation based solely on pollen findings remains uncertain, and we currently lack supportive evidence from other proxies to substantiate that domesticated plants were part of the main subsistence strategy in the 3rd millennium cal BCE.

2.2. Stable isotopes from human bone collagen

The collagen stable isotope analysis (SIA; raw data reported in electronic supplementary material, table S5) contains all the individuals directly dated to the 3rd millennium cal BCE from Estonia ($n = 10$) and Latvia ($n = 17$). Besides these, the Abora I site includes directly dated individuals ($n = 13$), 20 specimens found as loose human bones ($n = 11$) and intact burials ($n = 9$) all found from the settlement layer dated to *ca* 3400–1900 cal BCE [49,50]. The dataset also entails an infant and adult female from the Selgas double grave, and burial 307 from Zvejnieki due to their characteristic traits for CWC burials. This results in a total of 49 individuals, with either single or multiple skeletal elements being analysed, and thus represented by 65 samples of both adult and non-adult values. Since the collagen yield together with carbon and nitrogen percentage inclusion was inconsistently reported in previous studies, we use the C:N ratio falling between 3.1 and 3.5 in this study (electronic supplementary material, table S5) as a quality criterion for collagen preservation [51,52].

The human bone collagen SIA values display considerable heterogeneity among the 3rd millennium cal BCE populations (figure 3a), with values ranging from -24.9‰ to -15.7‰ for $\delta^{13}\text{C}$, and $+8.9\text{‰}$ to $+17.8\text{‰}$ for $\delta^{15}\text{N}$. The statistical comparison of CMC, CWC and Abora I SIA values shows significant differences (Mann–Whitney pairwise test all p -values < 0.05), but due to the small number of CMC samples this must be taken with caution, whereas the difference holds for the Abora I and CWC individuals. The CWC burials have a narrower range of values ($\delta^{13}\text{C}$ range -23.5‰ to -19.1‰ , $\delta^{15}\text{N} + 8.9\text{‰}$ to $+11.8\text{‰}$; see electronic supplementary material, table S4), which is indicative of protein intake from terrestrial sources. The consumption of terrestrial foods is indirectly evidenced by the overlapping paired AMS dates of terrestrial animals and humans from the Ardu II and Sope B I burials (electronic supplementary material, figure S1), showing that humans do not have apparent ages caused by a reservoir offset. Multiple samples are available for three CWC individuals—Ardu I, Kunila I and Sope B I—which permits us to track the dietary protein intake changes in an individual's lifetime. For Sope B I and Kunila I, the dietary change is negligible (less than 1‰), but Ardu I displays elevated $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in adulthood (based on comparing SIA results from molars and the rib; figure 3a).

In contrast to the CWC individuals, there is considerable intra-site variability in the SIA values among the 33 individuals from Abora I. Here the $\delta^{13}\text{C}$ values range from -24.8‰ to -20.5‰ , and $\delta^{15}\text{N}$ from $+9.7\text{‰}$ to $+17.8\text{‰}$, whereas the remarkably high $\delta^{15}\text{N}$ value ($+17.8\text{‰}$) for burial 81 belongs to a non-adult aged 1 ± 2 months [50] and is indicative of weaning. There is no significant chronological or burial practice (crouched versus supine position versus loose human bones) based difference in the Abora I cohort (Mann–Whitney pairwise test p -values ranging from 0.14 to 0.98). From the CMC group, the Tamula individuals had higher $\delta^{15}\text{N}$ values and lower $\delta^{13}\text{C}$ values indicating a reliance on freshwater resources, whereas the Tamula XXII individual demonstrated low lifetime intra-individual dietary changes. The HFG female from Naakamäe (Saaremaa island) with a maximum date range between 2910 and 2490 cal BCE (reservoir effect (RE) corrected, 95.4% probability [32,53]), stands out




Figure 3. Summary of biomolecular dietary results from the 3rd millennium cal BCE northeast Baltic. (a) plot of human bone collagen stable isotope data, (b) with an example of a typical CWC burial, Ardu I. Note that the Ardu I individual is displayed with connected data points showing intra-individual dietary change (the higher $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values are from adulthood). Data derive from this study and previous publications [12,18,32,50,53–59]. (c) $\delta^{13}\text{C}$ values of mid-chain-length fatty acids ($\text{C}_{16:0}$ and $\text{C}_{18:0}$) extracted from Estonian and Latvian (pre-)3rd millennium cal BCE CMC (i), CWC (ii) and other 3rd millennium cal BCE regional wares from Abora I (iii) with full aquatic biomarkers indicated by filled symbols. The reference ranges presented in C (i–iii), calculated at 68% confidence, were derived from data obtained from the tissues of modern animals, mostly from the Baltic region [60–62].

for her high $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values, being the only individual in our dataset with a clear focus on marine foodstuffs.

2.3. Pottery lipids

A total of 178 samples from CMC and CWC pottery from 18 sites in Estonia and Latvia were subject to organic residue analysis (ORA). Of these, 111 samples are from CMC and 67 from CWC vessels. This dataset is further supplemented by 15 Porous and six Lubāna-type ceramic samples from Latvia, resulting in a total of 199 samples, of which 113 are reported here for the first time. The biomarker and isotope results of ORA from all samples are reported in electronic supplementary material, table S6, whereas the results from previous publications [25,63,64] were included in the dataset only if both biomarker and compound specific isotope data was available.

To trace early farming through ORA, $\delta^{13}\text{C}$ measurements from the two major fatty acids, $\text{C}_{16:0}$ and $\text{C}_{18:0}$, are highly informative for distinguishing freshwater and marine organisms, porcine and ruminant fats, and most importantly dairy products [65–67]. As domesticated and wild ruminants are isotopically inseparable, the detection of dairy fats remains the major indicator of exploiting domesticated ruminant species and their products.

A Mann–Whitney *U*-test for the isotopic measurements of the $\delta^{13}\text{C}_{16:0}$ and $\delta^{13}\text{C}_{18:0}$ fatty acids in the CMC ($n = 93$) and CWC ($n = 64$) samples shows differences between the two groups (p -value < 0.029 for both), with the average values for $\delta^{13}\text{C}_{16:0}$ and $\delta^{13}\text{C}_{18:0}$ ranging between $-28.6 \pm 2.5\text{‰}$ and $-28.0 \pm 2.6\text{‰}$ for the CMC, and $-29.4 \pm 2.3\text{‰}$ and $-29.8 \pm 3.0\text{‰}$ for the CWC samples. The $\delta^{13}\text{C}$ results (figure 3c) from CWC, show the clear inclusion of dairy products in seven (potentially eight) CWC vessels, being the earliest evidence of dairy fats in the northeast Baltic. This number should be considered as a minimum, as mixing dairy with other products might result in isotopic values outside the dairy ellipses, also into the ruminant region [68]. Notably, all the dairy samples are from the north and

northeast part of Estonia, with none from the inland sites sampled. There is one sample from Abora I plotting between the dairy and ruminant ellipses, but it also includes the full suite of aquatic biomarkers, which might hint at mixing dairy products with other substances with more depleted $\delta^{13}\text{C}_{16:0}$ values.

Despite the first detection of dairy fats in this region, aquatic biomarkers from either marine or freshwater organisms indicate that different aquatic products were processed in both CMC (40 samples out of 93) and CWC vessels (22 samples out of 64) as well as in other regional wares (15 samples out of 18; figure 3c). These specific biomarkers include ω -(*o*-alkylphenyl)alkanoic acids with carbon atoms ranging from C_{16} to $\text{C}_{20/22}$, formed during the heating of long-chain polyunsaturated fatty acids of aquatic organisms, together with one or more of the isoprenoid fatty acids (phytanic, pristanic and 4,8,12-trimethyltridecanoic (TMTD)) [69,70], with further support by the ratios of 3S,7R,11R,15-phytanic acid (SRR) and 3R,7R,11R,15-phytanic acid (RRR) diastereomers [71]. Hence we have clear evidence for initial dairy production/storage, but also for the consumption of aquatic substances, suids and ruminants throughout the 3rd millennium cal BCE.

3. Discussion

Comparison of different dietary proxies—zooarchaeological, archaeobotanical, SIA and ORA—provides a more cohesive picture of the arrival, nature and spread of early farming in the northeast Baltic. In contrast to the northwest Baltic [72], the scarce macrobotanical and pollen data show an absence of considerable crop cultivation in the northeast Baltic during the 3rd millennium cal BCE. Plant consumption here was rather tuned towards wild species [16,73], although tentative experimenting with crop cultivation cannot be entirely excluded [74]. The oldest directly dated cereal grains in the northeast Baltic are from the 2nd millennium cal BCE [39,72,75,76], while crop cultivation at that time is also supported by continuous *Cerealia* pollen records [77,78] and the establishment of the first field systems [21].

Early farming in the northeast Baltic was based on animal husbandry, pioneered by CWC communities who introduced domesticates during the first half of the 3rd millennium cal BCE, and is comparable to the dates for early domesticates from the southeast Baltic [79]. In Finland, the earliest cattle find is from the 2nd millennium cal BCE [80], although a goat hair found from the Perttulanmäki CWC burial might also support an earlier arrival time of domesticates [81]. However, domesticated animals start to dominate the faunal record in the northeast Baltic only around 1000 cal BCE [12,30,34,82].

The introduction of domesticates is unlikely to have transformed the economies of everyone living in the northeast Baltic during the 3rd millennium cal BCE. The individuals afforded CWC burial form an isotopically narrow and homogeneous group (figure 3a) most probably relying on terrestrial resources, similar to CWC individuals from Lithuania and Central Europe [2,83]. By contrast, the other contemporaneous human remains display very diverse foodways (figure 3a) from strongly aquatic diets, mixed aquatic/terrestrial diets and more terrestrial orientated diets.

At the finer scale of culinary practices, pottery ORA shows the combination of domesticated, wild terrestrial and aquatic resources among the CWC populations (figure 3c). Dairy was identified in seven CWC samples (*ca* 10% of analysed CWC samples) from four coastal/estuarine sites, whereas in at least two of those the consumption of other (wild) resources, including aquatic, is evident. Hence CWC populations in the northeast Baltic were relying on a mixed economy with parallels from Lithuanian and Finnish 3rd millennium BCE contexts [6,24,26,84]. Similar tendencies have been also highlighted in the development of early farming in the northwest Baltic [85–87] and Central Europe [88].

The variations in 3rd millennium cal BCE subsistence systems outlined in this study might have different analytical, regional and socio-cultural explanations. The pottery ORA reflects the dietary practices of wider communities, while SIA shows the dietary history of single individuals (CWC burials) preserved to this day. We can hypothesize that the discrepancy between the CWC vessels' ORA (mixed economy) and CWC burial SIA results (mostly terrestrial diet) might point to different dietary availability and/or preferences among different CWC groups.

The individuals from the traditional CWC burials display a rather homogeneous range of SIA values, the vast majority of domesticated animal bones have been found in CWC burial contexts, while dairy products are detected in CWC vessels from a limited number of coastal/estuarine sites. Therefore, given that the pottery ORA and human bone collagen SIA data are contemporaneous, one possible explanation is that in the introduction phase, domesticated species and their secondary products might have been more accessible to certain subgroups of CWC people, who were afforded a specific

burial rite. Alternatively, we might be witness to analytical constraints, as it must be acknowledged that bulk SIA cannot trace the limited consumption of aquatic foods [89]. The inclusion of smaller proportions of aquatic products into the diets of the individuals in CWC burials could be refined with compound specific SIA of amino acids from bone collagen in the future [90]. Finally, the discrepancy between the pottery ORA and human SIA values might also derive from specific culinary practices, i.e. aquatic products being preferably processed in ceramic vessels, whereas a wider range of other foodstuffs were both procured and consumed without the use of ceramic containers.

Nevertheless, CWC pottery ORA shows that the abundant wild resources of northern latitudes offered optimal conditions for practising a mixed economy during the Mid-Holocene. Indeed, the exploitation of wild resources was common among CWC populations from the very beginning [91], with several CWC burials in the northeast Baltic including wild animal bones as well [12,79,92]. The Ardu I CWC individual whose burial customs and childhood diet follow the characteristic CWC traditions, serves as an intriguing example of dietary conversion from fully terrestrial to the mixed economy and a shift towards consuming aquatic, possibly marine resources, in his adulthood (figure 3*n*). The CWC early farmers were adaptive to different subsistence strategies, which might have facilitated their movement into northerly latitudes, whereas the natural conditions in the northeast Baltic probably provided favourable conditions for larger inclusion of forager subsistence elements into their livelihood.

There was no transition from HFG to farming in the 3rd millennium cal BCE northeast Baltic. The introduction of domesticates had little if any impact on local HFG groups, who continued to rely on wild economies and show no signs of early farming through ORA of pottery or SIA of human bone (figure 3). This proves the coexistence of parallel worlds, where farming and/or mixed economy-based CWC groups resided together with the foragers for nearly a full millennium. Such a socio-economic divide is perhaps also reflected by the ancient genomic data: the Tamula (south Estonia) and Naakamäe 3rd millennium cal BCE individuals following forager diets are affiliated with HFGs [17], whereas the CWC populations show different genetic ancestry, originating from/related to the steppe region [18,56]. Similar cultural but also genetic divergence between foragers and early farmers has been established in other parts of Europe across the so-called Neolithic transition [93–96]. However, the proportions of CWC and local CMC populations among the 3rd millennium cal BCE in the northeast Baltic need further investigations to determine the extent of population turnover prior to *ca* 2000 cal BCE [19].

The northeast Baltic 3rd millennium cal BCE data provides important insights for acknowledging the complexity for HFG–farmer transitions. There was no single or linear pathway in the change of subsistence systems, and we are very far from the hierarchical and ‘all-or-nothing’ conceptualization when it comes to early farming practices [97–99]. In fact, the whole concept of ‘transition’ is erroneous in the northeast Baltic context, as local HFGs remained true to their forager lifeways, which existed in parallel with (segregated) farming pioneering, yet mixed economy CWC populations. There is probably no total replacement of foraging by farming as the major source of food production in northern latitudes (beyond the 56th parallel north), and future research should rather address the economic dialogues between the wild and domesticated, as well as tackle the prolonged adaptation and experimenting processes with several success–failure episodes over the following millennia.

4. Methods

4.1. Organic residue analysis: sample preparation

Majority of samples were prepared and analysed at the University of Tartu Archemy laboratory and University of York BioArCh facility. Food crust adhered to pottery surfaces were removed using clean scalpels. Ceramic powder samples were drilled using clean drill bits, discarding the upper surface powder to avoid contamination. Lipids were extracted and derivatized (methylated) from *ca* 1 g of ceramic powder and 20 mg of foodcrusts using acid-catalysed methylation following previously reported procedure [61]. The samples were analysed with gas chromatography–flame ionization detector (GC-FID) for lipid quantification, gas chromatography–mass spectrometry (GC-MS) for general biomolecular characterization, and GC-MS in selected ion monitoring (SIM) mode targeting aquatic biomarkers and GC–combustion–isotope ratio mass spectrometry (GC-C-IRMS) to determine the $\delta^{13}\text{C}$ values of $\text{C}_{16:0}$ and $\text{C}_{18:0}$ fatty acids.

A subset of samples (sample nos. 186–199 reported in electronic supplementary material, table S6) was analysed at the University of Bristol, School of Chemistry using the previously reported

simultaneous extraction and methylation protocol [100], followed by the analysis with high temperature (HT) GC-FID, GC-MS and GC-C-IRMS.

4.2. Gas chromatography–flame ionization detector analysis

GC-FID analysis was conducted at the York BioArCh facility (University of York) using Agilent 7890 B Series GC and DB1-HT polyimide-coated fused silica column (15 m × 320 μm × 0.1 μm; J&W Scientific, Folsom, CA, USA). Injected sample size was 1 μl. The splitless injector was used at 300°C with helium carrier gas. The temperature was set at 100°C for 2 min, with the gradient of 20°C min⁻¹ up to 325°C maintained for 3 min. Data were acquired using CHEMSTATION software.

In Bristol, the analyses were undertaken using an Agilent Technologies 7890A GC system and 100% dimethyl polysiloxane-fused silica column (15 m × 320 μm × 0.1 μm; Restek, Rxi-HT). Injected sample size was 1 μl. The on-column injector was used with helium carrier gas. The temperature programme was set at 50°C for 2 min, followed by a gradient of 10°C min⁻¹ up to 350°C, followed by a 10 min isothermal hold. Data were acquired using HP ChemStation software (Rev. B.03.02 [341] Agilent Technologies).

4.3. Gas chromatography–mass spectrometry analysis

GC-MS was conducted at the Institute of Chemistry (University of Tartu) with Agilent 7890A Series gas chromatograph and Agilent 5975C Inert XL mass-selective detector with a DB5-MS (5%-phenyl)-methylpolysiloxane column (30 m × 0.25 mm × 0.25 μm). Injected sample size was 1 μl. The splitless injector and interface were maintained at 300°C and 280°C, respectively, helium was used as the carrier gas at a constant flow. The GC column was inserted directly into the ion source of the mass spectrometer. The temperature programme was set as follows: 50°C for 2 min, thereafter a gradient of 10°C min⁻¹ up to 325°C and kept there for 14.5 min. Compounds were identified with Agilent Chemstation software using also NIST mass spectral library.

Additional dedicated GC-SIM method was used for detecting aquatic biomarkers as reported in previous studies [61,101]. For this GC-MS equipped with a DB-23 (50%-cyanopropyl)-methylpolysiloxane column (60 m × 0.25 mm × 0.25 μm; J&W Scientific, Folsom, CA, USA) was used. Injected sample size was 1 μl. The oven temperature was set at 50°C for 2 min with an increase of the temperature up to 100°C (10°C min⁻¹). Thereafter, the temperature was raised by 4°C min⁻¹ to 140°C, then by 0.5°C min⁻¹ to 160°C and, finally, by 20°C min⁻¹ to 250°C where it was maintained for 10 min. The carrier gas used was helium with a flow rate of 2.4 ml min⁻¹. The SIM mode was set to target characteristic aquatic biomarkers' ions groups: *m/z* 74, 87, 213, 270 for 4,8,12-trimethyltridecanoic acid (TMTD), *m/z* 74, 88, 101, 312 for pristanic acid, *m/z* 74, 101, 171, 326 for phytanic acid and *m/z* 74, 105, 262, 290, 318, 346 for the detection of *ω*-(*o*-alkylphenyl) alkanolic acids of carbon lengths C₁₆–C₂₂ (APAA_{16–22}). Additionally, two phytanic acid diastereomers (3S,7R,11R,15-phytanic acid or SRR and 3R,7R,11R,15-phytanic acid or RRR) were separated for the calculation of the percentage of SRR in total phytanic acid (SRR%) by integrating the *m/z* 101 ion [71].

GC-MS analysis in Bristol was performed using a Thermo-Finnigan Trace MS and introduced via a PTV injector set to splitless mode onto an HP-1 fused silica capillary column (100% dimethylpolysiloxane, 50 m × 0.32 mm × 0.1 μm; Agilent). Injected sample size was 1 μl, and helium was used as a carrier gas at a constant flow. The temperature programme was set as follows: 50°C for 2 min, thereafter gradient of 10°C min⁻¹ up to 300°C where it was held for 10 min. The MS was operated in electron ionization (EI) mode at 70 eV, and acquired between *m/z* 50 and 650 at 1.7 scans s⁻¹ in full scan mode. Data were acquired and analysed using XCalibur (v. 1.2).

4.4. Gas chromatography–combustion–isotope ratio mass spectrometry analysis

GC-C-IRMS analysis was conducted on acid-extracted (methylated) samples to determine the ¹³C/¹²C ratio in two most abundant fatty acids (C_{16:0} and C_{18:0}). At the York BioArCh facility, the samples were analysed with two instruments: Delta V Advantage isotope ratio mass spectrometer (Thermo Fisher, Bremen, Germany) linked to a Trace Ultra 1310 gas chromatograph (Thermo Fisher) with a GC Isolink II interface (CuO combustion reactor held at 850°C); and with Isoprime 100 (Isoprime, Cheadle, UK) linked with Agilent 7890B Series Gas Chromatograph (Agilent Technologies, Cheadle, Cheshire, UK) with Isoprime GC5 interface (Isoprime Cheadle, UK, with CuO combustion reactor held at 850°C). Parallel acquisition of the molecular data was achieved by deriving a small part of the flow to an ISQ

mass spectrometer (Thermo Fisher) and Agilent 5975C inert XL mass-selective detector (MSD) equipped with a quadrupole mass analyser (Agilent technologies, Cheadle, Cheshire, UK), respectively.

All fatty acid methyl esters (FAMES) samples were diluted with *n*-hexane, and subsequently 1 μl of each sample was injected into a DB-5MS UI fused-silica column (PN 122-5562UI; 60 m \times 250 μm \times 0.25 μm ; J&W Scientific technologies, Folsom, CA, USA). The split/splitless injector was operated in splitless mode. The temperature was set at 50°C for 0.5 min and raised by 25°C min^{-1} to 175°C, then raised by 8°C min^{-1} to 325°C where it was held for 20 min. Ultra-high-purity-grade helium was used as the carrier gas at constant flow (2.0 ml min^{-1}). All samples were measured in duplicates.

Eluted products were combusted to CO_2 and ionized in the mass spectrometer by electron impact. Ion intensities of m/z 44, 45 and 46 were monitored in order to automatically compute the $^{13}\text{C}/^{12}\text{C}$ ratio of each peak in the extracts. Computations were made with Isodat (v. 3.0; Thermo Fisher) and IonOS/LyticOS software (Isoprime, Cheadle, UK) and were based on comparisons with a repeatedly measured standard reference gas (CO_2). The results from the analysis are reported in parts per thousand (‰) relative to an international standard (V-PDB) and corrected for the carbon atom added during methylation using a mass balance equation [102].

At Bristol the samples were analysed with Isoprime 100 (Isoprime, Cheadle, UK) linked with Agilent 7890A Series GC (Agilent Technologies, Cheadle, Cheshire, UK) via an Isoprime GC5 interface (Isoprime Cheadle, UK; with furnace temperature of 850°C). The FAMES were diluted with *n*-hexane and injected (1 μl) into an HP-1 fused silica column with a dimethyl polysiloxane stationary phase (50 m \times 0.32 mm \times 0.17 μm ; Agilent). The temperature programme was set at 40°C for 2 min and raised 10°C min^{-1} to 300°C where it was held for 10 min. Helium was used as a carrier gas at constant flow (2.0 ml min^{-1}). All analyses were conducted in duplicate. Following conversion to CO_2 over copper oxide pellets, ion intensities of m/z 44, 45 and 46 were monitored to determine $^{12}\text{C}/^{13}\text{C}$, relative to the repeatedly measured standard reference CO_2 gas. Instrument performance was determined using an in-house FAME mixture ($\text{C}_{11:0}$, $\text{C}_{13:0}$, $\text{C}_{16:0}$, $\text{C}_{21:0}$, $\text{C}_{23:0}$) with known $\delta^{13}\text{C}$ values, with an instrument precision of $\pm 0.3\text{‰}$, determined via repeated analysis between samples. Data processing was carried out using IonVantage (v. 1.5.6.0, Isoprime). $\delta^{13}\text{C}$ values are reported in parts per thousand (‰) relative to the international standard (V-PDB), and were corrected for the methyl group added during derivatization via a mass balance equation [102].

4.5. Zooarchaeology by mass spectrometry analysis

ZooMS analysis was carried out following previously published protocols [103,104]. Bone chips were demineralized in 0.5 M HCl and rinsed in 50 mM ammonium bicarbonate. The demineralized bone chip was then gelatinized at 65°C for 1 h and the resulting supernatant was treated with 0.4 μg trypsin (Thermo Scientific Pierce Trypsin Protease). Enzymatic digestion took place at 37°C for 18 h. The incubated samples were concentrated and desalted using C18 ZipTips (Thermo Scientific Pierce C18 Tips) and eluted in a final solution of 50 μl of 50% acetonitrile and 0.1% trifluoroacetic acid (TFA). Then 0.5 μl of the resulting solution was mixed with 0.5 μl of α cyano-4-hydroxycinnamic acid solution (10 mg ml^{-1} in 50% acetonitrile and 0.1% trifluoroacetic acid) and allowed to crystallize.

Samples were analysed at the Max Planck Institute for the Science of Human History using a Bruker AutoFlex LRF Speed. The resulting spectra were peak picked with a signal to noise ratio of 3.5 after baseline correction, smoothing and deisotoping with the default parameters and analysed with flexAnalysis 3.4 (Bruker Daltonics) and mMass software [105]. The spectra were compared against a reference library of known peptide markers [36,103,106,107]. The resulting spectra are provided (see electronic supplementary material, Data ZooMS files) and peptide markers are reported following standard nomenclature [108] (see electronic supplementary material, table S2).

4.6. Collagen stable isotope analysis

The SIA results presented in this paper derive mostly from published materials and thus are conducted at different laboratories with different pretreatment protocols for collagen extraction. The extraction methods and hardware used for analysis are described in the papers referred to in electronic supplementary material, table S5.

The samples analysed in this study (burials from Sope, Ardu, Kunila) were analysed at the Archemy Lab, University of Tartu (UT). Collagen was extracted using the modified Longin method following the previously reported procedure [109]. Bone/tooth dentine samples were demineralized in 0.25 M

hydrochloric acid (HCl) at room temperature for 48 h. The demineralized samples were rinsed with deionized water, which was replaced with 0.01 M HCl solution, to be placed in the oven (58°C) for 16 h for gelatinization. The resulting solution was filtered with Whatman nitrocellulose membrane filters with pore size 5 µm to remove any insoluble residue and freeze-dried. Stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotopic values were measured with an automated carbon and nitrogen elemental analyser isotope ratio mass spectrometer (EA-IRMS) in the Department of Geology, UT. Samples were weighed as duplicates into tin capsules (approx. 1.0–1.2 mg) and combusted in a Thermo Flash HT EA with introduction of separated N_2 and CO_2 to a Delta V plus via a ConFlo IV interface. The data were calibrated against international standards from IAEA (for nitrogen IAEA N-1, $\delta^{15}\text{N}_{\text{AIR}} = +0.4\text{‰}$, IAEA N-2, $\delta^{15}\text{N}_{\text{AIR}} = +20.3\text{‰}$, USGS25, $\delta^{15}\text{N}_{\text{AIR}} = -30.4\text{‰}$, and for carbon IAEA CH3, $\delta^{13}\text{C}_{\text{VPDB}} = -24.72\text{‰}$, and IAEA CH6, $\delta^{13}\text{C}_{\text{VPDB}} = -10.449\text{‰}$). The results are expressed using the delta notation in per thousand (‰) [110] relative to the international marine limestone VPDB standard for carbon and AIR for nitrogen. The long-term stability error for the isotope ratio measurements determined from repeated measurements of international and laboratory standards was better than $\pm 0.2\text{‰}$ (1 s.d.) for nitrogen, and $\pm 0.1\text{‰}$ (1 s.d.) for carbon.

4.7. Accelerator mass spectrometry dating

Approximately 1 g of sample was removed from the animal bones and submitted to AMS radiocarbon dating at the CHRONO Centre in Queen's University Belfast. The obtained AMS dates were calibrated with the OxCal 4.4.4., using the IntCal20 atmospheric calibration curve [31] and rounded by 10.

Ethics. The study includes solely archaeological material from Estonia and Latvia, with agreements and sampling protocols arranged in collaboration with the museum collections.

Data accessibility. The data are provided in electronic supplementary material [111].

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors' contributions. E.O.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, visualization, writing—original draft; M.T.: conceptualization, data curation, formal analysis, investigation, visualization, writing—original draft, writing—review and editing; K.J.: conceptualization, data curation, formal analysis, investigation, writing—original draft, writing—review and editing; E.R.: conceptualization, data curation, formal analysis, investigation, visualization, writing—original draft, writing—review and editing; A.P.: data curation, investigation, writing—review and editing; L.L.: data curation, resources, validation, writing—review and editing; A.L.: formal analysis, investigation, methodology, resources, validation, writing—review and editing; J.L.: data curation, formal analysis, investigation; S.B.: data curation, formal analysis, methodology, validation, writing—review and editing; S.C.: data curation, formal analysis; L.V.: data curation, formal analysis, investigation, visualization, writing—review and editing; V.V.: data curation, resources; D.L.: data curation, investigation, resources; G.Z.: data curation, resources; L.C.: conceptualization, data curation, investigation, methodology, resources, supervision, writing—review and editing; V.H.: resources, writing—review and editing; M.R.: data curation, formal analysis, writing—review and editing; Ł.P.: writing—review and editing; H.K.R.: conceptualization, data curation, writing—review and editing; K.N.: conceptualization, data curation, investigation, resources, visualization, writing—original draft, writing—review and editing; C.H.: conceptualization, funding acquisition, investigation, project administration, resources, writing—review and editing; O.E.C.: conceptualization, data curation, resources, supervision, writing—original draft, writing—review and editing; A.K.: conceptualization, data curation, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. This research was funded by an Estonian Research Council Personal Research Grant (grant no. PSG492) (E.O., E.R., M.T., K.J. and S.C.); a Swedish Collegium for Advanced Study Pro Futura Scientia Fellowship (E.O.); a Latvian Science Council project no. Izp-20202-0032 'Subsistence strategies and the first demographic transition at the Lake Lubāns wetland: Abora Late Neolithic settlement case study' (G.Z., V.V. and D.L.); an Alfred Kordelin Foundation funded project 'Estonian Corded Ware identity and its manifestation in a hunter-gatherer environment' (grant no. 210084) (K.N.); and a Polish National Science Centre project no. 2021/43/B/HS3/01995 'Subsistence strategies and their transformations in the Late and Final Neolithic of East-Central Europe' (Ł.P.). This project has received funding from the European Union HORIZON Coordination and Support Actions under grant agreement no. 101079396 and from Innovate UK grant no. 10063975. The project was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 695539; The Innovation, Dispersal and Use of Ceramics in northwest Eurasia).

Acknowledgements. We are very grateful to Auli Bläuer (University of Turku), Aija Macāne (University of Helsinki, University of Latvia), Eduards Plankājs (University of Latvia), Giedrė Piličiauskienė (Vilnius University), Laszlo Bartosiewicz (Stockholm University) and Daniel Makowiecki (Nicolaus Copernicus University) for information on the references for the first domesticates in the Baltic Sea region. We also wish to thank the local collections—Tallinn

University Archaeological Research Collections, University of Tartu archaeological collections, and the National History Museum of Latvia—for providing access to and permitting us to sample the archaeological material. We are grateful to Dr Helen Talbot (University of York) for her assistance with the GC-C-IRMS analysis.

References

- Moreau L. 2020 Social inequality before farming? Multidisciplinary approaches to the study of social organization in prehistoric and ethnographic hunter-gatherer-fisher societies. Cambridge, UK: McDonald Institute for Archaeological Research. (doi:10.17863/CAM.60627)
- Pospieszny Ł. 2015 Freshwater reservoir effect and the radiocarbon chronology of the cemetery in Żąbie, Poland. *J. Archaeol. Sci.* **53**, 264–276. (doi:10.1016/j.jas.2014.10.012)
- Cramp LJE *et al.* 2019 Regional diversity in subsistence among early farmers in Southeast Europe revealed by archaeological organic residues. *Proc. R. Soc. B* **286**, 20182347. (doi:10.1098/rspb.2018.2347)
- Nordqvist K. 2018 The Stone Age of north-eastern Europe 5500–1800 calBC: bridging the gap between the East and the West. PhD thesis, University of Oulu, Finland.
- Brami M, Winkelbach L, Schulz I, Schreiber M, Blöcher J, Diekmann Y, Burger J. 2022 Was the fishing village of Lepenski Vir built by Europe's first farmers? *J. World Prehist.* **35**, 109–133. (doi:10.1007/s10963-022-09169-9)
- Pääkkönen M, Holmqvist E, Bläuer A, Evershed RP, Asplund H. 2020 Diverse economic patterns in the north Baltic Sea region in the Late Neolithic and Early Metal periods. *Eur. J. Archaeol.* **23**, 4–21. (doi:10.1017/aea.2019.39)
- Barker G. 2006 *The agricultural revolution in prehistory: why did foragers become farmers?* Oxford, UK: Oxford University Press. (doi:10.1093/oso/9780199281091.001.0001)
- Zeder MA. 2011 The origins of agriculture in the Near East. *Curr. Anthropol.* **52**, S221–S235. (doi:10.1086/659307)
- Pinhasi R, Fort J, Ammerman AJ. 2005 Tracing the origin and spread of agriculture in Europe. *PLoS Biol.* **3**, e410. (doi:10.1371/journal.pbio.0030410)
- Vasks A. 2015 *No medniekiem un zvejniekiem līdz lopkopjiem un zemkopjiem: latvijas aizvēstures senākais posms (10 500.-1.g. pr. Kr.)*. Riga, Latvia: Zinātne.
- Kriiska A. 2009 The beginning of farming in the Eastern Baltic area. In *East European plain on the eve of agriculture* (eds PM Dolukhanov, Graeme R. Sarson, AM Shukurov), pp. 159–179. Oxford, UK: Archaeopress.
- Lõugas L, Kriiska A, Maldre L. 2007 New dates for the Late Neolithic Corded Ware Culture burials and early husbandry in the east Baltic region. *Archaeofauna* **16**, 21–31.
- Piličiauskas G, Jankauskas R, Piličiauskienė G, Dupras T. 2017 Reconstructing Subneolithic and Neolithic diets of the inhabitants of the SE Baltic coast (3100–2500 cal BC) using stable isotope analysis. *Archaeol. Anthropol. Sci.* **9**, 1421–1437. (doi:10.1007/s12520-017-0463-z)
- Piličiauskas G, Jankauskas R, Piličiauskienė G, Craig OE, Charlton S, Dupras T. 2017 The transition from foraging to farming (7000–500calBC) in the SE Baltic: a re-evaluation of chronological and palaeodietary evidence from human remains. *J. Archaeol. Sci. Rep.* **14**, 530–542. (doi:10.1016/j.jasrep.2017.06.004)
- Piličiauskas G, Kisieliene D, Piličiauskienė G. 2017 Deconstructing the concept of Subneolithic farming in the southeastern Baltic. *Veget. Hist. Archaeobot.* **26**, 183–193. (doi:10.1007/s00334-016-0584-9)
- Heyd V. 2022 The bumpy road towards farming in Finland – a short-essay re-appraisal. In *Oodeja Mikalle: Juhlakirja professori Mika Laventolle hänen täyttäväänsään 60 vuotta Odes to Mika: Festschrift for Professor Mika Lavento on the occasion of his 60th birthday* Оды Мике: Юбилейный сборник посвященный 60-летию профессора Мики Лавенто. (eds P Halinen, V Heyd, K Mannermaa), pp. 17–25. Helsinki, Finland: Archaeological Society of Finland.
- Mittnik A *et al.* 2018 The genetic prehistory of the Baltic Sea region. *Nat. Commun.* **9**, 442. (doi:10.1038/s41467-018-02825-9)
- Saag L *et al.* 2017 Extensive farming in Estonia started through a sex-biased migration from the steppe. *Curr. Biol.* **27**, 2185–2193.e6. (doi:10.1016/j.cub.2017.06.022)
- Allentoft ME *et al.* 2022 Population genomics of Stone Age Eurasia. *bioRxiv*. (doi:10.1101/2022.05.04.490594)
- Ahola M, Heyd V. 2020 The northern way: graves and funerary practices in Corded Ware Finland. *Praehist. Zeitsch.* **95**, 78–111. (doi:10.1515/pz-2020-0002)
- Kriiska A, Lang V, Mäesalu A, Tvauri A, Valk H. 2020 *Eesti ajalugu I. Eesti esiaeg*. Tartu, Estonia: Tartu Ülikooli ajaloo ja arheoloogia instituut.
- Furholt M. 2014 Upending a 'totality': re-evaluating Corded Ware variability in Late Neolithic Europe. *Proc. Prehist. Soc.* **80**, 67–86. (doi:10.1017/ppr.2013.20)
- Macãne A, Nordqvist K. 2021 More than just Zvejnieki: an overview of Latvian Stone Age burials. *Eur. J. Archaeol.* **24**, 299–323. (doi:10.1017/aea.2020.64)
- Heron C, Craig OE, Luquin A, Steele VJ, Thompson A, Piličiauskas G. 2015 Cooking fish and drinking milk? Patterns in pottery use in the southeastern Baltic, 3300–2400 cal BC. *J. Archaeol. Sci.* **63**, 33–43. (doi:10.1016/j.jas.2015.08.002)
- Piličiauskas G *et al.* 2020 Fishers of the Corded Ware culture in the Eastern Baltic. *Acta Archaeol.* **91**, 95–120. (doi:10.1111/j.1600-0390.2020.12223.x)
- Robson HK, Skipitytė R, Piličiauskienė G, Lucquin A, Heron C, Craig OE, Piličiauskas G. 2019 Diet, cuisine and consumption practices of the first farmers in the southeastern Baltic. *Archaeol. Anthropol. Sci.* **11**, 4011–4024 (doi:10.1007/s12520-019-00804-9)
- Kriiska A, Nordqvist K. 2021 Estonian Corded Ware culture (2800–2000 cal. BC): defining a regional group in the eastern Baltic. In *Yamnaya interactions* (eds V Heyd, G Kulcsár, B Preda-Bălănică), pp. 463–485. Budapest, Hungary: Archaeolingua.
- Lõugas L. 2006 Animals as subsistence and bones as raw material for settlers of prehistoric Zvejnieki. In *Back to the origin* (eds L Larsson, I Zagorska), pp. 75–89. Lund, Sweden: Lund University.
- Zagorska I. 2006 Radiocarbon chronology of the Zvejnieki burials. In *Back to the origin: new research in the Mesolithic-Neolithic Zvejnieki cemetery and environment, northern Latvia* (eds L Larsson, I Zagorska), pp. 91–113. Stockholm, Sweden: Almqvist & Wiksell International.
- Vasks A, Kalniņa L, Ritums R. 1999 The introduction and pre-Christian history of farming in Latvia. In *The development of the Baltic Sea and Stone Age Settlement in the Pärnu Area of Southwestern Estonia. Environmental and cultural history of the Eastern Baltic Region. PACT, 57* (eds U Miller, T Hackens, V Lang, A Raukas, S Hicks), pp. 291–304. Rixensart, Belgium: Pact Belgium.
- Reimer PJ *et al.* 2020 The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP). *Radiocarbon* **62**, 725–757. (doi:10.1017/RDC.2020.41)
- Lõugas L, Lidén K, Nelson DE. 1996 Resource utilisation along the Estonian coast during the Stone Age. In *Coastal Estonia: recent advances in environmental and cultural history. PACT 51* (eds T Hackens, S Hicks, V Lang), pp. 359–369. Rixensart, Belgium: Council of Europe.
- van Vuure T. 2014 Aurochs *Bos primigenius* Bojanus, 1827. In *Ecology, evolution and behaviour of wild cattle: implications for conservation* (eds J Burton, M Melletti), pp. 240–254. Cambridge, UK: Cambridge University Press.
- Paaver K, Птавер К. 1965 *Формирование териофауны и изменчивость млекопитающих Прибалтики в голоцене*=K. L. Paaver, *Development of teriofauna and variability of mammals of the Pribaltic territory in the holocen*. Tartu, Estonia: Akademiya nauk ESSR.
- Lepiksaar J. 1986 The Holocene history of theriofauna in Fennoscandia and Baltic countries. In *Nordic late quaternary biology and*

- ecology (ed. L-K Königsson), pp. 51–70. Uppsala, Sweden: Societas Upsaliensis pro Geologia Quaternaria.
36. Buckley M, Collins M, Thomas-Oates J, Wilson JC. 2009 Species identification by analysis of bone collagen using matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* **23**, 3843–3854. (doi:10.1002/rcm.4316)
 37. Rowley-Conwy P, Albarella U, Dobney K. 2012 Distinguishing wild boar from domestic pigs in prehistory: a review of approaches and recent results. *J. World Prehist.* **25**, 1–44. (doi:10.1007/s10963-012-9055-0)
 38. Rasiņš A, Tauriņa M. 1983 Pārskats par Latvijas PSR arheoloģiskajos izrakumos konstatētajām kultūru un nezaļu sēklām. *Arheoloģija un etnogrāfija* **14**, 152–176.
 39. Griķpēdis M, Matuzevičute GM. 2018 A review of the earliest evidence of agriculture in Lithuania and the earliest direct AMS date on cereal. *Eur. J. Archaeol.* **21**, 264–279. (doi:10.1017/ea.2017.36)
 40. Vassar A. 1939 Iru linnapära. In *Muistse eesti linnused: 1936–1938. a. Uurimiste tulemused* (ed. H Moora), pp. 53–100. Tartu, Estonia: Õpetatud Eesti Selts.
 41. Poska A, Saarse L, Veski S. 2004 Reflections of the pre- and early-agrarian human impact in the pollen diagrams of Estonia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **209**, 37–50. (doi:10.1016/j.palaeo.2003.12.024)
 42. Poska A, Saarse L. 2006 New evidence of possible crop introduction to north-eastern Europe during the Stone Age. *Veget. Hist. Archaeobot.* **15**, 169–179. (doi:10.1007/s00334-005-0024-8)
 43. Githumbi E *et al.* 2022 European pollen-based REVEALS land-cover reconstructions for the Holocene: methodology, mapping and potentials. *Earth System Sci. Data* **14**, 1581–1619. (doi:10.5194/essd-14-1581-2022)
 44. Josefsson T, Ramqvist PH, Hörnberg G. 2014 The history of early cereal cultivation in northernmost Fennoscandia as indicated by palynological research. *Veget. Hist. Archaeobot.* **23**, 821–840. (doi:10.1007/s00334-014-0446-2)
 45. Lahtinen M, Rowley-Conwy P. 2013 Early farming in Finland: was there cultivation before the Iron Age (500 BC)? *Eur. J. Archaeol.* **16**, 660–684. (doi:10.1179/1461957113Y.00000000040)
 46. Beug H-J. 2004 *Leitfaden der pollenbestimmung für mitteleuropa und angrenzende gebiete*. Munich, Germany: Friedrich Pfeil.
 47. Behre K-E. 2007 Evidence for Mesolithic agriculture in and around central Europe? *Veget. Hist. Archaeobot.* **16**, 203–219. (doi:10.1007/s00334-006-0081-7)
 48. Albert BM, Innes JB. 2020 On the distinction of pollen grains of early varieties of *Hordeum* from *Glyceria* species: addressing the early cereal cultivation problem in palynology. *Palynology* **44**, 369–381. (doi:10.1080/01916122.2019.1589593)
 49. Loze I. 2008 *Lubāna ezera mitrāja neolīta dzintars un tā apstrādes darbnīcas*. Riga, Latvia: Latvijas vestures institūta apgads.
 50. Legzdīņa D, Zariņa G. 2023 Living on the climate edge suitable for early farming: the complex subsistence strategies of the Late Neolithic people at the Abora site in Latvia. *Proc. Latvian Acad. Sci. Sect. B. Nat. Exact Appl. Sci.* **77**, 1–14. (doi:10.2478/prolas-2023-0001)
 51. DeNiro MJ. 1985 Postmortem preservation and alteration of *in vivo* bone collagen isotope ratios in relation to palaeodietary reconstruction. *Nature* **317**, 806–809. (doi:10.1038/317806a0)
 52. van Klinken GJ. 1999 Bone collagen quality indicators for palaeodietary and radiocarbon measurements. *J. Archaeol. Sci.* **26**, 687–695. (doi:10.1006/jasc.1998.0385)
 53. Tõrv M. 2018 *Persistent practices: a multi-disciplinary study of hunter-gatherer mortuary remains from c. 6500–2600 cal. BC, Estonia*. Kiel, Germany: Wachholtz, Murmann Publishers.
 54. Zagorska I. 2000 The art from Zvejnieki burial ground, Latvia. *Acta Academiae artium Vilnensis* **20**, 79–92.
 55. Eriksson G, Lõugas L, Zagorska I. 2003 Stone Age hunter–fisher–gatherers at Zvejnieki, northern Latvia. *Before Farming* **2003**, 1–25. (doi:10.3828/bfarm.2003.1.2)
 56. Rasmussen S *et al.* 2015 Early divergent strains of *Yersinia pestis* in Eurasia 5,000 years ago. *Cell* **163**, 571–582. (doi:10.1016/j.cell.2015.10.009)
 57. Meadows J, Bērziņš V, Legzdīņa D, Lūbke H, Schmöcke U, Zagorska I, Zariņa G. 2018 Stone-age subsistence strategies at Lake Burtnieks, Latvia. *J. Archaeol. Sci. Rep.* **17**, 992–1006. (doi:10.1016/j.jasrep.2016.03.042)
 58. Henderson RC, Zariņa G, Czermak A, Schulting RJ, Henderson PA, Legzdīņa D, Zagorska I, Lee-Thorp J. 2022 Life histories at stone age Zvejnieki based on stable isotope profiles of tooth dentine. *J. Archaeol. Sci. Rep.* **44**, 103496. (doi:10.1016/j.jasrep.2022.103496)
 59. Varul L, Tõrv M, Lõugas L, Kriiska A. In press. Corded Ware Culture (2800–2000 calBC) inhumations in Estonia, Northern Baltics. *Est. J. Archaeol.*
 60. Dudd SN. 1999 *Molecular and isotopic characterisation of animal fats in archaeological pottery*. Bristol, UK: University of Bristol.
 61. Courel B *et al.* 2020 Organic residue analysis shows sub-regional patterns in the use of pottery by Northern European hunter–gatherers. *R. Soc. Open Sci.* **7**, 192016. (doi:10.1098/rsos.192016)
 62. Pääkkönen M, Evershed RP, Asplund H. 2020 Compound-specific stable carbon isotope values of fatty acids in modern aquatic and terrestrial animals from the Baltic Sea and Finland as an aid to interpretations of the origins of organic residues preserved in archaeological pottery. *J. Nordic Archaeol. Sci.* **19**, 3–19.
 63. Spataro M, Oras E, Lucquin A, Bērziņš V. 2021 Hunter-fisher-gatherer pottery production and use at the Neolithic shell-midden of Rīņņukalns, Latvia. *Antiquity* **95**, 1446–1463. (doi:10.15184/aqy.2021.127)
 64. Piličauskas G, Skipitytė R, Oras E, Lucquin A, Craig OE, Robson HK. 2023 The globular amphora culture in the eastern Baltic: new discoveries. *Acta Archaeol.* **92**, 203–227. (doi:10.1163/16000390-20210037)
 65. Dudd SN, Evershed RP. 1998 Direct demonstration of milk as an element of archaeological economies. *Science* **282**, 1478–1481. (doi:10.1126/science.282.5393.1478)
 66. Evershed RP, Dudd SN, Copley MS, Mutherjee A. 2002 Identification of animal fats via compound specific $\delta^{13}C$ values of individual fatty acids: assessments of results for reference fats and lipid extracts of archaeological pottery vessels. *Docum. Praehist.* **29**, 73–96. (doi:10.4312/dp.29.7)
 67. Copley MS, Berstan R, Dudd SN, Docherty G, Mukherjee AJ, Straker V, Payne S, Evershed RP. 2003 Direct chemical evidence for widespread dairying in prehistoric Britain. *Proc. Natl Acad. Sci. USA* **100**, 1524–1529. (doi:10.1073/pnas.0335955100)
 68. Hendy J *et al.* 2018 Ancient proteins from ceramic vessels at Çatalhöyük West reveal the hidden cuisine of early farmers. *Nat. Commun.* **9**, 4064. (doi:10.1038/s41467-018-06335-6)
 69. Hansel FA, Copley MS, Madureira LAS, Evershed RP. 2004 Thermally produced ω -(*o*-alkylphenyl)alkanoic acids provide evidence for the processing of marine products in archaeological pottery vessels. *Tetrahedron Lett.* **45**, 2999–3002. (doi:10.1016/j.tetlet.2004.01.111)
 70. Cramp LJE, Evershed RP. 2014 Reconstructing aquatic resource exploitation in human prehistory using lipid biomarkers and stable isotopes. In *Treatise on geochemistry: archaeology and anthropology* (eds HD Holland, KK Turekian), pp. 319–339. Oxford, UK: Elsevier.
 71. Lucquin A, Colonese AC, Farrell TFG, Craig OE. 2016 Utilising phytanic acid diastereomers for the characterisation of archaeological lipid residues in pottery samples. *Tetrahedron Lett.* **57**, 703–707. (doi:10.1016/j.tetlet.2016.01.011)
 72. Vanhanen S, Gustafsson S, Ranheden H, Björck N, Kemell M, Heyd V. 2019 Maritime hunter-gatherers adopt cultivation at the farming extreme of Northern Europe 5000 years ago. *Sci. Rep.* **9**, 4756. (doi:10.1038/s41598-019-41293-z)
 73. Vanhanen S, Kriiska A, Nordqvist K. 2023 Corded Ware Culture plant gathering at the Narva-Jõesuu IIB settlement and burial site in Estonia. *Environ. Archaeol.*, 1–13. (doi:10.1080/14614103.2023.2216531)
 74. Alenius T, Gerasimov D, Sapelko T, Ludikova A, Kuznetsov D, Golyeva A, Nordqvist K. 2020 Human-environment interaction during the Holocene along the shoreline of the Ancient Lake Ladoga: a case study based on palaeoecological and archaeological material from the Karelian Isthmus, Russia. *The Holocene* **30**, 1622–1636. (doi:10.1177/0959683620941071)
 75. Piličauskas G, Vengalis R, Minkevičius K, Kisieliene D, Ežerinskis Ž, Šapolaitė J, Skipitytė R, Robson HK. 2021 The earliest evidence for crop cultivation during the Early Bronze Age in the southeastern Baltic. *J. Archaeol. Sci. Rep.* **36**, 102881. (doi:10.1016/j.jasrep.2021.102881)
 76. Lempiäinen T. 1999 Prehistoric cultivation in Finland - macrofossil evidence of cultivated cereals. In *The development of the Baltic Sea and Stone Age Settlement in the Pärnu Area of Southwest Estonia. Environmental and cultural history of the Eastern Baltic Region. PACT*, 57 (eds U Miller, T Hackens, V Lang, A

- Raukas, S Hicks), pp. 353–355. Rixensart, Belgium: Pact Belgium.
77. Lahtinen M, Oinonen M, Tallavaara M, Walker JWP, Rowley-Conwy P. 2017 The advance of cultivation at its northern European limit: process or event? *Holocene* **27**, 427–438. (doi:10.1177/0959683616660164)
 78. Antanaitis-Jacobs I, Stančikaitė M. 2004 The impact of the economic activities of Stone and Bronze Ages populations on their environment according to the archaeobotanical evidence. *Lietuvos Archeologija* **25**, 251–266.
 79. Piličiauskas G. 2018 *Virvelinės keramikos kultūra lietuvoje 2800–2400 cal BC*. Vilnius, Lithuania: Lietuvos Istorijos Institutas.
 80. Bläuer A, Korkeakoski-Väisänen K, Arppe L, Kantanen J. 2013 Bronze Age cattle teeth and cremations from a monumental burial cairn in Selkäkangas, Finland: new radiocarbon dates and isotopic analysis. *Eston. J. Archaeol.* **17**, 3–23. (<https://www.ceeol.com/search/article-detail?id=178330>)
 81. Ahola M, Kirkinen T, Vajanto K, Ruokolainen J. 2018 On the scent of an animal skin: new evidence on Corded Ware mortuary practices in Northern Europe. *Antiquity* **92**, 118–131. (doi:10.15184/aqy.2017.188)
 82. Löugas L. 1994 Subfossil vertebrate fauna of Asva site, Saaremaa. *Mammals. Stilis* **5**, 71–93.
 83. Antanaitis-Jacobs I, Richards M, Daugnora L, Jankauskas R, Ogrinc N. 2009 Diet in early Lithuanian prehistory and the new stable isotope evidence. *Archaeol. Baltica* **12**, 12–30.
 84. Cramp LJ *et al.* 2014 Neolithic dairy farming at the extreme of agriculture in northern Europe. *Proc. R. Soc. B* **281**, 1–9. (doi:10.1098/rspb.2014.0819)
 85. Lidén K, Eriksson G, Nordqvist B, Götherström A, Bendixen E. 2004 The wet and the wild followed by the dry and tame: or did they occur at the same time? Diet in Mesolithic - Neolithic southern Sweden. *Antiquity* **78**, 23–33. (doi:10.1017/S0003598X00092899)
 86. Craig OE *et al.* 2011 Ancient lipids reveal continuity in culinary practices across the transition to agriculture in Northern Europe. *Proc. Natl Acad. Sci. USA* **108**, 17 910–17 915. (doi:10.1073/pnas.1107202108)
 87. Robson HK, Saul H, Steele VJ, Meadows J, Otto Nielsen P, Fischer A, Heron CP, Craig OE. 2021 Organic residue analysis of Early Neolithic 'bog pots' from Denmark demonstrates the processing of wild and domestic foodstuffs. *J. Archaeol. Sci.* **36**, 102829. (doi:10.1016/j.jasrep.2021.102829)
 88. Casanova E *et al.* 2022 Dating the emergence of dairying by the first farmers of Central Europe using ¹⁴C analysis of fatty acids preserved in pottery vessels. *Proc. Natl Acad. Sci. USA* **119**, e2109325118. (doi:10.1073/pnas.2109325118)
 89. Milner N, Craig OE, Bailey GN, Pedersen K, Andersen SH. 2004 Something fishy in the Neolithic? A re-evaluation of stable isotope analysis of Mesolithic and Neolithic coastal populations. *Antiquity* **78**, 9–22. (doi:10.1017/S0003598X00092887)
 90. Soncin S *et al.* 2021 High-resolution dietary reconstruction of victims of the 79 CE Vesuvius eruption at Herculaneum by compound-specific isotope analysis. *Sci. Advances* **7**, eabg5791. (doi:10.1126/sciadv.abg5791)
 91. Artmenko, И=Артеменко ИИ. 1967 *Племена Верхнего и Среднего Поднепровья в эпоху бронзы*. Москва: Наука.
 92. Grasis N. 2007 The Skaistkalnes Selgas Double Burial and the Corded Ware/ Rzucewo Culture: a model of the culture and the development of burial practices. *Lietuvos Archeologija* **31**, 39–70.
 93. Bollongino R, Nehlich O, Richards MP, Orschiedt J, Thomas MG, Sell C, Fajkosová Z, Powell A, Burger J. 2013 2000 years of parallel societies in Stone Age Central Europe. *Science* **342**, 479–481. (doi:10.1126/science.1245049)
 94. Lazaridis I *et al.* 2014 Ancient human genomes suggest three ancestral populations for present-day Europeans. *Nature* **513**, 409–413. (doi:10.1038/nature13673)
 95. González-Fortes G *et al.* 2017 Paleogenomic evidence for multi-generational mixing between neolithic farmers and Mesolithic hunter-gatherers in the Lower Danube Basin. *Curr. Biol.* **27**, 1801–1810.e10. (doi:10.1016/j.cub.2017.05.023)
 96. Mathieson I *et al.* 2018 The genomic history of Southeastern Europe. *Nature* **555**, 197–203. (doi:10.1038/nature25778)
 97. Smith BD. 2001 Low-level food production. *J. Archaeol. Res.* **9**, 1–43. (doi:10.1023/A:1009436110049)
 98. Bulbeck D. 2013 The transition from foraging to farming in prehistory and 'ethnography'. *World Archaeol.* **45**, 557–573. (doi:10.1080/00438243.2013.821668)
 99. Denham T, Donohue M. 2022 Mapping the middle ground between foragers and farmers. *J. Anthropol. Archaeol.* **65**, 101390. (doi:10.1016/j.jaa.2021.101390)
 100. Correa-Ascencio M, Evershed RP. 2014 High throughput screening of organic residues in archaeological potsherds using direct acidified methanol extraction. *Anal. Methods* **6**, 1330–1340. (doi:10.1039/C3AY41678J)
 101. Dolbunova E *et al.* 2023 The transmission of pottery technology among prehistoric European hunter-gatherers. *Nat. Hum. Behav.* **7**, 171–183. (doi:10.1038/s41562-022-01491-8)
 102. Rieley G. 1994 Derivatization of organic compounds prior to gas chromatographic-combustion-isotope ratio mass spectrometric analysis: identification of isotope fractionation processes. *Analyst* **119**, 915–919. (doi:10.1039/AN9941900915)
 103. Welker F *et al.* 2016 Palaeoproteomic evidence identifies archaic hominins associated with the Châtelperronian at the Grotte du Renne. *Proc. Natl Acad. Sci. USA* **113**, 11 162–11 167. (doi:10.1073/pnas.1605834113)
 104. Brown S, Hebestreit S, Wang N, Boivin N, Douka K, Richter K. 2020 Zooarchaeology by Mass Spectrometry (ZooMS) for bone material – acid insoluble protocol. (doi:10.17504/protocols.io.bf43jqyn)
 105. Strohal M, Hassman M, Košata B, Kudiček M. 2008 mMass data miner: an open source alternative for mass spectrometric data analysis. *Rapid Commun. Mass Spectrom.* **22**, 905–908. (doi:10.1002/rcm.3444)
 106. Buckley M, Whitcher Kansa S, Howard S, Campbell S, Thomas-Oates J, Collins M. 2010 Distinguishing between archaeological sheep and goat bones using a single collagen peptide. *J. Archaeol. Sci.* **37**, 13–20. (doi:10.1016/j.jas.2009.08.020)
 107. Buckley M, Kansa SW. 2011 Collagen fingerprinting of archaeological bone and teeth remains from Domuztepe, South Eastern Turkey. *Archaeol. Anthropol. Sci.* **3**, 271–280. (doi:10.1007/s12520-011-0066-z)
 108. Brown S, Douka K, Collins MJ, Richter KK. 2021 On the standardization of ZooMS nomenclature. *J. Proteom.* **235**, 104041. (doi:10.1016/j.jprot.2020.104041)
 109. Brown TA, Nelson DE, Vogel JS, Southon JR. 1988 Improved collagen extraction by modified Longin method. *Radiocarbon* **30**, 171–177. (doi:10.1017/S0033822200044118)
 110. McKINNEY CR, McCREA JM, Epstein S, Allen HA, Urey HC. 1950 Improvements in mass spectrometers for the measurement of small differences in isotope abundance ratios. *Rev. Sci. Instrum.* **21**, 724–730. (doi:10.1063/1.1745698)
 111. Oras E *et al.* 2023 Parallel worlds and mixed economies: multi-proxy analysis reveals complex subsistence systems at the dawn of early farming in the northeast Baltic. Figshare. (doi:10.6084/m9.figshare.c.6845558)