



## OPEN ACCESS

## EDITED BY

Fernanda Dorea,  
Food and Agriculture Organization of the  
United Nations (Headquarters), Italy

## REVIEWED BY

Ravikanth Reddy Poonooru,  
University of Missouri, United States  
Giorgia Angeloni,  
Experimental Zooprophyllactic Institute of the  
Venezie (IZSVe), Italy  
Jean Perchet,  
World Organisation for Animal Health, France

## \*CORRESPONDENCE

Margarida Arede  
✉ margarida.decastro@uab.cat

RECEIVED 26 February 2023

ACCEPTED 30 August 2023

PUBLISHED 22 September 2023

## CITATION

Arede M, Beltrán-Alcrudo D, Aliyev J,  
Chaligava T, Keskin I, Markosyan T, Morozov D,  
Oste S, Pavlenko A, Ponea M, Starciuc N,  
Zdravkova A, Raizman E, Casal J and  
Allepuz A (2023) Examination of critical factors  
influencing ruminant disease dynamics in the  
Black Sea Basin.  
*Front. Vet. Sci.* 10:1174560.  
doi: 10.3389/fvets.2023.1174560

## COPYRIGHT

© 2023 Arede, Beltrán-Alcrudo, Aliyev,  
Chaligava, Keskin, Markosyan, Morozov, Oste,  
Pavlenko, Ponea, Starciuc, Zdravkova, Raizman,  
Casal and Allepuz. This is an open-access  
article distributed under the terms of the  
[Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).  
The use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in this  
journal is cited, in accordance with accepted  
academic practice. No use, distribution or  
reproduction is permitted which does not  
comply with these terms.

# Examination of critical factors influencing ruminant disease dynamics in the Black Sea Basin

Margarida Arede<sup>1\*</sup>, Daniel Beltrán-Alcrudo<sup>2</sup>, Jeyhun Aliyev<sup>3</sup>,  
Tengiz Chaligava<sup>4</sup>, Ipek Keskin<sup>5</sup>, Tigran Markosyan<sup>6</sup>,  
Dmitry Morozov<sup>7</sup>, Sarah Oste<sup>8</sup>, Andrii Pavlenko<sup>2</sup>, Mihai Ponea<sup>9</sup>,  
Nicolae Starciuc<sup>10</sup>, Anna Zdravkova<sup>11</sup>, Eran Raizman<sup>2</sup>, Jordi Casal<sup>1</sup>  
and Alberto Allepuz<sup>1</sup>

<sup>1</sup>Departament de Sanitat i Anatomia Animals, Facultat de Veterinària, Universitat Autònoma de Barcelona, Barcelona, Spain, <sup>2</sup>Food and Agriculture Organization of the United Nations (FAO), Regional Office for Europe and Central Asia, Budapest, Hungary, <sup>3</sup>Food Safety Agency of the Republic of Azerbaijan, Baku, Azerbaijan, <sup>4</sup>Veterinary Department, National Food Agency, Ministry of Environmental Protection and Agriculture of Georgia, Tbilisi, Georgia, <sup>5</sup>Veterinary Control Central Research Institute, Ministry of Agriculture and Forestry, Ankara, Türkiye, <sup>6</sup>Scientific Centre for Risk Assessment and Analysis in Food Safety Area, Ministry of Agriculture, Nubarashen, Yerevan, Armenia, <sup>7</sup>Vitebsk State Academy of Veterinary Medicine, Vitebsk, Belarus, <sup>8</sup>University Institute of Technology Nancy-Brabois, Lorraine University, Villers-lès-Nancy, France, <sup>9</sup>National Sanitary Veterinary and Food Safety Authority, Bucharest, Romania, <sup>10</sup>Faculty of Veterinary Medicine, State Agrarian University of Moldova, Chisinau, Moldova, <sup>11</sup>Bulgarian Agency for Food Safety, Sofia, Bulgaria

**Introduction:** Ruminant production in the Black Sea basin (BSB) is critical for national economies and the subsistence of rural populations. Yet, zoonoses and transboundary animal diseases (TADs) are limiting and threatening the sector. To gain a more comprehensive understanding, this study characterizes key aspects of the ruminant sector in nine countries of the BSB, including Armenia, Azerbaijan, Belarus, Bulgaria, Georgia, Moldova, Romania, Türkiye, and Ukraine.

**Methods:** We selected six priority ruminant diseases (anthrax, brucellosis, Crimean Congo haemorrhagic fever (CCHF), foot-and-mouth disease (FMD), lumpy skin disease (LSD), and peste des petits ruminants (PPR)) that are present or threaten to emerge in the region. Standardized questionnaires were completed by a network of focal points and supplemented with external sources. We examined country and ruminant-specific data such as demographics, economic importance, and value chains in each country. For disease-specific data, we analysed the sanitary status, management strategies, and temporal trends of the selected diseases.

**Results and discussion:** The shift from a centrally planned to a market economy, following the collapse of the Soviet Union, restructured the ruminant sector. This sector played a critical role in rural livelihoods within the BSB. Yet, it faced significant challenges such as the low sustainability of pastoralism, technological limitations, and unregistered farms. Additionally, ruminant health was hindered by informal animal trade as a result of economic factors, insufficient support for the development of formal trade, and socio-cultural drivers. In the Caucasus and Türkiye, where diseases were present, improvements to ruminant health were driven by access to trading opportunities. Conversely, European countries, mostly disease-free, prioritized preventing disease incursion to avoid a high economic burden. While international initiatives for disease management are underway in the BSB, there is still a need for more effective local resource allocation and international partnerships to strengthen veterinary health capacity, protect animal health and improve ruminant production.

## KEYWORDS

Black Sea, surveillance and control, ruminants, transboundary animal diseases, zoonoses

## 1. Introduction

Livestock production is critical for the subsistence of rural populations as a source of food, income, transportation, hides, and fertilizers, contributing to 40% of the agricultural economy worldwide (1). However, in recent decades, there has been a surge and spread of endemic and exotic diseases affecting livestock (2, 3), which significantly impact the sector and threaten public health and welfare (4). This surge has been intensified by several factors, including the high increase in international trade of animals and animal products (3), rise in intensive farming driven by higher market demands for animal protein and increasing middle-class purchasing power (5–7), changes in land use (8), shifts in migration and tourism patterns (9), and the effects of climate change (9).

TADs such as foot-and-mouth disease (FMD), lumpy skin disease (LSD), and peste des petits ruminants (PPR), along with zoonoses, particularly anthrax, brucellosis (*Brucella abortus* and *Brucella melitensis*), and Crimean Congo haemorrhagic fever (CCHF) (Table 1) are diseases that are either threatening ruminants or emerging in the Black Sea Basin (BSB). Ruminant production is the most important livestock subsector in most countries in the region, ensuring food security for rural populations and contributing significantly to national economies (16–25).

Nevertheless, key aspects linked with the dynamics of these diseases in the region remain poorly understood. Knowledge gaps include disease geographic coverage and prevalence, morbidity and mortality rates, economic impact, and risk factors influencing their spread and persistence. These gaps arise from weaknesses in a country's veterinary management programmes, which can be associated with lack of human resources (authorities, veterinarians and technicians) to sustain them, inadequate government funding for agriculture or livestock sectors, limited surveillance coverage (26), insufficient legislative action, and lack of support for implementing biosecurity measures (27). As a result, disease reporting is delayed, incomplete or biased, leading to ineffective responses to disease outbreaks (28, 29). These challenges are more pronounced in rural areas of lower to middle-income countries, as the BSB, where social inequality persists. In these regions, livestock, particularly ruminants, are ubiquitous and critical for livelihoods, and animal diseases hinder food security and the sector's development.

This study characterizes ruminant production and its importance around the BSB and describes the disease status and management efforts (i.e., surveillance and control activities) for the selected ruminant diseases (Table 1) in nine countries of the region (i.e., Armenia, Azerbaijan, Belarus, Bulgaria, Georgia, Moldova, Romania, Türkiye, and Ukraine). It also explores the most relevant factors that may influence the incursion and spread of these diseases in the region.

TABLE 1 Overview of the studied diseases.

		Disease	Agent	Main domestic host (s)	Transmission	Vaccine availability
Zoonoses	Bacterial	Anthrax (10)	<i>Bacillus anthracis</i>	All mammals	Contact with <i>B. anthracis</i> spores	Yes
		Brucella (11)	<i>Brucella abortus</i>	Cattle	Direct/indirect contact	Yes
	<i>Brucella melitensis</i>		Sheep and goats			
		CCHF (12)	CCHF virus • g. Orthonairovirus • f. Nairoviridae	Cattle, sheep, and goats	Tick-borne	No
TADs	Viral	FMD (13)	FMD virus • g. Aphthovirus • f. Picornaviridae	Cattle, sheep, goats, and swine	Direct/indirect contact	Yes
		LSD (14)	LSD virus • g. Capripoxvirus • f. Poxviridae	Cattle	Arthropod vector	Yes
		PPR (15)	Small ruminant morbillivirus • g. Morbillivirus • f. Paramixoviridae	Sheep and goats	Direct contact	Yes

g.: genus, f.: family.

## 2. Materials and methods

The current paper is a component of the GCP/GLO/074/USA project, which contributes to the broader “Global Framework for the Progressive Control of Transboundary Animal Diseases (GF-TADs)” initiative. This project targets nine countries located around the BSB, namely Armenia, Azerbaijan, Belarus, Bulgaria, Georgia, Moldova, Romania, Türkiye, and Ukraine. Herein, Armenia, Azerbaijan, and Georgia are referred to as “Caucasus,” when the statement is true for the three countries, and Türkiye is referred to as either “Thrace” or “Anatolia” when specific differences apply to each of the regions.

The primary focus of the project is on six diseases that are relevant for the region: anthrax, brucellosis, CCHF, FMD, LSD, and PPR. Consequently, this study focused on domestic ruminants (cattle, sheep and goats), which are the animal species most impacted by these diseases. These species are also interchangeably referred to as large ruminants (LR) and small ruminants (SR).

A report template was designed to collect information from each of the participating countries ([Supplementary material S1](#)). This document was developed by four authors of this paper (AA, DB-A, JC, and MA) as a semi-structured questionnaire. The selection of topics was based on the project’s objectives and aimed at addressing knowledge gaps in the BSB about the ruminant sector and the impact of the selected diseases. The initial version of the document was presented and shared with respondents from the nine participating countries during a virtual meeting. The final version of the report template accounted for edits and suggestions provided by the participants.

The report template was divided into two sections. The first section focused on the ruminant demographics, types of ruminant production, national and international trade, livestock markets, slaughterhouses, seasonal movements, and value chains. The second section focused on the six targeted diseases, requesting information on disease status, recent outbreaks, surveillance and control activities, awareness campaigns, and research activities in place.

Moreover, each report template requested information in two formats: narrative answers (e.g., description of a system or production type) and quantitative data in a database format (e.g., Excel datasheet). In some cases, quantitative data could complement descriptive information. To have high-quality figures, we requested the highest level of detail (e.g., the number of smallholder farms at the smallest administrative level) and, when applicable, exact locations (e.g., georeferenced locations of a livestock market). Further instructions prompted respondents to refer to additional documents like local veterinary authority national reports and national publications (i.e., grey literature).

One focal point (FP) of each participating country was appointed by FAO to answer the report template and collect country-specific information. FPs were carefully selected based on previous collaborations, the quality of their work, their expertise in the ruminant sector and selected diseases, and access to the data necessary for further analyses. FPs were based in each respective country and were working (or had recently worked) within relevant national institutions (e.g., veterinary services, food safety authorities, or national laboratories), during data collection. All nine FPs are co-authors of this paper.

FPs received the report template via email, filled it in with preliminary information, and iteratively and upon request, added further detail, following a back-and-forth exchange of emails and virtual meetings. Data collection was carried out by the FPs in collaboration with local peers, and all activities were coordinated with national authorities to request and obtain approval for data sharing. Data collection took place between October 2020 and December 2021.

Descriptive information and quantitative data were obtained and analysed from completed report templates. Then, data were assessed, and specific topics were selected to examine in this paper. To complement data on these topics, information was sourced from national reports and websites of the Food and Agriculture Organization of the United Nations (FAO), the World Organisation for Animal Health (WOAH), and the World Bank. To assess the economic importance of ruminant production for each country, we sourced data for the gross production value (GPV) of the main domestic production species from FAOSTAT (30). To find the proportional contribution of ruminant GPV to each country, we divided GPV for cattle, sheep, and goats, by the total GPV for all domestic species in 2020. Finally, ruminant distribution maps for ruminant populations (31–33) were sourced from FAO-NSAL (FAO’s Livestock Information, Sector Analysis and Policy) branch.

Quantitative data was managed, cleaned, harmonized, and collated in Microsoft Office Excel (2019), RStudio® (34), and analysed and visualised in Quantum GIS (35) and RStudio® (34).

## 3. Results

Selected topics from the nine participating countries were organized into two sections following the structure of the report template: (1) study region and ruminant-specific information, and (2) disease-specific information.



### 3.1. Study region and ruminant-specific information

#### 3.1.1. Study region

The main political changes and affiliations from countries of the study region between 1988 and 2021 are illustrated in [Figure 1](#). [Supplementary material S2](#) summarizes data for human and ruminant demographics, relevant economic indicators, and other characteristics of ruminant production. In 2020, most countries were classified as upper-middle-income economies, with the exceptions of Ukraine and Romania, which had a lower-middle-income economy and a high-income economy, respectively (36). The median GDP *per capita* of each region in 2020 was \$4,547 USD, ranging from \$3,725 USD in Ukraine to \$12,896 USD in Romania. For livestock production indicators, Moldova and Bulgaria had the lowest contribution to agricultural GDP at 23%, while Belarus had the highest at 57% (30). The proportion of ruminant GPV (per total domestic production species) ranged between 23% in Moldova to 92% in Belarus (30). Further details about this indicator are supplied in [Supplementary material S3](#).

#### 3.1.2. Ruminant demographics

The ruminant distribution varied significantly throughout the study region, both for LR and SR. LR heads ranged from 159,000 in



Moldova to 18 million in Türkiye, whereas SR heads were lowest in Belarus (148,000) and highest in Türkiye (54 million). Figure 2 illustrates the spatial distribution for LR and SR in the region and shows higher abundance of LR in Belarus, certain regions of Türkiye, western Georgia, and Azerbaijan, and higher number of SR in parts of Türkiye (Thrace and southeast Anatolia), Romania, and Azerbaijan. Additionally, the figures highlight lower LR populations in Ukraine, Moldova, southern Romania, and northern Bulgaria, and lower SR populations in Belarus, Ukraine, Moldova, and northern Bulgaria.

### 3.1.3. Production types

Countries classified ruminant production types using distinct terminology. To allow for comparisons, production types were grouped based on herd size and commercial purpose into smallholder and commercial farms, as defined in Supplementary material S6. In the Caucasus and Romania, over 90% of cattle farms were smallholdings, while Belarus had the highest proportion of cattle production in commercial herds. Across the

entire region, more than 75% of herds keeping sheep and goats were smallholdings.

### 3.1.4. Animal identification and registration systems

Most countries in the BSB had established National Animal Identification and Traceability Systems (NAITS). In contrast, the Caucasus had NAITSs under development, but not yet fully implemented at the time of data collection. In Azerbaijan, this system was being developed through a European Commission (EC) framework. It entered a regional pilot stage in late 2021 and began a country-wide phased implementation over 2022 (37). In Armenia, the Centre of Agribusiness and Rural Development (CARD), with support from the Austrian Development Agency (ADA) (38, 39), developed and conducted a pilot of its NAITS in the cattle sector in January 2022. Similarly, in Georgia, after a 5-year project supported by FAO and financed by the ADA and the Swiss Agency for Development and Cooperation (SDC) (40), the system was launched nationwide in February 2022 (40).



### 3.1.5. National trade of live ruminants

Recordings of live animal movements were linked to the existence of a NAITS in each country. Therefore, most countries in the region recorded these movements within a national centralized database. Each registration included information regarding the individual identification of the animal and the farm of origin, the destination farm, and a veterinary health report issued by an official veterinarian.

Conversely, Georgia did not have a recording system for animal movements. In Armenia and Azerbaijan, movements between provinces were registered, but the record consisted solely of a paper-based veterinary health certificate. These records were issued by official veterinarians and archived in regional divisions. There were no centralised databases for recording live animal movements in these three countries.

Live animal movements were characterized by a seasonal pattern that is not detailed in this paper. Nevertheless, it is worth noting that these movements significantly surged during cultural-religious celebrations such as Novruz, Kurban Bayram, and Ramadan Bayram in Azerbaijan and Türkiye, in which animals are transported to cities to be ritually slaughtered. Similarly, Easter and St George's Day in Bulgaria and Romania were also preceded by an increase in live animal movement due to the traditional consumption of mutton.

Furthermore, it is important to highlight that livestock trade was closely linked to animal density in each region, the demand for animal protein in densely populated areas, the location of slaughterhouses, and specific commercial partnerships with regions or countries. For example, in Georgia, ruminant trade primarily occurred from west to east due to the high exports to Azerbaijan. As for Bulgaria, the southern regions, where LR and SR production was more intense, also had an increased movement of ruminants. Moreover, in Türkiye, ruminants were moved from small to large provinces, and more specifically from east to west and north to south of the country.

#### 3.1.5.1. Livestock markets

The role of livestock markets in live ruminant trade varied across the region. Azerbaijan and Türkiye run ten and 150 licensed live animal markets, respectively, which played a significant role in ruminant trade. During Kurban Bayram in these two countries, markets worked exceptionally to sustain the surge in animal movements. In Armenia, Georgia, and Bulgaria, these facilities existed but were not as relevant for animal trade. In Belarus, official markets for live ruminant trade were absent, instead occasional fairs and exhibitions were held at the district level and on a small scale. In the same country, ruminant trade for breeding purposes occurred through state breeding companies. In Ukraine, smallholders used live animal markets for local ruminant trade.

#### 3.1.5.2. Seasonal movements

Pastoralism includes seasonal movements to pastures and can be sub-classified as nomadism, transhumance, or agropastoralism (definitions provided in [Supplementary material S6](#)). These practices are key to the seasonal sourcing of graze and water for livestock and were common across the study region. In Bulgaria, the Caucasus, Romania, and Türkiye, transhumant animals were moved to summer pastures, often found in mountainous areas, in spring and summer, and to lowland pastures or stables in autumn and winter. Migrating months had slight variations yearly depending on weather and pasture conditions. Georgia, Azerbaijan,

and Türkiye set up Veterinary Surveillance Points (VSP) along migration routes. These premises primarily focused on mass vaccination campaigns in Azerbaijan, but also served as rest points for supplying feed and water, as sanitary checkpoints for health status control, and anti-parasitic application in Georgia and Türkiye. The mingling of animals from various herds, regions or even neighbouring countries was common in seasonal pastures. Consequently, these animals were vaccinated either before going to pasture or during migration in VSPs. Furthermore, movements to seasonal pastures were recorded in centralized systems for movement control in Bulgaria, Romania ([41](#)) and Türkiye; however, these recordings, similarly to national movements, were not done in the Caucasus.

In Belarus, Moldova, and Ukraine, ruminants kept in smallholdings or smaller private farms in rural settings grazed seasonally in fields surrounding their holdings, in an agropastoral manner.

### 3.1.6. International trade of live ruminants

Partner trading countries with the BSB region are presented in the last two columns of [Supplementary material S2](#). International trade of live animals was done based on country partnerships, contingent on the trust in the exporting country's animal health capacity and/or the sanitary status for the main contagious zoonoses and TADs (at a specific time) ([42](#)). To guarantee disease freedom on entry into a country, imported live ruminants were accompanied by a health certificate validated by a veterinarian of the exporting country's competent authority. Particularly for the importation of live animals (and animal products) into the EU, the intra-EU trade, and EU exports of live animals, TRACES (Trade Control and Expert System) ([43](#)), an EC online platform, facilitates sanitary certification required for trade and centralizes trade information. Thus, Bulgaria and Romania along with other BSB countries exporting live animals or animal products into the EU, used this platform.


Similar to national live animal movements, international trade was influenced by cultural-religious events. Therefore, a surge in live animal imports preceded Kurban Bayram and Ramadan Bayram in Azerbaijan and Türkiye, and Easter and St George's Day in Bulgaria and Romania.

## 3.2. Disease-specific information

### 3.2.1. Disease status, surveillance, and control activities

[Figure 3](#) illustrates the country-level disease statuses for each selected disease. Countries self-classified their disease status as endemic, sporadic, or absent (definitions in [Supplementary material S6](#)). An *Absent* status was subclassified for brucellosis as "officially free" and for FMD as "officially free with or without vaccination" when WOAHP officially recognised these disease statuses.

In [Supplementary material S4](#), a table summarizes key details for the six studied diseases in each of the countries of the BSB. Moreover, temporal trends of disease outbreaks per country from 2010 to 2020 are shown in [Supplementary material S5](#). In this subsection of the results, we review the disease status and management practices applied in the region.



### 3.2.1.1. Anthrax

Anthrax was endemic or sporadic in all countries. All countries implemented passive surveillance and, upon suspicion, applied further clinical examinations, sampling, and testing. Due to the environmental nature of this disease, most national management programmes, in addition to guidelines for disease containment and carcass disposal, also included regulations for historically infected fields (e.g., signalling, fencing, digging restrictions, and awareness campaigns). Vaccination was compulsory for all ruminants in Azerbaijan, and Moldova, and all LR in Armenia. A risk-based vaccination approach was applied for all ruminants in Bulgaria, Georgia, Romania, Türkiye, and Ukraine, and exclusively for SR kept or moved to high-risk areas in Armenia. In Belarus, anthrax vaccination was not conducted.

### 3.2.1.2. Brucellosis

Brucellosis was endemic in the Caucasus and Türkiye, sporadic in Bulgaria, and absent in all other countries of the BSB. In 2012, Belarus was officially recognised by WOAHP as brucellosis-free, and to maintain this status, serosurveillance was conducted every three years. Moreover, surveillance was exclusively passive for Georgia, and active and risk-based in all other countries. In most of the BSB, passive surveillance for brucellosis was associated with the report and investigation of abortions in ruminants, which is a syndrome of this disease, but not exclusive to *Brucella* spp. infection. In Azerbaijan, Georgia, and Türkiye, vaccination for brucellosis was mandatory for all ruminants and performed at the same time as serosurveillance. Brucellosis vaccination was not part of the national veterinary control plan in Belarus, Bulgaria, Moldova, Ukraine, or Romania.

### 3.2.1.3. Crimean Congo haemorrhagic fever

CCHF was endemic in Georgia and Türkiye. These countries applied control measures upon outbreak identification, focusing on tick control and community awareness campaigns. These activities

comprised the application of acaricide sprays to ruminants, including during seasonal migrations from early spring to late autumn, and environmental tick elimination. Educational campaigns in Türkiye promoted contact restriction between livestock and wildlife, and tick management. These campaigns were included in the state budget at no cost to farmers. For the remaining countries, CCHF had never been reported in ruminants and there was no national surveillance programme in place. At the time of data collection, no licensed vaccine was available for CCHF in ruminants.

### 3.2.1.4. Foot-and-mouth disease

The WOAHP official FMD status varied between the two regions of Türkiye: Anatolia was classified as endemic, and Thrace held FMD-free status with vaccination. FMD was sporadic in Armenia, absent in Georgia and Azerbaijan, while in all other countries, WOAHP recognised the official status FMD-free without vaccination.

FMD surveillance was active in the Caucasus and Türkiye, as well as in regions of Bulgaria and Romania. The countries of the Caucasus were collaborating with EuFMD through the Progressive Control Pathway for Foot and Mouth Disease (PCP-FMD) to design and establish risk-based surveillance programmes. As part of these efforts, they implemented NSP (Non-Structural Protein) and SP (Structural Protein) serosurveys to evaluate the FMD virus circulation, seroconversion, and vaccination coverage. In regions bordering Thrace, Bulgaria conducted risk-based serosurveys on a sample of ruminants every three months. While in Romania, surveillance focused on clinical examination of LR and SR on high-density premises (e.g., live animal markets, exhibitions, ports, and airports), serosurveillance of all ruminants close to international borders, and SR upon their arrival from seasonal pastures.

In Türkiye, the FMD management programme in 2021 aimed to achieve FMD-free status without vaccination in Thrace and FMD-free status with vaccination in Anatolia by 2025 (44). In

Thrace, control measures comprised suspect FMD case culling, restrictions on live animal imports from Anatolia, and strict adherence to sanitary legislation. In Anatolia's southeast provinces bordering FMD-endemic countries, surveillance activities were enhanced and risk-based. Moreover, in case of an FMD outbreak, Türkiye conducted a field investigation, and vaccination, established a *cordon sanitaire*, animal quarantine, and thorough cleaning and disinfection, organized training, and awareness campaigns, and closely monitored all premises within a 10 km radius of the event.

FMD vaccination varied throughout the BSB. Türkiye vaccinated LR twice a year, and SR once a year only in Thrace. In case of an outbreak in Anatolia, SR were also vaccinated in established protection and surveillance zones. In Azerbaijan, LR were vaccinated twice a year (spring and autumn) and SR once a year, while Armenia, applied the same strategy only in high-risk areas. Since 2017, Georgia has conducted vaccination exclusively in high-risk areas, based on risk assessments, which considered seasonal migration, international borders with FMD-endemic countries, live animal markets, and informal trade.

#### 3.2.1.5. Lumpy skin disease

LSD was sporadic in Armenia, Azerbaijan, and Türkiye, and absent in all other countries. Surveillance activities varied: clinical examination was conducted in Belarus to a sample of LR in spring and summer, and in the six regions of Bulgaria bordering Thrace monthly. Georgia had active participatory surveillance, Türkiye implemented both active and passive surveillance activities, and all other countries only applied passive surveillance. Compulsory vaccination was practised nationwide in Azerbaijan, Bulgaria, and Türkiye, and in high-risk areas of Armenia and Georgia. Vaccination was not applied in Belarus, Moldova, Romania, or Ukraine.

#### 3.2.1.6. Peste des petits ruminants

PPR was endemic in Türkiye and absent in all other countries. In March 2021, Thrace was granted the classification of “PPR-protected area.” PPR surveillance varied across the BSB: Belarus did not conduct it, Moldova, Ukraine, and Anatolia exclusively applied passive surveillance, while Thrace and all other countries applied active surveillance. In Bulgaria, areas previously affected by PPR (2018 outbreak) implemented enhanced surveillance, and regions bordering Thrace applied risk-based serosurveillance on a sample of SR every two months. In Romania, active surveillance included clinical inspection of a sample of SR herds before and after pasture season.

Vaccination was implemented in Georgia, following the first PPR occurrence in 2016. In Türkiye vaccination was conducted, yet it ceased in Thrace after the region was granted a “PPR-protected area” classification in March 2021. This measure, coupled with strict live SR movement restrictions from Anatolia to Thrace, aimed at Thrace's application for WOAHP zonal freedom status in 2023. In Anatolia, PPR vaccines were applied to all newborn SR and unvaccinated adults.

## 4. Discussion

In this paper, we summarized the ruminant production sector and reviewed the sanitary status and management of six diseases affecting ruminants (anthrax, brucellosis, CCHF, FMD, LSD, and PPR) in the

BSB. Furthermore, we explored key factors contributing to the introduction and spread of these diseases in the region.

### 4.1. Post-Soviet Union reform

The fall of the Soviet Union caused a deterioration of public infrastructures and services across the former Soviet Union (FSU) and Communist Bloc countries, significantly affecting agricultural and livestock sectors (18, 45–48). In BSB countries, except Belarus, changes included the shift from collective and state-owned farms to private ownership, removal of government subsidies to the livestock sector (49), closure of large slaughterhouses (50), and depletion in resource allocation to veterinary services (51). Such factors left livestock production in the hands of unspecialized farmers, and unsupervised by veterinary services (52), resulting in increased disease incidence (52, 53). Thereafter, the region suffered a steep decline in the number of ruminants (49, 54) and, in some countries, as Ukraine and Belarus, a significant abandonment of agricultural lands (55). These abrupt structural changes were followed by a transition phase with gradual agricultural recovery and increasing productivity (56). Yet, rural poverty, particularly in the Caucasus and Moldova, persists and requires new and efficient policy measures that enable technological development and access to market channels and services (51). EU's farmer association model could aid smallholders of the FSU to actively engage to improve their marketing, input supplies, and support services (57).

### 4.2. Rural livelihoods and pastoralism

Pastoralism played a critical role in rural areas in most countries of the BSB (16–18, 20–25, 46, 58–60), creating a unique interdependence between ruminants, farmers, and the environment (61, 62). Preserving this practice is crucial, given its resilience to severe climates in arid and inhospitable areas, socio-cultural importance, and the potential opportunities brought to younger generations (62). However, its sustainability in the BSB is a matter of concern. Ageing rural farmers show reluctance to adopt new technologies and measures to improve animal production and health, and the mass migration of younger populations to urban centres leaves families without essential support for farming activities (57). Moreover, they have limited access to veterinary services also caused by ageing rural veterinarians, and difficulties in attracting young graduates due to low-income prospects and prevailing urban migration trends (57). These factors result in underperforming veterinary services (63) and high costs for disease management impeding improvements, even when advancements are made at higher levels (28). Solutions for these challenges need to be explored, as building private veterinary capacity and developing training programmes for veterinary paraprofessionals.

In addition, initiatives addressing pastoralism's limited sustainability and its associated risks to ruminant health and welfare are underway (64, 65). In Georgia (62), Türkiye, and Azerbaijan, VSPs were established along migration routes. In Armenia, the “Project Coordination Platform for Sustainable Management of Natural Grazing Lands – Pastures and Grasslands” was launched to address pasture management-related problems in the country (66). Internationally, the Pastoralist Knowledge Hub (PKH) by FAO aids



the development of synergies for dialogue and pastoralist development, while an extension of the Performance of Veterinary Services (PVS) (67) evaluation tool prioritizes finding solutions to control animal diseases in pastoralist areas (63). Collectively, these initiatives aim to foster and protect pastoralism while ensuring its sustainability.

### 4.3. Disease management and related factors

Disease management in the Caucasus and Türkiye was often inefficient. Nonetheless, improvements were being made in the field. Particularly in the Caucasus, the full operability of the NAITs is expected to make disease management programmes (68) and disease traceability (69) more efficient. As a result, these improvements will positively influence animal health and ruminant production, ultimately, leading to better trade opportunities and economic growth in these countries.

Sociocultural-religious events in Türkiye and Azerbaijan prompted the implementation of contingency plans and extraordinary measures, which, at times, proved inefficient in preventing disease introduction and spread. In fact, the epidemiological investigation conducted upon the PPR incursion to Bulgaria in July 2018 concluded that the high demand and resulting price difference of mutton between Bulgaria and Thrace during these festivals contributed to increased informal movements of people and animals (70).

In the BSB, only Türkiye reported the presence of all studied diseases. This can be attributed to its unique conditions, including a large ruminant population, vast geographical area with socio-economic disparities, and extensive rural regions. Moreover, its shared borders with six countries including Syria and Iraq, where social unrest leads to informal movement of people with their livestock, create a significant pathway for disease spread (71, 72). Recognising the high risk to animal and public health through this route, Türkiye introduced legislative acts for border control and supervision of the main roads (73, 74). These acts are open to amendment, and they aim to manage and identify informal/illegal trade for livestock and products of animal origin, along with enforcing animal culling. Following the implementation of these controls, a national report highlighted a significant reduction of nearly 95% and 50% of confiscated smuggled animals and animal products, respectively, caught during border controls conducted in 2011 and 2018. This demonstrates the successful impact of these actions (75).

EU countries, such as Bulgaria and Romania, had high resource allocation for disease management and prioritised the prevention of disease incursion to reduce economic losses. These countries followed harmonized live animal trade regulations set by the EC, enforcing additional control measures and trade restrictions (76) in the event of an exotic disease incursion. Therefore, responses to Bulgaria's FMD (2011), LSD (2016), and PPR (2018) outbreaks were quick and intensive (77). And LSD and FMD outbreaks resulted in an economic burden estimated at €8 million (78), and \$1.5 billion USD annually (79), respectively. To prevent disease re-emergence and further economic losses, disease management activities established upon these events, were still in place as of 2021.

Moreover, Thrace's proximity to Europe and shared borders with the EU through Bulgaria and Greece, prompted the establishment of partnership programmes between the EU and Türkiye. These

initiatives involved significant investments to curb disease introduction and spread into Europe, while also promoting trade opportunities (80, 81). Therefore, FMD and PPR statuses varied between Thrace and Anatolia, leading to distinct classifications by WOA, along with distinct approaches for disease management and movement control in these two regions.

### 4.4. The exception of CCHF

Türkiye and Georgia were the only countries in the BSB reporting the presence of CCHF in ruminants. In spite of this, past studies identified CCHF virological or serological evidence and the presence of competent vectors in most countries of the study region, except for Belarus (12, 82–84), while CCHF human cases were also notified in Bulgaria and Türkiye (84). Non-reporting of CCHF in ruminants was linked to two factors. Firstly, its exclusion from national veterinary programmes resulted in the absence of routine official surveys, and secondly, the subclinical nature of the disease in these species allows it to circulate unnoticed (85–87). Nevertheless, domestic ruminants play an important role in the epidemiology of the disease as they are involved in its vector life cycle (83) and amplification and spread of the virus (88, 89). Moreover, ruminant CCHFV antibody titers correlate with virus presence in a region (84), as well as human disease incidence (82). Given CCHF's public health threat, including potential human incurred deaths, the prudent course of action is to include the disease in national veterinary programmes. This would ensure regular disease monitoring and prompt response to any reported cases.

### 4.5. Current initiatives

Achieving effective disease management requires not only efficient resource allocation for national disease preparedness and response but also promoting collaborations with other countries and unions. An initiative that strengthens regional alliances for TADs management is the GF-TADs, a joint FAO and WOA effort, created to support capacity building and the establishment of disease management programmes based on regional priorities (77). GF-TADs' priority diseases in the BSB include brucellosis, FMD, LSD, and PPR. Under this initiative, the Global Strategy for the Control and Eradication of PPR aims to control and eradicate PPR and strengthen veterinary services (90, 91). Additionally, FAO's PCP-FMD guides endemic countries in progressively managing FMD risks and reducing its impacts and viral circulation (92–94).

### 4.6. Limitations

Study limitations were linked to country-specific factors and data quality issues. Absent or not fully operable NAITs are likely to have affected data validity, and completeness is limited due to unregistered herds along with underreporting across the BSB. Underreporting is often linked to farmers' poor disease awareness, distrust in governmental authorities, risk of penalty or stigmatization, or at a higher level, lack of capacity to enforce regulations (95) and low transparency. Additionally, variability in data availability and spatial



resolutions between countries led to reduced accuracy of certain indicators or made it impossible to compare and examine others. Finally, data quality might have been affected by resource reallocation during the COVID-19 pandemic, which partly coincided with the two-year data collection period.

## 4.7. The armed conflict in Ukraine

The armed conflict in Ukraine, starting in February 2022, had a significant impact on its livestock sector. Since its beginning, the conflict led to decreased agricultural production due to land abandonment, animal losses from death or forced slaughtering, and reduced demand for meat and milk due to mass emigration (96). It has disrupted the accessibility to veterinary services, vaccines, medication (97), and critical inputs, such as feed and fodder (96), compromising disease prevention and control, and increasing the risk of stress, malnourishment, and susceptibility to disease in livestock. Moreover, amongst security issues, unavailability of consumables and equipment, and competing urgent priorities, appropriate carcase disposal became challenging. These effects are expected to reshape ruminant demographics, its associated production sector, and value chains, particularly in front-line regions. International cooperation is vital to address the consequences on livestock health and revive the sector post-conflict. Guidelines aiming to support the livelihoods of livestock-keeping communities in humanitarian emergencies that affect livestock are in place (98, 99), being used to alleviate the consequences of the presented conflict.

## 5. Conclusion

This study provides a comprehensive overview of the ruminant production sector and the management of six major diseases of concern in the BSB. By examining the effects of the post-soviet reform, the importance of pastoralism, differences in disease management and countries' response to disease incursion, as well as the influence of cultural events and political affiliations on live animal trade, we have gained a valuable understanding of how these different factors work together to determine disease dynamics in the region.

Unlike the other studied diseases, CCHF was not included in veterinary management plans, and not surveyed in ruminants across the region, presenting a public health threat. Furthermore, the armed conflict in Ukraine starting after data collection will likely have a significant impact on ruminant production and animal disease emergence in this country, with potential spread to neighbouring countries.

Finally, despite recent developments in veterinary infrastructures, including the implementation of NAITs in the Caucasus, substantial support from international agencies and targeted initiatives for ruminant disease management, the need to improve animal health persists, particularly in rural and remote regions. A thorough understanding of the primary challenges, needs, and constraints faced by smallholders in each specific country context is essential. Establishing priorities and closely assessing them in collaboration with farmers, national stakeholders, and international agencies, will aid in identifying opportunities for

more effective disease management strategies contributing to alleviating and preventing future outbreak scenarios. These considerations go hand in hand with providing incentives for rural development, by seeking financial aid, efficiently allocating financial and human resources, and most importantly ensuring the sustainability of the implemented strategies.

## Data availability statement

The datasets presented in this article are not readily available for public access due to restrictions imposed by the governmental entities of the nine participating countries. The data that support the findings of this study were used under license from these entities and are subject to confidentiality and legal constraints. Requests to access the datasets should be directed to [margarida.decastro@uab.cat](mailto:margarida.decastro@uab.cat).

## Ethics statement

Ethical review and approval was not required for the study of animals in accordance with the local legislation and institutional requirements.

## Author contributions

AA, DB-A, JC, and MA developed the report template and contributed to the conception and design of the study. JA, TC, IK, TM, DM, AP, MP, NS, and AZ were hired as country FPs to fill in the report template and provide quantitative data and descriptive information for the project. SO wrote a document summarizing preliminary information. MA performed the data management, descriptive analysis and visualization materials, and wrote the first draft of the manuscript. All authors contributed to the article and approved the submitted version.

## Funding

The project GCP/GLO/074/USA "Global Framework for the Progressive Control of Transboundary Animal Diseases (GF-TADs)" was funded by the United States Department of Defense, Defense Threat Reduction Agency (DTRA). The content of the information does not necessarily reflect the position or the policy of the Federal Government of the United States, and no official endorsement should be inferred.

## Acknowledgments

The authors acknowledge the national authorities from the nine countries for providing all the data and documents on ruminant demographics and diseases. We also acknowledge Giuseppina Cinardi from FAO-NSAL for her contribution in generating GLW 4 maps for ruminants distribution. These maps, presented in Figure 2, were adjusted to incorporate data obtained during the project.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fvets.2023.1174560/full#supplementary-material>

## References

1. FAO. The state of food and agriculture. Livestock in the balance; (2009). Available at: <https://www.fao.org/3/a-i0680e.pdf>
2. Berezowski J, Lindberg A, Ward M. Surveillance against the odds: addressing the challenges of animal health surveillance. *Prev Vet Med.* (2015) 120:1–3. doi: 10.1016/j.PREVETMED.2015.02.020
3. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in emerging infectious diseases. *Nature.* (2008) 451:990–3. doi: 10.1038/nature06536
4. Cartin-Rojas A. Transboundary animal diseases and international trade In: V Bobek (editor), *International trade from economic and policy perspective*. InTech; (2012) 143–166.
5. Rushton J. Improving the use of economics in animal health – challenges in research, policy and education. *Prev Vet Med.* (2017) 137:130–9. doi: 10.1016/j.PREVETMED.2016.11.020
6. Popkin BM, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev.* (2012) 70:3–21. doi: 10.1111/j.1753-4887.2011.00456.x
7. HLPE. *Sustainable agricultural development for food security and nutrition: what roles for livestock? A report by the High Level Panel of Experts on Food Security and Nutrition of the Committee on World Food Security*, Rome. (2016).
8. Magouras I, Brookes VJ, Jori F, Martin A, Pfeiffer DU, Dürr S. Emerging zoonotic diseases: should we rethink the animal–human interface? *Front Vet Sci.* (2020) 7:748. doi: 10.3389/fvets.2020.582743/BIBTEX
9. Myers SS. Planetary health: protecting human health on a rapidly changing planet. *Lancet.* (2017) 390:2860–8. doi: 10.1016/S0140-6736(17)32846-5
10. World Organization for Animal Health (WOAH). 3.1.1 Anthrax. In: WOAH, ed. *Terrestrial Manual 2023*. 12th ed. WOAH; (2023). Available at: [https://www.woah.org/fileadmin/Home/eng/Health\\_standards/tahm/3.01.01\\_ANTHRAX.pdf](https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.01.01_ANTHRAX.pdf)
11. World Organization for Animal Health (WOAH). 3.1.4 Brucellosis (infection with *Brucella abortus*, *B. melitensis* and *B. suis*). In: WOAH, ed. *Terrestrial Manual 2022*. 11th ed. WOAH; (2022). Available at: [https://www.woah.org/fileadmin/Home/eng/Health\\_standards/tahm/3.01.04\\_BRUCELLOSIS.pdf](https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.01.04_BRUCELLOSIS.pdf)
12. World Organization for Animal Health (WOAH). 3.1.5 Crimean–Congo haemorrhagic fever. In: WOAH, ed. *Terrestrial Manual 2023*. 12th ed. WOAH; (2023). [https://www.woah.org/fileadmin/Home/eng/Health\\_standards/tahm/3.01.05\\_CCHF.pdf](https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.01.05_CCHF.pdf)
13. World Organization for Animal Health (WOAH). 3.1.8 Foot and mouth disease (infection with foot and mouth disease virus). In: WOAH, ed. *Terrestrial Manual 2022*. 11th ed. WOAH; (2022). [https://www.woah.org/fileadmin/Home/eng/Health\\_standards/tahm/3.01.08\\_FMD.pdf](https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.01.08_FMD.pdf)
14. World Organization for Animal Health (WOAH). 3.4.12 Lumpy skin disease. In: WOAH, ed. *Terrestrial Manual 2023*. 12th ed. WOAH; (2023). [https://www.woah.org/fileadmin/Home/eng/Health\\_standards/tahm/3.04.12\\_LSD.pdf](https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.04.12_LSD.pdf)
15. World Organization for Animal Health (WOAH). 3.8.9 Peste des petits ruminants (infection with small ruminant morbillivirus). In: WOAH, ed. *Terrestrial Manual 2021*. 10th ed. WOAH; (2021). [https://www.woah.org/fileadmin/Home/eng/Health\\_standards/tahm/3.08.09\\_PPR.pdf](https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.08.09_PPR.pdf)
16. Dreve V, Călin I, Bazgă B. Analysis on the evolution of Romanian sheep and goat sector after EU accession. *Scientific Papers Series D. Animal Science* (2016) 59.
17. FAO. *Smallholders and family farms in Georgia – country study report 2019* Budapest (2020). doi: 10.4060/ca9822en
18. Punjabi, M. Developing the sheep value chain in Azerbaijan – Vision 2025. Rome: FAO. (2020).
19. Rukhkyan L. *Country report on the state of the Armenian animal genetic resources*. Food and Agriculture Organization. (2003). Available at: <https://www.fao.org/3/a1250e/annexes/CountryReports/Armenia.pdf>
20. General Directorate of Agricultural Research and Policies. *Turkey Country Report on Farm Animal Genetic Resources*. Food and Agriculture Organization. (2004). Available at: <https://www.fao.org/3/a1250e/annexes/CountryReports/Turkey.pdf>
21. Sen O, Ruban S, Getya A, Nesterov Y. Current state and future outlook for development of the milk and beef sectors in Ukraine. In: A Kuipers, G Keane and A Rozstalnyy, eds. *Cattle Husbandry in Eastern Europe and China*. EAAP Scien. Wageningen Academic Publishers; (2014):169–80.
22. Tarashevych A. *Ukraine – livestock and products annual, 2019 USDA Foreign Agricultural Service report*. Kyiv: Office of Agricultural Affairs. (2019). Available at: [https://apps.fas.usda.gov/newgainapi/api/report/downloadreportbyfilename?filename=Livestock%20and%20Products%20Annual\\_Kiev\\_Ukraine\\_8-30-2019.pdf](https://apps.fas.usda.gov/newgainapi/api/report/downloadreportbyfilename?filename=Livestock%20and%20Products%20Annual_Kiev_Ukraine_8-30-2019.pdf)
23. Kuznetsova A. Prospects for the Development of the Dairy Industry in the Republic of Belarus and in the Russian Federation. In: *Hradec Economic Days*. University of Hradec Kralove, (2020).
24. Stankov K. Economic efficiency analysis of dairy cattle farms in Bulgaria. *Trakia J Sci.* (2015) 13:226–32. doi: 10.15547/tjs.2015.s.01.038
25. FAO. *Smallholders and family farms in the Republic of Moldova – country study report 2019*. Budapest (2020).
26. Randolph TF, Schelling E, Grace D, Nicholson CF, Leroy JL, Cole DC, et al. Invited review: role of livestock in human nutrition and health for poverty reduction in developing countries. *J Anim Sci.* (2007) 85:2788–800. doi: 10.2527/jas.2007-0467
27. Can MF, Altuğ N. Socioeconomic implications of biosecurity practices in small-scale dairy farms. *Vet Q.* (2014) 34:67–73. doi: 10.1080/01652176.2014.951130
28. Goutard FL, Binot A, Duboz R, Rasamoelina-Andriamanivo H, Pedrono M, Holl D, et al. How to reach the poor? Surveillance in low-income countries, lessons from experiences in Cambodia and Madagascar. *Prev Vet Med.* (2015) 120:12–26. doi: 10.1016/j.PREVETMED.2015.02.014
29. Worsley-Tonks KEL, Bender JB, Deem SL, Ferguson AW, Fèvre EM, Martins DJ, et al. Strengthening global health security by improving disease surveillance in remote rural areas of low-income and middle-income countries. *Lancet Glob Health.* (2022) 10:e579–84. doi: 10.1016/S2214-109X(22)00031-6
30. FAO. FAOSTAT. Published (2022). Available at: <https://www.fao.org/faostat/en/#data/QV> (Accessed July 5, 2022).
31. Gilbert, M, Cinardi, G, Da Re, D, Wint, WGR, Wisser, D, and Robinson, TP. Global sheep distribution in 2015 (5 minutes of arc), *Harvard Dataverse, V1.* (2022). doi: 10.7910/DVN/VZ0YHM
32. Gilbert, M, Cinardi, G, Da Re, D, Wint, WGR, Wisser, D, and Robinson, TP. 2022, Global goats distribution in 2015 (5 minutes of arc), *Harvard Dataverse, V1.* (2022). doi: 10.7910/DVN/YYG6ET
33. Gilbert, M, Cinardi, G, Da Re, D, Wint, WGR, Wisser, D, and Robinson, TP. Global cattle distribution in 2015 (5 minutes of arc), *Harvard Dataverse, V1.* (2022). doi: 10.7910/DVN/LHBICE
34. RStudio Team. *RStudio: Integrated development environment for R*. Boston, MA: RStudio, PBC, (2022). Available at: <http://www.rstudio.com/>
35. QGIS.org. QGIS Geographic Information System. Open Source Geospatial Foundation Project. (2022). <http://qgis.org>
36. World Bank Open Data. GDP per capita (current US\$) – Armenia, Azerbaijan, Georgia, Belarus, Bulgaria, Moldova, Romania, Ukraine, Turkey | World Bank. Data. (2022). Available at: <https://data.worldbank.org/indicator/NY.GDP.PCAP.CD?locations=AM-AZ-GE-BY-BG-MD-RO-UA-TR-ES-PT> (Accessed March 24, 2022).
37. Trend news agency. Azerbaijan plans to apply animal identification and registration system soon. *Aznews.* (2021). Available at: <https://www.azernews.az/business/182785.html> (Accessed October 2, 2022).
38. Center for Agribusiness and Rural Development (CARD). Holding and animal identification and registration in Armenia. CARD, Yerevan, Armenia.

- (2018). Available at: <http://card.am/en/categories/3/projects/14> (Accessed October 2, 2022).
39. Austrian Development Agency. Holding and animal identification and registration (HAI&R) in Armenia: Preparation – strengthening of the animal health and animal breeding and administrative capacities. Vienna, Austria: Austrian Development Agency, (2018). Available at: <https://www.entwicklung.at/en/projects/detail-en/holding-and-animal-identification-and-registration-hair-in-armenia-preparation-strengthening-of-the-animal-health-and-animal-breeding-and-administrative-capacities> (Accessed October 2, 2022).
40. FAO. *FAO in Georgia - Georgia's National Animal Identification and Traceability System implemented successfully*. (2022). Available at: <https://www.fao.org/georgia/news/detail-events/es/c/1470714> (Accessed May 24, 2022).
41. EC. REGULATION (EC) no 1760/2000 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 17 July 2000 – establishing a system for THE identification and registration of bovine animals and regarding the labelling of beef and beef Products and repealing Council Regulation (EC) no 820/97. Available at: <https://faolex.fao.org/docs/pdf/eur20924.pdf>; (2020).
42. World Organization for Animal Health (WOAH). *Handbook on Import Risk Analysis for Animals and Animal Products*, Volume 1. 2nd ed. (WOAH, ed.). WOAH; (2010). [https://rr-africa.woah.org/wp-content/uploads/2018/03/handbook\\_on\\_import\\_risk\\_analysis\\_-\\_oie\\_-\\_vol\\_1.pdf](https://rr-africa.woah.org/wp-content/uploads/2018/03/handbook_on_import_risk_analysis_-_oie_-_vol_1.pdf)
43. EC. TRACES - About TRACES (trade control and expert system). European Commission website (2023). Available at: [https://food.ec.europa.eu/animals/traces\\_en#more-information](https://food.ec.europa.eu/animals/traces_en#more-information) (Accessed August 15, 2023).
44. EuFMD vlearning. Fifth online training course for foot and mouth disease research in Turkish. EuFMD. (2021). Available at: [https://eufmdlearning.works/pluginfile.php/85731/mod\\_page/content/16/FITC\\_TUR\\_Opening\\_Webinar.pdf](https://eufmdlearning.works/pluginfile.php/85731/mod_page/content/16/FITC_TUR_Opening_Webinar.pdf)
45. Meurs M, Bogushev A. Forward to the past? Agricultural restructuring in Bulgaria. *Méditerranée*. (2008) 110:93–104. doi: 10.4000/mediterranee.545
46. Department of Animal Husbandry, Agriculture M of, Armenia R of. *The National Strategy for sustainable use and development of farm animal genetic resources in the Republic of Armenia* Routledge (2013).
47. Csaki C, Lerman Z, Nucifora A. Land Reform and Private Farming in Moldova - EC4NR Agriculture Policy Note No. 9. Washington, DC: World Bank. (1997). Available at: [https://www.researchgate.net/publication/304936403\\_Land\\_Reform\\_and\\_Private\\_Farming\\_in\\_Moldova](https://www.researchgate.net/publication/304936403_Land_Reform_and_Private_Farming_in_Moldova) (Accessed May 27, 2022)
48. Vasile E, Bălan M, Mitran D, Croitoru I. The restructuring of Romanian agriculture. *Rom Agric Res*. (2011) 28:263–70.
49. Economic Research Service/USDA. *The agricultural sector before and after the breakup of the Soviet Union*; (2002).
50. Torgerson PR, Oguljahan B, Muminov AE, Karaeva RR, Kuttubaev OT, Aminjanov M, et al. Present situation of cystic echinococcosis in Central Asia. *Parasitol Int*. (2006) 55:S207–12. doi: 10.1016/j.parint.2005.11.032
51. Zjalic, M. Remarks and recommendations of the workshop. In K. J. Peters, A. Kuipers, M. G. Keane and A. Dimitriadou (Eds.), *The cattle sector in Central and Eastern Europe - Developments and opportunities in a time of transition*. Wageningen: Wageningen Academic Publishers. (2009). 175–8.
52. Hotez PJ, Alibek K. Central Asia's hidden burden of neglected tropical diseases. *PLoS Negl Trop Dis*. (2011) 5:e1224. doi: 10.1371/JOURNAL.PNTD.0001224
53. Kralacik I, Abdullayev R, Asadov K, Ismayilova R, Baghirova M, Ustun N, et al. Changing patterns of human Anthrax in Azerbaijan during the post-soviet and preemptive livestock vaccination eras. *PLoS Negl Trop Dis*. (2014) 8:e2985. doi: 10.1371/JOURNAL.PNTD.0002985
54. Rozstalnyy A, Hoffmann I, Mack S. Dairy sector challenges and perspectives in central and Eastern Europe In: *The cattle sector in Central and Eastern Europe: Developments and opportunities in a time of transition*, vol. 10 (2009). 17.
55. Lesiv M, Schepaschenko D, Moltchanova E, Bun R, Dürauer M, Prishchepov AV, et al. Spatial distribution of arable and abandoned land across former Soviet Union countries. *Sci Data*. (2018) 5:1–12. doi: 10.1038/sdata.2018.56
56. Lerman, Z. Privatisation and Changing Farm Structure in the Commonwealth of Independent States. In: S. Gomez y Paloma, S. Mary, S. Langrell and P Ciaian. (eds) *The Eurasian Wheat Belt and Food Security*. Cham: Springer. (2017).
57. Millns J. *Agriculture and rural cooperation examples from Armenia, Georgia and Moldova – FAO-REU report*. Published online (2013).
58. SZUCS T. *Detailed assessment report on the cattle farming conditions in Azerbaijan*. FAO Regional Office for Europe and Central Asia. (2020).
59. Kyssa, I, Lutaye, D, Halubets, L, Babenkov, V, Yakubets, Y, and Popov, M. 14. Efficiency, cooperative aspects and prospects for the dairy and beef sectors in Belarus. In: A Kuipers, A Rozstalnyy and G Keane, eds. *Cattle Husbandry in Eastern Europe and China*. Wageningen: Wageningen Academic Publishers, (2014), 181–190.
60. Nykolyuk, O., Pyvovar, P., Chmil, A., Bogonos, M., Topolnycky, P., Cheban, I., et al. *Agricultural markets in Ukraine: current situation and market outlook until 2030*, EUR 30874 EN, Publications Office of the European Union, Luxembourg, (2021).
61. FAO. *Livestock and agroecology. How they can support the transition towards sustainable food and agriculture*. Food and Agriculture Organization. Rome. (2018) Available at: <https://www.fao.org/publications/card/fr/c/18926EN/>
62. Eloit, M. Editorial: Pastoralism and sanitary challenges. *Bull l'OEI*. (2018) 2018:1–3. doi: 10.20506/bull.2018.2.2863
63. Zinsstag, J, Abakar, MF, Ibrahim, M, Tschopp, R, Crump, L, Bonfoh, B, et al. Cost-effective control strategies for animal and zoonotic diseases in pastoralist populations. *WOAH. Rev Sci Tech*. (2016) 35:673–81. doi: 10.20506/rst.35.2.2548
64. Calkins CM, Scasta JD. Transboundary animal diseases (TADs) affecting domestic and wild African ungulates: African swine fever, foot and mouth disease, Rift Valley fever (1996–2018). *Res Vet Sci*. (2020) 131:69–77. doi: 10.1016/j.rvsc.2020.04.001
65. Daversa DR, Fenton A, Dell AI, Garner TWJ, Manica A. Infections on the move: how transient phases of host movement influence disease spread. *Proc R Soc B Biol Sci*. (1869) 2017:20171807. doi: 10.1098/rspb.2017.1807
66. Ayzvayan V. Study on pasture management issues and their causality in the Republic of Armenia. Strategic Development Agency. Yerevan, Republic of Armenia. (2019). Available at: <https://arot.am/wp-content/uploads/2020/09/Study-on-Pasture-Management-Issues-English-pages-2-81.pdf>
67. WOAH. PVS pathway – OIE tool for the evaluation of performance of veterinary services. (OIE PVS Tool), 7th Ed. WOAH, Paris, France. (2019). Available at: <https://www.woah.org/en/what-we-offer/improving-veterinary-services/pvs-pathway/> (Accessed October 17, 2022)
68. WOAH. General principles on identification and traceability of live animals, chapter 4.2 In: *Terrestrial animal health code* (2007)
69. Congressional Research Service. *Animal identification and traceability: overview and issues*. Published online 2010.
70. De Clercq K, Cetre-Sossah C, Métras R. Mission of the community veterinary emergency team to Bulgaria – PPR incursion (16–21 July 2018), EC presentation by CVET experts and DG SANTE officials. (2018). Available at: [https://ec.europa.eu/food/sites/food/files/animals/docs/reg-com\\_ahw\\_20180919\\_pres\\_ppr\\_cvet\\_bul.pdf](https://ec.europa.eu/food/sites/food/files/animals/docs/reg-com_ahw_20180919_pres_ppr_cvet_bul.pdf)
71. Angeloni G, Guardone L, Buono N, Lazzaro E, Ivascu CM, Rizk AA, et al. A methodology to identify socio-economic factors and movements impacting on ASF and LSD in rural and insecure areas (Poster). In: *Epizone 14th Annual Meeting*, Barcelona: New Horizons, New Challenges, (2022).
72. Grace D, Little P. Informal trade in livestock and livestock products. *Rev Sci Tech*. (2020) 39:183–92. doi: 10.20506/RST.39.1.3071
73. Ministry of Food Agriculture and Livestock General Directorate of Food and Control. 5996 law on veterinary services plant health food and feed agricultural Laws; (2010). Available at: <http://www.lawsturkey.com/law/5996-law-on-veterinary-services-plant-health-food-and-feed> (Accessed October 24, 2022).
74. Ministry of Food Agriculture and Livestock General Directorate of Food and Control. Circular on the liquidation of illegal livestock and animal products no. 2012/06; Ankara, Türkiye: Ministry of Food Agriculture. (2012). Available at: [https://webcache.googleusercontent.com/search?q=cache:Y130AL\\_E-vwJ:https://www.tarimorman.gov.tr/Belgeler/Mevzuat/Genelgeler/kacak\\_canli\\_hayvan\\_hayvansal\\_urunlern\\_tasfiyesi.pdf&cd=1&hl=en&ct=clnk&gl=es](https://webcache.googleusercontent.com/search?q=cache:Y130AL_E-vwJ:https://www.tarimorman.gov.tr/Belgeler/Mevzuat/Genelgeler/kacak_canli_hayvan_hayvansal_urunlern_tasfiyesi.pdf&cd=1&hl=en&ct=clnk&gl=es) (Accessed October 24, 2022).
75. Ministry of Agriculture and Forestry. Preventing the entry of smuggled animals and animal products across borders. *2018 Activity Report*. Ankara, Türkiye: Ministry of Agriculture and Forestry. (2018) p 47. Available at: [https://www.tarimorman.gov.tr/SGB/Belgeler/Bakanl%20i%20%20k\\_Faaliyet\\_Raporlar%20i/2018%20ACTIVITY%20REPORT.pdf](https://www.tarimorman.gov.tr/SGB/Belgeler/Bakanl%20i%20%20k_Faaliyet_Raporlar%20i/2018%20ACTIVITY%20REPORT.pdf)
76. EC. Live animal movements: Movements within the union and entry into the EU. European Commission website. (2022). Available at: [https://ec.europa.eu/food/animals/live-animal-movements\\_en](https://ec.europa.eu/food/animals/live-animal-movements_en) (Accessed May 24, 2022).
77. FAO-WOAH. The Global Framework for the Progressive Control of Transboundary Animal Diseases (GF-TADs); (2004). <https://www.gf-tads.org/>
78. Casal J, Allepuz A, Miteva A, Pite L, Tabakovsky B, Terzievski D, et al. Economic cost of lumpy skin disease outbreaks in three Balkan countries: Albania, Bulgaria and the former Yugoslav Republic of Macedonia (2016–2017). *Transbound Emerg Dis*. (2018) 65:1680–8. doi: 10.1111/TBED.12926
79. Knight-Jones TJD, Rushton J. The economic impacts of foot and mouth disease – what are they, how big are they and where do they occur? *Prev Vet Med*. (2013) 112:161–73. doi: 10.1016/j.pvetmed.2013.07.013
80. Motta P, Garner G, Hövari M, Alexandrov T, Bulut A, Fragou IA, et al. A framework for reviewing livestock disease reporting systems in high-risk areas: assessing performance and perceptions towards foot and mouth disease reporting in the Thrace region of Greece, Bulgaria and Turkey. *Transbound Emerg Dis*. (2019) 66:1268–79. doi: 10.1111/TBED.13143
81. EFSA Panel on Animal Health and Welfare. Scientific opinion on foot-and-mouth disease in Thrace. *EFSA J*. (2012) 10:4. doi: 10.2903/J.EFSA.2012.2635
82. Fillâtre P, Revest M, Tattevin P. Crimean-Congo hemorrhagic fever: an update. *Med Mal Infect*. (2019) 49:574–85. doi: 10.1016/j.medmal.2019.09.005
83. Spengler JR, Bergeron É, Rollin PE. Seroepidemiological studies of Crimean-Congo hemorrhagic fever virus in domestic and Wild animals. *PLoS Negl Trop Dis*. (2016) 10:e0004210. doi: 10.1371/JOURNAL.PNTD.0004210
84. Mertens M, Schuster I, Sas MA, Vatanserver Z, Zubalek Z, Güven E, et al. Crimean-Congo hemorrhagic fever virus in Bulgaria and Turkey. *Vector Borne Zoonotic Dis*. (2016) 16:619–23. doi: 10.1089/VBZ.2016.1944

85. Whitehouse CA. Crimean-Congo hemorrhagic fever. *Antivir Res.* (2004) 64:145–60. doi: 10.1016/J.ANTIVIRAL.2004.08.001
86. Ergönül Ö. Crimean-Congo haemorrhagic fever. *Lancet Infect Dis.* (2006) 6:203–14. doi: 10.1016/S1473-3099(06)70435-2
87. Hoogstraal H. The epidemiology of tick-borne Crimean-Congo hemorrhagic fever in Asia, Europe, and Africa. *J Med Entomol.* (1979) 15:307–417. doi: 10.1093/JMEDENT/15.4.307
88. Zeller HG, Cornet JP, Camicas JL. Experimental transmission of Crimean-Congo hemorrhagic fever virus by west African wild ground-feeding birds to *Hyalomma marginatum rufipes* ticks. *Am J Trop Med Hyg.* (1994) 50:676–81. doi: 10.4269/AJTMH.1994.50.676
89. Bente DA, Forrester NL, Watts DM, McAuley AJ, Whitehouse CA, Bray M. Crimean-Congo hemorrhagic fever: history, epidemiology, pathogenesis, clinical syndrome and genetic diversity. *Antivir Res.* (2013) 100:159–89. doi: 10.1016/J.ANTIVIRAL.2013.07.006
90. FAO, WOA. Global strategy for the control and eradication of PPR; (2015). Available at: <http://www.fao.org/3/I4460E/i4460e.pdf> (Accessed October 3, 2022).
91. Legnardi M, Raizman E, Beltran-Alcrudo D, Cinardi G, Robinson T, Falzon LC, et al. Peste des Petits ruminants in central and eastern Asia/West Eurasia: epidemiological situation and status of control and eradication activities after the first phase of the PPR global eradication Programme (2017–2021). *Animals (Basel).* (2022) 12:2030. doi: 10.3390/ANI12162030
92. FAO, WOA, GF-TADs. The progressive control pathway for foot and mouth disease control (PCP-FMD). Published online 2018. Available at: <https://www.fao.org/3/a-an390e.pdf>
93. Basiladze V. The PCP-FMD progress in Georgia and how it advances FAST control. In: EUDMD OS22; (2022). Available at: <https://www.slideshare.net/eufmd1/v-basiladze-the-pcpfmd-progress-in-georgia-and-how-it-advances-fast-control>
94. Aliyeva T, Aliyev J, Hajiyeva A, Vatani M, Ch Suleymanova. Comparison of fmd serosurveillance results in azerbaijan during 2016 – 2019. In: EufMD OS20; (2020).
95. Halliday J, Daborn C, Auty H, Mtema Z, Lembo T, Bronsvort BMC, et al. Bringing together emerging and endemic zoonoses surveillance: shared challenges and a common solution. *Philos Trans R Soc Lond B Biol Sci.* (2012) 367:2872–80. doi: 10.1098/RSTB.2011.0362
96. FAO. *Ukraine: Impact of the war on agricultural enterprises, findings of a Nationwide survey of agricultural enterprises with land up to 250 hectares, January–February 2023*; (2023).
97. FAO. *Impact of the Ukraine-Russia conflict on global food security and related matters under the mandate of the food and agriculture Organization of the United Nations* FAO (2022).
98. FAO In: P Ankers, S Bishop, S Mack and K Dietze, editors. *Livestock-related interventions during emergencies*, vol. 18: FAO Animal Production and Health Manual (2016)
99. Jode, H, and Watson, C. (eds.) *LEGS - Livestock Emergency Guidelines and Standards* 3rd ed. Rugby, UK: Practical Action Publishing. (2023).