

This is an Accepted Manuscript for Parasitology. This version may be subject to change during the production process. DOI: 10.1017/S0031182023000902

Molecular and morphological studies on *Contracaecum rudolphii* A and *C. rudolphii* B in great cormorants (*Phalacrocorax carbo sinensis*) from Italy and Israel

Monica Caffara<sup>1#</sup>, Perla Tedesco<sup>1#</sup>, Nadav Davidovich<sup>2</sup>, Silva Rubini<sup>3</sup>, Valentina Luci<sup>1</sup>, Alessia Cantori<sup>1</sup>, Patrycja Anna Glogowski<sup>1</sup>, Maria Letizia Fioravanti<sup>1</sup> Andrea Gustinelli<sup>1</sup>

<sup>1</sup> Department of Veterinary Medical Sciences (DIMEVET), *Alma Mater Studiorum* University of Bologna, Italy; <sup>2</sup> Israeli Veterinary Services, Israel; <sup>3</sup> Experimental Zooprophyllactic Institute of Lombardy and Emilia Romagna, Brescia, Italy

**Corresponding author:** Monica Caffara, E-mail: [monica.caffara@unibo.it](mailto:monica.caffara@unibo.it)

# these authors contributed equally

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

## Abstract

The distribution of parasites is shaped by a variety of factors, among which the migratory movements of their hosts. Israel has a unique position to migratory routes of several bird species leaving Europe to winter in Africa, however detailed studies on the parasite fauna of birds from this area are scarce. Our study investigates occurrence and distribution of sibling species among *Contracaecum rudolphii* complex in *Phalacrocorax carbo sinensis* from Italy and Israel, to acquire further information on the geographical range of these species to gain deeper knowledge on the ecology of these parasites and their bird host. A total of 2383 *Contracaecum* were collected from the gastric mucosa of 28 great cormorants (18 from Israel and 10 from Italy). A subsample was processed for morphological analyses in light and scanning electron microscopy (SEM), and for molecular analyses through amplification and sequencing of the ITS rDNA and the *cox2* mtDNA, and through PCR-RFLP.

All the 683 *Contracaecum* subjected to molecular identification belonged to *C. rudolphii* s.l., (300 *C. rudolphii* A and 383 *C. rudolphii* B). SEM micrographs provided, for the first time, details of taxonomic structures in male specimens from both sibling species, and the first SEM characterization of *C. rudolphii* B.

This work presents the first data on the occurrence of sibling species of *C. rudolphii* in Israel and provides additional information on the distribution of *C. rudolphii* A and B in Italy, confirming the high prevalence and intensity of infection observed in *Ph. carbo sinensis* from other Italian areas.

**Keywords:** *Contracaecum rudolphii* A; *C. rudolphii* B; taxonomy; *Phalacrocorax carbo sinensis*; Israel; Italy.

## Introduction

Members of the species complex *Contracaecum rudolphii* parasitize mainly cormorants (family Phalacrocoracidae) worldwide, with six sibling species displaying characteristic geographic distribution and host preference. Particularly, *C. rudolphii* A has been described in the great cormorant *Phalacrocorax carbo sinensis* (Mattiucci *et al.*, 2002; 2020; Amor *et al.*, 2020; Carmeno *et al.*, 2022; Cammilleri *et al.*, 2023), in the European shag *Ph. aristotelis aristotelis* (Abollo *et al.*, 2001) and *Ph. aristotelis desmarestii* (Roca-Geronès *et al.*, 2023) from Europe; *C. rudolphii* B parasitizes *Ph. carbo sinensis* (Mattiucci *et al.*, 2002; 2020; Amor *et al.*, 2020; Carmeno *et al.*, 2022; Cammilleri *et al.*, 2023) from Europe; *C. rudolphii* C is reported in the double-crested cormorant *Ph. auritus* from the USA (D'Amelio *et al.*, 2007), while *C. rudolphii* D and *C. rudolphii* E are reported in *Ph. carbo* and *Ph. varius* from Australia (Shamsi *et al.*, 2009). Moreover *C. rudolphii* F is reported in the brown pelican *Pelecanus occidentalis* (family Pelecanidae) from the Gulf of Mexico (D'Amelio *et al.*, 2012).

The distribution of these sibling species is shaped by the feeding ecology and migratory movements of their definitive hosts. Cormorants are piscivorous birds, feeding on a wide variety of marine, brackish and freshwater fish. The continuous ingestion of fish paratenic hosts results, in many occasions, in massive infections in cormorants, which are sometimes associated with severe gastric lesions (Rokicki *et al.*, 2011). In European cormorants, the sibling species *C. rudolphii* A and B are reported, with *C. rudolphii* sp. A having a life cycle more adapted to brackish and marine ecosystems, while *C. rudolphii* B occurring mainly in freshwater ecosystems (Mattiucci *et al.*, 2020; Roca-Geronès *et al.*, 2023). In central and eastern Europe, both species are found, often in mixed infections, in *Ph. carbo sinensis* (Mattiucci *et al.*, 2002), which are thought to feed in different environments during their migration (Frederiksen *et al.*, 2018). The great cormorant *Ph. carbo sinensis* is a cosmopolitan species widely distributed in all continents (Battisti *et al.*, 2008; Davidovich *et al.*, 2023 in

press) and its diet is essentially represented by fish.

In Italy, larval stages of *C. rudolphii s.l.* are reported from a number of fish species; particularly *C. rudolphii* A is found in fish from brackish and marine ecosystems, including European seabass *Dicentrarchus labrax* (Paggi *et al.*, 1998; Culurgioni *et al.*, 2014; Guardone *et al.*, 2020; Mattiucci *et al.*, 2020), gilthead sea bream *Sparus aurata* (Guardone *et al.*, 2020), common sole *Solea solea* and, gobies *Gobius niger* and *G. paganellus* (Culurgioni *et al.*, 2014); several studies also document its occurrence in European eel *Anguilla anguilla* from brackish waters and coastal lagoons (Paggi *et al.*, 1998; Culurgioni *et al.*, 2014; Dezfuli *et al.*, 2016; Mattiucci *et al.*, 2020). Conversely, *C. rudolphii* B has been reported in fish from freshwater environments, including chub *Squalius cephalus*, barbel *Barbus barbus*, goldfish *Carassius carassius* and big-scale sand smelt *Atherina boyeri* sampled from river and lake systems in central Italy (Mattiucci *et al.*, 2020), and in common bream *Abramis brama* and in European carp *Cyprinus carpio* from other parts of Europe (Molnár *et al.*, 2019). Interestingly, the absence of *C. rudolphii* B in fish sampled in brackish water was recently reported (Mattiucci *et al.*, 2020), which strengthens the hypothesis that this sibling species has a life cycle adapted to freshwater ecosystems.

With respect to adult stages, parasitological data on the occurrence of sibling species of *C. rudolphii* in great cormorants are available from different areas, particularly from brackish and freshwater ecosystems of northeastern and central Italy (Li *et al.*, 2005; Mattiucci *et al.*, 2020), from coastal brackish water ponds in Sardinia (Amor *et al.*, 2020), from freshwater ecosystems of the pre-Alpine area (Carmeno *et al.*, 2022), and from coastlines and the waterways of different regions in southern Italy (Cammilleri *et al.*, 2023).

Geographically, Israel has a unique position with respect to migratory routes of many bird species that winter in Africa, with tens of thousands of piscivorous birds also staying and over-wintering in Israel (Nemtzov, 2002). The only *Contracaecum* species described so far in

Israel in birds are *C. micropapillatum*, *C. quadripapillatum*, *C. gibsoni* and, *C. multipapillatum* E (Caffara *et al.*, 2023); larval stages of *C. quadripapillatum* and *C. multipapillatum* E have also been described in fish (Davidovich *et al.*, 2022; 2023), while *C. multipapillatum s.l.* has been reported by Smirnov *et al.* (2021). To the author's knowledge, no information on the distribution of sibling species of *C. rudolphii* in piscivorous birds or other intermediate/paratenic hosts from Israel are available so far.

This study aimed at investigate the occurrence and distribution patterns of *C. rudolphii* A and B in *Ph. carbo sinensis* collected in Italy and Israel, to acquire further information about the geographical range of these sibling species and to gain deeper knowledge on the ecology of these parasites and their bird host.

## Materials and Methods

### *Contracaecum* sampling

Two thousand three hundred eighty-three nematodes of the genus *Contracaecum* were collected from the gastric mucosa of 28 great cormorant (*Phalacrocorax carbo sinensis*): 18 birds were shot from seven localities in Israel under permits of the Israel Nature and Parks Authority 2020/42659 and 2021/42855; while 10 birds were found dead from 3 localities in Italy. In Figure 1 (Fig. 1A, B) the number of birds collected in each locality of both Countries are reported.

The nematodes were washed in saline and preserved in 70% ethanol for morphological and molecular analyses. For some adults, the anterior and posterior portions were preserved in 10% neutral buffered formalin for SEM.

### *Molecular study*

For molecular analysis, genomic DNA was extracted from 683 adult males by a fast DNA

extraction method using Chelex®100 (Sigma-Aldrich, Darmstadt, Germany) (Caffara *et al.*, 2023). The ITS rDNA was amplified with primers NC5\_f (5'-GTAGGTGAACCTGCGGAAGGATCATT-3') and NC2\_r (5'-TTAGTTTCTTCCTCCGCT-3') (Zhu *et al.*, 1998) and then 10 µl were digested with the restriction endonucleases *MspI* (*C. rudolphii* cut= 700-300 bp) and then with *NsiI* to distinguish *C. rudolphii* A (cut = ~ 850-49 bp) and B (cut = uncut) (modified from Zhu *et al.*, 2007). A fragment of the *cox2* mtDNA was also amplified with primers 211\_f (5'-TTTTCTAGTTATATAGATTGRTTTYAT-3') and 210\_r (5'-CACCAACTCTTAAAATTATC-3') of Mattiucci *et al.* (2008) following the same protocol. From 53 specimens, both the ITS rDNA and *cox2* mtDNA were sequenced with an ABI 3730 DNA analyzer (StarSEQ, Mainz, Germany) after purification by Nucleo-Spin Gel and PCR Clean-up (Mackerey-Nagel, Düren, Germany). The DNA trace files were assembled with Contig Express (VectorNTI Advance 11 software, Invitrogen, Carlsbad, CA, USA), and the consensus sequences of the ITS rDNA and *cox2* mtDNA were compared with published data by BLAST tools (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Multiple sequence alignments were performed using BioEdit 7.2.5 (Hall, 1999), p-distance and maximum-likelihood (ML) tree (GTR+G+I substitution model for ITS, bootstrap of 1,000 replicates) were obtained using MEGA 7 (Kumar *et al.*, 2016). The ITS1 and ITS2 rDNA sequences were concatenated (after deleting the 5.8S rDNA) and used to build a ML tree together with the sequences of *Contracecum* spp. reported by Mattiucci *et al.* (2020) and *Ascaris suum* (MH030604) as outgroup. The *cox2* mtDNA gene was also aligned with the sequences reported by Mattiucci *et al.* (2020), plus *Pseudoterranova ceticola* (DQ116435) and *Anisakis pegreffii* (MT912471) as outgroups. The phylogenesis was performed by the Bayesian analysis (BI) with MrBayes 3.2.7a software (Ronquist *et al.*, 2012), with GTR+G model, four heated Markov chains runs for 200000 generations with sampling frequency set at 500, discarding the first 25% of the samples from the cold chain. Posterior probabilities were estimated to assess support for each

branch (significant support > 0.90). To infer the population genetics of *C. rudolphii* A and *C. rudolphii* B, 52 sequences newly generated plus 89 sequences of the two siblings from Italy, Spain and one from Poland, retrieved from GenBank were aligned and analyzed by DnaSP V6.12.03 (Rozas *et al.*, 2017): number of haplotypes (hn), diversity of haplotypes (hd), private haplotype (ph), and nucleotide diversity were determined. TCS network of haplotypes was constructed by PopART (Clement *et al.*, 2000).

The sequences generated in this study have been deposited in GenBank under accession numbers OR263194-OR263246 (ITS rDNA) and OR269666-OR269717 (*cox2* mtDNA).

### *Morphological study*

For morphological study 93 males and 13 females, randomly selected, were observed under a dissection microscope to evaluate gross morphology and to record total length (TL), then by light microscope (Leica Microsystems, Wetzlar, Germany) with the aid of a digital Nikon DS-Fi1 camera and image-acquisition software (Nikon Nis-Elements D3.0). The central part of the worms, devoid of taxonomic informative features was removed for DNA extraction. Anterior and posterior portions of the parasite body were clarified in Amman's lactophenol to measure internal structures. Morphometric analysis was carried out following Yamaguti (1935), Hartwich (1964), and Baruš *et al.* (1978).

For scanning electron microscopy (SEM), anterior and posterior portions of male and female specimens of genetically identified *C. rudolphii* A and *C. rudolphii* B were dehydrated through a graded ethanol series, dried in hexamethyldisilazane, sputter-coated with gold palladium, and observed using a Phenom XL G2 Desktop SEM (Thermo Fisher Scientific, Eindhoven, The Netherlands) operating at 5 kV.

## Results

### *Molecular analyses*

All the 683 *Contracaecum* subjected to PCR-RFLP confirmed they belong to *C. rudolphii s.l.* and, 300 were members of the sibling species *C. rudolphii* A (*Nsi*I = 840-49 bp) while 383 were *C. rudolphii* B (*Nsi*I = uncut) (Fig. 2). Table 1 report the distribution of the two-sibling species together with the sampling locality for both countries under study. In most cases we detected mixed infections between the two-sibling species, especially in cormorants from Italy.

All the sequences of the ITS rDNA of 53 specimens were of good quality and the BLAST search returned 99-100% similarity with *C. rudolphii* A (20 specimens) and *C. rudolphii* B (33 specimens). Among each sibling species the sequences were identical to each other and showed a p-distance of 0.2% between A and B-F and 0.1-0.2% between B and A-F. Interestingly the alignment of the siblings A and B newly obtained in the present study together with the other sibling species in the *C. rudolphii* complex showed the presence of an indel represented by an insertion of “GTTCGTGTG” in all but not in *C. rudolphii* B. The ML tree showed a well resolved branch (99%) with a cluster containing *C. rudolphii s.l.*, with *C. rudolphii* B basal to all the rest of the sibling species of the complex (Fig. 3).

Concerning the *cox2* mtDNA gene, 52 sequences were newly generated (20 *C. rudolphii* A and 32 *C. rudolphii* B). Among *C. rudolphii* A the p-distance was 0-0.2%, while among *C. rudolphii* B was 0-0.4%; between the two siblings were 0.8-0.9% as with *C. rudolphii* F (the only *cox2* available). Moreover, the alignment of the two siblings showed the presence of several transitions: 4 A/G, 11 G/A, 1 C/T and 1 T/C (*C. rudolphii* A / *C. rudolphii* B), that are the only variations observed between the two siblings. The BLAST search gave 99-100% similarity with the two siblings respectively. The BI inference phylogenetic tree obtained indicate that *C. rudolphii* A and *C. rudolphii* B form two distinct clades with high probability values (Fig. 4) including the same siblings retrieved from GB (MK496476 and MK496482



respectively) and with *C. rudolphii* F more related to *C. rudolphii* B. Similarly, the other *Contracecum* species form a well-supported clade separated from the abovementioned siblings.

Regarding the genetic diversity between the two siblings, the number of haplotypes obtained among 141 sequences (83 *C. rudolphii* A + 58 *C. rudolphii* B) from Italy, Israel, and Spain (plus one from Poland) was 106, of which 92 were private haplotypes with a haplotype's diversity of 0.99 ( $\pm 0.003$ ) and a nucleotide diversity of 0.044 ( $\pm 0.0013$ ). Concerning the haplotypes frequency, the most represented is the Hap12 of the *C. rudolphii* A haplogroup shared by 7 specimens from all the geographical regions (Italy, Israel, and Spain). The second most representative group of haplotypes are Hap17-18 and 80, all belonging to *C. rudolphii* A haplogroup, containing 5 mixed geographical haplotypes each except the latter (Hap80) composed only by specimens from Sardinia. Finally, the haplotypes 3 and 14 contain *C. rudolphii* B from Italy and Israel. In any case the two siblings (A and B) never mixed together. Analyzing the genetic diversity among each sibling, we observed in *C. rudolphii* A 60 haplotypes out of 83 sequences analyzed, most of them private, with a haplotype's diversity of 0.984 ( $\pm 0.006$ ) and a nucleotide diversity of 0.0105 ( $\pm 0.00062$ ); while among *C. rudolphii* B haplotypes were 46 out of 58 sequences with a similar haplotype's diversity of  $0.985 \pm 0.008$  and a nucleotide diversity of 0.0170 ( $\pm 0.0014$ ). The distribution of the haplotypes frequencies between the two siblings is reported in Fig. 5 (see also supplementary material, S1 and S2) as a complex web of haplotypes composed by a double system of star-like network: one star encloses the haplogroup A while the other contain the haplogroup B, separated by 26 mutation. No clear geographical differentiation in the distribution of the haplogroups has been detected.

### *Morphological descriptions*

Morphological analysis allowed to identify all adult males (n = 93) and females (n = 13)

collected as *C. rudolphii* s.l. Morphometric features of specimens identified as *C. rudolphii* A and *C. rudolphii* B by PCR-RFLP and sequencing are reported in Table 2.

Main morphological details of *C. rudolphii* s.l. are as follows.

Adults with transversely striated cuticle, more marked at anterior end, forming a conspicuous cephalic collar (Figs. 6A,B; 7C,D; 8A,B; 9A); three well developed lips; two pyriform cephalic papillae on dorsal lip (Figs. 6A; 7C), one cephalic papilla on each subventral lip (Figs. 6B; 8A; 9A,B); interlabia well developed, with large base and bilobed tip (Fig. 8B); excretory pore opening at base of ventral interlabium (Fig. 6A); oesophagus muscular, with small globular ventricle; ventricular appendix directed posteriorly; intestinal cecum two to three times longer than ventricular appendix, directed anteriorly.

Males with tail conical, curved at tip (Figs. 6F, 8D), with 27–40 pairs of pre-cloacal papillae, forming 2 subventral lines, 2 pairs of paracloacal papillae, 2 pairs of distal subventral papillae and 2 pairs of distal sublateral papillae, 1 pair of phasmids (Figs. 6F, 8D, F); spicules sub-equal, folded, with longitudinal alae and pointed tips (Figs. 6D,E; 8D,E).

Females larger than males; vulva around second quarter of body length (Figs. 7E; 9E); tail conical (Figs. 7F; 9F), with rounded tip. Eggs subspherical.

## Discussion

The lack of morphological features useful to discriminate among the *C. rudolphii* A and *C. rudolphii* B confirms the need of coupling the traditional parasitological observations with the molecular approach based at least on two molecular markers, e.g. ITS rDNA and *cox2* mtDNA to reach the correct identification.

All adults collected from the *Ph. carbo sinensis* sampled in Italy and Israel belong to *Contraecaecum rudolphii* s.l. confirming great cormorant as the main definitive host for this complex of species in EU (Mattiucci *et al.*, 2020; Amor *et al.*, 2020; Carmeno *et al.*, 2022;

Cammilleri *et al.*, 2023) as well as in Israel, where *C. rudolphii s.l.* had never been described.

The molecular analyses (PCR-RFLP and sequencing) performed on 683 adults from 28 *Ph. carbo sinensis* (10 from Italy and 18 from Israel) allowed the identification of the sibling species *C. rudolphii* A and *C. rudolphii* B, often in mixed infections (Table 1). The co-occurrence of both species in the same bird host has been already reported in other studies (Szostakowska and Fagerholm, 2007; Szostakowska *et al.*, 2012; Amor *et al.*, 2020; Mattiucci *et al.*, 2020; Carmeno *et al.*, 2021; Cammilleri *et al.*, 2023); similar distributions have been observed also in our study. In Italian specimens, the two siblings showed a similar proportion (184 A vs 195 B), while the birds from Israel were more frequently infected with *C. rudolphii* B (116 A vs 188 B) with only four birds out of 18 examined showing a mixed infection.

The phylogenetical analyses of both genetic markers showed a well-supported separation between *C. rudolphii* A and *C. rudolphii* B, with *C. rudolphii* B more closely related to *C. rudolphii* F than to *C. rudolphii* A in the *cox2* (Amor *et al.*, 2020; Mattiucci *et al.*, 2020; Roca-Geronès *et al.*, 2023), but not in ITS rDNA. Unfortunately, except for *C. rudolphii* F, no *cox2* mtDNA sequences are available for the other siblings (C, D, E) included in *C. rudolphii* complex to better clarify the relationship among the complex even by this more evolving gene.

Concerning the population structure, we observed a high genetic variability as demonstrated by the high numbers of haplotypes in both *C. rudolphii* A (83 sequences/60 haplotypes) and *C. rudolphii* B (58 sequences/46 haplotypes) populations; similar results were obtained also by Amor *et al.* (2020) in Sardinia (Italy) for *cox2* mtDNA (*C. rudolphii* A: n. haplotypes 33 out of 158,  $hd=0.985$ , nucleotide diversity=0.013; *C. rudolphii* B: nucleotide diversity 7/22,  $hd=0.952$ , nucleotide diversity =0.021). In Spain, Roca-Geronès *et al.* (2023) analyzed only *C. rudolphii* A, that was considered as single population due to the low genetic diversity (n. haplotypes 40/56,  $hd=0.969$ , nucleotide diversity=0.00681). In the same study, the comparisons with the sequences of *C. rudolphii* A from 2 Italian areas showed similar values

as obtained in the present study (Tyrrhenian: n. haplotypes 80/110,  $hd=0.986$ , nucleotide diversity  $=0.00871$ ; western Sardinia: n. haplotypes 28/33,  $hd=0.979$ , nucleotide diversity  $=0.0117$ ). The haplotypes diversity in any case reflect the geographical distributions of the sequences analyzed (Amor *et al.*, 2020; Roca-Geronès *et al.*, 2023). Amor *et al.* (2020) speculate about the low genetic diversity among the two siblings, suggesting that the high infection rates in definitive hosts slow down the genetic drift, but also it could be due to the parasites dispersal model mediated by the host dispersal dynamic. The migrations of the definitive hosts and the population structure could influence the genetic structure of this Anisakidae, as reported by Cipriani *et al.* (2022) for *Anisakis* spp.

The SEM observations allowed for the first time a detailed characterization of the external features of *C. rudolphii* B, and added further morphological information about *C. rudolphii* A, which was already examined using SEM by Abollo *et al.* (2001). Particularly, we provide the first detailed SEM characterization of the spicule morphology in both sibling species, with special reference to their folded appearance and to the morphology of the distal extremity.

Overall, compared to other siblings of the species complex (Abollo *et al.*, 2001; Shamsi *et al.*, 2009) and to *C. rudolphii* s.l. (Amato *et al.*, 2006; Li *et al.*, 2013) previously analyzed by SEM, our specimens show a similar external morphology of anterior and posterior extremity. The pattern of post-cloacal papillae in adult males and the conspicuous cuticular collar in both males and females are particularly distinctive characters that support the identification of *C. rudolphii* s.l.

Generally, cormorants from central Italy showed a similar proportion of adult stages of *C. rudolphii* A and *C. rudolphii* B while the ones from Israel showed a higher proportion of *C. rudolphii* B. It has been suggested that the different feeding ecology and wintering behavior of different populations of *Ph. carbo sinensis*, could be one of the “drivers” of the differential

spatial distribution of *C. rudolphii* A and *C. rudolphii* B in the different aquatic ecosystems, i.e. brackish/marine and freshwater environments, respectively (Mattiucci *et al.*, 2020). Moreover, abiotic factors related to early stages of the parasites, have been supposed to contribute to the differential occurrence of the two sibling species in the two aquatic ecosystems (Moravec *et al.*, 2009; Mattiucci *et al.*, 2020).

Recent work suggested the possible role of the migration routes of wintering populations of cormorants in the Mediterranean Sea in influencing the distribution and genetic structure of *C. rudolphii* (Roca-Geronès *et al.*, 2023). Cormorants migrate from Europe and winter in Israel during November-March, forming large colonies along the Mediterranean and Red Sea coasts and at inland streams and wetlands (Nemtsov, 2008). Large numbers of *C. rudolphii* A could be acquired by cormorants feeding in coastal areas during their migration across the Mediterranean Sea. Furthermore, most of Israel's wetlands are exploited for fish farming, mainly in freshwater fishponds; these fish are particularly susceptible to predation by piscivorous birds during migration stopovers (Nemtsov, 2002) and could provide opportunities for maintaining the life cycle of *C. rudolphii* B. In our study, all the great cormorants from both countries have been sampled in winter; interestingly, in two cases the birds from Israel were sampled in an area very close to the sea but they were infected only by *C. rudolphii* B, therefore we could hypothesize that they arrived in these wintering sites already parasitized. To date, larval stages of *C. rudolphii* have never been reported in marine/brackish/freshwater fish from Israel. This could either be due to biotic/abiotic factors unfavorable to the development of *C. rudolphii* larvae, or to difficulties in the detection of small larval stages (much smaller than those of other *Contracaecum* spp.) during field inspections. Despite their adaptation to different aquatic ecosystems, larvae of both *C. rudolphii* A and *C. rudolphii* B have generally shown a low specificity for their fish host (Mattiucci *et al.*, 2020), therefore the availability of suitable paratenic hosts does not seem to be a limiting factor for the successful establishment of the

parasite life cycle.

This work presents the first data on the occurrence of sibling species of *C. rudolphii* in Israel and provides additional data on the distribution of *C. rudolphii* A and B in Italy, confirming the high prevalence and intensity of infection observed in *Ph. carbo sinensis* from other Italian areas. Based on these results, we stress the importance of carrying out targeted investigations aimed at evaluating the occurrence of *C. rudolphii* in marine and freshwater fish from Israel, to understand the epidemiology of the parasite in this important wintering stopover.

### **Supplementary material.**

The supplementary material for this article can be found at DOI: .

### **Data availability.**

The DNA sequences generated in this study have been deposited on the public database GenBank under accession numbers OR263194-OR263246 (ITS rDNA) and OR269666-OR269717 (*cox2* mtDNA).

### **Acknowledgements.**

The authors thanks prof. Roberta Galuppi and dr. Giorgia Bignami of the DIMEVET for their help in the collection of samples. We also thank to Omer Ben-Asher and Ziv Riklis from the GIS Unit within the Ministry of Agriculture and Rural Development for the preparation of the maps.

### **Author's contribution.**

MC and PT wrote the main manuscript text, carried out all the analyses and prepared all the figures. AC, VL and PAG provided the technical support for SEM and molecular analyses. ND

and SR carried out the sampling and necropsies. AG and MLF revised the manuscript. All authors reviewed the manuscript and approved the final version.

**Financial support.**

This study was supported by the Israeli Veterinary Services and by the University of Bologna, Italy. Scanning Electron Microscopy acquired thanks to the Department of Excellence Project 2018-2022 funded by the Italian Ministry of Education, Universities and Research.

**Competing interests.**

The authors declare there are no conflicts of interest.

**Ethical standards.**

Not applicable.

Accepted Manuscript

## References

- Abollo, E, Gestal, C and Pascual, S** (2001) Anisakid infection in the European shag *Phalacrocorax aristotelis aristotelis*. *Journal of Helminthology* **75**, 209-214. <https://doi.org/10.1079/JOH200051>.
- Amato, JFR, Monteiro, CM and Amato, SB** (2006) *Contracaecum rudolphii* Hartwich (Nematoda, Anisakidae) from the Neotropical Cormorant, *Phalacrocorax brasilianus* (Gmelin) (Aves, Phalacrocoracidae) in southern Brazil. *Revista Brasileira de Zoologia* **23**, 1284-1289. <https://doi.org/10.1590/S0101-81752006000400046>.
- Amor, A, Farjallah, S, Piras, MC, Burreddu, C, Garippa, G and Merella, P** (2020) New insights into the coexistence of *Contracaecum rudolphii* A and *Contracaecum rudolphii* B (Nematoda: Anisakidae) in *Phalacrocorax carbo sinensis* from Sardinia: genetic variability and phylogenetic analysis. *Parasitology* **147**, 1538-1551. <https://doi.org/10.1017/S0031182020001341>.
- Baruš, V, Sergeeva, TP, Sonin, MD and Ryzhikov, KM** (1978) Helminths of fish-eating birds of the Palaearctic Region: Nematoda. In: Ryšavý, B and Ryzhikov, KM (eds), Helminths of Fish-Eating Birds. Springer: Dordrecht, p 318. <https://doi.org/10.1007/978-94-009-9972-5>.
- Battisti, C, Taffon, D and Giucca, F** (2008) Atlante degli uccelli nidificanti, Gangemi Editore: Roma. p. 190.
- Caffara, M, Tedesco, P, Davidovich, N, Locke, SA, Gustinelli, A, King, R, Nuytten, M, Nuzzo, M and Fioravanti, ML** (2023) Advancing understanding of the taxonomy and diversity of the genus *Contracaecum* in the great white pelican (*Pelecanus onocrotalus*). *Parasitology Research* **122**, 315-331. <https://doi.org/10.1007/s00436-022-07732-z>.
- Cammilleri, G, D'Amelio, S, Ferrantelli, V, Costa, A, Buscemi, MD, Castello, A, Bacchi, E, Goffredo, E, Mancini, ME and Cavallero, S** (2023) Identification of *Contracaecum*



*rudolphii* (Nematoda: Anisakidae) in Great Cormorants *Phalacrocorax carbo sinensis* (Blumenbach, 1978) from Southern Italy. *Veterinary Science* **10**, 194. <https://doi.org/10.3390/vetsci10030194>.

**Carmeno, C, Rusconi, A, Castelli, M, Prati, P, Bragoni, R, Santoro, A, Postiglione, U, Sasser, S and Olivieri, E** (2022) Molecular identification of *Contracaecum rudolphii* A and B (Nematoda: Anisakidae) from cormorants collected in a freshwater ecosystem of the pre-alpine area in Northern Italy. *Veterinary Parasitology: Regional Studies and Reports* **27**, 100674. <https://doi.org/10.1016/j.vprsr.2021.100674>.

**Cipriani, P, Palomba, M, Giulietti, L, Marcer, F, Mazzariol, S, Santoro, M, Aco Alburquerque, R, Covelo, P, López, A, Santos, MB, Pierce, GJ, Brownlow, A, Davison, NJ, McGovern, B, Frantzis, A, Alexiadou, P, Højgaard, DP, Mikkelsen, B, Paoletti, M, Nascetti, G, Levsen, A, and Mattiucci, S** (2022) Distribution and genetic diversity of *Anisakis* spp. in cetaceans from the Northeast Atlantic Ocean and the Mediterranean Sea. *Scientific Reports* **12**, 13664. <https://doi.org/10.1038/s41598-022-17710-1>.

**Clement, M, Posada, D and Crandall, K** (2000) TCS: a computer program to estimate gene genealogies. *Molecular Ecology* **9**, 1657-1659. <https://doi.org/10.1046/j.1365-294x.2000.01020.x>.

**Culurgioni, J, Sabatini, A, De Murtas, R, Mattiucci, S and Figus, V** (2014) Helminth parasites of fish and shellfish from the Santa Gilla Lagoon in southern Sardinia, Italy. *Journal of Helminthology* **88**, 489–498. <https://doi.org/10.1017/S0022149X13000461>.

**D'Amelio, S, Cavallero, S, Dronen, NO, Barros, NB and Paggi L** (2012) Two new species of *Contracaecum* Railliet & Henry, 1912 (Nematoda: Anisakidae), *C. fagerholmi* n. sp. and *C. rudolphii* F from the brown pelican *Pelecanus occidentalis* in the northern Gulf of Mexico. *Systematic Parasitology* **81**, 1-16. <https://doi.org/10.1007/s11230-011-9323->

[X](#).

**D'Amelio, S, Barros, NB, Ingrosso, S, Fauquier, DA, Russo, R and Paggi, L** (2007) Genetic characterization of members of the genus *Contracaecum* (Nematoda: Anisakidae) from fish-eating birds from west-central Florida, USA, with evidence of new species. *Parasitology* **134**, 1041–1051. <https://doi.org/10.1017/S003118200700251X>.

**Davidovich, N, Tedesco, P, Caffara, M, Yasur-Landau, D, Gustinelli, A, Drabkin, V, Minkova, E, Aflalo, O, Morick, D and Fioravanti, ML** (2022) Morphological description and molecular characterization of *Contracaecum* larvae (Nematoda: Anisakidae) parasitizing market-size hybrid tilapia (*Oreochromis aureus* x *Oreochromis niloticus*) and red drum (*Sciaenops ocellatus*) farmed in Israel. *Food and Waterborne Parasitology* **26**, e00147. <https://doi.org/10.1016/j.fawpar.2022.e00147>

**Davidovich, N, Tedesco, P, Caffara, M, Luci, V, Cantori, A, Morick, D, Fioravanti, ML and Gustinelli, A** (2023) Low prevalence of *Contracaecum* third-stage larvae parasitizing sea of Galilee fisheries: 1-year survey after 57 years of no information. *Food and Waterborne Parasitology* **32**, e00204. <https://doi.org/10.1016/j.fawpar.2023.e00204>.

**Dezfuli, B.S., Manera, M., Bosi, G., DePasquale, J.A., D'Amelio, S., Castaldelli, G and Giari, L** (2016) *Anguilla anguilla* intestinal immune response to natural infection with *Contracaecum rudolphii* A larvae. *Journal of Fish Diseases* **39**, 1187–1200. <https://doi.org/10.1111/jfd.12455>.

**Frederiksen, M, Korner-Nievergelt, F, Marion, L and Bregnballe, T** (2018) Where do wintering cormorants come from? Long-term changes in the geographical origin of a migratory bird on a continental scale. *Journal of Applied Ecology* **55**, 1–14. <https://doi.org/10.1002/ece3.4348>.

**Guardone, L, Susini, F, Castiglione, D, Ricci, E, Corradini, C, Guidi, A and Armani, A**

- (2020) Ascaridoid nematode larvae in wild gilthead seabream (*Sparus aurata*) and European seabass (*Dicentrarchus labrax*) caught in the Tyrrhenian Sea (Western Mediterranean Sea): a contribute towards the parasitological risk assessment on two commercially important fish species. *Food Control*. 118, 107377. <https://doi.org/10.1016/j.foodcont.2020.107377>.
- Hall, T.A** (1999) BioEdit: A User-Friendly Biological Sequence Alignment Editor and Analysis Program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**, 95-98
- Hartwich, G** (1964) Revision der Vogelparasitischen Nematoden Mitteleuropas II. Die Gattung *Contracaecum* Railliet and Henry, 1912 (Ascaridoidea). *Mitteilungen aus dem Zoologischen Museum in Berlin* **40**:15–53.
- Israeli Veterinary Services** (2017) Pre-marketing control of locally grown edible fish: procedure. <https://www.gov.il/he/departments/policies/moag-pro-082> (in Hebrew)
- Kumar, S, Stecher, G and Tamura, K** (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7 for bigger datasets. *Molecular Biology and Evolution* **33**:1870-1874. <https://doi.org/10.1093/molbev/msw054>.
- Li, AX, D'Amelio, S, Paggi, L, He, F, Gasser, RB, Lun, ZR, Abollo, E, Turchetto, M and Zhu, XQ** (2005). Genetic evidence for the existence of sibling species within *Contracaecum rudolphii* (Hartwich, 1964) and the validity of *Contracaecum septentrionale* (Kreis, 1955) (Nematoda: Anisakidae). *Parasitology Research* **96**, 361-366. <https://doi.org/10.1007/s00436-005-1366-y>.
- Li, L, Xu, Z and Zhang, LP** (2013) Further studies on *Contracaecum spasskii* Mozgovoi, 1950 and *C. rudolphii* Hartwich, 1964 (*sensu lato*) (Ascaridida: Anisakidae) from piscivorous birds in China. *Systematic Parasitology* **84**, 225-236. <https://doi.org/10.1007/s11230-012-9399-y>.
- Li, L, Gibson, DI and Zhang, LP** (2016) An annotated catalogue of the ascaridoid nematode

parasites of Chinese vertebrates. *Systematic Parasitology* **93**,1-35.  
<https://doi.org/10.1007/s11230-015-9617-5>.

**Mattiucci, S, Turchetto, M, Brigantini, F and Nascetti, G** (2002) On the occurrence of the sibling species of *Contracaecum rudolphii* complex (Nematoda: Anisakidae) in cormorants (*Phalacrocorax carbo sinensis*) from Venice and Caorle lagoons: Genetic markers and ecological studies. *Parassitologia*. **44**, 105.

**Mattiucci, S, Sbaraglia, GL, Palomba, M, Filippi, S, Paoletti, M, Cipriani, P and Nascetti, G** (2020) Genetic identification and insights into the ecology of *Contracaecum rudolphii* A and *C. rudolphii* B (Nematoda: Anisakidae) from cormorants and fish of aquatic ecosystems of Central Italy. *Parasitology Research* **119**, 1243–1257.  
<https://doi.org/10.1007/s00436-020-06658-8>.

**Molnár, K, Székely, C, Baska, F, Müller, T, Zuo, S, Kania, PW, Nowak, B and Buchmann, K** (2019) Differential survival of 3rd stage larvae of *Contracaecum rudolphii* type B infecting common bream (*Abramis brama*) and common carp (*Cyprinus carpio*). *Parasitology Research* **118**, 2811-2817. <https://doi.org/10.1007/s00436-019-06441-4>.

**Moravec, F** (2009) Experimental studies on the development of *Contracaecum rudolphii* (nematode: Anisakidae) in copepod and fish paratenic hosts. *Folia Parasitologica* **56**, 185–193. <https://doi.org/10.14411/fp.2009.023>.

**Nemtsov, SC** (2002) Management of Wildlife-Human Conflicts in Israel: A wide variety of vertebrate pest problems in a difficult and compact environment. Proc. 20th Vertebr. Pest Conf. (Timm, R.M., Schmidt, R.H. Eds.) Published at Univ. of Calif., Davis. 2002. Pp. 348-353. <https://doi.org/10.5070/V420110190>.

**Nemtsov, SC** (2008) Israel-Ukraine cooperation for experimental management of a shared overabundant population of Great Cormorants (*Phalacrocorax carbo*). Proc. 23rd Vertebr. Pest Conf. (Timm, R.M., Madon, M.B. Eds.) Published at Univ. of Calif., Davis.

2008. Pp. 108-112. <https://doi.org/10.5070/V423110628>.

**Paggi L, D'Amelio S, Ortis M, Mattiucci S, Mariniello L, Di Cave D and Orecchia P**

(1998) Metazoi parassiti di pesci di valore economico delle lagune costiere del Parco Nazionale del Circeo. *Biologia Marina Mediterranea* **5**, 1593–1599

**Roca-Geronès, X, Fisa, R, Montoliu, I, Casadevall, M, Tobella, C, Bas, JB, Palomba, M**

**and Mattiucci S** (2023) Genetic diversity of *Contraecaecum rudolphii* sp. A (Nematoda: Anisakidae) parasitizing the European Shag *Phalacrocorax aristotelis desmarestii* from the Spanish Mediterranean coast. *Frontiers in Veterinary Science* **10**, 1122291. <https://doi.org/10.3389/fvets.2023.1122291>.

**Rokicki, J, Sołtysiak, Z, Dziekońska-Rynko, J and Borucińska, J** (2011) Pathology

associated with *Contraecaecum rudolphii* (Nematoda: Anisakidae) infection in the great cormorant *Phalacrocorax carbo* (L. 1758). *Helminthologia* **48**, 28-35. <https://doi.org/10.2478/s11687-011-0006-6>.

**Ronquist, F, Teslenko, M, van der Mark, P, Ayres, DL, Darling, A, Höhna, S, Larget, B,**

**Liu, L, Suchard MA and Huelsenbeck, JP** (2012) MRBAYES 3.2: Efficient Bayesian phylogenetic inference and model selection across a large model space. *Systematic Biology* **61**, 539-542. <https://doi.org/10.1093/sysbio/sys029>.

**Rozas, J, Ferrer-Mata, A, Sánchez-DelBarrio, JC, Guirao-Rico, S, Librado, P, Ramos-**


**Onsins, SE and Sánchez-Gracia, A** (2017) DnaSP v6: DNA Sequence Polymorphism Analysis of Large Datasets. *Molecular Biology and Evolution* **34**, 3299-3302. <https://doi.org/10.1093/molbev/msx248>.

**Shamsi, S, Norman, R, Gasser, R and Beveridge, I** (2009) Redescription and genetic


characterization of selected *Contraecaecum* spp. (Nematoda: Anisakidae) from various hosts in Australia. *Parasitology Research* **104**, 1507-1525. <https://doi.org/10.1007/s00436-009-1357-5>.

- Smirnov, M, Ofek, T and Ron, TB** (2021) Forty years of fluctuation in the occurrence of two various metazoan parasites: *Contraecum multipapillatum* (Nematoda) and Digenean metacercaria N1 in warm water fish culture in the Israeli farms. *Israeli Journal of Aquaculture* **73**, 1-6. <https://doi.org/10.46989/001c.21408> .
- Szostakowska, B and Fagerholm, HP** (2007) Molecular identification of two strains of third-stage larvae of *Contraecum rudolphii Sensu Lato* (Nematoda: Anisakidae) from fish in Poland. *Journal of Parasitology* **93**, 961–964. <https://doi.org/10.1645/GE-1100R.1>.
- Szostakowska, B, Myjak, P and Kur, J** (2002) Identification of anisakid nematodes from the Southern Baltic Sea using PCR-based methods. *Molecular and Cellular Probes* **16**, 111–118. <https://doi.org/10.1006/mcpr.2001.0391>.
- Yamaguti, S** (1935) Studies on the helminth fauna of Japan. Part 12. Avian nematodes, I. *Journal of Zoology* **6**, 403–431.
- Zhu, X, Gasser, RB, Podolska, M and Chilton, NB** (1998) Characterisation of anisakid nematodes with zoonotic potential by nuclear ribosomal DNA Sequences. *International Journal for Parasitology* **28**, 1911–1921. [https://doi.org/10.1016/S0020-7519\(98\)00150-7](https://doi.org/10.1016/S0020-7519(98)00150-7).
- Zhu, XQ, D'Amelio, S, Gasser, RB, Yang, TB, Paggi, L, He, F, Lin, RQ, Song, HQ, Ai, L and Li, AX** (2007) Practical PCR tools for the delineation of *Contraecum rudolphii* A and *Contraecum rudolphii* B (Ascaridoidea: Anisakidae) using genetic markers in nuclear ribosomal DNA. *Molecular and Cellular Probes* **21**, 97–102. <https://doi.org/10.1016/j.mcp.2006.08.004>.


## Legends to figures



**Figure 1.** Maps of Israel (A) and Italy (B) with detail of the sampling localities together with the numbers of *Phalacrocorax carbo sinensis* collected from each locality.




**Figure 2.** PCR-RFLP pattern of *C. rudolphii* A and *C. rudolphii* B after digestion with *MspI* (*C. rudolphii* s.l.: 700-300 bp) and *NsiI* (*C. rudolphii* A: uncut; *C. rudolphii* B: 850-49 bp).




**Figure 3.** Maximum-Likelihood tree based on the concatenated ITS1-ITS2 rDNA sequences showing the relationship between *C. rudolphii* A (condensed, containing 20 newly generated sequences: GenBank accession numbers OR263194, OR263197-OR263205, OR263223-24, OR263226-34, plus two concatenated reference sequences AJ634782+AJ634785 of *C. rudolphii* A, Li *et al.*, 2005), *C. rudolphii* B (condensed, containing 33 sequences, GB acc. n. OR263195-96, OR263202, OR263206-22, OR263225, OR263235-46, plus two concatenated reference sequences AJ634783+AJ634786 of *C. rudolphii* B, Li *et al.*, 2005) described in the present study (in bold) and the congeneric *Contraecaecum* species. The tree is drawn to scale, with branch length measured in the number of substitutions per site.







**Figure 4.** Bayesian Inference (BI) tree based on the *cox2* mtDNA sequences showing the relationship between *C. rudolphii* A (condensed, containing 20 sequences, GB acc. n. OR269671, OR269673, OR269678-79, OR269681-88, OR269690-92, OR269705-09, plus one reference sequence MK496476 of *C. rudolphii* A, Mattiucci *et al.*, 2020), *C. rudolphii* B (condensed, containing 32 sequences, GenBank accession numbers: OR269666-70, OR269672, OR269674-77, OR269680, OR269689, OR269693-OR269704, OR269710-17, plus one reference sequence MK496482 of *C. rudolphii* B, Mattiucci *et al.*, 2020) described in the present study (in bold) and the congeneric *Contracaecum* species. The posterior probability is reported for each branch.




**Figure 5.** TCS network of haplotypes recorded in the present study, with indications of only the haplotypes detected at least 5 times. All the rest of haplotypes are reported in the supplementary figure S1 and table S1.




**Figure 6.** SEM micrographs of *C. rudolphii* A adult male. (A) Apical view of anterior end showing the dorsal lip (dl) and two ventral lips (vl) separate by interlabia, and the excretory pore opening at the base of ventral interlabium (arrow). (B) Lateral view of anterior end. (C) Anterior end with detail of the amphid. (D) Caudal end with everted spicules. (E) Spicules with detail of the spicule tip. (F) Caudal end showing the pattern of post-cloacal papillae.



**Figure 7.** SEM micrographs of *C. rudolphii* B adult female. (A) Ventral view of anterior end. (B) Detail of amphid. (C) Dorsal view of anterior end. (D) Apical view of anterior end. (E) Detail of vulva. (F) Caudal end.



**Figure 8.** SEM micrographs of *Contracaecum rudolphii* B, adult male. (A) Lateral view of anterior end. (B) Subapical view of anterior end, showing interlabia with bilobed tip. (C) Detail of lip edge. (D) Caudal end showing post cloacal papillae and section of spicules. (E) Caudal end with everted spicules and detail of the spicule tip. (F) Detail of caudal end showing post cloacal papillae (pcp) and phasmid (ph).



**Figure 9.** SEM micrographs of *Contracaecum rudolphii* B, adult female. (A) Lateral view of anterior end. (B) Detail of ventral lip. (C) Lateral view of anterior portion. (D) detail of amphid. (E) Detail of vulva. (F) Caudal end.

**Table 1.** Distribution of the adults of *C. rudolphii* A and *C. rudolphii* B collected from great cormorant and the number of specimens identified by molecular methods, together with the sampling locality and years of sampling for both countries under study.

Birds ID	Locality	Sampling Years	Adults/mol ID	<i>C. rudolphii</i> A	<i>C. rudolphii</i> B
<b>Israel</b>					
PC-01	Neve Ur	Dec 2020	114/55	1	54
PC-02	Gesher	Dec 2020	70/33	10	23
PC-03			59/28	6	22
PC-04			6/6	6	0
PC-05			95/50	50	0
PC-06			38/9	6	3
PC-07	Misgav-Baraam	Nov 2021	45/10	10	0
PC-08			12/10	10	0
PC-09			64/10	0	10
PC-10	Tirat Zvi	Dec 2021	111/10	0	10
PC-16			21/16	16	0
PC-17			116/10	0	10
PC-18			82/10	0	10
PC-11	Maoz Haim	Dec 2021	49/15	1	14
PC-14			219/10	0	10
PC-12	Lohamei HaGeta'ot	Dec 2021	2/2	0	2
PC-13	Maagan Michael	Dec 2021	14/10	0	10
PC-15			49/10	0	10
<b>Total</b>			<b>1166/304</b>	<b>116</b>	<b>188</b>
<b>Italy</b>					
IT-01	Emilia Romagna (FE)	Feb 2021	51/24	11	13
IT-02			29/13	5	8
IT-03			47/17	16	1
IT-04			93/51	47	4
IT-05			18/11	7	4
IT-06			9/3	1	2
IT-07			118/60	14	46
IT-08			82/36	17	19
IT-09	Tuscany (AR)	Feb 2022	161/140	50	90
IT-10	Emilia Romagna (BO)	Dic 2021	31/24	16	8
<b>Total</b>			<b>639/379</b>	<b>184</b>	<b>195</b>

**Table 2.** Morphometric features of mature male and female specimens of *C. rudolphii* A and *C. rudolphii* B from *Ph. carbo sinensis* from present study. Measurements are in  $\mu\text{m}$  unless otherwise stated.

	Males		Females	
	<i>C. rudolphii</i> A	<i>C. rudolphii</i> B	<i>C. rudolphii</i> A	<i>C. rudolphii</i> B
<b>TL</b>	13 - 28 (22.3 $\pm$ 3.5) mm	11 - 32 (20.6 $\pm$ 5.1) mm	26 - 39 (30.8 $\pm$ 4.9)	16 - 34 (24.7 $\pm$ 6.4)
<b>Oe</b>	1906 - 4190.7 (3451.2 $\pm$ 529.6)	1614.9 - 6051.5 (3299.1 $\pm$ 660.3)	3281.4 - 4188.3 (3910.3 $\pm$ 371.3)	2091.9 - 4360.7 (3200.5 $\pm$ 895.2)
<b>IC</b>	1222.1 - 3374.7 (2551.6 $\pm$ 473.3)	1130 - 4731 (2436.3 $\pm$ 525)	2700 - 3471.5 (3061.2 $\pm$ 290.4)	1530.8 - 3545.4 (2601.5 $\pm$ 875.5)
<b>VA</b>	644 - 1443.1 (1070.4 $\pm$ 205.8)	511.4 - 1456.7 (931.2 $\pm$ 187.2)	1096.9 - 1506.4 (1229.9 $\pm$ 182)	667.5 - 1326 (1011.1 $\pm$ 239.7)
<b>NR</b>	298.4 - 786.6 (593.7 $\pm$ 118.6)	274.9 - 743.2 (534.1 $\pm$ 100.9)	502.2 - 604.4 (563.6 $\pm$ 39.2)	401 - 618.1 (504.4 $\pm$ 86.7)
<b>Ta</b>	163.9 - 365.9 (240.9 $\pm$ 42.1)	160.1 - 336.4 (239 $\pm$ 34.4)	346.5 - 416.4 (377.1 $\pm$ 25.7)	303.5 - 398.7 (345.4 $\pm$ 29.3)
<b>RSp</b>	5429.3 - 10217.5 (8054.5 $\pm$ 1169.2)	2876.6 - 10580.7 (7527.3 $\pm$ 1780.9)	-	-
<b>LSp</b>	5128.2 - 9985.9 (7755.9 $\pm$ 1206.6)	3316.8 - 10091.5 (7449.9 $\pm$ 1687.2)	-	-
<b>Vu</b>	-	-	9260.3 - 12807.7 (11469.3 $\pm$ 1456.1)	7243.7 - 11133.5 (8914.2 $\pm$ 1346.3)

Abbreviations TL = total length; Oe = oesophagus length; IC = intestinal caecum length; VA = ventricular appendix length; NR = distance of nerve ring from anterior end; Ta = tail length; RSp = right spicule length; LRp = left spicule length; Vu = distance of vulva from anterior end.

Accepted Manuscript