



Turkish nurses' burnout levels, job satisfaction, and mental symptoms during the COVID-19 pandemic

Nuray Turan ^{1*}, Arzu Erkoç ², Çiğdem Beki ³, Türkinaz Aştı ⁴, İsmet Kırpınar ⁵

¹ Department of Fundamentals of Nursing Istanbul University, Faculty of Nursing, Istanbul, Türkiye

² Department of Medical Nursing, Florence Nightingale, Faculty of Nursing, Istanbul University-Cerrahpasa, Istanbul, Türkiye

³ Health Sciences University, Başakşehir Çam and Sakura City Hospital, Istanbul, Türkiye

⁴ Department of Nursing, Bezmialem University, Faculty of Health Sciences, Istanbul, Türkiye

⁵ Head of Psychiatry, Bezmialem Vakıf University, Faculty of Medicine, Istanbul, Türkiye

* **Corresponding author:** Nuray Turan, Istanbul University Nursing Faculty, Fatih-Istanbul, Türkiye.

Email: nkaraman@istanbul.edu.tr

Received: 2 December 2022 **Revised:** 1 March 2023 **Accepted:** 18 April 2023 **e-Published:** 30 June 2023

Abstract

Background: Nurses involved in healthcare delivery during the Coronavirus disease 2019 (COVID-19) pandemic experience higher levels of burnout, job satisfaction, and mental symptoms than other healthcare professionals did.

Objectives: This study examined nurses' burnout level, job satisfaction, and mental symptoms during the COVID-19 pandemic.

Methods: A descriptive, correlational study was conducted on 262 nurses working at a foundation university hospital in Istanbul between July and December 2021. The data were collected by using the Nurse Information Form, Maslach Burnout Inventory, Minnesota Job Satisfaction Scale, and Mental Symptom Checklist. Data analysis was performed using the Mann-Whitney U test, Kruskal-Wallis H test, Spearman correlation, and Bonferroni post hoc test.

Results: Nurses' Burnout, Job Satisfaction, and Mental Symptom average scores were 70.36 ± 11.00 , 2.80 ± 0.49 , and 1.20 ± 0.79 , respectively. A statistically significant negative and weak relationship was found between burnout, and job satisfaction average scores ($r = -0.380$; $P < 0.001$). Age, marital status, and income level were associated with burnout, job satisfaction, and mental symptoms ($P < 0.05$).

Conclusion: Nurses had a moderate level of burnout, a low level of job satisfaction, and a high level of mental symptoms.

Keywords: COVID-19, Burnout, Job satisfaction, Mental symptoms, Nurses.

Introduction

Burnout can reduce job performance, and increase physical and mental health problems such as depressive affection, restlessness, and concentration impairment, which will lead to absenteeism, job dissatisfaction, and job leave. [1-3]

Changes in healthcare settings during the Coronavirus disease 2019 (COVID-19) pandemic to maintain their dynamicity and efficiency may also cause varying levels of depression, anxiety, and stress for healthcare providers, particularly nurses. [2-6] Nurses' feelings of job satisfaction, mental symptoms, and burnout profoundly impact the quality of nursing care. [7] COVID-19, due to its unknown nature and high contagiousness, can cause high levels of

stress, mental symptoms and tension, and, finally, burnout in nurses, affecting their job satisfaction. Several studies investigated burnout, job satisfaction, or mental symptoms in nurses. [3,7-9] However, studies that evaluated these variables together in Turkish nurses during the COVID-19 pandemic are not available. Therefore, the question that arises is whether there is a relationship between job burnout, job satisfaction, and depression in Turkish nurses.

Objectives

This study examined the burnout level, job satisfaction, and mental symptoms of Turkish nurses during the COVID-19 pandemic.

Methods

Study design and participants

A cross-sectional study was conducted between July and December 2021. A total of 390 nurses working in the internal and surgical and adult inpatient services of a foundation university hospital in Istanbul, Türkiye, constituted the population of the study. As a result of the power analysis using the G*Power 3.0.10 program; with 0.90 power, 0.5 margins of error, and $d=0.202$ effect size, a total of at least 262 samples were found to be sufficient. The inclusion criteria included the active working of nurses in internal, surgical, and adult inpatient services.

Data collection instruments

The Nurse Information Form consisted of questions to determine the socio-demographic characteristics of nurses.

Maslach Burnout Inventory (MBI)^[10] measures the burnout status of individuals working in different occupational groups. Its Turkish translation^[11] consists of 22 items under three dimensions: Emotional Exhaustion (EE), Depersonalization (DP), and Feeling of Personal Accomplishment (PA). The items are rated on a seven-point Likert-type scale ranging between 0 and 6. The increase in EE and DP scores indicates a high level of burnout, and the increase in the PA score indicates a low level of burnout. Ergin found Cronbach- α coefficients for the three dimensions as 0.83 for Emotional Exhaustion, 0.65 for Depersonalization, and 0.72 for Personal Accomplishment.^[11] In this study, the Cronbach- α coefficient was found to be 0.722.

Minnesota Job Satisfaction Scale (MJSS)^[12] measures job satisfaction. The MJSS without reverse questions consists of 20 items and has features that determine the internal, external, and general satisfaction levels. General Satisfaction covers all items of the scale. Items are rated on a five-point Likert type scale between 1 and 5. The Cronbach- α coefficient was reported 0.91.^[13] In this study, the Cronbach- α coefficient was found to be 0.821.

The Mental Symptom Checklist (SCL-90-R) is a psychiatric checklist tool used in determining the psychiatric symptoms, and the level of challenge or stress experienced by individuals.^[14] The Turkish validity and reliability study of the scale was conducted by Dağ.^[15] It contains 90 self-reported items and each item is scored between 0 and 4. The increase in the General Symptom Level, which is the general average score of the scale, indicates the increase in the distress of the individual's psychiatric symptoms and is the highest index receivable from the scale. While a score greater than 1 indicates a

mental problem, a score between 0.5 and 1 indicates a moderate problem, and a score less than 0.5 indicates that there is no problem. Three general scores can be calculated from the scale. In the original study of the scale, the Cronbach- α coefficient was 0.97.^[16] In this study, the Cronbach- α coefficient was found to be 0.97.

Data collection

The data were collected from nurses recruited to the study by convenience sampling and by visiting hospital units three days a week between 08:00 and 16:00. Nurse Information Form, Minnesota Job Satisfaction Scale, and Mental Symptom Checklist were applied to each nurse. It took approximately 10-15 minutes to collect data from each nurse.

Ethical considerations

Before starting the study, permission was obtained from the researchers who developed the scale to use the scales. The study was approved by the Ministry of Health General Directorate of Health Services (approval code: T16_54_25, issued at 202-05-17), institutional permission from a foundation university hospital and ethical approval from the clinical research ethics committee (approval code: 8725) were obtained. The researcher explained to the nurses the purpose, content, and what was expected from them. The principles of the Declaration of Helsinki were followed in the conduct of the study.

Data analysis

IBM SPSS Statistics 24 (IBM SPSS, Türkiye) was used for statistical analysis. Non-parametric methods were used for the measurement values that did not conform the normal distribution. The Mann-Whitney U test was used for the comparison of measurement values of two independent groups, and the Kruskal-Wallis H test was used for the comparison of measurement values of three or more independent groups. The Bonferroni correction was applied for binary comparisons of the variables that differed significantly for three or more groups. The Spearman correlation coefficient examined the relationship between measurement values without normal distribution. P values <0.05 were considered significant.

Results

The mean age of the nurses was 26.11 ± 6.55 years, 79.4% of them were female, 79% were married, 33.5% were in the 23-24 age group, and the income of 53.1% of them did not meet their expenses. About 53.6% of nurses were medical vocational high school graduates, 61.5% worked in their

current unit for an average of 1-4 years, 46.2% worked in the profession for an average of 1-4 years, and 48.9% did not provide care in the family.

Nurses' average burnout score was 70.36±11.00, average job satisfaction score was 2.80±0.49, and their average mental symptom score was 1.20±0.79 [Table 1]. A statistically significant negative correlation was found between nurses' burnout and job satisfaction scores (r=-0.380; p<0.001). A significant positive correlation was found between nurses' burnout and mental symptom scores (r=0.423; p<0.001). A significant negative correlation was found between nurses' job satisfaction and mental symptom scores (r=-0.408; p<0.001).

Mental symptom average scores of those aged 22 and lower and those in the age group of 23-34, and 25-26 were significantly higher compared to those aged 27 and over (p=0.022). A significant negative relationship was found between age and mental symptom scores (r=-0.159; p=0.010). The mental symptom average scores of the single nurses were significantly different from married individuals (p<0.001). The mental symptom average scores of nurses whose income did not meet their expenses were significantly higher than that of those whose income met their expenses (p<0.001) [Table 2].

Discussion

Our nurses had moderate burnout. Nurses' burnout has been shown to be increased by heavy workloads during the pandemic, posing a significant threat to members of the medical team.^[17] Kaya and İşler Dalğıç^[18] found the burnout level was moderate during the pandemic. In the study conducted by Liu and Aungsuroch,^[19] it was determined that the work-related stress of nurses affected burnout moderately. Evidence show that long weekly working hours, fear of infection, extreme physical fatigue, and difficulty in working in full protective equipment^[20] may lead to physical and mental burnout in nurses. Nurses in this study also had a low level of job satisfaction. Working in pandemic units brings along a significant

workload for nurses, which causes nurses to have physical and mental difficulties. While this finding supports the increase in burnout in nurses, it emphasizes the need for close monitoring of nurses' job satisfaction, improving working conditions and providing psychological support. It was determined that the mental symptom levels of the nurses were generally high. Healthcare workers are exposed to both physical and psychological stress during the COVID-19 pandemic.^[21,22] A study showed that the pandemic negatively affected the mental health of nurses and that these effects continued over time.^[22] Brouwer et al.,^[23] and Zhang et al.,^[24] similarly reported that nurses experienced mental problems during the COVID-19 pandemic. Our findings along with these studies reflect the moderate to high levels of job burnout and low job satisfaction among nurses.

Burnout and job satisfaction in nurses showed a reverse correlation. It seems that the physical, mental, and social disorders to which nurses are exposed in the working environment, especially during the COVID-19 pandemic has an effect on job satisfaction and burnout.^[25,26]

Nurses' burnout levels increased as their mental symptom levels increased. Similarly, nurses' burnout levels decreased as their mental symptom levels decreased. Since nursing is a profession based on relationships with people, it needs mentally healthy professionals more than other professions. The result of the study is compatible with the literature^[9,27,28] and signifies the fact that nurses should be mentally healthy in order to fulfill their duties and responsibilities.

The mental symptom levels of nurse were correlated with their job satisfaction. Job satisfaction can be defined as how the workers perceive that their needs are met by their job.^[17] The COVID-19 pandemic has globally led to unprecedented pressure on healthcare workers, especially nurses.^[20] According to recent studies, some nurses have developed mental symptoms and decreased job satisfaction, while facing the COVID-19 pandemic.^[28,29] However, the knowledge on the impact of COVID-19 on nurses' mental health and job satisfaction is still incipient.

Table 1. Distribution of Nurses' Maslach Burnout, Minnesota Job Satisfaction, and Mental Symptom Checklist Scores (N=262)

| Scale | Mean± SD | Median | Min.-Max. |
|----------------------------------|-------------|--------|-----------|
| Maslach Burnout Inventory | 70.36±11.00 | 69.0 | 29.0-98.0 |
| Minnesota Job Satisfaction Scale | 2.80±0.49 | 2.8 | 1.1-3.6 |
| Mental Symptom Checklist | 1.20±0.79 | 1.1 | 0.0-4.0 |

Min.: Minimum. Max.: Maximum. Mean: Mean. SD: Standard deviation

Table 2. Comparison of Maslach Burnout, Minnesota Job Satisfaction, and Mental Symptom Checklist scores according to individual characteristics of nurses ^a (N=262)

| Variable | N | Maslach Burnout | | Minnesota Job Satisfaction | | Mental Symptom Checklist | |
|--------------------------------|-----|-----------------------|-------------|----------------------------|-----------|--------------------------|-----------|
| | | Mean± SD | Median | Mean± SD | Median | Mean± SD | Median |
| Age classes | | | | | | | |
| ≤22 ⁽¹⁾ | 71 | 68.70±12.18 | 69.0 [15.0] | 2.84±0.51 | 2.9 [0.7] | 1.32±0.90 | 1.3 [1.6] |
| 23-24 ⁽²⁾ | 88 | 71.39±10.52 | 71.0 [13.8] | 2.77±0.44 | 2.8 [0.4] | 1.26±0.81 | 1.2 [1.4] |
| 25-26 ⁽³⁾ | 35 | 69.20±11.82 | 68.0 [11.0] | 2.76±0.53 | 2.8 [0.7] | 1.26±0.70 | 1.3 [1.3] |
| ≥27 ⁽⁴⁾ | 68 | 71.37±9.80 | 70.0 [11.8] | 2.80±0.50 | 2.9 [0.8] | 0.96±0.65 | 0.9 [0.8] |
| | | F=1.113 | | χ ² =1.120 | | χ ² =9.602 | |
| | | p=0.344 | | p=0.772 | | p=0.022 | |
| Gender | | | | | | | |
| Female | 208 | 70.60±10.93 | 69.0 [13.0] | 2.80±0.47 | 2.8 [0.6] | 1.22±0.81 | 1.1 [1.4] |
| Male | 54 | 69.46±11.32 | 70.0 [12.3] | 2.80±0.56 | 2.9 [0.9] | 1.10±0.74 | 1.0 [1.1] |
| | | Z=-0.136 | | Z=-0.222 | | Z=-0.879 | |
| | | p=0.892 | | p=0.824 | | p=0.380 | |
| Marital status | | | | | | | |
| Married | 55 | 70.56±10.16 | 70.0 [14.0] | 2.89±0.41 | 2.9 [0.6] | 0.86±0.60 | 0.8 [0.8] |
| Single | 207 | 70.31±11.24 | 69.0 [13.0] | 2.77±0.50 | 2.8 [0.6] | 1.29±0.82 | 1.2 [1.4] |
| | | Z=-0.256 | | Z=-1.532 | | Z=-3.532 | |
| | | p=0.798 | | p=0.126 | | P<0.001 | |
| School graduated | | | | | | | |
| Medical vocational high school | 96 | 70.07±11.38 | 69.0 [13.0] | 2.88±0.49 | 2.9 [0.6] | 1.23±0.81 | 1.2 [1.4] |
| Associate degree | 66 | 71.21±10.61 | 71.5 [14.3] | 2.81±0.45 | 2.8 [0.8] | 1.20±0.86 | 1.0 [1.3] |
| Undergraduate and higher | 100 | 70.08±10.97 | 69.0 [12.0] | 2.70±0.52 | 2.7 [0.6] | 1.16±0.73 | 1.1 [1.1] |
| | | F=0.262 | | χ ² =5.559 | | χ ² =0.262 | |
| | | p=0.770 | | p=0.062 | | p=0.877 | |
| Income level | | | | | | | |
| Expenses are met | 123 | 69.61±9.26 | 69.0 [11.0] | 2.83±0.45 | 2.9 [0.6] | 1.03±0.74 | 0.8 [1.2] |
| Expenses are not met | 139 | 71.03±12.32 | 70.0 [14.0] | 2.77±0.52 | 2.8 [0.8] | 1.35±0.81 | 1.3 [1.2] |
| | | Z=-1.274 | | Z=-1.065 | | Z=-3.290 | |
| | | p=0.203 | | p=0.287 | | P<0.001 | |
| Person cared | | | | | | | |
| No one | 128 | 70.34±10.14 | 70.0 [13.0] | 2.77±0.47 | 2.8 [0.6] | 1.17±0.77 | 1.1 [1.2] |
| Mother and/or father | 77 | 70.25±12.74 | 69.0 [13.5] | 2.73±0.54 | 2.8 [0.7] | 1.39±0.87 | 1.5 [1.6] |
| Child/children | 34 | 72.35±9.57 | 70.0 [12.3] | 2.93±0.43 | 3.0 [0.6] | 1.01±0.62 | 1.0 [0.9] |
| Other | 23 | 67.91±11.48 | 72.0 [18.0] | 2.99±0.42 | 3.0 [0.6] | 0.98±0.78 | 0.8 [1.1] |
| | | χ ² =2.150 | | χ ² =3.234 | | χ ² =7.379 | |
| | | p=0.542 | | p=0.357 | | p=0.061 | |

^a“ANOVA” test (F-table value) statistics were used for the comparison of three or more independent groups in the data without normal distribution. “Mann-Whitney U” test (Z-table value) was used for the comparison of measurement values of two independent groups in the data without normal distribution, and “Kruskall-Wallis H” test (χ²-table value) statistics were used for the comparison of three or more independent groups.

In the present study, younger nurses showed higher mental symptoms. It can be attributed to the fact that nurses who have just started their profession and started to work at an early age, are concerned about meeting new people and beginning relationships and many unknown environments. We also found that the mental symptom scores of the single nurses were higher compared to married individuals. Being married is generally

considered a protective factor for mental health.^[28] According to the income status, it was determined that the mental symptom average scores of nurses whose income did not meet their expenses were significantly higher than those whose income met their expenses. When the socioeconomic status of nurses is low, they are deprived of many opportunities such as cultural and social activities, an adequate lifestyle, which causes nurses to experience

various feelings of frustration and be exposed to high levels of stress.^[30,31] In this context, it can be stated that income status has a direct effect on the mental health of nurses.

This study has some limitations. The use of self-report scales is among the limitations. Future studies can employ qualitative methods to assess nurses' job satisfaction, burnout, and mental status. Furthermore, since the study was conducted in only one institution, the results can only be generalized to this institution. We did not have data about the variables before the COVID-19 pandemic, so it is difficult to relate the burnout, job satisfaction, and mental health of nurses to the pandemic.

Conclusions

Nurses had a moderate level of burnout, a low level of job satisfaction, and a high level of mental symptom. While a negative relationship was found between nurses' burnout and job satisfaction, a positive relationship was found between their burnout and mental symptom levels, and a negative relationship was found between their job satisfaction and mental symptom levels. Significant differences were found between the mental symptom levels of nurses according to their age, marital status, and income level. It may be recommended to adjust working conditions to reduce burnout and increase job satisfaction in nurses during the COVID-19. Nurse authorities also should evaluate nurses' burnout level, job satisfaction, and mental status at certain intervals, support nurses psychologically in hospitals, and increase the practices that protect nurses' mental health.

Acknowledgment

The authors would like to thank the nurses who participated in this study in the intensive care unit.

Competing interests

The authors declare that they have no competing interests.

Abbreviations

Coronavirus disease 2019: COVID-19;
Maslach Burnout Inventory: MBI;
Minnesota Job Satisfaction Scale: MJSS;
Mental Symptom Checklist: SCL-90-R.

Authors' contributions

All authors read and approved the final manuscript. All authors take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Role of the funding source

None.

Availability of data and materials

The data used in this study are available from the corresponding author on request.

Ethics approval and consent to participate

The study was conducted in accordance with the Declaration of Helsinki. The study was approved by the Ministry of Health General Directorate of Health Services (2020-05-17T16_54_25), institutional permission from a foundation university hospital and ethical approval from the clinical research ethics committee (24.07.2021/8725) were obtained.

Consent for publication

By submitting this document, the authors declare their consent for the final accepted version of the manuscript to be considered for publication.

References

1. Arpacioğlu MS, Baltacı Z, Ünübol B. Burnout, fear of COVID-19, depression, occupational satisfaction levels and related factors in healthcare professionals in the COVID-19 pandemic. *Cukurova Med J* 2021;46:88-100.
2. Wu Y, Wang J, Luo C, Hu S, Lin X, Anderson AE, et al. A comparison of burnout frequency among oncology physicians and nurses working on the front lines and usual wards during the COVID-19 epidemic in Wuhan, China. *J Pain Symptom Manage* 2020;60:e60-e65. doi:10.1016/j.jpainsymman.2020.04.008 PMID:32283221 PMCID:PMC7151285
3. Hassanvandi S, Mohammadi M, Shahyad S. Predicting the severity of covid-19 anxiety based on sleep quality and mental health in health care workers. *Novel Clin Med* 2022; 1(4): 184-191. doi:10.22034/ncm.2022.344007.1048
4. Fallahi CR, Mitchell MT, Blau JJ, Daigle CD, Rodrigues HA, Deleo L. Burnout, personal and secondary trauma among mental health care professionals during the COVID-19 pandemic. *Minerva Psychiatry* 2022;63:219-230. doi:10.23736/S2724-6612.21.02192-8
5. Kaya B. Effects of pandemic on mental health. *J Clin Psy* 2020; 23: 123-124. doi:10.5505/kpd.2020.64325
6. Aydın A, Akay B, Baydemir S. Burnout and factors affecting it in the nursing students. *J DU Health Sci Inst* 2017;7:19-23.
7. Ertekin PŞ, Bilgiç D, Demirel G, Akyüz MB, Karatepe C, Sevim D. Relationship between burnout and life satisfaction of university students in the health field. *TAF Prev Med Bull* 2015;14;284-292. doi:10.5455/pmb.1-1417432935
8. Kılıç M, İpekçi NN, Dokur M, Kaya S. Determination of burnout levels of the nurses working at Kilis state hospital. *J Anatolia Nurs*

- Health Sci 2015;18:1-8. doi:10.17049/ahsbd.69979
9. Heidarijamebozorgi M, Jafari H, Sadeghi R, Sheikhbardsiri H, Kargar M, Amiri Gharaghani M. The prevalence of depression, anxiety, and stress among nurses during the coronavirus disease 2019: A comparison between nurses in the frontline and the second line of care delivery. *Nurs Midwifery Stud* 2021;10:188-193. doi:10.4103/nms.nms_103_20
 10. Maslach C, Jackson SE. The measurement of experienced burnout. *J Occup Behav* 1981;2: 99-113. doi:10.1002/job.4030020205
 11. Ergin C. Burnout in doctors and nurses and adaptation of Maslach Burnout Scale. National Psychology Congress, 22th September 1992 Ankara, Türkiye.
 12. Weiss DJ, Dawis RV, England GW. Manual for the Minnesota Satisfaction Questionnaire. *Minnesota Stud Vocational Rehabil* 1967; 22:120-124. doi:10.1037/t05540-000
 13. Teleş, M. Validity and Reliability of the Turkish Version of the General Work Stress Scale. *J Nurs Manag* 2021;29:710-720. doi:10.1111/jonm.13211 PMID:33174261
 14. Derogatis LR. SCL-90: Administration, Scoring and Procedure Manual-1 for the revised version, Baltimore MD: Johns Hopkins Univ 1977.
 15. Dağ Y. Reliability and validity of the symptom screening list (SCL-90-R) for university students. *Turk Psikiyatri Derg* 1991;2:5-12.
 16. Koğar H. The validity and reliability study of the Symptom Checklist (SCL-90): Mokken Scale Analysis. *Turkish Psychological Guidance J* 2019;9:689-705.
 17. Soto-Rubio A, Giménez-Espert MDC, Prado-Gascó V. Effect of emotional intelligence and psychosocial risks on burnout, job satisfaction, and nurses' health during the COVID-19 Pandemic. *Int J Environ Res Public Health* 2020;17:7998. doi:10.3390/ijerph17217998 PMID:33143172 PMCid:PMC7663663
 18. Kaya A, İşler Dalgıç A. Examination of job satisfaction and burnout status of pediatric nurses: A cross-sectional and correlational study using online survey research in Türkiye. *Perspect Psychiatr Care* 2021;57:800-808. doi:10.1111/ppc.12617 PMID:32924165
 19. Liu Y, Aunguroch Y. Work stress perceived social support, self-efficacy, and burnout among Chinese registered nurses. *J Nurs Manag* 2019;27:1445-1453. doi:10.1111/jonm.12828 PMID:31306524
 20. Denning M, Goh ET, Tan B, Kanneganti A, Almonte M, Scott A, et al. Determinants of burnout and other aspects of psychological wellbeing in healthcare workers during the COVID-19 pandemic: A multinational cross-sectional study. *Plos One* 2021;16: e0238666. doi:10.1371/journal.pone.0238666 PMID:33861739 PMCid:PMC8051812
 21. Chen Q, Liang M, Li Y, Guo J, Fei D, Wang L, et al. Mental health care for medical staff in China during the COVID-19 outbreak. *Lancet Psychiatry* 2020;7:e15-e16. doi:10.1016/S2215-0366(20)30078-X PMID:32085839
 22. Lu W, Wang H, Lin Y, Li L. Psychological status of medical workforce during the COVID-19 pandemic: a cross-sectional study. *Psychiatry Res* 2020;288:112936. doi:10.1016/j.psychres.2020.112936 PMID:32276196 PMCid:PMC7195354
 23. Brouwer KR, Walmsley LA, Parrish EM, McCubbin AK, Welsh JD, Braido CEC, et al. Examining the associations between self-care practices and psychological distress among nursing students during the COVID-19 pandemic. *Nurse Educ Today* 2021;100: 104864. doi:10.1016/j.nedt.2021.104864 PMID:33744816 PMCid:PMC7946540
 24. Zhang S, Wang J, Xie F, Yin D, Shi Y, Zhang M, et al. A cross-sectional study of job burnout, psychological attachment, and the career calling of Chinese doctors. *BMC Health Serv Res* 2020;20: 193. doi:10.1186/s12913-020-4996-y PMID:32164684 PMCid:PMC7068889
 25. Duarte I, Teixeira A, Castro L, Marina S, Ribeiro C, Jácome C, et al. Burnout among Portuguese healthcare workers during the COVID-19 pandemic. *BMC Public Health* 2020; 20:1885. doi:10.1186/s12889-020-09980-z PMID:33287794 PMCid:PMC7720923
 26. Danacı E, Koç Z. The association of job satisfaction and burnout with individualized care perceptions in nurses. *Nurs Ethics* 2020; 27:301-315. doi:10.1177/0969733019836151 PMID:30966862
 27. Doğan S, Erdoğan C, Çakmak R, Kızılaslan D, Çiftçi B, Karaslan P. Reviewing work-related strain, perceived stress and job satisfaction among the nurses working at new type 2019 corona virus services. *Göbeklitepe Int J Health Sci* 2021;4:16-25.
 28. Yu X, Zhao Y, Li Y, Hu C, Xu H, Zhao X, et al. Factors associated with job satisfaction of frontline medical staff fighting against COVID-19: A cross-sectional study in China. *Front Public Health* 2020;8:426. doi:10.3389/fpubh.2020.00426 PMID:32850610 PMCid:PMC7417651
 29. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw Open* 2020;3:e203976. doi:10.1001/jamanetworkopen.2020.3976 PMID:32202646 PMCid:PMC7090843
 30. Oral M, Karakurt N. Evaluation of the relationship between burnout and psychological symptoms among staff working with individuals with disabilities and elderly. *Cyp Turk J of Psychiatry and Psychol* 2021;3:123-131.
 31. Özel Y, Türkleş S, Erdoğan S. Investigation of mental state in university students. *J Academic Nurs* 2021;6:220-228. doi:10.5222/jaren.2020.41033

How to Cite this Article:

Turan N, Erkoç A, Beki Ç, Aştı T, Kırpınar İ. Turkish nurses' burnout levels, job satisfaction, and mental symptoms during the COVID-19 pandemic. *Nurs Midwifery Stud*. 2023;12(2): 104-109. doi:10.48307/NMS.2023.175271