



## Effect of precision land leveling and permanent raised bed planting on soil properties, input use efficiency, productivity and profitability under maize (*Zea mays*) – wheat (*Triticum aestivum*) cropping system

R K NARESH<sup>1</sup>, R S RATHORE<sup>2</sup>, PARVINDER KUMAR<sup>3</sup>, S P SINGH<sup>4</sup>, ADESH SINGH<sup>5</sup> and U P SHAHI<sup>6</sup>

Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut, Uttar Pradesh

Received: 16 April 2012; Revised accepted: 20 March 2014

### ABSTRACT

Precision land leveling with permanent raised bed planting with recommended dose of NPK can be used to improve crop yield, water and nutrient use efficiency over the existing traditional land leveling with flat beds planting with recommended dose of NPK practices. The objective of present study was to establish an understanding of maize (*Zea mays* L.) -wheat (*Triticum aestivum* L.) rotation yield and input use efficiency can be improved and how land leveling and crop establishment practices can be modified to be more efficient in water use through precision conservation crop management techniques. A farmers participatory field experiment was conducted during 2009-2011 in the jurisdiction of Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut, UP. Multi crop planter with inclined plate seed metering device machine were given to the farmers and crops were sown on permanent raised beds in maize-wheat cropping system. The data collected from the farmers participatory field experiment showed that there was about 20.4% (295.8 mm/ha for wide beds, i.e. 107 cm furrow centre gap) water saving and about 16.5% (310.3 mm/ha for narrow beds, i.e. 37 cm furrow centre gap) with grain yield increase about 13.5% (5.13 and 4.44 tonnes/ha) for wheat crop and 11.8% (4.33 and 3.82 tonnes/ha) for maize crop with precision land leveling raised bed planting compared to traditional land leveling with flat beds planting. The agronomic efficiency (AE) of N (23.4 and 30.4 kg grain/kg N for maize and wheat) and uptake of N, P and K (103.85, 25.6 and 110.7 kg/ha for maize and 112.95, 19.49 and 112.96 kg/ha for wheat) were significantly improved under precision land leveling with raised bed planting technique compared to other practices.

**KeyWords:** Input use efficiency, Permanent raised beds, Precision land leveling, Productivity, Profitability

Presently, 50% of the human population relies on nitrogen (N) fertilizer for food production. The world today uses around 83 million metric tonnes of N, which is about a 100-fold increase over the last 100 years (Ladha *et al.* 2013). About 60% of global N fertilizer is used for producing the world's three major cereals: rice, wheat, and maize. Projections estimate that 50 to 70% more cereal grain will be required by 2050 to feed 9.3 billion people (Ladha *et al.* 2013). Stagnating yield and declining input use efficiency in irrigated maize (*Zea mays* L.) -wheat (*Triticum aestivum* L.) of the Western Indo-Gangetic Plain

(WIGP) coupled with diminishing availability of water for agriculture is a major concern of food security in South Asia. Achieving sustainable food security is a major challenge considering the growing population with changing diets and a degrading resource scenario. Maize or corn is an important food and feed crop and is one of the most versatile, high yielding food crops of the world. In South Asia with limited availability of water, and aberration in temperature on account of climate change, it is becoming increasingly important that issues concerning it be addressed. Maize lends itself to various applications and has a high yield potential and wide adaptability across regions, seasons and altitudes, making it suited in different cropping and farming systems. Its demand is increasing on account of the shift towards animal based diets and expansion of the bio-fuel industry. Raised beds were introduced to rice-wheat systems of the Indo-Gangetic Plain (IGP) in the mid 1990s, initially for wheat, inspired by the success of irrigated maize-wheat on permanent raised beds (PRB) in Mexico (Sayre and Hobbs 2004). Since then, many advantages of growing wheat on beds have been reported, including increased

<sup>1</sup> Associate Professor, Department of Agronomy (e mail: r.knaresh@yahoo.com), <sup>2</sup> DDG (e mail: rajvir\_ddg@yahoo.co.in), UPCAR, Lucknow, Uttar Pradesh; <sup>3</sup> Research Associate (e mail: balyanpk@yahoo.com), Department of Agrimet, CCS Haryana Agricultural University, Hisar; <sup>4</sup> Assistant Professor (e mail: drpsingh75@gmail.com), Department of Soil Science, <sup>5</sup> Assistant Professor (e mail: adeshsingh29@rediffmail.com), Department of Agronomy, <sup>6</sup> Assistant Professor (e mail: upshahi@gmail.com), Department of Soil Science

yields, reduced lodging, opportunities for mechanical weeding and improved fertiliser placement, irrigation water savings, reduced waterlogging, reduced seed rate and opportunities for intercropping (Ram *et al.* 2005, Naresh *et al.* 2011). Around the same time that PRB were being proposed, an unprecedented revolution in adoption of 'zero-till' (direct-drilled) wheat after rice was underway across the IGP (Malik *et al.* 2004). Majority of the maize-wheat growers in the western Uttar Pradesh practice surface irrigation either through flood or check basin methods. The light textured soils under undulating topography leads to uneven distribution of water, which limits the availability of water and nutrients to the crop plants. Undulated crop fields when managed with flood irrigation, also lead to within field spatial variability in grain production owing to leaching of certain nutrients due to excess water at lower elevations and in-adequate availability of irrigated water at higher elevations. Naresh *et al.* (2011) reported that wheat and maize are planted in many parts of the world on beds and bed planting on an average saved 29% of water as compared to flat beds.

Land leveling is a precursor to good agronomic, soil and crop management practices and the levelness of the land surface has significant influence on all the farming operations. Jat *et al.* (2006) rated the development of laser technology for precision land leveling as second only to breeding of high yielding varieties. The soil moisture status throughout the field governed by its levelness has great influence not only on farming operations but also the yield and input use efficiency. The leveling of land for achieving higher resource use efficiency is not a new technique but the way in which it is done is not up to the mark as frequent patches of dikes and ditches stretched over a minimum workable distance are created even with best effort by conventional leveling practices. Undulated land hampers the seedbed preparation, seed placement, germination and also requires heavy draught for machines, which leads to consumption of more energy, and ultimately to more cost of production and low productivity levels. Improvement in operational efficiency, weed control efficiency, water use efficiency, nutrient use efficiency, crop productivity and economic returns (Naresh *et al.* 2011) and environmental benefits (Jat *et al.* 2006) have been reported as a result of precision land leveling when compared to traditional practice of land leveling. The objective of this study was to evaluate the effect of precision land leveling and permanent raised beds planting on soil properties, input use efficiency, productivity and profitability under maize-wheat cropping system on a sandy loam soil of western Uttar Pradesh.

#### MATERIALS AND METHODS

An experiment was conducted on maize-wheat rotation in three districts (Meerut, Ghaziabad and Saharanpur) in farmers participatory mode in the jurisdiction of Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut (Uttar Pradesh), India, (28°40'07"N to 29°28'11"N, 77°28'14"E to 77° 44 '18"E) during 2009-10 to 2011-12

and was designed as a farmer-managed with a single replicate, repeated over many farmers. Therefore, the experimental design was Randomized Block Design in which farmer as a replicate/block commencing with *khariif* in 2009. The climate of the area is semi-arid, with an average annual rainfall of 805 mm 75–80% of which is received during July to September, minimum temperature of 4°C in January, maximum temperature of 41–45°C in June, and relative humidity of 67–83% during the year. The soils are generally sandy loam to loam in texture and low to medium in organic matter content. Soil with a bulk density of 1.48 Mg/m<sup>3</sup>, pH =7.9, total C=8.3 g/kg, total N =0.83 g/kg, Olsen P =28 mg/kg, and K =128 mg/kg. Groundwater pumping was the predominant method of irrigation in Western UP. The plots consisted of seven crop establishment treatments and details is given as follows: T<sub>1</sub> - Precision land leveling with wide raised Beds with recommended dose of NPK (PL WB + RNPK), T<sub>2</sub> - Traditional land leveling with wide raised Beds with recommended dose of NPK (TL WB + RNPK), T<sub>3</sub> - Precision land leveling with narrow raised Beds with recommended dose of NPK (PL NB + RNPK), T<sub>4</sub> - Traditional land leveling with narrow raised Beds with recommended dose of NPK (TL NB + RNPK), T<sub>5</sub> - Precision land leveling with flat Beds with recommended dose of NPK (PL FB + RNPK), T<sub>6</sub> - Traditional land leveling with flat Beds with recommended dose of NPK (TL FB + RNPK), T<sub>7</sub> - Traditional land leveling with flat beds without NPK (Control/conventional practices) (TL FB + N<sub>0</sub> P<sub>0</sub> K<sub>0</sub>).

For laser-assisted precision land leveling, the land was first plowed at the optimum moisture level (field capacity) with a harrow/cultivator for pulverization and was leveled using a laser-equipped drag scrapper (TrimbleTM, USA) with an automatic hydraulic system attached to a 50-60 HP tractor. Before running the laser leveler, the field was surveyed at 3-m distance to record the elevation and the elevation points were averaged to know the desired elevation for leveling the field. The average elevation value was entered into the digital control box for controlling the scrapper at the desired elevation point (Naresh *et al.* 2011) and the tractor was run across the field till the desired elevation was achieved throughout the field. For the traditional land leveling treatment, the field was first ploughed as described above and was leveled using an iron plank attached to a tractor and was dragged across the land surface.

After ploughing with a harrow/cultivator for pulverization of the field at the optimum moisture level, an iron scraper attached to the tractor was moved on the land surface on a visual elevation level. After the cuts and fills of soil, a wooden planker attached to the tractor was moved across the field to smooth the land surface.

HM 10 maize cultivar was seeded in the last week of June and wheat cultivar PBW 343 was seeded on 7, 9 and 9 November 2009, 2010 and 2011, respectively. A seed rate of 20 kg/ha for maize and 80 kg/ha was used in treatments where wheat was seeded on beds, and 100 kg/ha was used

in the rest of the treatments. The bed furrow width at top was kept at 37 cm:30 cm having one seed rows for maize and three seed rows for wheat and the depth of the furrow was kept at 15 cm for narrow beds and furrow width at top was kept at 107 cm:30 cm having two seed rows for maize and six seed rows for wheat and the depth of the furrow was kept at 12 cm for wide beds. The plant population was maintained equal in flat as well as raised bed planting.

Irrigation water was applied using polyvinyl chloride pipes of 15-cm diameter and the amount of water applied to each plot was measured using a water meter (Dasmesh Co., India). The quantity of water applied and the depth of irrigation were computed using the following equations:

$$\text{Quantity of water applied (L)} = F \times t \quad (1)$$

$$\text{Depth of water applied (mm)} = \{L / A / 1000\} \quad (2)$$

where F is flow rate (l/s), t is time (s) taken during each irrigation and A is area of the plot (m<sup>2</sup>). Rainfall data was recorded using a rain gauge installed within the meteorological station. The total amount of water (input water) applied was computed as the sum of water received through irrigation and rainfall (I+R). Water productivity (WP<sub>I+R</sub>) (kg grains/m<sup>3</sup> of water) was computed as follows (Humphreys *et al.* 2008): WP<sub>I+R</sub> = grain yield (kg/ha) / [irrigation water applied (m<sup>3</sup>) + rainfall received by the crop (m<sup>3</sup>)]/ha.

Soil samples were collected at the start of the experiment from 0 to 15-cm soil depth using an auger of 5-cm diameter. Each sample was a composite from three locations within a plot. The freshly collected soil samples were mixed thoroughly, air-dried, crushed to pass through a 2-mm sieve and stored in sealed plastic jars before analysis. Olsen P (0.5 M NaHCO<sub>3</sub> extractable) and NH<sub>4</sub>OAc-extractable K were analyzed using the methods described by [Page *et al.* 1982], respectively. Soil organic C was analyzed by the Walkley and Black method (Page *et al.* 1982). The samples for determination of soil physical properties (soil aggregates, mean weight diameter of aggregates) were collected at the start of the experiment and after the harvest of each crop. Soil aggregation and mean weight diameters of aggregates were analyzed using the wet-sieving method (Yoder method). Bulk density was measured to a depth of 20-cm at intervals of 5-cm soil depth using the core-ring method and one core

per stratus of each plot was collected and the samples were oven-dried for 48 h at 105°C, weighed and bulk density calculated according to (Blake and Hartge 1986). Soil moisture by gravimetric method (Jalota *et al.* 1998), soil strength by cone penetrometer. The bulk density were measured at the onset of the experiment and after the 3 years of study.

The plants measured for growth and yield were used for analyzing the N, P and K content in grain and straw. The grain and straw samples were dried at 70°C in a hot air oven. The dried samples were ground in a stainless steel Wiley Mill. The N content in grain and straw were determined by digesting the samples in sulfuric acid (H<sub>2</sub>SO<sub>4</sub>), followed by analysis of total N by Kjeldahl method (Page *et al.* 1982) using a Kjeltec autoanalyser. The P content (grain and straw) was determined by vanadomolybdo-phosphoric yellow colour method and the K content both in grain and straw was analysed in di-acid (HNO<sub>3</sub> and HClO<sub>4</sub>) digests by Flame Photometric method (Page *et al.* 1982). The uptake of the nutrients was calculated by multiplying the nutrient content (%) by respective yield (kg/ha) and was divided by 100 to get the uptake values in kg/ha. The uptake in grain and straw was summed to get the total uptake of nutrients/ha. In general, four terms are used in relation to NUE. These are: Agronomic Efficiency (AE), Recovery Efficiency (RE), Physiological Efficiency (PE), and Partial Factor Productivity of Fertilizers (PFP). The following expressions are used for determining these:

Agronomic efficiency = kg grain yield increase per kg applied AE = (Y-Y<sub>0</sub>)/F AE

Partial factor productivity = kg grain yield per kg N applied PFP = Y/F = (Y<sub>0</sub>/F) + AE

Recovery efficiency ('apparent') = kg increase in crop uptake per kg nutrient applied RE = (U-U<sub>0</sub>)/F

Physiological efficiency = kg yield increase per kg increase in crop uptake PE = (Y-Y<sub>0</sub>)/(U-U<sub>0</sub>)

## RESULTS AND DISCUSSION

### Soil properties

Tillage significantly affected the soil significant variations after three crop cycles the soil physical properties

Table 1 Effect of crop establishment on bulk density, water stability of aggregates, clod breaking strength and soil organic carbon (%) etc. soil properties under maize-wheat cropping system after 03 year's of experimentation in 0-15 cm

Treatment	Bulk density (Mg m <sup>-3</sup> )	Water stable	Aggregate	Clod breaking	Soil organic carbon (%)	Field capacity		Permanant witing	
		aggregates >0.25 mm (%)	porosity (%)	strength (kPa)		(% moisture)		point (% moisture)	
						0-5 cm	5-20 cm	0-5 cm	5-20 cm
T <sub>1</sub>	1.44	82.8	43.2	204.8	0.63	31	32	13	11
T <sub>2</sub>	1.46	79.0	40.8	332.9	0.58	29	30	11	10
T <sub>3</sub>	1.45	81.9	42.7	235.6	0.61	30	31	12	11
T <sub>4</sub>	1.48	72.9	40.2	367.5	0.55	29	29	11	10
T <sub>5</sub>	1.49	80.3	41.3	289.7	0.59	29	30	12	11
T <sub>6</sub>	1.50	66.7	39.6	418.7	0.54	28	29	11	10
T <sub>7</sub>	1.55	59.1	36.2	423.8	0.52	28	29	11	09
CD (P=0.05)	0.09	5.3	1.74	95.3	0.53**				

Table 2 Effect of crop establishment on available N, available P and available K (kg/ha) under maize-wheat cropping system after 03 year's of experimentation in 0-15 cm

Treatment	Available N (kg/ha)	Available P (kg/ha)	Available K (kg/ha)
T <sub>1</sub>	261.5	13.7	247.0
T <sub>2</sub>	250.1	12.1	241.8
T <sub>3</sub>	259.3	13.5	245.9
T <sub>4</sub>	249.0	11.9	240.7
T <sub>5</sub>	258.1	13.2	244.3
T <sub>6</sub>	243.4	11.8	240.2
T <sub>7</sub>	139.5	8.6	232.5
CD (P=0.05)	6.7	0.40	2.15

in bulk density, water stable aggregates, aggregate porosity, clod breaking strength, organic carbon, available N, P and K were recorded due to different treatments (Table 1). The bulk density did not vary significantly due to land leveling however, planting techniques had significant influence and it was significantly reduced under raised bed planting compared to flat sowing irrespective of the land leveling practice. This was attributed mainly due to more pore spaces created in the beds through modified land configuration by accumulations the topsoil. Bed planting provides natural opportunity to reduce compaction by confining traffic to the furrow bottoms (Govaerts *et al.* 2006). The soil organic carbon content in top soil (0-15 cm) was increased significantly due to raised bed planting compared to flat sowing planting mostly because of localized deposition of more fertile top soil on beds under altered land configuration than flat planting. Available nitrogen, phosphorus and potassium status of soil analyzed after harvest of third wheat crop showed significant variation due to different treatments (Table 2). Maximum available N, P and K content in soil was recorded under PLWB being at par with TLWB but were significantly superior to all other treatments. Further, flat planting either on precision or traditional leveling were at par with each other at similar fertility levels.

#### Nutrient uptake

Total (grain + straw) uptake of nutrients (N, P, K) analyzed at crop maturity varied significantly due to land leveling and crop establishment techniques. Maximum uptake of total N was recorded with PLWB which was significantly higher over all other treatments (Table 3). Similar to nitrogen, maximum uptake of total P was also recorded in PLWB which was at par to PLNB but it was significantly higher over rest of the treatments (Table 3). The total K uptake by the crop though at par, under precision land leveling irrespective of the planting technique (i.e PLWB, PLNB and PLFB) but significantly higher over rest of the treatments (Table 3). The higher amount of uptake of nutrients under precision leveling and raised bed planting techniques was associated with higher biomass accumulation under these treatments, which led to higher amount of uptake of these nutrients. The higher nutrient uptake in

Table 3 Effect of various tillage and establishment techniques on total N, P and K uptake in maize-wheat rotation

Crop establishment	Total N uptake (Kg/ha)		Total P uptake (kg/ha)		Total K uptake (kg/ha)	
	Maize	Wheat	Maize	Wheat	Maize	Wheat
T <sub>1</sub> - PLWB + RNPk	103.85	112.95	25.6	19.49	110.7	112.96
T <sub>2</sub> - TLWB + RNPk	99.75	107.56	22.7	16.61	104.3	108.27
T <sub>3</sub> - PLNB + RNPk	99.75	109.95	24.8	17.62	106.8	110.35
T <sub>4</sub> - TLNB + RNPk	93.05	103.80	20.3	14.85	098.2	096.36
T <sub>5</sub> - PLFB + RNPk	94.95	104.65	23.9	15.06	102.6	105.40
T <sub>6</sub> - TLFB + RNPk	89.85	100.45	17.5	13.31	81.4	083.56
T <sub>7</sub> - TLFB + N <sub>0</sub> P <sub>0</sub> K <sub>0</sub>	53.70	58.70	13.7	7.66	57.3	051.18
CD (P=0.05)	8.95	8.02	1.21	1.15	3.97	4.63

precision leveling with raised beds is mainly due to less leaching loss of nutrients and availability of sufficient moisture for mineralization of native as well as applied nutrients. The higher uptake efficiency of nutrients depends on a myriad of factors including nutrient availability due to favourable soil biota under precision leveling with raised beds compared to precision leveling with flat beds. These findings are in agreement with Jat *et al.* (2006) and Walker *et al.* (2003).

#### Nutrient use efficiency

The agronomic as well as recovery efficiency of applied nutrients was in general higher in precision leveling permanent wide beds. Efficiencies tended to be lower in traditional leveling flat beds (TLFB) than precision leveled wide or narrow raised beds treatments. Efficiencies on raised beds consistently increased but the differences between precision leveled and traditional leveled permanent raised beds were not always significant for the maize-wheat crop cycles with the same level of nutrient application (Table 4).

#### Agronomic efficiency (AE)

The agronomic efficiency (AE) of applied nutrients as unit grain production per unit of applied nutrients after deducting the soil supplying capacity was calculated for all the treatments. The AE of applied N was significantly higher under precision leveling with wide raised bed treatment compared to other treatments. The efficiency of the nutrient under PLWB+RNPk, and PLNB+RNPk was at par but significantly superior to PLFB+RNPk. The efficiency under PLFB+RNPk, TLNB+NPK and TLFB+RNPk were at par but significantly inferior to TLWB+RNPk. While traditional leveling without NPK (TLFB+N<sub>0</sub>P<sub>0</sub>K<sub>0</sub>) had significantly lower AE than all other treatments Table 4.



Table 4 Estimates of N use efficiency, i.e. Agronomic Efficiency of N (AEn), Recovery Efficiency of N (REn), Physiological Efficiency of N (Pen) and Partial Factor Productivity of N (PFPn) in maize-wheat rotation

Crop establishment	AEn (kg grain/kg N)		REn (%)		Pen		PFPn	
	Maize	Wheat	Maize	Wheat	Maize	Wheat	Maize	Wheat
T <sub>1</sub> - PLWB + RNPk	23.4	30.4	46.3	49.2	47.8	52.8	70.3	83.7
T <sub>2</sub> -TLWB + RNPk	21.6	28.5	40.7	42.3	39.4	43.5	55.8	64.2
T <sub>3</sub> - PLNB + RNPk	22.1	28.7	43.8	46.6	42.6	50.6	62.4	68.4
T <sub>4</sub> -TLNB + RNPk	20.3	27.2	39.3	38.4	37.7	41.6	52.1	47.7
T <sub>5</sub> - PLFB + RNPk	20.4	27.6	41.6	41.3	40.4	46.7	59.5	50.2
T <sub>6</sub> - TLFB + RNPk	19.7	26.2	36.7	32.3	33.8	39.8	44.6	39.6
T <sub>7</sub> . TLFB + N <sub>0</sub> P <sub>0</sub> K <sub>0</sub>	16.8	18.4	26.4	28.7	24.3	21.4	20.7	31.3
CD (P=0.05)	3.3	2.6	5.9	3.7	2.9	3.3	4.8	3.6

*Recovery efficiency (RE)*

Data on true recovery efficiency (REnt) shows that in maize the values ranged from 26.4 to 46.3%, while in wheat these ranged from 28.7 to 49.7%. Maize-wheat cropping system is a new rising system for the backbone of food security in India and the values of apparent recovery efficiency of N (REn) of maize and wheat experiment are in Table 4. Data in Table 4 clearly show that the values of all the terms associated with N use efficiency (NiUE) declined in all traditional leveling plots. At N levels similar to those in Table 4, values of all NiUE terms in maize were lower in compared to the wheat. On the other hand, values of all terms of NiUE showing that in experiment N is more efficiently utilized for wheat than maize. Thus in maize there is considerable scope to increase NiUE. Precision leveling irrespective of planting technique exerted significant effect on RE-N. The RE-N under PLWB+RNPk was significantly higher over all other treatments. Further, the recovery efficiency under TLLFB+RNPk also improved significantly compared to TLNB+RNPk, TLFB+RNPk and

TLFB+N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> (Table 4).

*Water application and water productivity*

The input water application includes the irrigation water applied and the rain water during the maize-wheat season of 2009-10 to 2011-12. The water application in maize-wheat system was remarkably lower with permanent wide and narrow beds compared to other practices (Table 5 and 6). The higher irrigation water application in maize-wheat system under traditional leveling treatments as compared to precision leveled plots. The precision leveled plots savings in water use in raised beds with recommended dose of NPK were 11.5% to 20.5% in maize and 14.1% to 26.7% in wheat as compared to traditional leveled flat beds with recommended dose of fertilizer treatment (T<sub>6</sub>). Water productivity under permanent beds was higher as compared to other tillage and crop establishment techniques and lowest system water productivity was recorded with traditional leveled flat beds. Bhushan *et al.* (2007) revealed that the saving in irrigation water with raised bed planting technique

Table 5 Water productivity and profitability of maize and wheat rotation under various tillage and establishment techniques

Crop establishment	Water productivity (kg yield m <sup>-3</sup> water )						Net profit ( ₹/ha )					
	Maize			Wheat			Maize			Wheat		
	2009	2010	2011	2009-10	2010-11	2011-12	2009	2010	2011	2009-10	2010-11	2011-12
T <sub>1</sub> - PLWB + RNPk	1.31	1.42	1.53	1.75	1.82	2.06	23 580	24 680	25 375	23 585	23 875	24 560
T <sub>2</sub> -TLWB + RNPk	1.16	1.17	1.17	1.54	1.52	1.50	22 320	22 495	23 310	21 525	21 650	21 985
T <sub>3</sub> - PLNB + RNPk	1.24	1.33	1.39	1.57	1.69	1.78	22 765	23 570	24 125	22 350	22 790	23 275
T <sub>4</sub> -TLNB + RNPk	1.10	1.11	1.12	1.41	1.40	1.36	21 550	22 200	23 050	21 300	21 435	21 525
T <sub>5</sub> - PLFB + RNPk	1.10	1.15	1.22	1.40	1.45	1.52	20 910	21 375	21 890	21 750	21 975	22 430
T <sub>6</sub> - TLFB + RNPk	1.01	1.00	1.01	1.25	1.19	1.19	19 500	19 760	20 300	20 050	19 610	19 250
T <sub>7</sub> . TLFB + N <sub>0</sub> P <sub>0</sub> K <sub>0</sub>	0.60	0.52	0.49	0.76	0.70	0.59	12 650	11 275	10 725	11 950	10 990	9 725

Table 6 Maize and wheat water application (mm/ha) and crop yield (t/ha) in laser-leveled and traditionally leveled field under different tillage and crop establishment methods

Crop establishment	Maize						Wheat					
	2009		2010		2011		2009 - 10		2010 - 11		2011 - 12	
	Water	Yield	Water	Yield	Water	Yield	Water	Yield	Water	Yield	Water	Yield
T <sub>1</sub> - PLWB + RNPk	325	4.25	310	4.40	300	4.60	295	5.15	285	5.20	260	5.35
T <sub>2</sub> -TLWB + RNPk	345	4.00	350	4.05	360	4.20	305	4.70	310	4.72	320	4.80
T <sub>3</sub> - PLNB + RNPk	330	4.10	320	4.25	312	4.35	315	4.95	295	5.00	290	5.15
T <sub>4</sub> -TLNB + RNPk	355	3.90	360	4.00	370	4.15	330	4.65	335	4.68	345	4.70
T <sub>5</sub> - PLFB + RNPk	360	3.95	350	4.02	345	4.20	340	4.75	330	4.77	320	4.85
T <sub>6</sub> - TLFB + RNPk	370	3.75	375	3.80	385	3.90	355	4.45	370	4.42	375	4.45
T <sub>7</sub> - TLFB + N <sub>0</sub> P <sub>0</sub> K <sub>0</sub>	385	2.30	390	2.05	397	1.95	360	2.75	365	2.55	380	2.25
CD (P=0.05)		0.21		0.29		0.37		0.24		0.23		0.31

was more under traditional leveling as in this technique water moves in furrows only. Laser assisted precision land leveling can reduce evaporation and percolation losses from wheat by enabling faster irrigation times and by eliminating depressions and therefore ponding of water in depressions.

#### Grain yields

The crop yield data from 2009-2012 (Table 6) showed that the higher grain yields of maize occurred in precision land leveling permanent wide beds with recommended dose of NPK. Yields on raised beds consistently increased from year 1 to year 3 in laser leveling with recommended NPK, but the differences between laser leveling permanent wide raised beds and permanent narrow raised beds were not always significant for the three maize-wheat crop cycles. Precision leveling with residue retain increase the yield by 8.5% to 10.9% in maize and 10.1% to 13.1% in wheat as compared to conventional seeding. This is an extremely important finding in relation to practical management of such systems by farmers. Data pertaining to crop yield parameters of wheat (Table 6) showed significant variation due to land leveling and planting techniques during the study years. The yield level, in general, under all the treatments was little higher during year 3 compared to year 1 and year 2. This was attributed mainly due to more sunshine hours cross the season in year 3 compared to year 1 and year 2. Also, the minimum temperature during flowering season was higher during year 1 and year 2 compared to year 3 which limits the reproductive period and responsible for lower yields of wheat. Grain yield of wheat varied significantly due to laser leveling permanent wide raised beds with recommended dose of NPK (PLWB+RNPk) techniques and significantly higher yield levels of

5.15, 5.20 and 5.35 tonnes/ha were recorded under (PLWB+RNPk) during year 1, year 2 and year 3, respectively compared to other treatments. The increase in grain yield with (PLWB+RNPk) was 8.3%, 8.7% and 9.3% during year 1, year 2 and year 3, respectively whereas the corresponding increase under flat bed planting was recorded at 6.3%, 7.3% and 8.2%. The yield under permanent wide raised beds traditional land leveling with recommended dose of NPK (TLWB+RNPk) and laser leveling permanent narrow raised beds with recommended dose of NPK (PLNB+RNPk) did not vary significantly during the years because productive tillers, length of spike and number of grains/spike are almost same. Further, with the same level of land leveling and different levels of planting techniques, the wheat yield varied remarkably. Raised bed showed 5.9, 6.4 and 7.6% yield advantage over flat bed planting under precision leveling during year 1, year 2 and year 3, respectively whereas, the corresponding increase in yield under traditional leveling was recorded at 4.7%, 5.9% and 6.3%. It showed that the raised bed planting technique is more advantageous under precisely leveled fields. Significantly higher yield of maize-wheat was recorded with precision land leveling as it takes care of maintaining near homogeneity by way of cut and fill and also tillage (Hassan *et al.* 2005, Borrell and Garside 2004). These findings are in agreement with Gupta and Sayre (2007), Rajput and Patel (2004), Naresh *et al.* (2011) who summarized the finding of multi-location trials across IGP and reported higher yield of wheat with raised beds compared to flat sowing.

#### Profitability

The net income through maize was higher with precision leveling permanent raised beds with recommended dose of

NPK followed by PLNB + RNPK > TLWB + RNPK > TLNB + RNPK and PLFB + RNPK and the lowest being recorded with TLFB+N<sub>0</sub>P<sub>0</sub>K<sub>0</sub> (Table 5). The lower net income with the conventional tillage was due to the cost on preparing the field. Profitability of wheat was remarkably higher with precision leveling permanent wide raised beds practices due to higher productivity and less cost of production compared to conventional tillage practices (T<sub>7</sub>) treatment.

### CONCLUSIONS

Over the past decade, researchers in association with farmers and entrepreneurs have been trying to overcome the problems of depleting water resources, diminishing input use efficiency, declining farm profitability, and deteriorating soil health by developing, evaluating and refining conservation and precision agriculture-based resource-conserving technologies for the maize-wheat rotation in the western Uttar Pradesh. Precision land leveling and permanent beds planting with recommended dose of NPK have a tremendous potential for improving the use efficiency of natural as well as externally applied resources. Many new opportunities based on precision land leveling and permanent beds planting with recommended dose of NPK have appeared to give stimulus to the productivity through a more sustainable pace of natural resource use in maize-wheat based cropping system. Taken together, these practices can rise productivity, cut costs, save water and soils, reduce use of external inputs, foster greater agro-ecosystem diversity and generate employment. Lignified residual straw and roots added more organic matter and nutrients into the soils under permanent raised beds, resulting in increased nutrient uptake by the crops. Crop yields on beds with precision land leveling rose by about 11.8% for maize and 13.5% for wheat over a 3-year cycle compared with conventional tillage on the flat beds. Future thrust of research are: Conservation tillage practices like precision land leveling and permanent bed planting utilizes more judiciously the plant available water than the conventional tillage when the other factors are similar. Precision-Conservation Agriculture (PCA) based crop management solutions seem to be promising options to sustain the maize-wheat systems of South Asia on a long-term basis.

### ACKNOWLEDGEMENTS

The authors grateful to the Director Research of the Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut, Uttar Pradesh, India for providing facilities and encouragement. Financial assistance by the Uttar Pradesh Council of Agricultural Research, Lucknow under the Resource Conservation Technologies for Sustainable Development of Agriculture project is duly acknowledged.

### REFERENCES

Blake G R and Hartge K H. 1986. Bulk density. (In) *Methods of Soil Analysis, Part I*, pp 363–75. Klute A, Campbell G S, Jackson R D, Mortland M M, Nielsen D R (Eds). ASA and SSSA, Madison, WI, USA.

- Bhushan L, Ladha J K, Gupta R K, Singh S, Padre A T, Saharawat Y S, Gathala M and Pathak H. 2007. Saving of water and labour in a rice-wheat system with no-tillage and direct seeding technologies. *Agronomy Journal* **99**(5): 1288–96.
- Borrell A and Garside A. 2005. Early work on permanent raised beds in tropical and subtropical Australia focusing on the development of a rice-based cropping system. (In) *Evaluation and performance of permanent raised bed cropping systems in Asia, Australia and Mexico*. Roth C H, Fischer R A and Meisner C A (Eds). ACIAR Proceedings No. 121, pp 120–8.
- Gupta R and Sayre K D. 2007. Conservation agriculture in South Asia. *Journal of Agricultural Sciences* **145**: 207–14.
- Govaerts B, Sayre K D, Ceballos-Ramirez J, Luna-Guido M L, Limon-Ortega A, Deckers J and Den-dooen L. 2006. Conventionally tilled and permanent raised beds with different crop residue management: Effect on soil C and N dynamics. *Plant and Soil* **280**: 143–55.
- Hassan I, Hussain Z and Akbar G. 2005. Effect of permanent raised beds on water productivity for irrigated maize-wheat cropping system. (In) *Evaluation and performance of permanent raised bed cropping systems in Asia, Australia and Mexico*. Roth C H, Fischer R A and Meisner C A (Eds). ACIAR Proceedings No. 121, pp 59–65.
- Humphreys E, Kukal S S, Amanpreet Kaur, Thaman S, Yadav S, Yadvinder Singh, Balwinder Singh, Timsina J, Dhillon S S, Prashar A and Smith D J. 2008. Permanent beds for rice-wheat in Punjab, India. Part 2: water balance and soil water dynamics. (In) *Permanent Beds and Rice-residue Management for Rice-wheat Systems in the Indo-Gangetic Plain*. Humphreys E and Roth C H (Eds).
- Jalota S K, Khera R and Ghuman B S. 1998. *Methods in Soil Physics*, pp 65–7. Narosa Publishing House, New Delhi.
- Jat M L, Chandana P, Sharma S K, Gill M A and Gupta R K. 2006. Laser land leveling—A precursor technology for resource conservation. Rice-Wheat Consortium Technical Bulletin Series 7, Rice-Wheat Consortium for the Indo-Gangetic Plains, New Delhi.
- Jat M L, Gupta Raj, Saharawat Y S and Khosla Raj. 2011. Layering precision land leveling and furrow irrigated raised bed planting: Productivity and input use efficiency of irrigated bread wheat in Indo-Gangetic plains. *American Journal of Plant Sciences* **2**: 1–11.
- Ladha, Jagdish K, Pathak, Himanshu, Timothy J, Krupnik J Six and Chris van Kessel. 2013. Efficiency of fertilizer nitrogen in cereal production: Retrospects and prospects. *Advances in Agronomy* **87**: 85–156.
- Malik R K, Yadav A, Gill G S, Sardana P, Gupta R K and Piggan C. 2004. Evolution and acceleration of no-till farming in rice-wheat system of the Indo-Gangetic Plains. (In) *4<sup>th</sup> International Crop Science Congress*, held in Brisbane, Australia from September 26 to October 1, 73 p.
- Naresh R K, Gupta, Raj, K, Satya Prakash, Kumar, Ashok, Singh Madhvendra and Misra A K. 2011. Permanent beds and rice-residue management for rice-wheat systems in the North West India. *International Journal of Agricultural Sciences* **7**(2): 429–39.
- Naresh R K, Gupta, Raj, K, Kumar, Ashok, Satya Prakesh, Tomar S S, Singh, Adesh, Rathi R C, Misra A K and Singh, Madhvendra. 2011. Impact of laser leveler for enhancing water productivity in Western Uttar Pradesh. *International Journal of Agricultural Engineering* **4**(2): 133–47.
- Page A L, Miller R H and Keeney D R (Eds). 1982. *Methods of*

- Soil Analysis. Part 2. Chemical and Microbiological Properties*, 2<sup>nd</sup> ed. Agron. Monogr. 9, ASA, CSSA and SSSA, Madison, WI.
- Rajput T B S and Patel Neelam. 2004. Effect of land leveling on irrigation efficiencies and wheat yield. *Journal of Soil and Water Conservation* **3**: 86–96.
- Ram H, Singh Yadvinder, Timsina J, Humphreys E, Dhillion S S, Kumar K and Kaler D S. 2005. *Evaluation and performance of permanent raised bed cropping systems in Asia, Australia and Mexico*. Roth C H, Fischer R A and Meisner C A (Eds). ACIAR Proceedings No. 121, pp 1–58.
- Sayre K D and Hobbs P R. 2004. The raised-bed system of cultivation for irrigated production conditions. (In) *Sustainable Agriculture and the International Rice-wheat System*, pp 337–55.
- Walker T W, Kingery W L, Street Joe E, Lox M S, Oldham J L, Gerard P D and Aan F X. 2003. Rice yield and soil chemical properties as affected by precision land leveling in Alluvial soils. *Agronomy Journal* **95**: 1 483–8.
- Yoder R E. 1936. A direct method of aggregate analysis and study of the physical nature of erosion losses. *Journal of American Society of Soil Science* **28**: 337–51.