

**Title: Improvements in sperm motility following low or high intensity dietary interventions in men with obesity.**

**Authors:** Aditi Sharma<sup>1\*</sup>, Nikoleta Papanikolaou<sup>1\*</sup>, Sara Abou Sherif<sup>1</sup>, Anastasia Dimakopolou<sup>1</sup>, Thilipan Thaventhiran<sup>1</sup>, Cara Go<sup>1</sup>, Olivia Holtermann Entwistle<sup>1</sup>, Adrian Brown<sup>2</sup>, Rong Luo<sup>1</sup>, Rama Jha<sup>1</sup>, Anavi Prakash<sup>1</sup>, Dalia Khalifa<sup>1</sup>, Hannah Lewis<sup>1</sup>, Sruthi Ramaraju<sup>1</sup>, Anthony R Leeds<sup>3</sup>, Harvinder Chahal<sup>1</sup>, Sanjay Purkayastha<sup>4</sup>, Ralf Henkel<sup>1</sup>, Sukhbinder Minhas<sup>5</sup>, Gary Frost<sup>1</sup>, Waljit S Dhillon<sup>1</sup>, Channa N Jayasena<sup>1</sup>

**\*Co-first authors**

1. Department of Metabolism, Digestion and Reproduction, Imperial College, 6th Floor Commonwealth building, 150 Du Cane Road, London, UK, W12 0NN
2. Centre for Obesity Research, University College London, London, UK.
3. Clinical Research Unit, Parker Institute, Frederiksberg Hospital, Copenhagen, Denmark.
4. Department of General and Bariatric Surgery, Imperial College Healthcare NHS Trust, St. Mary's Hospital, London, UK.
5. Department of Urology, Imperial College Healthcare NHS Trust, Charing Cross Hospital, Fulham Palace Road, Hammersmith, London, UK.

**Correspondence and reprint requests:**

Dr. Channa N. Jayasena

6th Floor Commonwealth building

Imperial College Faculty of Medicine

Hammersmith Hospital

1 Du Cane Road

2 London W12 0NN

3 UK.

4 E-mail: [c.jayasena@imperial.ac.uk](mailto:c.jayasena@imperial.ac.uk)

5 ORCID: 0000-0002-2578-8223

6

7 **Funding:** National Institute of Health and Care Research (NIHR) Post-Doctoral Fellowship (PDF-2017-10-  
8 098). The Section of Endocrinology and Investigative Medicine is funded by grants from the MRC, NIHR and  
9 is supported by the NIHR Biomedical Research Centre (BRC) Funding Scheme and the NIHR/Imperial Clinical  
10 Research Facility. The views expressed are those of the authors and not necessarily those of the above-  
11 mentioned funders, the NHS, the NIHR, or the Department of Health. The following authors have grant funding  
12 as follows: AS, Imperial College Healthcare Charity Fellowship; CNJ, NIHR Post-Doctoral Fellowship &  
13 Imperial BRC. Cambridge Weight Plan® Ltd, Corby, Northants, UK, and LighterLife®, Harlow, Essex, UK,  
14 provided food products free of charge, without condition, and without any other involvement in the study.  
15 LogixX Pharma, Theale, Berkshire, UK, sponsored the flow cytometer and established the TUNEL assay to  
16 determine sperm DNA fragmentation.

17

18 **Role of funder:** NIHR funded the study, but was not involved in its design, conduct or analysis. Cambridge  
19 Weight Plan® Ltd, Corby, Northants, UK, and LighterLife®, Harlow, Essex, UK, provided food products free  
20 of charge, without condition, and without any other involvement in the study. LogixX Pharma sponsored the  
21 flow cytometer but was not involved in its design or conduct.

22

**Conflicts of interest:** CNJ holds a researcher-led grant from LogixX Pharma Ltd and is an Editorial Board member for the Journal of Endocrinology and Metabolism. AB reports honoraria from Novo Nordisk, Johnson & Johnson, OHID and Obesity UK outside the submitted work and on the Medical Advisory Board and shareholder of Reset Health Clinics Ltd. ARL receives fees as chair of TDMR Europe. RH is employed by LogixX Pharma.

**Key words:** Male reproduction, Male fertility, Obesity, Spermatogenesis, Weight loss, Sperm quality

## ABSTRACT

**Introduction:** Obesity increases risks of male infertility, but bariatric surgery does not improve semen quality. Recent uncontrolled studies suggest that low energy diet (LED) improves semen quality. Further evaluation within a randomized, controlled setting is warranted.

**Methods:** Men with obesity (18-60 years) with normal sperm concentration (normal-count) (n=24) or oligozoospermia (n=43) were randomised 1:1 to either 800Kcal/day LED for 16-weeks, or control, brief dietary intervention (BDI) with 16-weeks' observation. Semen parameters were compared at baseline and 16-weeks.

**Results:** Mean age of normal-count men was 39.4±6.4 in BDI and 40.2±9.6 years in LED group. Mean age of oligozoospermic men was 39.5±7.5 in BDI and 37.7±6.6 years in LED group. LED caused more weight loss compared with BDI in normal-count (14.4 vs 6.3kg; P<0.001) and oligozoospermic men (17.6 vs 1.8kg; P<0.001). Compared with baseline, in normal-count men total motility (TM) increased 48±17% to 60±10% (P<0.05) after LED, and 52±8% to 61±6% (P<0.0001) after BDI; progressive motility (PM) increased 41±16% to 53±10% (P<0.05) after LED, and 45±8% to 54±65% (P<0.001) after BDI. In oligozoospermic men compared with baseline: TM increased 35%[26] to 52%[16] (P<0.05) after LED, and 43%[28] to 50%[23] (P=0.0587) after BDI; PM increased 29%[23] to 46%[18] (P<0.05) after LED, and 33%[25] to 44%[25] (P<0.05) after BDI.

1 No differences in post-intervention TM or PM were observed between LED and BDI groups, in normal-count  
2 or oligozoospermic men.

3 **Conclusion:** LED or BDI may be sufficient to improve sperm motility in men with obesity. The effects of  
4 paternal dietary intervention on fertility outcomes require investigation.

## 6 **Introduction**

7 Infertility is defined as the inability to conceive after 12 months of regular, unprotected intercourse.(1) Poor  
8 sperm quality is the most common indication for assisted reproductive technologies (ART) such as in vitro  
9 fertilisation (IVF), and global sperm counts are declining (2). IVF is an effective yet prohibitively expensive  
10 treatment for male infertility which is unavailable to most couples and public healthcare systems worldwide (3).  
11 Gonadotrophin therapy can restore fertility in men with hypogonadotropic hypogonadism (HH) (4). However,  
12 there is currently no pharmacological therapy able to improve semen quality in the majority of men with poor  
13 semen quality without HH. Novel and affordable therapies are therefore needed to improve sperm quality in  
14 couples affected by male infertility.

15 Body mass index (BMI) is inversely correlated with sperm quality in men (5). Forty percent of all men  
16 investigated for infertility are reported to be living with overweight or obesity (6). Bariatric surgery is the most  
17 efficacious treatment for obesity, but does not improve, and may even worsen semen quality in men (7,8). This  
18 suggests that extreme, rapid weight loss may have some negative effects on sperm quality (9), at least in the  
19 short term. Recently published, uncontrolled interventional studies have reported improvements in sperm  
20 concentration during intensive programmes of dietary weight loss in men with obesity (10,11) suggesting that  
21 lifestyle interventions are a potential, novel therapy for male infertility. However, the extent of dietary weight  
22 loss required to improve semen quality in men with obesity, has not been investigated previously.

1 Male infertility is a marker of comorbidities including cardiovascular disease, diabetes mellitus and  
2 osteoporosis (12,13). Since available treatments for infertile men are limited, the possible contribution of  
3 obesity in men with no other risk factors for infertility requires investigation.

4 We conducted randomised, controlled studies comparing the effects of high- versus low-intensity dietary  
5 interventions on semen parameters in men with obesity. Men with obesity and either normal-count (Study A) or  
6 oligozoospermia (Study B) were randomised to either a 16-week low energy diet (LED), or a single, brief  
7 dietary intervention (BDI) according to National Health Service (NHS) guidelines with 16 weeks of  
8 observation.

## 10 **Materials and Methods**

11 **Governance and study recruitment:** The studies were granted ethical approval by the London-Queen Square  
12 Research Ethics Committee (Registration number 18/LO/0376) and conducted in accordance with the principles  
13 of the Declaration of Helsinki. Participants were recruited through local online and paper advertisements,  
14 clinics, Andrology Department at Hammersmith Hospital, Imperial College Healthcare NHS Trust and primary  
15 care clinics within North-West London, UK. Participants were invited to a screening visit to ensure they were  
16 eligible based on study's inclusion and exclusion criteria. Inclusion criteria consisted of men aged between 18-  
17 60 years with a BMI  $\geq 30$  kg/m<sup>2</sup> and normal sperm count or oligozoospermia (sperm concentration  $< 15 \times 10^6$ /ml)  
18 for study A and B, respectively. Exclusion criteria were as follows: any medical condition likely to affect  
19 testicular function such as chronic or acute systemic illnesses, undescended testes, significant smoking history,  
20 medications with adverse effect on sperm, clinical evidence of varicocele, azoospermia. All participants  
21 provided written informed consent prior to randomisation. Twenty-four men completed study A and 43 men  
22 completed study B.

**Protocol:** The participants were assessed twice (screening visit and visit 1) before they were commenced on the dietary intervention (Supplemental Figure 1 (14)). During the two visits, baseline measurements of body composition, and semen analysis were performed. Eligibility for enrolment was assessed during the screening visit, based on study's inclusion and exclusion criteria. Baseline semen parameters used later in data analysis were averaged from the two semen samples for each participant. During visit 1, all participants were randomised in a 1:1 ratio to either formula LED or BDI. LED products consisted of soups, shakes, and bars. In the first 12 weeks, participants randomised to LED were provided diet products providing 800 Kcal/day (i.e. four Cambridge Weight Plan® products/day in study A, and four LighterLife® products combined with 400 ml of semi-skimmed milk/day in study B) to achieve weight loss (weight loss phase); in the remaining 4 weeks, participants were gradually reintroduced to food by replacing one LED product every 1-2 weeks with low carbohydrate, high protein meals. Participants randomised to BDI were provided a single, brief (10 minute) intervention consisting of NHS standard dietary recommendations based on 'The Eatwell Guide' (NHS Eatwell) (15) which was then reinforced at every visit over the 16-week period. All participants attended either 4 (study A) or 5 (study B) subsequent visits until 16 weeks following randomisation. All men were advised to limit physical activity to resistance training of 30 minutes three times a week, and to maintain an exercise log throughout the study.

**Clinical and biochemical parameters:** body weight and composition (percentage and absolute values of fat mass, lean mass, and water) were measured using bioelectrical impedance analysis using the Tanita MC-780MA P (Tanita Corp., Tokyo, Japan). Height was recorded during the screening visit. BMI was calculated with the formula (weight (kg) / height (m)<sup>2</sup>). Waist circumference was measured with a tape measure placed halfway between the end of ribs and superior iliac crests.

Morning fasting blood samples (up to 11 am) were performed at screening and final visit for measurements of luteinizing hormone (LH), follicle-stimulating hormone (FSH), testosterone and sex hormone-binding globulin (SHBG). These were analysed in the clinical biochemistry department of Charing Cross Hospital using Abbott

1 ARCHITECT, an automated immunoassay platform under UKNEQAS accreditation. Reference ranges for  
2 males were as follows: LH, 2-12 IU/L; FSH, 1.7-8 IU/L; testosterone, 10-30 nmol/L; SHBG, 15-55 nmol/L.

3 Semen samples were collected on-site in a designated private room in the Andrology Department of  
4 Hammersmith Hospital. Semen handling and manual semen analysis were performed according to the 5th  
5 Edition of the WHO Manual for the Laboratory Examination and Processing of Human Semen (16). All  
6 samples were analysed by experienced biomedical scientists within a specialist hospital laboratory accredited by  
7 the UK National External Quality Accreditation Service (NEQAS). We used manual semen analysis which is  
8 the gold standard method recommended by WHO manual, 5<sup>th</sup> edition (16). For this reason, computer-assisted  
9 sperm analysis (CASA) was not used for this study. Sperm motility was determined as the percentage of  
10 progressive motile, non-progressive motile, and immotile spermatozoa by scoring at least 200 spermatozoa/slide  
11 (16). Normal sperm morphology was examined on Papanicolaou pre-stained slides, using strict criteria (17).  
12 The total motile sperm count was calculated by multiplying the concentration by the volume and the fraction of  
13 motile sperm. The lower reference limit for sperm parameters according to WHO manual, 5<sup>th</sup> edition was as  
14 follows: semen volume, 1.5 ml; total sperm count, 39 million/ejaculate; sperm concentration, 15 million/ml;  
15 total motility, 40%; progressive motility, 32%; morphology, 4% (16). The interobserver coefficient of variation  
16 (CV) are provided here: progressive motility, 6.92%; non-progressive motility, 23.95 %; immotile: 6.67%;  
17 morphology: 17.43%.

18 Semen samples were also assessed for oxidative stress using two markers, namely sperm DNA fragmentation  
19 index (DFI) and Reactive Oxygen Species (ROS). Sperm DFI was measured with an established TUNEL  
20 (terminal uridine nick-end labelling) assay using the Apo-Direct™ kit (Pharmingen, San Diego, CA, USA)  
21 according to previously described protocol (18). TUNEL assay directly measures single- and double-strand  
22 DNA breaks using the enzyme terminal deoxynucleotidyl transferase (TdT) to catalyse the attachment of  
23 florescent labels or deoxyribonucleotides to the 3'-hydroxyl 'free ends' of single and double DNA breaks  
24 ('nicks') (19). The fluorescence, which is proportional to the number of strand breaks, is then quantified using

1 flow cytometry (BD Accuri C6 Plus Flow Cytometer; Becton Dickinson, San Jose, CA, USA). The more DNA  
2 strand break sites are present, the more labels are incorporated within a cell.

3 ROS was measured using an established in-house validated chemiluminescence assay (20). In brief, 400µL neat  
4 (native) semen mixed with 100µL working solution containing luminol. Each sample was vortexed to evenly  
5 disperse the samples before taking luminometer readings (GloMax; Promega Corporation; Madison, WI, USA).  
6 For each negative and positive controls, and specimen assay, 10 readings were taken every minute for ten  
7 minutes and the means were used. Chemiluminescence was expressed as mean relative light units per second  
8 (RLU/sec). ROS value was calculated via the following formula:

$$9 \text{ ROS} = \frac{\text{Mean seminal sample chemiluminescence} - \text{Negative control chemiluminescence}}{\text{Sperm concentration}}$$

10 The reference range for semen ROS was <3.8 RLU/sec/million sperm (20). In-house validation was performed  
11 daily to ensure consistent positive and negative calibration.

### 13 **Statistical analysis and sample size**

14 Hakonsen et al reported that weight loss increased total sperm count by 193 million (95% confidence interval  
15 45-341) (10). Based on this data and, assuming that NHS dietary advice would have no effect on semen  
16 parameters, we estimated that 10 subjects would be required to detect a significant increment in semen  
17 parameters during LED compared with a non-effective comparator, with 80% power ( $\alpha=0.05$ , two-sided).  
18 Accounting for dropouts, we aimed to recruit 12 per group in study A. Sample size for Study B was increased  
19 due to the greater standard deviation in semen parameters in men with oligozoospermia compared with normal-  
20 count and was based on previous unpublished pilot data. Quantitative data were assessed for normality using the  
21 Shapiro-Wilk Normality test. Data are presented as mean  $\pm$  standard deviation if normally distributed and  
22 median [interquartile range] if not normally distributed. Comparisons between the groups were calculated as  
23 independent samples t-test for normally distributed data; not normally distributed data was compared using a



1 Mann Whitney U-Test. Paired samples t-test and Wilcoxon signed rank test for parametric and non-parametric  
2 data, respectively, were carried out to assess the differences at baseline and end of intervention. Categorical data  
3 was compared using a Chi Squared or Fisher's Exact test. Correlations were calculated using Pearson's  
4 correlation coefficient or Spearman's rank correlation coefficient for normally or not normally distributed data,  
5 respectively.

6 Outcome data following the end of intervention were analysed using Analysis of Covariance (ANCOVA), with  
7 the post intervention data as the response, and the baseline measurements as an adjusting covariate. Fitting  
8 fixed effects in the regression model for study, group, and the interaction between these two terms was  
9 performed to examine for group differences (LED vs. BDI) between the two studies. Where an interaction was  
10 shown to be statistically significant, this was retained in the model and the effects of the intervention were  
11 quantified separately for each study. Outcomes meeting the assumptions of the linear regression model were  
12 analysed on the original scale of measurement. Outcomes with a strongly positively skewed distribution were  
13 analysed on the log scale. In all cases,  $P < 0.05$  was considered statistically significant. Statistical analyses were  
14 performed using Prism v.9 (GraphPad Software Inc, La Jolla, CA, USA) and Stata v.15.1 (StataCorp.  
15 2017. Stata Statistical Software: Release 15. College Station, TX: StataCorp LLC).

## 17 **Results**

18 *Study A. Effects of weight loss on semen parameters in men with obesity and normal-count sperm.*

19 The mean age did not differ between study groups ( $39.4 \pm 6.4$  years, BDI;  $40.2 \pm 9.6$  years, LED;  $p=0.84$ ).  
20 Baseline BMI was not significantly different between the groups ( $36.2 \pm 2.4$  kg/m<sup>2</sup>, BDI;  $34.8 \pm 3.8$  kg/m<sup>2</sup>,  
21 LED;  $p=0.29$ ). Baseline semen characteristics were also similar between the two groups, except for total sperm  
22 count and total motile count (Table 1). Remaining baseline demographic and anthropometric characteristics are  
23 shown in Supplemental Table 1 (14).

1 Men in the LED group lost a mean of nearly 3-fold more weight compared with the BDI group during the 16-  
2 week study period ( $6.3 \text{ kg} \pm 6.4$ , BDI;  $17.6 \text{ kg} \pm 7.7$ , LED;  $p < 0.001$ ) (Supplemental Figure 2A- (14)). Sperm  
3 concentration at the end of intervention did not differ significantly between the BDI and LED groups.  
4 Furthermore, sperm concentration did not increase significantly from baseline following either BDI or LED  
5 (Figure 1 A, B, C).

6 Total motility (TM) at the end of the intervention did not differ significantly between the BDI and LED groups.  
7 However, TM increased significantly following both BDI ( $61 \pm 6\%$  vs.  $52 \pm 8\%$ ;  $p < 0.0001$ ) and LED ( $60 \pm$   
8  $10\%$  vs.  $48 \pm 17\%$ ;  $p < 0.05$ ) compared with baseline (Figure 1 D, E, F)

9 Progressive motility (PM) at the end of intervention did not differ between the BDI and LED groups. However,  
10 PM increased significantly following both BDI ( $54 \pm 6\%$  vs.  $45 \pm 8\%$ ;  $p < 0.001$ ) and LED ( $53 \pm 10\%$  vs.  $41 \pm$   
11  $16\%$ ;  $p < 0.05$ ) compared with baseline (Figure 1 G, H, I).

12 In addition to conventional semen parameters, we investigated the effect of weight loss on sperm DNA  
13 fragmentation in men with obesity and normal-count. Sperm DNA fragmentation was significantly lower in the  
14 LED group compared with BDI group at the end of the intervention period (DFI:  $18.3\%$  (6.8) BDI;  $4.1\%$  (8),  
15 LED;  $p < 0.001$ ) (Figure 2A). Regression analysis also showed mean DFI in LED to be lower by 9.5 units  
16 compared to DFI in BDI group (95% CI:  $-16.4$ ,  $-2.5$ ;  $p = 0.009$ ) (Table 2). In men with obesity during LED  
17 intervention, the DFI decreased, but this was non-significant (Figure 2B). There was a significant inverse  
18 correlation ( $r = -0.56$ ,  $p = 0.005$ ) between weight loss and DFI values at the end of the intervention (Figure 2D),  
19 but the correlation between reduction in DFI with weight loss was non-significant (Supplemental Figure 3 (14)).

20 Three men randomised to LED intervention had asthenospermia. After excluding these from the analysis,  
21 statistically significant increase in sperm motilities during LED were no longer observed in the remaining  
22 participants. The findings in sperm concentration and DFI remained unchanged (Supplemental Figure 4 (14)).  
23

1 *Study B. Effects of weight loss on semen parameters in men with obesity and oligozoospermia.*

2 Baseline characteristics including demographics, anthropometric, metabolic and hormonal parameters are  
3 shown in Table 1 and Supplemental Table 2 (14). Mean weight loss in LED group was significantly higher  
4 compared to BDI group (14.4 Kg  $\pm$  5.3, LED vs. 1.8 kg  $\pm$  4.2, BDI;  $p < 0.0001$ ) (Supplemental Figure 2B (14)).

5 Baseline semen parameters did not differ between the two groups except for sperm morphology (Table 2).  
6 Sperm concentration at the end of intervention did not differ significantly between the BDI and LED groups or  
7 change significantly from baseline following either BDI or LED (Figure 3 A, B, C). The degree of weight loss  
8 did not correlate with semen quality (Supplemental Table 3 (14)).

9 TM at the end of intervention did not differ significantly between the BDI and LED groups. TM increased  
10 significantly following the LED intervention (52% [16] vs. 35% [26],  $p < 0.05$ ), though TM following the BDI  
11 was non-significant (50% [23] vs. 43% [28],  $p = 0.0587$ ) (Figure 3 D, E, F).

12 PM at the end of intervention did not differ significantly between the BDI and LED groups. However, PM  
13 increased significantly following both the BDI (44% [25] vs. 33% [25];  $P < 0.05$ ) or LED (46% [18] vs. 29%  
14 [23];  $p < 0.05$ ) groups compared with baseline (Figure 3 G, H, I).

15 At baseline, approximately half of the participants in both interventions had reduced TM and PM as defined by  
16 the 2010 WHO reference range (16) (TM  $< 40\%$  in 11/20, LED group and 12/23, BDI group; PM  $< 32\%$  in  
17 12/20, LED group and 11/23, BDI group). In a subgroup analysis, in men with asthenospermia TM and PM at  
18 the end of intervention were significantly higher with LED compared to BDI. There were no differences at the  
19 end of intervention in the men with normal sperm motilities at baseline (Supplemental Figure 5 (14)).

20 Sperm DFI at the end of intervention did not differ significantly between the BDI and LED groups or change  
21 significantly from baseline following either BDI or LED (Supplemental Figure 6 (14)). We observed no  
22 statistically significant difference in semen ROS between the two groups at the end of intervention (ROS in  
23 RLU/sec/million sperm: 5.04 (36.5) BDI; 13.02 (36.14), LED;  $p = 0.55$ ). In a subgroup of men with elevated

1 ROS at baseline (validated threshold  $\leq 3.8$ ), 36% men normalised their ROS following the formula LED  
2 compared with 12.5% with the BDI ( $p=0.34$ ) (data not provided).

### 3 *Combining the two studies*

4 The distinction between the two studies were based on the sperm concentration WHO 2010 cut off value (16).  
5 We combined the data from the two studies as participants received the same dietary interventions per group  
6 and were followed up for the same period. Data were adjusted for baseline measurements. We observed no  
7 difference in sperm concentration, TM, PM, semen volume, sperm count, total motile count, and morphology  
8 between the two intervention groups (Table 2). We additionally investigated for possible differences according  
9 to BMI severity. The results did not change after adjusting for baseline BMI. (Supplemental Table 4 (14)).

10 We analysed levels of serum reproductive hormones from both studies. No differences between serum  
11 reproductive hormone levels were observed between BDI and LED groups at the end of the study. In the LED  
12 group, levels of serum testosterone and SHBG increased significantly from baseline to end of intervention. In  
13 the BDI group, levels of serum LH reduced significantly from baseline to end of study. No other before-after  
14 hormonal changes were observed (Figure 4). Analysis of reproductive hormones for each study separately are  
15 shown in Supplemental Figures 7 & 8 (14). Baseline reproductive hormone profile is shown in Table 1.

### 17 **Discussion**

18 A recent meta-analysis concluded that reported mean sperm counts in both high and low / middle-income  
19 countries, have halved during the last 50 years (21). In the absence of drug therapies, it is important to develop  
20 simple and affordable interventions to improve semen quality, and therefore male fertility. We expected that an  
21 intensive programme of dietary weight loss would be superior to a control, brief dietary intervention for  
22 improving semen quality; surprisingly, similar improvements in sperm motility were observed following both  
23 dietary interventions. However, high intensity diet had a more favourable effect in improving sperm motilities  
24 in men with asthenospermia. Similarly, when men with asthenospermia were excluded, there were no

1 significant increases in TM and PM during LED intervention. This suggests that improvements may be more  
2 pronounced in men with asthenospermia. Our data therefore supports the growing view that dietary  
3 interventions may potentially be a method of improving fertility outcomes in couples in whom the male partner  
4 has obesity. However, we also suggest for the first time that even mild dietary interventions may be sufficient to  
5 improve semen quality, which may greatly increase the accessibility of fertility treatment for couples currently  
6 unable to access support.

7 Global rates of obesity have tripled since 1975 (22). Obesity may impair semen quality through several  
8 mechanisms including suppressed hypothalamic gonadotrophin releasing hormone (GnRH) secretion, increased  
9 aromatisation of androgens to oestrogens, insulin resistance, oxidative and heat stress within the testes (23–25).  
10 However, the role of male obesity in semen parameters remains less clear (26,27). The paucity of studies  
11 investigating the role of weight loss on sperm parameters and ART outcomes creates an evidence gap of  
12 uncertainty. This may reflect the technical challenge of studying semen parameters which have high degrees of  
13 biological variation. Furthermore, the relationship between weight loss and semen parameters may be complex,  
14 as demonstrated by lack of improvement of semen parameters with bariatric surgery (28). This may reflect that  
15 either significant energy restriction or inflammatory effects during substantial weight loss have negative effects  
16 on male reproductive function. We therefore hypothesised that a smaller degree of weight loss could ameliorate  
17 the detrimental effects of obesity on semen quality, without the impairments of semen quality observed during  
18 greater absolute weight loss. Our protocol for weight loss and sample size were based on pilot studies  
19 suggesting that a threshold of 12kg of weight loss would significantly improve semen quality in men with  
20 obesity. Independent ethical review recommended that participants with obesity randomised to the control  
21 intervention should receive the level of dietary support available to the general population. Therefore, we  
22 provided a brief (10 minutes) intervention which is openly available online from the NHS, providing advice on  
23 healthy eating (15). Participants randomised to the BDI arm lost on average 3-fold less weight compared with  
24 the LED arm, this however, did not result in greater improvements in sperm motility between the groups.  
25 Collectively, our data support the view that weight loss via dietary modification may improve semen quality in

1 men. We also suggest for the first time that a universally available public health intervention could also provide  
2 clinical benefit for improving reproductive potential in men with obesity. Although our initial results are  
3 promising, larger studies with pregnancy outcomes are required. Our data would require replication in a larger  
4 study powered to live birth outcomes; if confirmed, dietary intervention might have substantial future potential  
5 for improving reproductive health in couples with infertility, including those in low-to-middle income countries  
6 without access to assisted reproductive technologies.

7 The current study broadly supports a limited number of prior studies suggesting that dietary intervention may  
8 improve semen quality in men with obesity. Hakonsen et al. studied the effects of a 14-week residential diet and  
9 exercise programme in 27 men with a mean BMI 44kg/m<sup>2</sup>; improvements in total sperm count and semen  
10 volume were observed in the 10 participants with greatest weight loss (mean 25.4%) (10). In the recently  
11 published S-Lite study (11), men with obesity underwent a mean 16.5kg weight loss during 8-weeks of non-  
12 randomised LED intervention; total sperm count increased by 40% vs. baseline, and sperm concentration  
13 increased by 49% vs. baseline. All participants were then randomised to one of four 52-week intervention  
14 groups: placebo; exercise and placebo; GLP-1 agonist; both exercise and GLP-1 agonist. S-Lite failed to  
15 demonstrate differences in semen parameters among the intervention groups. However, post-hoc analysis  
16 suggested that improvements in semen parameters were sustained in men who maintained a median weight loss  
17 of more than 11.7kg at 52 weeks regardless of the randomised group. Mir et al. studied the effects of a 'healthy  
18 diet and exercise' programme in 105 men (mean BMI 33.2 kg/m<sup>2</sup>) and reported significant improvements in  
19 sperm morphology and PM with approximately 10% of BMI reduction (29). In summary, there is concordance  
20 among our and other studies that intensive forms of dietary intervention are associated with improved semen  
21 quality in men with obesity. However, our study is the first to suggest that a single, publicly available  
22 intervention may also be sufficient to improve semen quality in men with obesity. Larger studies would be  
23 needed to resolve whether LED has differential effects on semen quality compared with BDI in men with  
24 obesity.

1 Sperm DNA fragmentation is an important mechanism of sperm damage (30). Men with elevated DFI are  
2 known to have adverse fertility outcomes (30) and an increased risk of recurrent pregnancy loss (31–33). We  
3 observed a 4-fold improvement in sperm DFI for normal-count men with obesity during LED compared with  
4 BDI. This finding was replicated during regression analysis after adjusting for baseline value. This is of  
5 particular importance as seminal oxidative stress and sperm DNA fragmentation have widely been accepted as  
6 major causes of male infertility (34). In our study, participants losing more weight were more likely to have  
7 lower DFI sperm values by the end of the study. However, the correlation of weight loss with DFI reduction  
8 was not significant. Our results were not replicated in oligozoospermic men; this may reflect that DFI could not  
9 be measured in some participants owing to an insufficient sperm concentration required for the TUNEL assay.  
10 We noticed similar results with analysis of covariance. Our data are in concordance with two recent studies.  
11 Mir et al. observed improvements in sperm chromatin dispersion, an indirect marker of DFI, in men undergoing  
12 a 12-week diet and exercise programme (29). Faure et al. observed reductions in TUNEL DFI in six men after  
13 3-8 months of a dietary weight loss programme (35). However, Hakonsen et al failed to observe any significant  
14 improvements in sperm chromatin structure assay in men with morbid obesity undergoing weight loss and  
15 exercise (10). Hence, our study data suggest that in men with obesity, the degree of dietary weight loss may be  
16 correlated with degree of sperm DNA damage. Further studies would be needed to determine if different  
17 methods of weight loss would provide clinically significant differences in sperm DNA fragmentation which  
18 were sufficient to affect live birth outcomes in couples with male infertility. Seminal oxidative stress is a  
19 potential mediator of sperm damage and DNA fragmentation in men (36). Seminal ROS are correlated with  
20 reduced fertility outcomes (37,38) and recurrent pregnancy loss (39). One study of six men observed significant  
21 increase in seminal superoxide dismutase protein 2 levels, which is a ROS scavenger, with abdominal fat loss  
22 (35). We observed reductions in ROS during weight loss, but not significant differences between the study  
23 groups. Further studies are needed to investigate whether reductions in semen ROS represent a mechanism for  
24 improving semen quality in men during weight loss.

1 We additionally measured serum LH, FSH, testosterone and SHBG in men with normal-count and  
2 oligozoospermia. On combined analysis, serum testosterone and SHBG increased significantly following LED  
3 but not BDI. This is expected as it is widely known that weight loss either through dietary intervention or  
4 bariatric surgery results in rising of testosterone and SHBG (40–43). However, post-treatment levels were not  
5 different between the groups. Caloric restriction is known to suppress pulsatile GnRH / LH secretion (44); we  
6 are unable to explain why LH reduced significantly following BDI, but not following LED which provides  
7 more extensive weight loss.

8 This study utilised a randomised design and measured WHO and advanced semen parameters using validated  
9 assay techniques. A further strength of our study is its investigation of an effectively cost-free intervention  
10 available to the general public. Furthermore, unlike some published studies in this field, our study period was  
11 long enough to investigate effects of interventions on a complete cycle of spermatogenesis in men (range 42 to  
12 76 days) (45). The semen analyses were performed by biomedical scientists in the andrology laboratory who  
13 were not part of the study team and were blinded to the intervention given. However, all interventional studies  
14 of semen quality are limited by large, biological variations observed in semen parameters of men (46). This  
15 probably prohibited any associations of weight loss with semen quality. The study failed to detect any  
16 significant differences in its primary outcome (sperm concentration). Positive findings were shown regarding  
17 sperm motilities (secondary outcomes), but one needs to consider that these would have been underpowered.  
18 We cannot exclude that undetected differences between semen parameters may exist between men in the  
19 intervention groups. Additionally, the observed improvements in sperm motilities (but not in concentration)  
20 raise the possibility of regression to the mean. We attempted to mitigate this phenomenon at the design stage by  
21 including a randomly allocated placebo and intervention group and taking two baseline measurements for  
22 semen parameters (47). We cannot exclude though that the observed changes in sperm motility were by random  
23 chance. Many men were screened to find sufficient participants for the study. We speculate that the  
24 embarrassment of producing semen samples together with the social stigma of male infertility (48) may have  
25 contributed to the relative low number of patients who agreed to participate. We also cannot exclude the



1 possibility that participants may have voluntarily changed their exercise behavior during the study, and the  
2 potential effects this may have had on our results. Furthermore, it should be recognized that regular visits to see  
3 a medical team within a specialist hospital may have encouraged weight loss in the BDI group which might not  
4 be achieved within a community setting. Lastly, compliance with diet was not measured. Eight to sixteen weeks  
5 duration of LED or very low energy diets (VLED) results in 10-15% of weight loss (49,50). In our cohort, 25  
6 out of 32 men (78%) receiving LED achieved >10% weight loss and only 1 man had <8% weight loss  
7 indicating a moderate compliance with LED diet. Prior studies have reported 53%-94% of participants  
8 achieving the target of >10% weight loss with LED or VLED (51–53). Various factors have been deemed  
9 barriers to adherence to LED/ VLED meal replacement such as product unpalatability, unrealistic weight loss  
10 expectations, poor program accessibility, unforeseeable circumstances, and externalised weight-related stigma  
11 (54).

12 In conclusion, our data suggest that dietary interventions resulting in modest degrees of weight loss may be  
13 sufficient to improve sperm motilities in men living with obesity, thus having the potential to improve fertility  
14 in couples with male infertility. Current clinical guidelines for the management of infertility do not identify  
15 weight loss as a potential method of improving semen quality for male partners (55–57). Further studies are  
16 required to determine the feasibility, clinical- and cost-effectiveness of publicly available dietary intervention  
17 programme to improve fertility outcomes for couples with infertility. Such approaches may be particularly  
18 suited to geographical regions with both a high prevalence of obesity and limited fertility healthcare provisions.

19  
20 **Acknowledgments:** The studies were supported by Imperial NIHR Clinical Research Facility and the NIHR  
21 Clinical Research Network (CRN). We would like to particularly thank the staff at the Andrology department of  
22 Hammersmith hospital for their contribution to this study and Mr Paul Basset at Statsconsultancy Ltd for  
23 providing statistical support.

1 **Data availability:** Some or all datasets generated during and/or analyzed during the current study are not  
2 publicly available, but are available from the corresponding author on reasonable request.

#### 4 **References**

- 5 1. World Health Organization (WHO). International Classification of Diseases, 11th Revision (ICD-11).  
6 Geneva: WHO; 2018.
- 7 2. Human Fertilisation and Embryology Authority. Published online 2013. Accessed December 12, 2022.  
8 [www.hfea.gov.uk](http://www.hfea.gov.uk).
- 9 3. Ombelet W, Campo R. Affordable IVF for developing countries. *Reprod Biomed Online*.  
10 2007;15(3):257-265. doi:10.1016/S1472-6483(10)60337-9
- 11 4. Papanikolaou N, Luo R, Jayasena CN. Fertility Considerations in Hypogonadal Men. *Endocrinol Metab*  
12 *Clin North Am*. 2022;51(1):133-148. doi:10.1016/J.ECL.2021.11.009
- 13 5. Sermondade N, Faure C, Fezeu L, et al. BMI in relation to sperm count: an updated systematic review  
14 and collaborative meta-analysis. *Hum Reprod Update*. 2013;19(3):221-231.  
15 doi:10.1093/humupd/dms050
- 16 6. Punab M, Poolamets O, Paju P, et al. Causes of male infertility: a 9-year prospective monocentre study  
17 on 1737 patients with reduced total sperm counts. *Hum Reprod*. 2017;32(1):18-31.  
18 <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med14&NEWS=N&AN=27864361>
- 19 7. Legro RS, Kunesman AR, Meadows JW, et al. Time-related increase in urinary testosterone levels and  
20 stable semen analysis parameters after bariatric surgery in men. *Reprod Biomed Online*. 2015;30(2):150-  
21 156. doi:10.1016/J.RBMO.2014.10.014
- 22 8. di Frega AS, Dale B, Di Matteo L, Wilding M. Secondary male factor infertility after Roux-en-Y gastric  
23 bypass for morbid obesity: Case report. *Hum Reprod*. 2005;20(4):997-998.  
24 doi:10.1093/HUMREP/DEH707
- 25 9. Sermondade N, Massin N, Boitrelle F, et al. Sperm parameters and male fertility after bariatric surgery:  
26 three case series. *Reprod Biomed Online*. 2012;24(2):206-210. doi:10.1016/J.RBMO.2011.10.014
- 27 10. Håkonsen L, Thulstrup A, Aggerholm A, et al. Does weight loss improve semen quality and reproductive  
28 hormones? results from a cohort of severely obese men. *Reprod Health*. 2011;8(1):24. doi:10.1186/1742-  
29 4755-8-24
- 30 11. Andersen E, Juhl CR, Kjøller ET, et al. Sperm count is increased by diet-induced weight loss and  
31 maintained by exercise or GLP-1 analogue treatment: a randomized controlled trial. *Hum Reprod*.  
32 2022;37(7):1414-1422. doi:10.1093/HUMREP/DEAC096
- 33 12. Ferlin A, Garolla A, Ghezzi M, et al. Sperm Count and Hypogonadism as Markers of General Male  
34 Health. *Eur Urol Focus*. 2021;7(1):205-213. doi:10.1016/j.euf.2019.08.001
- 35 13. Latif T, Kold Jensen T, Mehlsen J, et al. Semen Quality as a Predictor of Subsequent Morbidity: A  
36 Danish Cohort Study of 4,712 Men With Long-Term Follow-up. *Am J Epidemiol*. 2017;186(8):910-917.  
37 doi:10.1093/aje/kwx067

14. Sharma A, Papanikolaou N, Abou Sherif S, et al. Data from: Supplemental materials for improvements in sperm motility following low or high intensity dietary interventions in men with obesity. Repository reference pending.
15. The Eatwell Guide - NHS. Accessed November 7, 2022. <https://www.nhs.uk/live-well/eat-well/food-guidelines-and-food-labels/the-eatwell-guide/>
16. WHO laboratory manual for the examination and processing of human semen. Accessed April 28, 2021. <https://www.who.int/publications/i/item/9789241547789>
17. Menkveld R, Stander FS, Kotze TJ, Kruger TF, van Zyl JA. The evaluation of morphological characteristics of human spermatozoa according to stricter criteria. *Hum Reprod*. 1990;5(5):586-592. doi:10.1093/oxfordjournals.humrep.a137150
18. Sharma R, Iovine C, Agarwal A, Henkel R. TUNEL assay-Standardized method for testing sperm DNA fragmentation. *Andrologia*. 2021;53(2):e13738. doi:10.1111/and.13738
19. Ribas-Maynou J, García-Peiró A, Fernández-Encinas A, et al. Comprehensive analysis of sperm DNA fragmentation by five different assays: TUNEL assay, SCSA, SCD test and alkaline and neutral Comet assay. *Andrology*. 2013;1(5):715-722. doi:10.1111/J.2047-2927.2013.00111.X
20. Vessey W, Perez-Miranda A, Macfarquhar R, Agarwal A, Homa S. Reactive oxygen species in human semen: Validation and qualification of a chemiluminescence assay. *Fertil Steril*. 2014;102(6):1576-1583.e4. doi:10.1016/J.FERTNSTERT.2014.09.009
21. Levine H, Jørgensen N, Martino-Andrade A, et al. Temporal trends in sperm count: a systematic review and meta-regression analysis of samples collected globally in the 20th and 21st centuries. *Hum Reprod Update*. Published online November 15, 2022. doi:10.1093/HUMUPD/DMAC035
22. Obesity and overweight. Accessed December 12, 2022. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
23. Davidson LM, Millar K, Jones C, Fatum M, Coward K. Deleterious effects of obesity upon the hormonal and molecular mechanisms controlling spermatogenesis and male fertility. *Hum Fertil (Camb)*. 2015;18(3):184-193. doi:10.3109/14647273.2015.1070438
24. Pasquali R. Obesity and androgens: facts and perspectives. *Fertil Steril*. 2006;85(5):1319-1340. doi:10.1016/J.FERTNSTERT.2005.10.054
25. Craig JR, Jenkins TG, Carrell DT, Hotaling JM. Obesity, male infertility, and the sperm epigenome. *Fertil Steril*. 2017;107(4):848-859. doi:10.1016/J.FERTNSTERT.2017.02.115
26. Campbell JM, Lane M, Owens JA, Bakos HW. Paternal obesity negatively affects male fertility and assisted reproduction outcomes: a systematic review and meta-analysis. *Reprod Biomed Online*. 2015;31(5):593-604. doi:10.1016/J.RBMO.2015.07.012
27. Salas-Huetos A, Maghsoumi-Norouzabad L, James ER, et al. Male adiposity, sperm parameters and reproductive hormones: An updated systematic review and collaborative meta-analysis. *Obes Rev*. 2021;22(1):e13082. doi:10.1111/OBR.13082
28. Lee Y, Dang JT, Switzer N, et al. Impact of Bariatric Surgery on Male Sex Hormones and Sperm Quality: a Systematic Review and Meta-Analysis. *Obes Surg*. 2019;29(1):334-346. doi:10.1007/s11695-018-3557-5
29. Mir J, Franken D, Andrabi SW, Ashraf M, Rao K. Impact of weight loss on sperm DNA integrity in obese men. *Andrologia*. 2018;50(4):e12957. doi:10.1111/AND.12957

- 1 30. Lewis SEM, Aitken RJ, Conner SJ, et al. The impact of sperm DNA damage in assisted conception and  
2 beyond: recent advances in diagnosis and treatment. *Reprod Biomed Online*. 2013;27(4):325-337.  
3 doi:10.1016/J.RBMO.2013.06.014
- 4 31. Carrell DT, Liu L, Peterson CM, et al. Sperm DNA fragmentation is increased in couples with  
5 unexplained recurrent pregnancy loss. *Arch Androl*. 2003;49(1):49-55. doi:10.1080/01485010290099390
- 6 32. McQueen DB, Zhang J, Robins JC. Sperm DNA fragmentation and recurrent pregnancy loss: a  
7 systematic review and meta-analysis. *Fertil Steril*. 2019;112(1):54-60.e3.  
8 doi:10.1016/J.FERTNSTERT.2019.03.003
- 9 33. Dai Y, Liu J, Yuan E, Li Y, Shi Y, Zhang L. Relationship Among Traditional Semen Parameters, Sperm  
10 DNA Fragmentation, and Unexplained Recurrent Miscarriage: A Systematic Review and Meta-Analysis.  
11 *Front Endocrinol (Lausanne)*. 2022;12. doi:10.3389/FENDO.2021.802632
- 12 34. Agarwal A, Said TM. Oxidative stress, DNA damage and apoptosis in male infertility: a clinical  
13 approach. *BJU Int*. 2005;95(4):503-507. doi:10.1111/j.1464-410X.2005.05328.x
- 14 35. Faure C, Dupont C, Baraibar MA, et al. In Subfertile Couple, Abdominal Fat Loss in Men Is Associated  
15 with Improvement of Sperm Quality and Pregnancy: A Case-Series. *PLoS One*. 2014;9(2):e86300.  
16 doi:10.1371/JOURNAL.PONE.0086300
- 17 36. Agarwal A, Makker K, Sharma R. Clinical relevance of oxidative stress in male factor infertility: an  
18 update. *Am J Reprod Immunol*. 2008;59(1):2-11. doi:10.1111/J.1600-0897.2007.00559.X
- 19 37. Bisht S, Faiq M, Tolahunase M, Dada R. Oxidative stress and male infertility. Published online 2017.  
20 doi:10.1038/nruirol.2017.69
- 21 38. Agarwal A, Parekh N, Selvam MKP, et al. Male Oxidative Stress Infertility (MOSI): Proposed  
22 Terminology and Clinical Practice Guidelines for Management of Idiopathic Male Infertility. *World J*  
23 *Mens Health*. 2019;37(3):296-312. doi:10.5534/WJM.H.190055
- 24 39. Jayasena CN, Radia UK, Figueiredo M, et al. Reduced Testicular Steroidogenesis and Increased Semen  
25 Oxidative Stress in Male Partners as Novel Markers of Recurrent Miscarriage. *Clin Chem*.  
26 2019;65(1):161-169. doi:10.1373/CLINCHEM.2018.289348
- 27 40. Stanik S, Dornfeld LP, Maxwell MH, Viosca SP, Korenman SG. The Effect of Weight Loss on  
28 Reproductive Hormones in Obese Men. *J Clin Endocrinol Metab*. 1981;53(4):828-832.  
29 doi:10.1210/JCEM-53-4-828
- 30 41. Strain G, Zumoff B, Rosner W, Pi-Sunyer X. The relationship between serum levels of insulin and sex  
31 hormone-binding globulin in men: the effect of weight loss. *J Clin Endocrinol Metab*. 1994;79(4):1173-  
32 1176. doi:10.1210/jcem.79.4.7962291
- 33 42. Kaukua J, Pekkarinen T, Sane T, Mustajoki P. Sex hormones and sexual function in obese men losing  
34 weight. *Obes Res*. 2003;11(6):689-694. doi:10.1038/oby.2003.98
- 35 43. Niskanen L, Laaksonen DE, Punnonen K, Mustajoki P, Kaukua J, Rissanen A. Changes in sex hormone-  
36 binding globulin and testosterone during weight loss and weight maintenance in abdominally obese men  
37 with the metabolic syndrome. *Diabetes Obes Metab*. 2004;6(3):208-215. doi:10.1111/j.1462-  
38 8902.2004.00335.x
- 39 44. Martin B, Golden E, Carlson OD, Egan JM, Mattson MP, Maudsley S. Caloric restriction: impact upon  
40 pituitary function and reproduction. *Ageing Res Rev*. 2008;7(3):209-224. doi:10.1016/j.arr.2008.01.002
- 41 45. Heller CG, Clermont Y. Spermatogenesis in Man: An Estimate of Its Duration. *Science (80- )*.

1 1963;140(3563):184-186. doi:10.1126/SCIENCE.140.3563.184

- 2 46. Schwartz D, Laplanche A, Jouannet P, David G. Within-subject variability of human semen in regard to  
3 sperm count, volume, total number of spermatozoa and length of abstinence. *Reproduction*.  
4 1979;57(2):391-395. doi:10.1530/JRF.0.0570391
- 5 47. Barnett AG, van der Pols JC, Dobson AJ. Regression to the mean: what it is and how to deal with it. *Int J*  
6 *Epidemiol*. 2005;34(1):215-220. doi:10.1093/ije/dyh299
- 7 48. Nachtigall RD, Becker G, Wozny M. The effects of gender-specific diagnosis on men's and women's  
8 response to infertility. *Fertil Steril*. 1992;57(1):113-121.
- 9 49. Johansson K, Neovius M, Hemmingsson E. Effects of anti-obesity drugs, diet, and exercise on weight-  
10 loss maintenance after a very-low-calorie diet or low-calorie diet: a systematic review and meta-analysis  
11 of randomized controlled trials. *Am J Clin Nutr*. 2014;99(1):14-23. doi:10.3945/ajcn.113.070052
- 12 50. Leeds AR. Formula food-reducing diets: A new evidence-based addition to the weight management tool  
13 box. *Nutr Bull*. 2014;39(3):238-246. doi:10.1111/nbu.12098
- 14 51. Scragg J, Hallsworth K, Taylor G, et al. Factors associated with engagement and adherence to a low-  
15 energy diet to promote 10% weight loss in patients with clinically significant non-alcoholic fatty liver  
16 disease. *BMJ open Gastroenterol*. 2021;8(1). doi:10.1136/bmjgast-2021-000678
- 17 52. Christensen P, Frederiksen R, Bliddal H, et al. Comparison of three weight maintenance programs on  
18 cardiovascular risk, bone and vitamins in sedentary older adults. *Obesity (Silver Spring)*.  
19 2013;21(10):1982-1990. doi:10.1002/oby.20413
- 20 53. Mathus-Vliegen EMH. Long-term maintenance of weight loss with sibutramine in a GP setting following  
21 a specialist guided very-low-calorie diet: a double-blind, placebo-controlled, parallel group study. *Eur J*  
22 *Clin Nutr*. 2005;59 Suppl 1:S31-8; discussion S39. doi:10.1038/sj.ejcn.1602172
- 23 54. Maston G, Franklin J, Hocking S, et al. Dietary adherence and program attrition during a severely  
24 energy-restricted diet among people with complex class III obesity: A qualitative exploration. *PLoS One*.  
25 2021;16(6):e0253127. doi:10.1371/journal.pone.0253127
- 26 55. Schlegel PN, Sigman M, Collura B, et al. Diagnosis and treatment of infertility in men: AUA/ASRM  
27 guideline part II. *Fertil Steril*. 2021;115(1):62-69. doi:10.1016/j.fertnstert.2020.11.016
- 28 56. Minhas S, Bettocchi C, Boeri L, et al. European Association of Urology Guidelines on Male Sexual and  
29 Reproductive Health: 2021 Update on Male Infertility. *Eur Urol*. 2021;80(5):603-620.  
30 doi:10.1016/j.EURURO.2021.08.014
- 31 57. Overview | Fertility problems: assessment and treatment | Guidance | NICE.

## 32 33 **Legends**

### 34 **Figure 1. Effects of dietary interventions on semen parameters in normal-count men**

35 Bar graphs of **A.** sperm concentration ( $58.5 \times 10^6 \pm 32.8$ , BDI;  $53.6 \times 10^6 \pm 36.4$ , LED;  $p=0.73$ ), **D.** total  
36 motility ( $61\% \pm 6$ , BDI;  $60\% \pm 10$ , LED;  $p=0.69$ ), and **G.** progressive motility ( $54\% \pm 6$ , BDI;  $53\% \pm 10$ , LED;  
37  $p=0.85$ ) at the end of the dietary intervention in BDI groups and LED groups.

38 Baseline and end of intervention, **B.** sperm concentration ( $p=0.12$ ), **E.** total motility ( $p=0.02$ ) and **H.** progressive  
39 motility ( $p=0.02$ ) in the LED group. Baseline and end of intervention, **C.** sperm concentration ( $p=0.59$ ), **F.** total  
40 motility ( $p < 0.0001$ ) and **I.** progressive motility ( $p < 0.001$ ) in the BDI group.

1 (A, D, G) Data are presented as mean  $\pm$  standard deviation. \*,  $p < 0.05$ ; \*\*\*,  $p < 0.001$

2 BDI: Brief dietary intervention; LED: Low energy diet

3

4 **Figure 2. Effects of dietary interventions on DNA fragmentation index in normal-count men** A.

5 Bar graph of DFI at the end of the dietary intervention between the 2 groups (18.3% (6.8); BDI, 4.1% (8), LED;  
6  $p < 0.001$ ). B. Difference in DFI before and after the dietary intervention in LED group ( $p = 0.23$ ). C. Difference  
7 in DFI before and after the dietary intervention in BDI group ( $p = 0.15$ ). D. Scatter plot of weight loss and DFI as  
8 measured at the end of the study across both groups ( $r = -0.56$ ,  $p = 0.005$ ).

9 (A) Data are given as median with interquartile range. \*\*\*,  $p < 0.001$

10 DFI: DNA Fragmentation Index; BDI: Brief dietary intervention; LED: Low energy diet

11

12 **Figure 3. Effects of dietary interventions on semen parameters in men with oligozoospermia** Bar  
13 graphs of A. sperm concentration ( $8.3 \times 10^6$  [18.3], BDI;  $3.4 \times 10^6$  [13.1], LED;  $p = 0.12$ ), D. total motility (50%  
14 [23], BDI; 52% [16], LED;  $p = 0.48$ ), and G. progressive motility (44% [25], BDI; 46% [18], LED;  $p = 0.67$ ) at  
15 the end of the dietary intervention in the two groups. Baseline and end of intervention, B. sperm concentration  
16 ( $p = 0.82$ ), E. total motility ( $p = 0.02$ ) and H. progressive motility ( $p = 0.01$ ) in LED group. Baseline and end of  
17 intervention, C. sperm concentration ( $p = 0.21$ ), F. total motility ( $p = 0.0587$ ) and I. progressive motility ( $p = 0.01$ )  
18 in BDI group.

19 (A, D, G) Data are given as median with interquartile range. \*,  $p < 0.05$

20 BDI: Brief dietary intervention; LED: Low energy diet

21

22 **Figure 4. Effects of dietary interventions on hormonal parameters after combining the two studies** Bar  
23 graphs of A. LH ( $3.4 \pm 1.5$ , BDI;  $3.5 \pm 1.4$ , LED;  $p = 0.81$ ), D. FSH ( $4.2 \pm 2.3$ , BDI;  $4.6 \pm 2.4$ , LED;  $p = 0.52$ ),  
24 G. Testosterone ( $13.3 \pm 3.7$ , BDI;  $14.8 \pm 5.8$ , LED;  $p = 0.19$ ), and J. SHBG ( $24 \pm 8$ , BDI;  $27 \pm 13$ , LED;  $p =$   
25  $0.30$ ) at the end of the dietary intervention in BDI groups and LED groups.

26 Baseline and end of intervention, B. LH ( $p = 0.70$ ), E. FSH ( $p = 0.26$ ), H. Testosterone ( $p < 0.0001$ ) and K. SHBG  
27 ( $p < 0.001$ ) in the LED group. Baseline and end of intervention, C. LH ( $p < 0.01$ ), F. FSH ( $p = 0.15$ ), I.  
28 Testosterone ( $p = 0.28$ ) and L. SHBG ( $p = 0.15$ ) in the BDI group.

29 (A, D, G, J) Data are presented as mean  $\pm$  standard deviation. \*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$

30 BDI: Brief dietary intervention; LED: Low energy diet

31

32 **Table 1: Baseline anthropometric, semen and reproductive hormonal parameters of men with obesity and normal-**  
33 **count sperm (Study A) and oligozoospermia (Study B) according to diet groups; LED vs. BDI.** Data presented as  
34 mean  $\pm$  standard deviation unless otherwise stated. <sup>1</sup>Total Motile Count (TMC) = Sperm concentration  $\times$  Sperm Volume  $\times$   
35 (Total Motility/100) <sup>2</sup>Contains missing values. \*Denotes where differences between LED and BDI groups were  
36 statistically significant at  $p < 0.05$ . LED, Low energy diet; BDI, Brief dietary intervention; BMI, Body mass index; DFI,  
37 DNA fragmentation index; ROS, Reactive oxygen species; RLU, Relative light units; DFI, DNA Fragmentation Index;  
38 IQR: Interquartile range; LH: Luteinizing hormone; FSH: Follicle stimulating hormone; SHBG: Sex hormone-binding  
39 globulin.

Demographics/Parameters	Dietary intervention groups		
	Study A	LED (n = 12)	BDI (n= 12)
<b>ANTHROPOMETRIC PARAMETERS</b>			
Age (years)	40.2 ± 9.6	39.4 ± 6.4	0.84
Weight (kg)	111.1 ± 12	114.1 ± 11.0	0.53
BMI (kg/m <sup>2</sup> )	35.3 ± 4.1	36.2 ± 2.4	0.52
<b>SEMEN PARAMETERS</b>			
Sperm Concentration (million/ml)	38.3 ± 15	54.2 ± 21.3	0.05
Semen Volume (ml)	3.3 ± 1.4	3.9 ± 1.1	0.28
Total Sperm Count (million/ejaculate) Median [IQR]	99.5 [76.3]	175.2 [72.3]	0.008*
Progressive Motility (%)	41 ± 16	45 ± 8	0.50
Total Motility (%)	48 ± 17	52 ± 8	0.51
Total Motile Sperm Count (million/ejaculate) <sup>1</sup> Median [IQR]	50.4 [62.4]	91.3 [36.1]	0.03*
Normal Morphology (%)	2 ± 1	2 ± 2	0.89
DFI (%) Median [IQR]	5.9 [22.1]	16.3 [10.5]	0.08
<b>REPRODUCTIVE HORMONE PROFILE</b>			
LH (IU/L)	3.2 ± 1.0	3.5 ± 1.7	0.64
FSH (IU/L)	4.2 ± 2.6	4.4 ± 1.9	0.83
Testosterone (nmol/L)	12.7 ± 4.2	12.3 ± 2.5	0.80
SHBG (nmol/L)	22 ± 8	25 ± 9	0.34
Study B	LED (n= 20)	BDI (n= 23)	P value
<b>ANTHROPOMETRIC PARAMETERS</b>			
Age (years)	37.7 ± 6.6	39.5 ± 7.5	0.40
Weight (kg)	116.8 ± 17.3	117.0 ± 21.9	0.90
BMI (kg/m <sup>2</sup> )	38 ± 4.0	38.0 ± 6.3	0.90
<b>SEMEN PARAMETERS</b>			
Sperm Concentration (million/ml) Median [IQR]	5.6 [6.0]	7.8 [8.9]	0.13
Sperm Volume (ml) Median [IQR]	3.4 [3.5]	2.7 [2.3]	0.54
Total Sperm Count (million/ejaculate) Median [IQR]	15.2 [14.6]	22.7 [28.8]	0.20
Progressive Motility (%) <sup>2</sup>	28.7 [23.0]	33.4 [25.9]	0.35

Median [IQR]			
<b>Total Motility (%)</b> <sup>2</sup> Median [IQR]	34.5 [26.5]	42.5 [28.0]	0.32
<b>Total Motile Count (million/ejaculate)</b> <sup>1,2</sup> Median [IQR]	5.0 [6.3]	9.6 [15.9]	0.08
<b>Morphology (%)</b> <sup>2</sup> Median [IQR]	0.0 [0.8]	1.0 [1.4]	0.03*
<b>ROS (RLU)</b> <sup>2</sup> Median [IQR]	8.5 [171.5]	3.6 [35.8]	0.10
<b>REPRODUCTIVE HORMONE PROFILE</b>			
<b>LH (IU/L)</b>	3.7 ± 0.9	4.3 ± 1.5	0.18
<b>FSH (IU/L)</b>	5.6 ± 2.9	5 ± 2.5	0.51
<b>Testosterone (nmol/L)</b>	12.6 ± 5.1	14.6 ± 5.0	0.21
<b>SHBG (nmol/L)</b>	21 ± 10	25 ± 10	0.27

**Table 2: Analysis for covariance and comparisons of outcomes from both studies.** For outcomes analysed on the original scale, the mean difference in outcome between groups is reported with a corresponding confidence interval. For outcomes analysed on the log scale, the ratio of values in the LED group relative to the BDI group is reported with a corresponding confidence interval. P-values indicating the significance of the group differences are reported in the final column.

\* Group difference expressed as outcome in LED group minus outcome in BDI group. Adjusted for outcome at baseline.

\*\* Group difference expressed as outcome in LED group relative to outcome in BDI group. Adjusted for outcome at baseline.

# Interaction not relevant as outcome measured for Study B only.

Outcome	Study by Group interaction p-value	Patients	Group difference Mean (95% CI) *	Group P-value
Total motility	0.42	All	2.4 (-3.6, 8.)	0.43
Progressive motility	0.62	All	2.8 (-3.3, 8.9)	0.36
Semen volume	0.81	All	0.44 (-0.12, 0.99)	0.12
DFI	<b>0.004</b>	Study A Study B	-9.5 (-16.4, -2.5) 4.8 (-1.6, 11.2)	<b>0.009</b> 0.14
Outcome	Study by Group interaction p-value	Patients	Group difference Ratio (95% CI) **	Group P-value
Sperm concentration	0.53	All	0.77 (0.40, 1.48)	0.43
Sperm count	0.79	All	0.87 (0.43, 1.74)	0.69
Total motile count	0.66	All	1.00 (0.52, 1.93)	0.99
Morphology	0.89	All	0.81 (0.60, 1.10)	0.17
ROS	#	Study B	0.77 (0.18, 3.36)	0.72



Figure 1




Figure 1  
181x216 mm ( x DPI)

Figure 2




Figure 2  
195x75 mm (x DPI)

1  
2  
3  
4

Figure 3




Figure 3  
179x202 mm ( x DPI)

Figure 4




Figure 4  
149x235 mm (x DPI)

1  
2  
3