

Research Article

# Increasing prevalence of overweight and obesity among Tanzanian women of reproductive age intending to conceive: evidence from three Demographic Health Surveys, 2004-2016

Amani I Kikula<sup>1,2,3</sup>, Aline Semaan<sup>1</sup>, Belinda Balandya<sup>4</sup>, Naku K Makoko<sup>5</sup>, Andrea B Pembe<sup>4</sup>, José L Peñalvo<sup>2</sup>, Lenka Beňová<sup>1</sup>

<sup>1</sup> Department of Public Health, Instituut voor Tropische Geneeskunde, <sup>2</sup> Global Health Institute, Faculty of Medicine and Health Sciences, University of Antwerp, <sup>3</sup> Obstetrics and Gynaecology, Muhimbili University of Health and Allied Sciences, <sup>4</sup> Department of Obstetrics and Gynaecology, Muhimbili University of Health and Allied Sciences, <sup>5</sup> National Institute for Medical Research

Keywords: Obesity, overweight, Pre-pregnancy, Tanzania, BMI

<https://doi.org/10.29392/001c.87443>

## Journal of Global Health Reports

Vol. 7, 2023

### Background

The prevalence of people who are overweight or obese is increasing globally, especially in low- and middle-income countries. High body mass index (BMI) among women of reproductive age is a risk factor for various adverse reproductive and pregnancy outcomes. This study aims to describe trends over time in the distribution of BMI among Tanzanian women of reproductive age intending to conceive between 2004/5 and 2015/16, and identify factors associated with high BMI.

### Methods

We used data on 20,819 women of reproductive age (15-49 years) intending to conceive who participated in the Tanzania Demographic and Health Surveys in 2004/5, 2010 and 2015/16. We estimated the prevalence of high BMI (being overweight [ $\geq 25$  to  $< 30$  kg/m<sup>2</sup>] and obesity [ $\geq 30$  kg/m<sup>2</sup>]) and trends in the prevalence of high BMI across the three surveys. Using survey-weighted multivariable logistic regression, we used the most recent 2015/16 survey data to identify factors associated with high BMI.

### Results

Median BMI increased from 21.7 kg/m<sup>2</sup> (inter-quartile range, IQR=19.9-24.1 kg/m<sup>2</sup>) in 2004/5 to 22.0 kg/m<sup>2</sup> (IQR=20.0-24.8 kg/m<sup>2</sup>) in 2010 to 22.7 kg/m<sup>2</sup> (IQR=20.4-26.0 kg/m<sup>2</sup>) in 2015/16. The prevalence of overweight women increased from 11.1% in 2004/5 to 15.8% in 2015 (P < 0.001). The prevalence of obesity increased from 3.1% in 2004/5 to 8.0% in 2015/16 (P < 0.001). Women in the highest wealth quintile had higher odds (adjusted odds ratio, aOR= 4.5; 95%CI 3.4-6.3, P < 0.001) of high BMI than women in the lowest quintile. The odds of high BMI were about four times greater (aOR=3.9; 95%CI=2.9-5.4, P < 0.001) for women 40-44 years compared to 20-24-year-olds. Women in the high-paying occupations had greater odds of high BMI than those working in agriculture (aOR=1.5; 95% CI=1.1-2.2, P=0.002). Women residing in the Southern zone had 1.9 (95%CI=1.5-2.5, P < 0.001) greater odds of high BMI than Lake zone residents.

### Conclusions

In Tanzania, high BMI affects almost 1 in 4 women of reproductive age who intend to conceive. This contributes to the burden of poor maternal and reproductive health outcomes. We recommend developing and implementing health-system strategies for addressing high BMI, tailored to the modifiable risk factors identified among women of reproductive age.

Since the early 1990s, the burden of being overweight and obese to morbidity and mortality in low- and middle-income countries (LMICs) is increasing more than that contributed by underweight.<sup>1</sup> Globally, adult obesity doubled

between 1980 to 2014, and the burden is higher among women (40%) compared to men (38%).<sup>2</sup> Obesity is causally linked to all major non-communicable diseases (NCDs), including type 2 diabetes mellitus, hypertension, cardiovas-

cular disease and some types of cancer.<sup>3</sup> In 2022, the World Health Organization (WHO) estimated that over three-quarters of global deaths were attributed to NCDs and most were in LMICs.<sup>4</sup> While obesity is not explicitly mentioned in the Sustainable Development Goals (SDGs), there are opportunities to develop actions against obesity under 13 of the 17 SDGs,<sup>5</sup> particularly under target 3.4, to reduce premature mortality from NCDs by one-third through prevention, treatment, promotion of mental health and well-being.

The increasing prevalence of obesity among adults globally has specific consequences for women of reproductive age, especially those planning to be or are currently pregnant. Being overweight or obese in the pre-pregnancy period poses an increased risk for subfertility or infertility<sup>6</sup> and adverse maternal and perinatal outcomes. For example, among Danish, Chilean and Chinese women, higher pre-pregnancy body mass index (BMI) was associated with higher odds of gestational diabetes, preeclampsia and operative delivery. Among newborns, there was a higher risk for macrosomia, birth injuries, stillbirth and pre-term birth.<sup>7,8</sup> These adverse associations persist beyond the neonatal period and could affect children's neuro-developmental milestones<sup>9</sup> and increase their risk for adult onset of cardiovascular diseases<sup>10</sup> and type 2 diabetes mellitus.<sup>11</sup>

Structural determinants for being overweight and obese among women of reproductive age include socio-economic factors such as lower wealth and lack of health insurance coverage.<sup>12</sup> Irrespective of the country's development level, women of older age and higher parity have higher odds of being overweight and obese.<sup>13</sup> Among women in high-income countries, being married,<sup>14</sup> lower education levels, smoking and having a lower-income job are factors that increase the risk of high pre-pregnancy BMI.<sup>15</sup> In LMICs, being wealthier, living in urban areas, and achieving higher education levels were associated with a higher risk of being overweight or obese.<sup>16</sup>

Maternal mortality ratio (MMR) in Tanzania was estimated at 238/100,000 livebirths in 2020.<sup>17</sup> Even though this represents a decline from 524/100,000 livebirths in 2015, it is still high, suggesting that the SDG of reducing MMR to less than 140/100,000 livebirths might not be attained by 2030.<sup>18</sup> Recently, over half of global avertable adult mortality is due to NCD with majority being clustered within LMICs.<sup>19</sup> With the ongoing global obstetric transition the contribution of NCDs to maternal mortality is expected to rise also,<sup>20</sup> highlighting a need to re-strategize and accommodate measures needed to consider and manage NCDs within maternal healthcare services.<sup>21</sup>

Tanzania is experiencing an increasing prevalence of overweight and obesity in the general adult population, with a high prevalence in urban areas of 30-37% in 2021.<sup>22, 23</sup> Existing evidence of the ongoing nutrition transition in rural areas of Tanzania suggests a likelihood of an increasing prevalence of being overweight (16%) and obese (6%) among rural residents similar to urban areas.<sup>24</sup> Among adult women, the prevalence of overweight and obesity doubled from 11.2% in 1991 to 21.6% in 2016.<sup>25</sup> Considering the association of overweight and obesity among

women of reproductive age with NCDs and the resulting reproductive health-related morbidity and mortality, we intend to study the evolution over time and factors associated with women's BMI using a robust nationally representative database to inform stakeholders of the potential need to develop and implement context-specific interventions relevant to this issue.

This study aimed to describe the change over time in the distribution of BMI among women of reproductive age intending to conceive in Tanzania between 2004/5 and 2015/16 and identify factors associated with high BMI in 2015/16.

## METHODS

### STUDY DESIGN

This is a cross-sectional study using secondary data. We used three consecutive Demographic and Health Surveys (DHS) in Tanzania to characterize changes in BMI distribution over time and analyzed factors associated with high BMI among women using data from the most recent survey.

### SETTING

We used the last three Tanzanian DHS conducted in 2004/5, 2010 and 2015/16. These nationally representative household-based surveys were conducted using standardized protocols with country-specific adaptations. Briefly, the survey uses multistage cluster sampling, with the initial stage involving the selection of clusters and the second stage selection of households. The survey questionnaires captured dimensions of reproductive and child health characteristics, anthropometric and nutritional status variables, individual and household socio-demographic characteristics.<sup>26-28</sup>

### POPULATION

Because of the importance of pre-conception BMI to maternal and perinatal outcomes, we included women of reproductive age (15 to 49 years) who resided in sampled households and were interviewed by the DHS and who reported wanting to have children in the next two years or within an undecided future period. We excluded women who had undergone female sterilization, were declared infertile, or reported not wanting to conceive in the future. We also excluded women who reported being pregnant at the time of the survey due to the physiological weight gain that occurs during pregnancy.

### OUTCOMES

Height and weight were measured as per DHS protocols by trained personnel. BMI was calculated by dividing body weight (kg) by squared height (m<sup>2</sup>). We used BMI as a continuous variable and classified it into categories. For women aged 20 years and older, we classified BMI into four categories: underweight (BMI  $\leq$  18.5kg/m<sup>2</sup>), normal weight (>18.5 to <25 kg/m<sup>2</sup>), overweight ( $\geq$ 25 to <30kg/m<sup>2</sup>) and obese ( $\geq$ 30kg/m<sup>2</sup>).<sup>29</sup> In the age group 15-19 years, BMI was classified as underweight (z-score  $\leq$  -2), normal weight (z

score  $>-2$  to  $<1$ ), overweight (z score  $\geq 1$  to  $< 2$ ) and obese (z score  $\geq 2$ ) using age-specific z-scores based on WHO guidelines.<sup>30</sup> We also constructed a combined category of 'high' BMI by combining women who were overweight or obese.

## COVARIATES

Variables theoretically associated with high BMI and available on the DHS included residence (rural or urban); zone of residence (Central, Lake, Western, Northern, Southern highland, South-west highlands, Southern, Eastern, and Zanzibar); household wealth classified by DHS using a quintile distribution into poorest, poorer, middle, more affluent and richest; health insurance coverage (yes or no); parity (0, 1-4 and  $\geq 5$ ); women's highest reached education level (no formal education, at least primary education, at least secondary education or university/college education); occupation at the time of survey (not working, unskilled manual, household and domestic, services, agricultural, skilled manual, professional /technical /managerial/clerical); 5-year age group at the time of the survey, and marital status at the time of the survey (married or living together or single/widowed/separated/divorced). The 2004/5 survey had no variable for health insurance coverage; four women had missing values for insurance coverage and 12 for occupation on the 2010 survey. The 2015/16 survey data contained no missing values in variables used in regression analysis.

## DATA ANALYSIS

We analyzed the data using Stata SE version 14.1 (Stata-Corp, College Station, TX, USA). The normal distribution of BMI as a continuous variable was assessed using the Shapiro-Wilk test, which indicated a significant departure from normality ( $P<0.001$ ). The distribution across the three surveys were described using median and Interquartile Range (IQR) by survey year/s. We used counts and percentages for categorical variables to describe the data across surveys. Differences of distribution (proportion) of the outcome (BMI) across the surveys were tested using Kruskal-wallis rank test. Using crude and multivariable logistic models, we used the most recent survey (2015/16) to estimate the association between the selected covariates and outcome (high BMI). We analyzed factors associated with high BMI as a binary variable (normal BMI and high BMI), excluding the underweight category since effect modifiers for underweight to normal weight differ from those of normal weight to overweight or obesity. We then considered BMI as an ordinal variable (normal weight, overweight, or obese). We conducted an ordinal multivariable logistic regression analysis to test whether the measure of the effect of the independent variable had different results when overweight and obesity were considered separately. The results are shown separately in the **Online Supplementary Document (Table S1)** with adjusted odds values similar to the binary multivariable logistic model. Normal weight was considered a reference category for multivariable logistic regression analysis.

We expressed the measure of effect in terms of crude and adjusted odds ratios (OR) with their 95% confidence interval (CI). We used the svyset command in STATA to adjust for the survey's individual sampling weights, clustering and stratification in all analyses. All tests with  $P<0.05$  were considered statistically significant.

## ETHICS APPROVAL

The DHS received government permission, used informed consent and assured respondents of confidentiality. We conducted secondary data analysis of the available anonymized data with permission from the DHS.

## RESULTS

### CHARACTERISTICS OF THE STUDY POPULATION

We included information on 6,731 women from the 2004/5 survey, 5,011 from 2010, and 9,077 from 2015/16. From each survey, women with flagged (missing and improbable) BMI values were excluded from the analysis ( $n=76$  in 2004/5,  $n=53$  in 2010 and  $n=87$  in 2015). A flowchart showing the distribution of women included in the analysis is in **Online Supplementary Document, Figure S1**. **Table 1** shows the women's characteristics across the three surveys. In all three surveys, about two-thirds of women resided in rural areas, while about a quarter of the women were from the Lake zone. Women in the lower or lowest wealth quintile decreased from 2004/5 to 2015/16 survey, while those with health insurance increased from 5.8% (2010) to 8.1% (2015). Lastly, the percentage of women with at least secondary education increased from 10.4% in 2004/5 to 28.8% in 2015.

### BMI CHANGES OVER TIME

Median BMI increased from 21.7 kg/m<sup>2</sup> (IQR=19.9 – 24.1 kg/m<sup>2</sup>) in 2004/5 to 22.0 kg/m<sup>2</sup> (IQR=20.0 – 24.8 kg/m<sup>2</sup>) in 2010 to 22.7 kg/m<sup>2</sup> (IQR=20.4 – 26.0 kg/m<sup>2</sup>) in 2015/16. The prevalence of normal BMI decreased from 80.2% (2004/5) to 71.6% (2010) and 71.3% (2015/16). The prevalence of overweight increased from 11.1% in 2004/5 to 14.5% in 2010 ( $P<0.001$ ) and 15.8% in 2015/16 ( $P<0.001$ ). The prevalence of obesity increased from 3.1% in 2004/5 to 8.0% in 2015/16 ( $P<0.001$ ) (**Figure 1**). The prevalence of high BMI (overweight and obesity combined) increased from 14.2% to 23.8% ( $P<0.001$ ) in the same period.

### FACTORS ASSOCIATED WITH HIGH BMI FROM THE 2015/6 SURVEY.

In univariate analysis, compared to nulliparous women, women with 2-4 children had almost three times greater odds of high BMI ( $P<0.001$ ) and those with health insurance coverage had 1.7 times higher odds of high BMI than those without. Women residing in urban areas had 2.7 greater odds ( $P<0.001$ ) of high BMI compared to those in rural areas, and those married or living with a partner had 1.8 greater odds of high BMI than those who were single or living alone ( $P<0.001$ ) (**Table 2**).

**Table 1. Characteristics of samples of women of reproductive age who intend to conceive by survey year**

Variable	2004/5 n= 6,731		2010 n= 5,011		2015/16 n= 9,077	
	Median BMI - 21.7 kg/m <sup>2</sup> (IQR 19.9 - 24.1 kg/m <sup>2</sup> )		Median BMI - 22.0 kg/m <sup>2</sup> (IQR 20.0 - 24.8 kg/m <sup>2</sup> )		Median BMI - 22.7 kg/m <sup>2</sup> (IQR 20.4 - 26.0 kg/m <sup>2</sup> )	
	n	% (95% CI)	n	% (95% CI)	n	% (95% CI)
<b>Age group</b>						
15 - 19	2,002	29.3 (27.9-30.8)	664	15.1 (13.8-16.6)	2,684	29.6 (28.4-30.9)
20 - 24	1,542	24.2 (23.0-25.4)	1,218	26.4 (24.9-27.9)	2,099	23.7 (22.7-24.7)
25 - 29	1,294	19.9 (18.7-21.2)	1,152	23.4 (22.0-24.8)	1,638	18.2 (17.33-19.2)
30 - 34	934	14.2 (13.2-15.2)	881	16.5 (15.2-17.8)	1,165	13 (12.2-13.9)
35 - 39	535	7.3 (6.5-8.1)	619	11.6 (10.5-12.8)	834	8.9 (8.2-9.6)
40 - 44	288	3.4 (2.9-3.9)	326	5.2 (4.5-6.1)	475	4.7 (4.2-5.3)
45 - 49	136	1.8 (1.4-2.2)	151	1.9 (1.5-2.3)	182	1.8 (1.5-2.3)
<b>Occupation status<sup>b</sup></b>						
Not working	1,772	21.8 (19.5-24.3)	890	15.4 (13.7-17.3)	2,656	27.8 (26.4-29.3)
Unskilled manual	571	8.1 (6.8-9.6)	724	14.3 (12.2-16.7)	1,389	15.6 (14.3-17.0)
Household and domestic Services	206	3.9 (2.9-5.2)	133	4.2 (2.2-3.7)	454	6.1 (5.4-6.9)
Agricultural	61	1.1 (0.6-1.8)	157	5.3 (2.6-3.9)	330	4.4 (3.9-5.1)
Skilled manual	3,711	60.7 (56.1-65.1)	2,676	54.8 (52.5-60.7)	3,482	38.6 (36.5-40.8)
Professional/technical /managerial/clerical	240	2.6 (2.0-3.3)	280	13.3 (4.0-5.8)	416	3.5 (3.0-4.1)
Professional/technical /managerial/clerical	170	1.9 (1.4-2.5)	146	2.6 (1.9-3.5)	350	3.9 (3.3-4.5)
<b>Zone of residence</b>						
Western	541	9.3 (6.7-13.1)	455	8.9 (7.0-12.9)	715	9.8 (8.6-11.0)
Northern	685	11.4 (8.6-14.9)	425	12.9 (8.9-13.3)	829	11.9 (10.4-13.6)
Central	756	10.8 (8.1-14.3)	505	10.2 (8.1-11.4)	767	9.6 (8.6-10.7)
Southern highlands	462	7.3 (5.1-10.2)	329	8.3 (5.5-10.3)	651	5.6 (5.0-6.3)
Southern	446	5.8 (4.1-8.2)	382	5.9 (5.4-8.9)	551	5.9 (5.2-6.7)
Southwest highlands	496	9.7 (6.2-14.8)	324	8.7 (5.8-11.5)	774	8.3 (7.3-9.4)
Lake	963	23.9 (18.9-29.8)	873	25.9 (21.6-29.5)	2,006	25.2 (23.7-26.8)
Eastern	769	18.3 (13.9-23.6)	670	15.9 (16.8-21.8)	1,080	20 (18.1-22.1)
Zanzibar	1,613	3.5 (2.7-4.6)	1,048	3.2 (2.4-3.2)	1,704	3.7 (3.4-3.9)
<b>Residence</b>						
Urban	1,688	29.4 (24.2-35.2)	1,296	30.2 (25.1-35.7)	2,977	38.4 (36.3-40.5)
Rural	5,043	70.6 (64.8-75.8)	3,715	69.8 (64.3-74.9)	6,100	61.6 (59.5-63.7)
<b>Household wealth quintile</b>						
Poorest	1,099	17.3	844	16.9	1,347	15.9

		(14.8-19.9)		(14.8-19.2)		(14.1-18.1)	
Poorer	1,243	19.4 (17.2-21.8)	953	19.1 (16.9-21.5)	1,406	16 (14.7-17.5)	
Middle	1,151	17.6 (15.6-19.9)	932	18.9 (17.0-20.9)	1,574	16.7 (15.3-18.2)	
Richer	1,498	18.3 (16.3-20.5)	1,096	20.9 (18.6-23.4)	2,162	21.3 (19.6-23.2)	
Richest	1,740	27.4 (22.8-32.6)	1,186	24.2 (20.5-28.4)	2,588	29.9 (27.6-32.5)	
<b>Health insurance coverage <sup>a</sup></b>							
Not covered	N/A	N/A	N/A	4,782	94.1 (92.9-95.1)	8,414	91.9 (90.8-92.8)
Covered	N/A	N/A	N/A	229	5.9 (4.9-7.0)	663	8.1 (7.2-9.2)
<b>Parity</b>							
0	2,425	33.5 (31.6-35.5)	787	17.3 (15.7-18.9)	3,230	34.3 (32.6-35.9)	
1	1,063	18.1 (16.9-19.5)	1,007	21.9 (20.4-23.4)	1,674	20.5 (19.4-21.6)	
2-4	2,266	35.8 (34.3-37.4)	2,287	46.1 (44.4-47.8)	3,006	34.2 (32.8-35.6)	
≥5	977	12.5 (11.2-14.1)	930	14.8 (13.4-16.3)	1,167	11.1 (10.2-12.1)	
<b>Marital status</b>							
Not married/ Not living together	2,755	38.5 (36.1-40.8)	1,274	28.1 (26.2-30.1)	4,139	44.9 (43.2-46.7)	
Married/ living with partner	3,976	61.6 (59.2-63.9)	3,737	71.9 (69.9-73.8)	4,938	55.1 (53.3-56.8)	
<b>Highest education level</b>							
No formal education	1,474	22.2 (19.1-25.6)	1,059	20.5 (18.2-22.9)	1,191	12.9 (11.7-14.4)	
Primary education	4,144	67.5 (64.7-70.2)	3,013	64.9 (62.7-67.1)	4,861	58.2 (56.7-59.7)	
Secondary or higher education	1,113	10.4 (8.7-12.3)	939	14.7 (12.8-16.7)	3,025	28.8 (27.1-30.6)	

N/A – Not applicable – in 2004/5, data for health insurance coverage was not collected, <sup>b</sup> n=4 missing from the 2010 data set, <sup>a</sup> n=12 missing from the 2010 dataset

After controlling for other factors in a multivariable model, parity, marital status, health insurance coverage, highest level of education, and residence were no longer significantly associated with high BMI. Older age group was associated with greater odds of high BMI. Women in the 40–44 age category had 3.9 greater odds (95% CI=2.9 – 5.4,  $P<0.001$ ) of high BMI compared to women 20–24 years. Women from the wealthiest households had nearly 5 times greater odds of high BMI than those in the lower wealth quintile (95% CI=3.4 - 6.3,  $P<0.001$ ). Women from the Southern and Southwestern zones of the country had higher odds (aOR=1.9, 95% CI=1.5 - 2.5,  $P<0.001$ ) of high BMI than women from the Lake zone. Women who did -skilled working-class jobs or those dedicated to household/domestic work had greater odds of high BMI than women working in agriculture.

## DISCUSSION

In this study, we analyzed a nationally representative data to show the changes in Tanzania for the prevalence of high

BMI among women intending to conceive over time. We found that between 2004/5 and 2015/16, the prevalence of high BMI increased from 14.2% to 23.8% due to an increase in the prevalence of both overweight and obesity BMI categories. In addition, the prevalence of underweight decreased over the time period under consideration. Women who were older, residing in the Southern zones, from wealthier households, to a higher skilled working class had greater odds of high BMI in 2015/16 than their counterparts.

The increasing prevalence of high BMI among women of reproductive age reflects the pattern seen among everyone in the general population. Globally, there is an increase in the prevalence of high BMI.<sup>31</sup> In Africa, the increase in the prevalence of high BMI has been related to other public health morbidities.<sup>32</sup> Specifically, Uganda has also witnessed a similar prevalence of high BMI among women of reproductive age.<sup>33</sup> In Dar es Salaam, Tanzania, a study showed that more than half of the women of reproductive age were overweight or obese.<sup>34</sup> Related studies conducted among people with similar demographic characteristics are limited. However, reviews in Europe and Africa on maternal

**Table 2. Crude and adjusted association using logistic regression analysis between socio-demographic factors and high BMI among sub-sample of non-underweight women, 2015/16 DHS (n=8,506)**

Variable	Study sample		Crude analysis			Multivariable analysis <sup>a</sup>		
	n <sup>b</sup>	% (95% CI)	cOR	95% CI	P	aOR	95% CI	P
<b>Age</b>								
15 - 19	2601	31.1 (29.9-32.4)	0.2	0.2-0.3	<0.001	0.2	0.2-0.3	0.001
20 - 24	1920	22.9 (21.9-24.0)	Ref			Ref		
25 - 29	1515	18.1 (17.2-19.1)	1.8	1.5-2.1	<0.001	1.8	1.5-2.3	0.001
30 - 34	1056	12.6 (11.8-13.5)	2.5	2.0-3.1	<0.001	2.8	2.2-3.6	0.001
35 - 39	731	8.7 (8.1-9.5)	2.6	2.0-3.2	<0.001	2.9	2.2-3.8	0.001
40 - 44	387	4.6 (4.1-5.2)	3.2	2.5-4.1	<0.001	3.9	2.9-5.4	0.001
45 - 49	154	1.8 (1.5-2.3)	2.2	1.5-3.3	<0.001	3.1	1.9-4.9	0.001
<b>Occupation status</b>								
Not working	2353	28.1 (26.7-29.6)	1.1	0.9-1.4	0.158	1.2	0.9-1.5	0.079
Unskilled manual	1314	15.7 (14.4-17.1)	3.3	2.7-3.9	<0.001	1.5	1.2-1.8	<0.001
Household/domestic	522	6.2 (5.4-7.1)	2.9	2.3-3.8	<0.001	1.7	1.2-2.3	0.001
Services	374	4.5 (3.9-5.1)	2.9	2.2-3.9	<0.001	1.2	0.9-1.7	0.231
Agricultural	3199	38.2 (36.1-40.4)	Ref			Ref		
Skilled manual	287	3.4 (2.9-3.9)	2.6	1.9-3.7	<0.001	1.3	0.8-1.9	0.275
Professional/technical/ managerial/clerical	317	3.8 (3.2-4.4)	5.7	4.2-7.7	<0.001	1.5	1.1-2.2	0.017
<b>Zone of residence</b>								
Western	818	9.8 (8.6-11.1)	1.1	0.9-1.5	0.43	1.3	0.9-1.7	0.053
Northern	984	11.8 (10.2-13.5)	2.8	2.1-3.6	<0.01	1.8	1.4-2.3	<0.001
Central	772	9.2 (8.2-10.4)	1.3	1.0-1.7	0.039	1.3	0.9-1.7	0.111
Southern Highlands	479	5.7 (5.1-6.4)	1.7	1.3-2.3	0.001	1.2	0.9-1.6	0.219
Southern	502	5.9 (5.3-6.8)	2.3	1.8-2.9	<0.001	1.9	1.5-2.5	<0.001
South west Highlands	712	8.5 (7.4-9.7)	2.1	1.5-2.9	<0.001	1.9	1.4-2.6	<0.001
Lake	2112	25.2 (23.6-26.9)	Ref			Ref		
Eastern	1688	20.2 (18.2-22.4)	3.4	2.8-4.2	<0.001	1.6	1.3-2.0	<0.001
Zanzibar	297	3.6 (3.3-3.9)	3.3	2.7-4.1	<0.001	1.8	1.4-2.2	<0.001
<b>Residence</b>								
Urban	3240	38.7 (36.6-40.9)	2.7	2.4-3.1	<0.001	1.1	0.9-1.4	0.159

Rural	5125	61.3 (59.1-63.4)	Ref			Ref			
<b>Wealth quintile</b>									
Poorest	1302	15.6 (13.7-17.6)	Ref			Ref			
Poorer	1326	15.9 (14.5-17.3)	1.7	1.3-2.2	<0.001	1.5	1.2-2.0	0.002	
Middle	1406	16.8 (15.4-18.3)	2.2	1.7-2.8	<0.001	1.9	1.4-2.5	0.001	
Richer	1792	21.4 (19.7-23.3)	3.6	2.7-4.5	<0.001	2.8	2.1-3.6	0.001	
Richest	2540	30.4 (27.9-32.9)	6.7	5.3-8.3	<0.001	4.6	3.4-6.3	0.001	
<b>Health insurance coverage</b>									
Not covered	7669	91.7 (90.6-92.7)	Ref			Ref			
Covered	696	8.3 (7.3-9.4)	1.7	1.4-2.1	<0.001	1.3	0.9-1.6	0.067	
<b>Parity</b>									
0	2924	34.9 (33.3-36.7)	Ref			Ref			
1	1714	20.5 (19.3-21.7)	2.1	1.7-2.5	<0.001	1.1	0.9-1.4	0.498	
2-4	2826	33.8 (32.4-35.2)	2.9	2.4-3.4	<0.001	1.0	0.8-1.3	0.859	
≥5	902	10.8 (9.9-11.7)	2.1	1.7-2.7	<0.001	0.9	0.6-1.2	0.422	
<b>Marital status</b>									
Not married/ Not living together	3817	45.6 (43.9-47.4)	Ref			Ref			
Married/ living with partner	4548	51.4 (52.6-56.2)	1.8	1.6-2.1	<0.001	1.2	0.9-1.4	0.137	
<b>Highest education level</b>									
No formal education	1039	12.4 (11.2-13.8)	Ref			Ref			
Primary education	4900	58.6 (57.0-60.1)	1.3	1.1-1.6	0.012	1.2	0.9-1.4	0.216	
Secondary or higher education	2425	29 (27.3-30.8)	1.6	1.3-2.0	<0.001	1.3	0.9-1.7	0.076	


cOR=crude Odds Ratio; aOR=adjusted Odds Ratio; <sup>a</sup> Multivariable logistic regression analysis, <sup>b</sup> Weighted counts

obesity (closely associated with pre-pregnancy BMI status) are on the rise.<sup>35,36</sup>

While drivers of high BMI globally vary, the global trend towards increased access to high-caloric foods, often ultra-processed with suboptimal nutritional profiles and inexpensive, which has led to the obesity epidemic in high-income countries<sup>37,38</sup> is also observed in LMICs through increased commercialization and penetrative food marketing.<sup>39</sup> What is common in both settings is the co-existence of calorie-dense diets with low energy expending activities as a result of industrialization with less manual labor or subsistence farming and advanced technology that promotes sedentary lifestyles.<sup>40,41</sup> Additionally, national-level policy drivers such as lower prices for high-calorie diets even in rural areas and their commercialization contribute to this increasing burden of overweight and obesity.<sup>42,43</sup>

Cultural perceptions can contribute to this growing burden; for example, the difference in an ideal body size in Japan<sup>43</sup> differs from the perceived African ideal body size.<sup>44,45</sup>

The increasing prevalence of high BMI in this study also highlights the co-existence of multiple forms of malnutrition among women of reproductive age in Tanzania. In addition to the rising prevalence of high BMI, women in the country also bear a high prevalence of anemia, reported to be at 57%<sup>46</sup> with an accompanying prevalence of folate deficiency at 26%<sup>47</sup> and of iron deficiency at 33%.<sup>26</sup> Maternal anemia is a risk factor for maternal mortality (the risk of maternal mortality doubles with maternal haemoglobin less than 7gm/dl).<sup>48</sup> While as the pre-pregnancy BMI increases from overweight to obesity, the odds of maternal death increase from just over two times to over three times those with normal BMI.<sup>49</sup> The compounding effect of ane-



**Figure 1. Changes of prevalence of BMI categories among women of reproductive age intending to conceive in Tanzania.**

mia and high BMI could be even more catastrophic to the well-being of women in Tanzania. This needs immediate action from the health system to protect the health of women and contribute to lowering the risk of maternal and perinatal mortality and morbidity in Tanzania.

Our results also show that older women living in relatively wealthier households and in a professional working class had greater odds of a high BMI. This can be hypothesized to be due to policies that enhance access to more high-energy foods and less energy-consuming activities, similar to what has been observed for individuals of such socio-economic positions in African countries, including Tanzania.<sup>16,33,50,51</sup> Women from high agricultural food-producing zones (the Southern and Southwest zones) had greater odds of having high BMI than those in the Lake zone. This could be due to these being high -producing zones in the country and having access to diverse food, which has been linked with high BMI in Tanzania.<sup>50</sup>

Having a high BMI before conception is linked to worse outcomes during and after pregnancy, including but not limited to gestational diabetes, or preeclampsia and an increased risk of type 2 diabetes mellitus.<sup>3</sup> On the healthcare service delivery, there have been improvements in access to quality health services in Tanzania. However, health facilities are still not well equipped to screen or manage NCD-related conditions relative to the capacity for managing communicable diseases.<sup>52</sup> This calls for immediate action to address this rising burden of nutritional morbidity (high BMI) as a factor contributing to maternal mortality and morbidity in Tanzania. Incorporating weight management interventions within antenatal care and during the post-natal period have been suggested to address and prevent

this burden and should be tailored to specific country contexts.<sup>53</sup>

Strengths of this study include the use of nationally representative, comparable data of high quality. We classified BMI based on the standard WHO classification of BMI. We also leveraged the information on the DHS employing multivariable models that partially account for the effects of other variables on increased BMI, although residual confounding cannot be ruled out. We were also somehow limited by the lack of data on women's intention to conceive in the near future. We acknowledge that women who intend to get pregnant might not be able to, and vice versa, women who did not intend to get pregnant (and were thus excluded from our study sample) might do so. Lastly, from 2016 to now, a lot may have changed, but the 2015/16 survey was the latest available TDHS data set. Still, these results are valid and help to highlight the need to accommodate high BMI-related reproductive health needs in the Tanzanian health system.

## CONCLUSIONS

The prevalence of high BMI among women of reproductive age who intend to conceive is increasing in Tanzania, in both rural and urban areas. In a country where communicable diseases and undernutrition have historically challenged maternal health, innovative strategies to improve access to healthy lifestyles will need to be implemented. We recommend developing context-specific and culturally sensitive policies and interventions for addressing BMI-related challenges in women of reproductive age in Tanzania. These can be incorporated into available systems – to improve screening and linkage to care on high BMI-re-



lated morbidities, promoting physical activity, and having a safe exercising environment for the community, particularly targeting women of reproductive age.

.....

#### DISCLAIMER

The views expressed in this manuscript are the author's opinion and do not represent the official opinion of the institutions the authors are affiliated.

#### ETHICS STATEMENT

The DHS received government permission, used informed consent and assured respondents of confidentiality. We conducted secondary data analysis of the available anonymized data with permission from the DHS.

#### DATA AVAILABILITY

Data were obtained from the DHS and are available from [https://www.dhsprogram.com/data/dataset\\_admin](https://www.dhsprogram.com/data/dataset_admin) with the permission of the DHS.

#### FUNDING

The APC was funded by Directorate-general Development Cooperation and Humanitarian Aid Belgium – Sandwich PhD; Sida Muhimbili University of Health and Allied Sciences, Reproductive and Child health sub-program –

funded the training course in the development of the manuscript.

#### AUTHORSHIP CONTRIBUTIONS

Conceptualization, A.K.; methodology, A.K., N.M., B.B., A.B.P., A.S, L.B, and J.L.P; formal analysis, A.K., L.B., A.S and J.L.P.; writing—original draft preparation, A.K. and N.M.; writing—review and editing final draft, A.K., L.B., A.S., J.L.P., B.B., N.M., and A.B.P. All authors have read and agreed to the published version of the manuscript.

#### DISCLOSURE OF INTEREST

The authors completed the Unified Competing Interest form at <http://www.icmje.org/disclosure-of-interest/> (available upon request from the corresponding author) and declare no conflicts of interest.

#### ADDITIONAL MATERIAL

This manuscript contains additional information in the **Online Supplementary Document**.

#### CORRESPONDENCE TO:

Amani Kikula, MD, MMed (Obstetrics and Gynaecology), PhD candidate; P.O. Box 60371 Dar es salaam, Ilala – 11103; Email address – [amanikikula@live.com](mailto:amanikikula@live.com).

Submitted: July 26, 2023 BST, Accepted: August 15, 2023 BST



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

## REFERENCES

1. James WPT. WHO recognition of the global obesity epidemic. *Int J Obes*. 2008;32(S7):S120-S126. doi:10.1038/ijo.2008.247
2. WHO. *Global Status Report on Noncommunicable Diseases 2014*.; 2014.
3. Aras M, Tchang BG, Pape J. Obesity and Diabetes. *Nurs Clin North Am*. 2021;56(4):527-541. doi:10.1016/j.cnur.2021.07.008
4. WHO. *Invisible Numbers: The True Extent of Noncommunicable Diseases and What to Do about Them*.; 2022. <https://www.who.int/publications/i/item/9789240057661>
5. Cooper K. *Obesity and the SDGs: An Opportunity Hidden in Plain Sight Obesity within the SDGs?*
6. Gautam D, Purandare N, Maxwell CV, et al. The challenges of obesity for fertility: A FIGO literature review. *Intl J Gynecology & Obste*. 2023;160(S1):50-55. doi:10.1002/ijgo.14538
7. Li MF, Ke JF, Ma L, et al. Maternal Pre-Pregnancy Obesity Combined With Abnormal Glucose Metabolism Further Increases Adverse Pregnancy Outcomes in Chinese Pregnant Women. *Front Endocrinol*. 2022;12. doi:10.3389/fendo.2021.754406
8. Zong X, Wang H, Yang L, et al. Maternal Pre-pregnancy Body Mass Index Categories and Infant Birth Outcomes: A Population-Based Study of 9 Million Mother-Infant Pairs. *Front Nutr*. 2022;9(789833). doi:10.3389/FNUT.2022.789833/FULL
9. Sanchez CE, Barry C, Sabhlok A, et al. Maternal pre-pregnancy obesity and child neurodevelopmental outcomes: A Meta-analysis. *Obes Rev*. 2017;19(4):464-484. doi:10.1111/obr.12643
10. Kereliuk SM, Dolinsky VW. Recent Experimental Studies of Maternal Obesity, Diabetes during Pregnancy and the Developmental Origins of Cardiovascular Disease. *Int J Mol Sci*. 2022;23(8):4467. doi:10.3390/ijms23084467
11. Pirkola J, Pouta A, Bloigu A, et al. Prepregnancy overweight and gestational diabetes as determinants of subsequent diabetes and hypertension after 20-year follow-up. *J Clin Endocrinol Metab*. 2010;95(2):772-778. doi:10.1210/jc.2009-1075
12. Siega-Riz AM. Prepregnancy obesity: Determinants, consequences, and solutions. *Advances in Nutrition*. 2012;3(1):105-107. doi:10.3945/an.111.001081
13. Boudet-Berquier J, Salanave B, Desenclos JC, Castetbon K. Sociodemographic factors and pregnancy outcomes associated with prepregnancy obesity: Effect modification of parity in the nationwide Epifane birth-cohort. *BMC Pregnancy Childbirth*. 2017;17(1). doi:10.1186/s12884-017-1456-
14. Tzotzas T, Vlahavas G, Papadopoulou SK, Kapantais E, Kaklamanou D, Hassapidou M. Marital status and educational level associated to obesity in Greek adults: Data from the National Epidemiological Survey. *BMC Public Health*. 2010;10(1):1-8. doi:10.1186/1471-2458-10-732/TABLES/3
15. Cheng TS, Ali N, Elbarazi I, et al. Sociodemographic determinants of prepregnancy body mass index and gestational weight gain: The Mutaba'ah study. *Obes Sci Pract*. 2021;8(3):308-319. doi:10.1002/osp4.573
16. Nglazi MD, Ataguba JEO. Overweight and obesity in non-pregnant women of childbearing age in South Africa: subgroup regression analyses of survey data from 1998 to 2017. *BMC Public Health*. 2022;22(1):1-18. doi:10.1186/S12889-022-12601-6/TABLES/4
17. WHO, UNFPA, WORLDBANK. *Trends in Maternal Mortality 2000 to 2020: Estimates*.; 2023. <https://www.who.int/reproductivehealth/publications/maternal-mortality-2000-2017/en/>
18. WHO. SDG Target 3.1 | Maternal mortality: By 2030, reduce the global maternal mortality ratio to less than 70 per 100 000 live births. Published online 2022. Accessed June 6, 2023. <https://www.who.int/data/gho/data/themes/topics/sdg-target-3-1-maternal-mortality>
19. Martinez R, Lloyd-Sherlock P, Soliz P, et al. Trends in premature avertable mortality from non-communicable diseases for 195 countries and territories, 1990–2017: a population-based study. *Lancet Glob Health*. 2020;8(4):e511-e523. doi:10.1016/s2214-109x(20)30035-8
20. Souza JP, Tunçalp Ö, Vogel JP, et al. Obstetric transition: the pathway towards ending preventable maternal deaths. *BJOG*. 2014;121:1-4. doi:10.1111/1471-0528.12735

21. Sarah B, Ramnarayanan D, Amin N. The Unseen Side of Pregnancy: Non-Communicable Diseases and maternal health. In: *Utopian Spaces of Modernism*. Palgrave Macmillan; 2020. doi:10.1057/9780230358300.0018
22. Pallangyo P, Mkojera ZS, Hemed NR, et al. Obesity epidemic in urban Tanzania: a public health calamity in an already overwhelmed and fragmented health system. *BMC Endocr Disord*. 2020;20(1). doi:10.1186/s12902-020-00631-3
23. Zubery D, Kimiywe J, Martin HD. Prevalence of Overweight and Obesity, and Its Associated Factors Among Health-care Workers, Teachers, and Bankers in Arusha City, Tanzania. *Diabetes Metab Syndr Obes*. 2021;14:455-465. doi:10.2147/dmso.s283595
24. Keding GB, Msuya JM, Maass BL, Krawinkel MB. Dietary patterns and nutritional health of women: the nutrition transition in rural Tanzania. *Food Nutr Bull*. 2011;32(3):218-226. doi:10.1177/156482651103200306
25. Tanzania | World Obesity Federation Global Obesity Observatory. [https://data.worldobesity.org/country/tanzania-213/?msclkid=b8bdd723ce8611ec889fee3c8988e857#data\\_trends](https://data.worldobesity.org/country/tanzania-213/?msclkid=b8bdd723ce8611ec889fee3c8988e857#data_trends)
26. TDHS. *Tanzania 2015-16 Demographic Health Survey and Malaria Indicator Survey*. Tanzania 2015-16 Demographic and Health Survey and Malaria Indicator Survey.; 2016:24.
27. NBS/Tanzania NB of S, Macro I. anzia Demographic and Health Survey 2010. Published 2011. <https://dhsprogram.com/publications/publication-FR243-DHS-Final-Reports.cfm>
28. NBS/Tanzania NB of S. Tanzania Demographic and Health Survey. *Published online*. Published online 2004. <https://www.measuredhs.com>
29. Weir CB, Jan A. BMI Classification Percentile And Cut Off Points. *StatPearls*. Published online June 27, 2022.
30. who. BMI-for-age (5-19 years). Published online 2022. <https://www.who.int/tools/growth-reference-data-for-5to19-years/indicators/bmi-for-age>
31. Ng M, Fleming T, Robinson M, et al. Global, regional and national prevalence of overweight and obesity in children and adults 1980-2013: A systematic analysis. *Lancet*. 2014;384(9945):766-781. doi:10.1016/s0140-6736(14)60460-8
32. WHO. Obesity | WHO | Regional Office for Africa. Published 2023. <https://www.afro.who.int/health-topics/obesity>
33. Yaya S, Ghose B. Trend in overweight and obesity among women of reproductive age in Uganda: 1995–2016. *Obes Sci Pract*. 2019;5(4):312-323. doi:10.1002/osp4.351
34. Mosha D, Paulo HA, Mwanyika-Sando M, et al. Risk factors for overweight and obesity among women of reproductive age in Dar es Salaam, Tanzania. *BMC Nutr*. 2021;7(1):1-10. doi:10.1186/S40795-021-00445-Z/TABLES/4
35. Onubi OJ, Marais D, Aucott L, Okonofua F, Poobalan AS. Maternal obesity in Africa: a systematic review and meta-analysis. *J Public Health*. 2015;38(3):e218-e231. doi:10.1093/pubmed/fdv138
36. Devlieger R, Benhalima K, Damm P, et al. Maternal obesity in Europe: where do we stand and how to move forward?: A scientific paper commissioned by the European Board and College of Obstetrics and Gynaecology (EBCOG). *Eur J Obstet Gynecol Reprod Biol*. 2016;201:203-208. doi:10.1016/j.ejogrb.2016.04.005
37. Swinburn B, Sacks G, Ravussin E. Increased food energy supply is more than sufficient to explain the US epidemic of obesity. *American Journal of Clinical Nutrition*. 2009;90(6):1453-1456. doi:10.3945/ajcn.2009.28595
38. Scarborough P, Burg MR, Foster C, et al. Increased energy intake entirely accounts for increase in body weight in women but not in men in the UK between 1986 and 2000. *Br J Nutr*. 2011;105(9):1399-1404. doi:10.1017/s0007114510005076
39. WHO. Obesity rising in Africa, WHO analysis finds | WHO | Regional Office for Africa. Published 2022. <https://www.afro.who.int/news/obesity-rising-africa-who-analysis-finds>
40. Fox A, Feng W, Asal V. What is driving global obesity trends? Globalization or “modernization”? *Global Health*. 2019;15(1):1-16. doi:10.1186/S12992-019-0457-Y/FIGURES/4
41. Swinburn BA, Sacks G, Hall KD, et al. The global obesity pandemic: shaped by global drivers and local environments. *Lancet*. 2011;378(9793):804-814. doi:10.1016/s0140-6736(11)60813-1
42. Sacks G, Veerman JL, Moodie M, Swinburn B. ‘Traffic-light’ nutrition labelling and ‘junk-food’ tax: a modelled comparison of cost-effectiveness for obesity prevention. *Int J Obes*. 2011;35(7):1001-1009. doi:10.1038/ijo.2010.228
43. Mytton O, Gray A, Rayner M, Rutter H. Could targeted food taxes improve health? *J Epidemiol Community Health*. 2007;61(8):689-694. doi:10.1136/jech.2006.047746

44. Naigaga DA, Jahanlu D, Claudius HM, Gjerlaug AK, Barikmo I, Henjum S. Body size perceptions and preferences favor overweight in adult Saharawi refugees. *Nutr J*. 2018;17(1):1-8. [doi:10.1186/S12937-018-0330-5/FIGURES/4](https://doi.org/10.1186/S12937-018-0330-5/FIGURES/4)
45. Chigbu CO, Aniebue UU, Berger U, Parhofer KG. Impact of perceptions of body size on obesity and weight management behaviour: a large representative population study in an African setting. *Journal of Public Health*. 2019;43(1):e54-e61. [doi:10.1093/pubmed/fdz127](https://doi.org/10.1093/pubmed/fdz127)
46. Sunguya BF, Ge Y, Mlunde L, Mpembeni R, Leyna G, Huang J. High burden of anemia among pregnant women in Tanzania: a call to address its determinants. *Nutr J*. 2021;20(1). [doi:10.1186/s12937-021-00726-0](https://doi.org/10.1186/s12937-021-00726-0)
47. Noor RA. Micronutrient Deficiencies among Vulnerable Populations in Tanzania: Evidence for Interventions and Programmatic Implications. Published online 2019.
48. Daru J, Zamora J, Fernández-Félix BM, et al. Risk of maternal mortality in women with severe anaemia during pregnancy and post partum: a multilevel analysis. *Lancet Glob Health*. 2018;6(5):e548-e554. [doi:10.1016/s2214-109x\(18\)30078-0](https://doi.org/10.1016/s2214-109x(18)30078-0)
49. Saucedo M, Esteves-Pereira AP, Pencolé L, et al. Understanding maternal mortality in women with obesity and the role of care they receive: a national case-control study. *Int J Obes*. 2020;45(1):258-265. [doi:10.1038/s41366-020-00691-4](https://doi.org/10.1038/s41366-020-00691-4)
50. Keding GB, Msuya JM, Maass BL, Krawinkel MB. Obesity as a public health problem among adult women in rural Tanzania. *Glob Health Sci Pract*. 2013;1(3):359-371. [doi:10.9745/ghsp-d-13-00082](https://doi.org/10.9745/ghsp-d-13-00082)
51. Ramonienė G, Maleckienė L, Nadišauskienė RJ, et al. Maternal obesity and obstetric outcomes in a tertiary referral center. *Medicina*. 2017;53(2):109-113. [doi:10.1016/j.medici.2017.03.003](https://doi.org/10.1016/j.medici.2017.03.003)
52. Ramaiya KL, Swai AM, Mutabingwa TK, Mwanri AW, Kagaruki GB. Capacity and capability of Tanzania health facilities to diagnose and manage diabetes mellitus in pregnancy. *Diabetes Res Clin Pract*. 2018;145:119-129. [doi:10.1016/j.diabres.2018.05.003](https://doi.org/10.1016/j.diabres.2018.05.003)
53. Dalrymple KV, Flynn AC, Relph SA, O’Keeffe M, Poston L. Lifestyle Interventions in Overweight and Obese Pregnant or Postpartum Women for Postpartum Weight Management: A Systematic Review of the Literature. *Nutrients*. 2018;10(11):1704. [doi:10.3390/nu10111704](https://doi.org/10.3390/nu10111704)

## SUPPLEMENTARY MATERIALS

### ONLINE SUPPLEMENTARY DOCUMENT

Download: <https://www.joghr.org/article/87443-increasing-prevalence-of-overweight-and-obesity-among-tanzanian-women-of-reproductive-age-intending-to-conceive-evidence-from-three-demographic-health/attachment/178660.docx>

---