

**Effects of task difficulty on performance and event-related bradycardia during preparation for action**

Henderson, Jennifer; Kavussanu, Maria; Gallicchio, Germano; Ring, Christopher

Psychology of Sport and Exercise

DOI:

[10.1016/j.psychsport.2023.102548](https://doi.org/10.1016/j.psychsport.2023.102548)

E-pub ahead of print: 07/10/2023

Version created as part of publication process; publisher's layout; not normally made publicly available

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Henderson, J., Kavussanu, M., Gallicchio, G., & Ring, C. (2023). Effects of task difficulty on performance and event-related bradycardia during preparation for action. *Psychology of Sport and Exercise*. <https://doi.org/10.1016/j.psychsport.2023.102548>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?


Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Journal Pre-proof

Effects of task difficulty on performance and event-related bradycardia during preparation for action

Jennifer Henderson, Maria Kavussanu, Germano Gallicchio, Christopher Ring



PII: S1469-0292(23)00172-3

DOI: <https://doi.org/10.1016/j.psychsport.2023.102548>

Reference: PSYSPO 102548

To appear in: *Psychology of Sport & Exercise*

Received Date: 16 May 2023

Revised Date: 1 September 2023

Accepted Date: 1 October 2023

Please cite this article as: Henderson, J., Kavussanu, M., Gallicchio, G., Ring, C., Effects of task difficulty on performance and event-related bradycardia during preparation for action, *Psychology of Sport & Exercise* (2023), doi: <https://doi.org/10.1016/j.psychsport.2023.102548>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Ltd.

**Effects of task difficulty on performance and
event-related bradycardia during preparation for action**

Jennifer Henderson¹, Maria Kavussanu¹, Germano Gallicchio², & Christopher Ring¹

¹ School of Sport, Exercise & Rehabilitation Sciences, University of Birmingham, Birmingham,
UK

² School of Sport, Health & Exercise Sciences, Bangor University, Bangor, UK

Correspondence to:

Prof Christopher Ring, School of Sport, Exercise & Rehabilitation Sciences, University of
Birmingham, Birmingham, UK. c.m.ring@bham.ac.uk +44 1214144275

Abstract

The slowing of heart rate prior to movement onset has been presented as a marker of task-related cognitive processing and linked with performance accuracy. Here we examined this event-related bradycardia and task performance as a function of task difficulty. Forty experienced golfers completed a series of golf putting conditions that manipulated task difficulty by varying target distance, target size, and surface contour. Performance was measured by the number of holed putts and finishing distance from the hole. Physiological activity was recorded throughout. Analyses confirmed that performance varied as a function of task difficulty, worsening with longer distances to target, smaller targets, and sloping paths to target. Task difficulty also impacted the cardiac response, including the rate of heart rate deceleration, change in heart rate, and heart rate at impact. These heart rate metrics were found to correlate with performance strongly, moderately, and weakly, respectively. In conclusion, heart rate deceleration in the moments preceding movement onset was affected by task difficulty. Features of this cardiac deceleration pattern were characteristic of successful performance. Our findings are discussed in terms of the role of cognitive and motor processes during the execution of complex motor skills.

**Effects of task difficulty on performance and
event-related bradycardia during preparation for action**

Journal Pre-proof

1 Abstract

2 The slowing of heart rate prior to movement onset has been presented as a marker of task-
3 related cognitive processing and linked with performance accuracy. Here we examined this event-
4 related bradycardia and task performance as a function of task difficulty. Forty experienced golfers
5 completed a series of golf putting conditions that manipulated task difficulty by varying target
6 distance, target size, and surface contour. Performance was measured by the number of holed
7 putts and finishing distance from the hole. Physiological activity was recorded throughout. Analyses
8 confirmed that performance varied as a function of task difficulty, worsening with longer distances
9 to target, smaller targets, and sloping paths to target. Task difficulty also impacted the cardiac
10 response, including the rate of heart rate deceleration, change in heart rate, and heart rate at
11 impact. These heart rate metrics were found to correlate with performance strongly, moderately,
12 and weakly, respectively. In conclusion, heart rate deceleration in the moments preceding
13 movement onset was affected by task difficulty. Features of this cardiac deceleration pattern were
14 characteristic of successful performance. Our findings are discussed in terms of the role of
15 cognitive and motor processes during the execution of complex motor skills.

16

1 Introduction

2 The sufficient and necessary allocation of attentional resources is a key feature of expertise and
3 performance excellence in self-paced sport skills (Abernethy *et al.*, 2007; Wulf, 2007). Physiological
4 recordings can tell us about preparatory cognitive and motor processes and thereby serve as
5 markers of movement preparation when executing and learning motor skills (Cooke, 2013). One
6 such marker is the transient bradycardia that can be observed before cued and uncued (self-paced)
7 movements. A short-term phasic pattern of heart rate deceleration is typical in the seconds
8 preceding self-paced acts, such as golf (Cooke *et al.*, 2010, 2011; 2014; Cotterill & Collins, 2005;
9 Moore *et al.*, 2012; Neumann & Thomas, 2009), pistol (Tremayne & Barry, 2001), and rifle
10 (Hatfield *et al.*, 1987; Hoffman & Street, 1992) shots. Despite consensus on the reproducibility of
11 this phenomenon, its interpretation remains open to debate. Preparatory bradycardia has been
12 interpreted as a marker of somatic quiescence by some (Obrist, 1968; Obrist *et al.*, 1969, 1972,
13 1974) and a marker of attentional focus by others (Lacey & Lacey, 1970, 1974, 1980). The current
14 study sought to contribute to this unresolved debate by examining cardiac activity and
15 performance outcomes in relation to task difficulty for a self-paced motor task.

16 Lacey and Lacey's (1970, 1974, 1980) *intake-rejection hypothesis* argues that the event-related
17 heart rate deceleration pattern reflects a relative shift in the allocation of attentional resources
18 whereby individuals preferentially process (intake) external events and ignore (reject) internal
19 events. In broad terms, it can be considered as representing a change in stimulus processing
20 towards exteroception and away from interoception. For instance, the bradycardia before overt
21 movement onset might be interpreted in terms of attentional focus, namely, a shift to external
22 from internal focus of attention. The *intake-rejection hypothesis* is grounded in evidence from
23 reaction time studies (Lacey, 1967; Lacey & Lacey, 1970) showing that heart rate decelerates in
24 the fixed foreperiod between the presentation of the ready signal and the presentation of the
25 imperative signal.

1 These original studies have been replicated and extended by studies of simple and complex
2 motor skill tasks. Heart rate was shown to decelerate between the warning signal and imperative
3 signal cueing participants to perform leg lifts, climb a flight of stairs, and pedal on a cycle
4 ergometer (Chase *et al.*, 1968; Stern, 1976). It is worth noting that the size of the heart rate
5 deceleration was relatively small in these cued reaction time studies, generally averaging less than
6 five beats per minute. In contrast, evidence from studies that examined self-paced sport skills have
7 reported greater falls in heart rate (Cooke *et al.*, 2010, 2011; 2014; Cotterill & Collins, 2005;
8 Cottyn *et al.*, 2008; Hatfield *et al.*, 1987; Hoffman & Street, 1992; Moore *et al.*, 2012; Neumann &
9 Thomas, 2009; Tremayne & Barry, 2001). For instance, experts exhibited more substantial and
10 profound heart rate deceleration (i.e., -12 bpm) before the onset of the backswing than novices
11 (i.e., -4 bpm) while performing a golf putting task (Neumann & Thomas, 2009). It has been
12 suggested that the superior performance of experts compared to novices may reflect differences
13 in the extent to which performers pay attention to their external environment (e.g., the hole, the
14 ball, and the path between the ball and hole) that may inform the programming of the movement
15 (i.e., the direction and extent of the swing of the putter), and, that this contributes to the more
16 pronounced drop in heart rate.

17 Obrist and colleagues (e.g., Obrist, 1968; Obrist *et al.*, 1970; 1974; 1969) challenged the
18 cognitive interpretation of the movement-related heart rate deceleration pattern. Instead, they
19 offered an interpretation based upon a *general inhibition hypothesis*, whereby decreases in heart rate
20 are attributable to lower peripheral muscle activity/metabolism driven by diminished efferent
21 central motor commands. Thus, cardiac-somatic coupling is considered an organising feature of
22 the cardiovascular and motor systems of the body under standard conditions, and, therefore, it is
23 likely that variations in heart rate are closely connected with variations in somatomotor activity.
24 By way of illustration, Howard *et al.* (1974) observed that variations in heart rate in the context of
25 classical aversive conditioning were closely related to peripheral (i.e., general activity) and central

1 (i.e., pyramidal tract activity) movement-related activity, thereby demonstrating how physical
2 demands can orchestrate the heart rate response to a motor task.

3 With the aim of contributing to the debate on the interpretation of heart rate deceleration
4 during preparation for action, the present study adopted a multi-method approach to explore the
5 effects of task difficulty on phasic heart rate during a golf putting task in experienced golfers. Based
6 on the assumption that increased task difficulty requires increased cognitive processing (e.g.,
7 Henry & Rogers 1960; Walters-Symons, 2018), and that increased task difficulty is associated with
8 greater heart rate deceleration (e.g., Coles, 1972; 1974; Coles & Duncan-Johnson, 1977), we
9 manipulated task demands by altering distance from the hole (target), hole size, and putting surface
10 curvature. Evidence that the degree of bradycardia preceding task-relevant movement onset (i.e.,
11 swinging the club) is related to task difficulty independently of muscle activity would provide
12 support for the *intake-rejection hypothesis* (Lacey & Lacey, 1970, 1974, 1980). In contrast, evidence
13 that the degree of bradycardia is related to task difficulty relative to changes in muscle activity
14 would provide support for the *general inhibition hypothesis* (e.g., Obrist, 1968; Obrist *et al.*, 1970;
15 1974; 1969).

17 **Methods**

18 ***Participants***

19 Male ($n = 31$) and female ($n = 9$) right-handed sport and exercise science students participated
20 in exchange for course credit. Participants ($M = 20.18$, $SD = 1.34$ years) were regular golfers with
21 on-course playing experience ($M = 17.34$, $SD = 14.39$ golf handicap). The protocol was approved
22 by the local research ethics committee and all participants provided informed consent. Power
23 calculations using GPower 3.1.9.7 (Faul, et al., 2007) software indicated that with a sample size of
24 40 the current study was powered at .80 to detect significant ($p < .05$) differences among the
25 conditions using repeated measures analyses of variance corresponding to a small-to-medium ($f =$
26 .19 to .20) effect size (Cohen, 1992). The current sample size also exceeded those recruited for

1 previous experiments that compared the effects of task manipulations on various outcome
2 measures in this context (see Introduction).

3 **Measures**

4 *Perceived difficulty.* Participants rated task difficulty on a 7-point Likert scale, anchored by 1 “not
5 at all difficult” and 7 “very difficult”.

6 *Performance.* The primary measure of performance was number of holed putts (out of 9). A
7 secondary measure of performance, mean radial error (i.e., arithmetic mean of the distances the
8 ball finished from the hole), indexed putting accuracy (Hancock et al., 1995). Distance of the ball
9 from the hole was recorded as the distance (cm) from the centre of the hole to the closest point
10 of the ball. The number of putts per condition was a compromise between the effectiveness of
11 each experimental manipulation (Cooke et al., 2011) and the reliability of the measurements
12 (Schweizer et al., 2020). On the one hand, on a one-off trial, the impact of a manipulation on the
13 performer has high effectiveness and high ecological validity whereas measurements have low
14 reliability and high variability (e.g., Woodman & Davis, 2008). On the other hand, with large
15 numbers of trials, the impact of a manipulation on the performer has low effectiveness and low
16 ecological validity whereas measurements have high reliability and low variability (e.g., Cooke et al.,
17 2014). Mindful of the influence of consecutive repetitions on the attenuation of motor preparatory
18 and control processes (Gallicchio & Ring, 2019), we designed 9 trials per condition.

19 *Kinematics.* A tri-axial accelerometer (ADXL337 Breakout, Cool Components) recorded putter
20 head acceleration in three planes: X, Y, and Z acceleration measured lateral, vertical, and back-
21 and-forth clubhead movements. Contact between the ball and putter was measured by an impact
22 sensor (Piezo Vibration Sensor, Measurement Specialties). The accelerometer and impact sensor
23 were attached to the bottom of the putter shaft. Movement kinematics were determined during
24 the time between the initiation of the downswing and impact with the ball. The Z-axis
25 (mediolateral), which is the primary movement in putting (Cooke et al., 2010, 2011; Maxwell et al.,
26 2003), was used to calculate measures of kinematic proficiency (Nelson, 1983; Stelmach et al.,

1 1989), namely, root mean square jerk (i.e., rate of change of acceleration) and smoothness (i.e.,
2 number of sign changes in jerk signal). Mean values for each kinematic variable were computed by
3 averaging values across all putts in each condition.

4 *Muscular activity.* Left forearm and right upper arm muscle activity was recorded using single
5 differential surface electrodes (DE 2.1, Delsys) and amplifier (Bagnoli-4, Delsys), with a ground
6 electrode attached on the collar bone. The left flexor carpi radialis and right biceps brachii muscles
7 have been implicated in putting (Smith *et al.*, 2000; Stinear *et al.*, 2006). EMG signals were amplified,
8 (Power 1401, CED), filtered (20-450 Hz), digitalized (2500 Hz), and recorded using Spike 2
9 software. Mean EMG amplitude (μV) was calculated during the entire condition (i.e., mostly resting
10 muscle activity between the first and ninth ball strike) and during the brief window (c. 500 ms)
11 just before the initiation of the upswing of the putter stroke (see Cooke *et al.*, 2010). We
12 computed the change in EMG (i.e., pre-initiation minus overall resting activity) to capture the
13 characteristic increase in muscle activity in the preparatory period (see Moore *et al.*, 2012).

14 *Cardiac activity.* An electrocardiogram was recorded using three silver/silver chloride spot
15 electrodes (Cleartrace, ConMed) in a modified chest configuration. The signal was amplified
16 (Bagnoli-4 Delsys), filtered (1-100 Hz), and digitalized at 2500 Hz with 16-bit resolution (Power
17 1401, CED) using bespoke software (Spike2, CED). R-wave peaks were identified and verified by
18 an interactive program. The R-R intervals were used to compute heart rate (bpm) for each 0.5 s
19 epoch, from 10 s before impact with the ball to 5 s post-impact to capture the heart rate
20 deceleration profile (Cooke *et al.*, 2014; Neuman & Thomas, 2009, 2011). T tests were used to
21 confirm the highest and lowest heart rate in the 10 s before impact for the group in each
22 condition. Heart rate at putter-ball impact corresponded with the lowest point of the heart rate
23 deceleration response. The heart rate change was computed as the difference between the mean
24 highest heart rate and mean lowest heart rate per condition for each participant. The rate of heart
25 rate deceleration was computed as the heart rate change described above divided by the time
26 between the two heart rate epochs.

1 **Task and conditions**

2 Participants performed a golf putting task. Similar tasks have been used in previous preparation
3 for action studies (e.g., Cooke *et al.*, 2014; Neuman & Thomas, 2009, 2011). To be successful in
4 this task, participants were expected to accurately plan and program both movement force and
5 direction. Accordingly, cognitive processes, such as external focus of attention, were likely in the
6 seconds before movement execution.

7 In the control condition, participants putted nine standard sized golf balls (Pro V1, Titleist) to a
8 standard sized golf hole (10.8 cm diameter), located 2 m away using a standard length (90 cm)
9 steel-shafted blade style putter (Sedona 2, Ping). The hole was centrally located 0.25 m from the
10 end of a flat 1.5 m x 5 m artificial putting surface (Augin Turfites) with a Stimpmeter reading of
11 4.27 m. The finishing position of each putt was marked by a dot sticker and the ball removed;
12 these dots were used to measure distance from the hole at the end of each condition.

13 Task difficulty was manipulated in three ways: distance from target, hole diameter, and surface
14 contour. In the distance from target conditions, participants putted to a hole located 3 m, 1 m and
15 0.5 m away. These distances were expected to be harder, easier, and much easier than control,
16 respectively. In the hole diameter conditions, participants putted to a hole with a diameter of 8.1
17 cm and 5.4 cm; these were 75% and 50% the size of a standard golf hole. These hole diameters
18 were expected to be harder and much harder than control, respectively. In the surface profile
19 conditions, participants putted to a hole with a right-to-left break and left-to-right break. These
20 sloping surfaces were expected to be harder and much harder than control (Carnegie *et al.*, 2020).
21 Aside from the manipulated factor, conditions were otherwise the same as control (see above).

22 **Procedure**

23 Participants attended a single testing session. Following preparation and instruction, they
24 completed nine practice putts. They completed eight conditions (described above), with order
25 counterbalanced using a Latin square (Williams, 1949). No technical putting instructions were
26 provided. Participants were instructed to putt at their own pace. They were told that performance

1 was based on the number of holed putts and radial error, and, therefore, they should try to hole
2 the putt and, if they miss the putt, to finish as close to the hole as possible. A £20 reward was
3 offered for the best overall performer. Standard scripted instructions were read out by the
4 experimenter. A ball was placed in position by the experimenter prior to each putt to ensure
5 participants always stood upright. Participants completed self-report measures using a tablet
6 computer after each condition creating a 3 min rest period between conditions.

7 ***Statistical analysis***

8 A series of condition multivariate analyses of variance (ANOVAs), followed by Student *t* tests,
9 were employed to examine how performance, psychological, physiological, and kinematic
10 measures changed with task difficulty compared to control. This analytic approach was also used
11 to establish the effects of task difficulty within the three sub-themes: distance from target, hole
12 diameter, and surface profile. Heart rate was subjected to a condition × epoch ANOVA. We
13 report multivariate statistics for these repeated measures ANOVAs to minimize the risk of
14 violating sphericity and compound symmetry assumptions (Vasey & Thayer, 1987). Partial eta-
15 squared (η_p^2) is reported as a measure of effect size, with values of .02, .12 and .26 indicating small,
16 medium, and large effect sizes, respectively (Cohen, 1992). Within-participant correlations were
17 computed between our measures of performance (holed putts, radial error) and each of the heart
18 rate metrics (rate of heart rate deceleration, change in heart rate, heart rate at impact) in each of
19 the eight conditions. These correlation coefficients were transformed using the Fisher Z-
20 transformation (Siegel & Castellan, 1956), and were then averaged, back-transformed, and
21 interpreted. The effect size of the back-transformed coefficients were evaluated using guidelines,
22 with values of .10, .30 and .50 reflecting small, medium, and large effect sizes, respectively (Cohen,
23 1992).

24

25 **Results**

26 ***Effects of distance-based task difficulty***

1 Separate 4 condition (0.5 m, 1 m, 2 m, 3 m) ANOVAs confirmed large-sized effects of distance
 2 from the hole on 12 out of 15 measures (Table 1). The increase in distance from the ball to the
 3 hole was associated with greater perceived difficulty, decreased putting performance, and higher
 4 kinematics while putting. Muscle activity increased during the preparatory phase but this increase
 5 did not vary as a function of task difficulty. Although the initial resting heart rate and heart rate at
 6 impact were broadly similar across difficulty conditions, the rate of heart rate deceleration and
 7 change in heart rate tended to be smaller with increasing distance from the hole (i.e., greater task
 8 difficulty).

9 Heart rate decelerated in the moments before movement onset (Figure 1A). This event-related
 10 bradycardia was confirmed by a 4 condition (0.5 m, 1 m, 2 m, 3 m) \times 21 epoch (-10, -9.5, -9.0
 11 0 s) polynomial contrast analysis. These contrasts yielded time-varying effects for epoch (linear =
 12 $F(1, 39) = 64.02, p < .001, \eta_p^2 = .63$; quadratic = $F(1, 39) = 118.91, p < .001, \eta_p^2 = .75$; cubic = $F(1,$
 13 $39) = 8.51, p = .006, \eta_p^2 = .18$), condition (linear = $F(1, 39) = 21.18, p < .001, \eta_p^2 = .35$), and
 14 condition by epoch (linear \times linear = $F(1, 39) = 18.34, p < .001, \eta_p^2 = .32$; linear \times quadratic = $F(1,$
 15 $39) = 11.41, p = .002, \eta_p^2 = .23$; linear \times cubic = $F(1, 39) = 18.73, p < .001, \eta_p^2 = .32$).

16 **Effects of target size-based task difficulty**

17 Separate 3 condition (100%, 75%, 50%) ANOVAs confirmed large-sized effects of target (hole)
 18 size on 8 out of 15 measures (Table 2). The decrease in hole size was associated with greater
 19 perceived difficulty, poorer putting performance, and lower kinematics while putting. Muscle
 20 activity increased during the preparatory phase, but this increase did not vary as a function of task
 21 difficulty. The heart rate metrics were mostly unchanged by target size-based task difficulty; the
 22 exception was that the rate of heart rate deceleration was larger with decreasing hole size (i.e.,
 23 greater task difficulty).

24 Heart rate deceleration in the moments before movement onset (Figure 1B) was confirmed by
 25 a 3 condition (100%, 75%, 50%) \times 21 epoch (-10, -9.5, -9.0 0 s) polynomial contrast analysis,
 26 which yielded time-varying contrast effects for epoch (linear = $F(1, 39) = 75.58, p < .001, \eta_p^2 = .65$;

1 quadratic = $F(1, 39) = 64.23, p < .001, \eta_p^2 = .62$). No condition or condition \times epoch contrasts
2 were found.

3 **Effects of contour-based task difficulty**

4 Separate 3 condition (straight, right-to-left break, left-to-right break) ANOVAs confirmed large-
5 sized effects of surface contour on 9 out of 15 measures (Table 3). Compared to straight putts on
6 a flat surface, breaking putts on a sloping surface, especially those with a left-to-right break, were
7 associated with greater perceived difficulty, poorer putting performance, and higher kinematics
8 while putting. Muscle activity increased during the preparatory phase but did not vary as a function
9 of task difficulty. The heart rate metrics were mostly unchanged by contour-based task difficulty;
10 the exception was that initial heart rate was marginally slower before executing breaking putts
11 (i.e., greater task difficulty).

12 Heart rate deceleration in the moments before movement onset (Figure 1C) was confirmed by
13 a 3 condition (straight, right-to-left break, left-to-right break) \times 21 epoch (-10, -9.5, -9.0 0 s)
14 polynomial contrast analysis. These contrasts yielded time-varying effects for epoch (linear = $F(1,$
15 $39) = 67.00, p < .001, \eta_p^2 = .63$; quadratic = $F(1, 39) = 82.35, p < .001, \eta_p^2 = .68$), and condition
16 (linear = $F(1, 39) = 7.83, p = .008, \eta_p^2 = .17$). No condition \times epoch contrast was observed.

17 **Relationships between cardiac metrics and task performance indices**

18 Within-participant correlations were performed between each cardiac metric (rate of heart
19 rate deceleration, change in heart rate, heart rate at impact) and the separate performance indices
20 (number of holed putts, radial error) across the eight task conditions using the average across the
21 9 putts for each condition (Table S1, Supplementary Materials). Each participant's Pearson's
22 correlation coefficient (e.g., between change in heart rate and radial error) was transformed using
23 the Fisher Z transformation, the average of these transformed coefficients was computed, the
24 average was back-transformed to a Pearson correlation coefficient, and the size of this coefficient
25 tested for linear independence (i.e., compared with 0 using a t test) and interpreted as small,
26 medium or large (Cohen, 1992). These analyses indicated that the rate of heart rate deceleration

1 was strongly related (i.e., large effect size) to the number of holed putts, $r(6) = -.58$, $p = .07$, and
2 radial error, $r(6) = .48$, $p = .11$. The change in heart rate was moderately related (i.e., medium
3 effect size) to holed putts, $r = -.28$, $p = .25$, and radial error, $r(6) = -.23$, $p = .29$. Heart rate at
4 impact was weakly related (i.e., small effect size) to holed putts, $r(6) = -.16$, $p = .35$, and radial
5 error, $r(6) = .11$, $p = .40$. In sum, task performance accuracy was better – more putts were holed
6 and putts finished closer to the hole – when heart rate decelerated faster.

7

8 **Discussion**

9 The current study sought to evaluate the *intake-rejection hypothesis* (Lacey & Lacey, 1970, 1974,
10 1980) and *general inhibition hypothesis* (Obrist, 1968; Obrist et al., 1970; 1974; 1969) as
11 explanations for the bradycardia response during preparation for action (Cooke, 2013). We
12 examined the effects of task difficulty on heart rate in the seconds preceding a complex self-paced
13 and goal-directed movement. Heart rate decelerated by about 12 beats per minute over a period
14 of about 8 s until club-ball impact (Figure 1). Subsequently, it then sped up over the next 6 s and
15 returned to and eventually surpassed initial values. This consistent pattern resembled that
16 reported previously in golf putting tasks (Boutcher & Zinsser, 1990; Cooke et al., 2014; Moore et
17 al., 2012; Neumann & Thomas, 2009, 2011). Our findings confirmed that task difficulty influenced
18 the rate but not the magnitude of heart rate deceleration. In particular, the magnitude of heart
19 rate deceleration and heart rate at impact were relatively invariant whereas heart rate began to fall
20 earlier and took longer to reach its minimum with increasing task difficulty (Figure 1). By
21 illustration, when the ball was closest to the hole and putting was easiest, heart rate only began to
22 fall 4 s before the club hit the ball compared to the typical 8 s in the other conditions.

23 Previous studies have found that the magnitude of the heart rate deceleration response is a
24 feature of expertise, being greater in experts than novices (Cooke et al., 2014; Neumann &
25 Thomas 2009, 2011). The current findings argue that the extent of this bradycardia is not sensitive
26 to task demands. Instead, the onset of the bradycardia was moderated by task difficulty. The *intake*

1 *rejection hypothesis* (Lacey & Lacey, 1970, 1974, 1980) would interpret these data as showing that
2 participants adopted an external focus of attention when preparing to putt a golf ball to a distant
3 target. Variations among task difficulty conditions in the rate of heart rate deceleration might
4 suggest that information in the external environment, such as required ball path and distance from
5 ball to hole, is processed and used to program movement parameters, such as direction and force
6 (Moore *et al.*, 2012; Requin *et al.*, 1991). Enhanced exteroceptive processing during preparation for
7 action, where novel features of the environment provide information about the task, may
8 therefore aid motor execution. Previous evidence has linked the onset of heart rate changes with
9 performance accuracy (e.g., Moore *et al.*, 2012; Neumann & Thomas, 2009; Tremayne & Barry,
10 2001). For instance, expert pistol shooters' heart rate deceleration began 3.5 s earlier on best
11 shots compared to worst shots (Tremayne & Barry, 2001). Accordingly, the slower rate of heart
12 rate deceleration observed in the current study suggests that attention shifts to process
13 information from the external environment depending on the difficulty of the motor task. More
14 difficult task demands, which required attention to begin earlier and to last longer, were associated
15 with earlier onset and more gradual slowing of heart rate. It should be noted that, in line with the
16 current findings, most previous golf putting studies have failed to find evidence that the absolute
17 change in heart rate was related to variations in putting performance. Instead, they found evidence
18 that the absolute change in heart rate was a characteristic of expertise. We speculate that this
19 consistent finding suggests that the absolute change in heart rate may simply reflect the well-honed
20 and practiced pre-shot routines of experts.

21 The *general inhibition hypothesis* (Obrist, 1968; Obrist *et al.*, 1970; 1974; 1969) explains changes
22 in cardiac activity in terms of accompanying changes in peripheral and/or central somatomotor
23 activity. In accordance with the *cardiac-somatic coupling* principle, decreases in cardiac activity
24 should be accompanied by concomitant decreases in muscle activity. Several issues need to be
25 considered here. First, the hypothesis would expect that the heart rate deceleration response in
26 the seconds before movement onset should be accompanied by reductions in muscle activity.

1 However, previous golf putting studies (Cooke *et al.*, 2014; Moore *et al.*, 2012) show that
2 decreases in heart rate in the seconds before the golf swing are accompanied by increases in
3 muscle activity in the upper and lower arms, which are explicitly required to control the putter
4 and execute the task. Similarly, we observed increased electromyographic activity in these arm
5 muscles (see Tables 1, 2 and 3). It should be noted that other muscles, such as postural,
6 oculomotor, or respiratory muscles, may have been deactivated during this preparatory period.
7 This is a possibility given evidence that the eyes are likely to fix on the ball and remain relatively
8 still in the period before movement onset in golf putting tasks (e.g., Moore *et al.*, 2012; Gallicchio
9 *et al.*, 2018; Gallicchio & Ring, 2020). Similarly, it is possible that participants varied their rate and
10 depth of breathing before putting (e.g., Neumann & Thomas, 2009). None of these muscles were
11 recorded in the current study and therefore this possibility cannot be discounted. Second, the
12 hypothesis would expect that the onset of the heart rate deceleration response should be
13 preceded by greater muscle activity. However, we found no differences among the eight task
14 conditions in background electromyographic activity in these arm muscles. This evidence would
15 therefore argue against this possibility. It should be conceded that we did not measure the activity
16 of other muscles, which may have been activated prior to the onset of the bradycardia.

17 **Conclusion**

18 The present study found that the rate but not the magnitude of heart rate deceleration in the
19 seconds preceding movement was sensitive to variations in task difficulty. These findings may
20 reflect the effects of the experimental manipulations on attentional (Lacey & Lacey, 1970, 1974,
21 1980) or motor (Obrist, 1968) processes. The current findings are better explained by the *intake-*
22 *rejection hypothesis* than the *general inhibition hypothesis*. It should be acknowledged that our
23 interpretation favouring the intake-rejection hypothesis is best supported by our findings from the
24 distance from the hole manipulations (Figure 1A, Table 1). The interpretation receives less support
25 from the size of the hole manipulation (Figure 1B, Table 2). Finally, the interpretation is not clearly
26 supported by the surface contour manipulation (Figure 1C, Table 3). Clearly, additional

1 manipulations, including task difficulty, and markers, such as cortical activity, are needed before we
2 can confidently decide which of the two competing hypotheses best account for cardiac
3 deceleration in preparation for action.

4

5 **References**

- 6 Abernethy, B., Maxwell, J.P., Masters, R.S.W., van der Kamp, J., & Jackson, R.C. (2007). Attentional
7 processes in skill learning and expert performance. In: G. Tenenbaum & R. C. Eklund (Eds.),
8 *Handbook of sport psychology* (pp. 245–263). Wiley.
- 9 Boutcher, S.H., & Zinsser, N.W. (1990). Cardiac deceleration of elite and beginning golfers during
10 putting. *Journal of Sport & Exercise Psychology*, *12*(1), 37-47. <https://doi.org/10.1123/jsep.12.1.37>
- 11 Carnegie, E., Marchant, D., Towers, S., & Ellison, P. (2020). Beyond visual fixations and gaze
12 behaviour. Using pupillometry to examine the mechanisms in the planning and motor
13 performance of a golf putt. *Human Movement Science*, *71*, 102622.
14 <https://doi.org/10.1016/j.humov.2020.102622>
- 15 Carroll, D., Rick Turner, J., & Hellawell, J. C. (1986). Heart rate and oxygen consumption during
16 active psychological challenge: The effects of level of difficulty. *Psychophysiology*, *23*(2), 174-181.
17 <https://doi.org/10.1111/j.1469-8986.1986.tb00613.x>
- 18 Chase, W. G., Graham, F. K., & Graham, D. T. (1968). Components of HR response in anticipation
19 of reaction time and exercise tasks. *Journal of Experimental Psychology*, *76*(4), 642-648.
20 <https://doi.org/10.1037/h0021288>
- 21 Cohen, J. (1992). A power primer. *Psychological Bulletin*, *112*(1), 155-159.
22 <https://doi.org/10.1037/0033-2909.112.1.155>
- 23 Coles, M. G. H. (1972). Cardiac and respiratory activity during visual search. *Journal of Experimental*
24 *Psychology*, *96*(2), 371-379. <https://doi.org/10.1037/h0033603>

- 1 Coles, M. G. H. (1974). Physiological activity and detection: The effects of attentional requirements
2 and the prediction of performance. *Biological Psychology*, 2(2), 113-125.
3 [https://doi.org/10.1016/0301-0511\(74\)90019-2](https://doi.org/10.1016/0301-0511(74)90019-2)
- 4 Coles, M. G., & Duncan-Johnson, C. C. (1977). Attention and cardiac activity: Heart rate
5 responses during a variable foreperiod, disjunctive reaction time task. *Biological Psychology*, 5(2),
6 151-158. [https://doi.org/10.1016/0301-0511\(77\)90010-2](https://doi.org/10.1016/0301-0511(77)90010-2)
- 7 Cooke, A. (2013) Readyng the head and steadying the heart: A review of cortical and cardiac
8 studies of preparation for action in sport. *International Review of Sport & Exercise Psychology*, 6(1),
9 122-138. <https://doi.org/10.1080/1750984X.2012.724438>
- 10 Cooke, A., Kavussanu, M., McIntyre, D., Boardley, I.D., & Ring, C. (2011). Effects of competitive
11 pressure on expert performance: Underlying psychological, physiological and kinematic
12 mechanisms. *Psychophysiology*, 48(8), 1146-1156. <https://doi.org/10.1111/j.1469-8986.2011.01175.x>
- 13 Cooke, A., Kavussanu, M., McIntyre, D., & Ring, C. (2010). Psychological, muscular and kinematic
14 factors mediate performance under pressure. *Psychophysiology*, 47(6), 1109-1118.
15 <https://doi.org/10.1111/j.1469-8986.2010.01021.x>
- 16 Cooke, A., Kavussanu, M., Gallicchio, G., Willoughby, A., McIntyre, D., & Ring, C. (2014).
17 Preparation for action: Psychophysiological activity preceding a motor skill as a function of
18 expertise, performance outcome, and psychological pressure. *Psychophysiology*, 51(4), 374-384.
19 <https://doi.org/10.1111/psyp.12182>
- 20 Cotterill, S., & Collins, D. (2005). Heart rate deceleration characteristics across shot types in golf.
21 *Journal of Sports Science*, 23, 173-174.
- 22 Cottyn, J., De Clercq, D., Crombez, G., & Lenoir, M. (2008). The role of preparatory heart rate
23 deceleration on balance beam performance. *Journal of Sport & Exercise Psychology*, 30(2), 159-
24 170. <https://doi.org/10.1123/jsep.30.2.159>

- 1 Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power
2 analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*,
3 39, 175-191. <https://doi.org/10.3758/BF03193146>
- 4 Gallicchio, G., Cooke, A., & Ring, C. (2018). Assessing ocular activity during performance of motor
5 skills use electroculography. *Psychophysiology*, 55(7), e13070. <https://doi.org/10.1111/psyp.13070>
- 6 Gallicchio, G., & Ring, C. (2019). Don't look, don't think, just do it! Toward an understanding of
7 alpha gating in a discrete aiming task. *Psychophysiology*, 56(3), e13298.
8 <https://doi.org/10.1111/psyp.13298>
- 9 Gallicchio, G., & Ring, C. (2020). The quiet eye effect: visual and postural-kinematic hypotheses.
10 *Sport, Exercise & Performance Psychology*, 9(1), 143-159. <https://doi.org/10.1037/spy0000162>
- 11 Hancock, G. R., Butler, M. S., & Fischman, M. G. (1995). On the problem of two-dimensional error
12 scores: measures and analyses of accuracy, bias, and consistency. *Journal of Motor Behavior*, 27(3),
13 241-250. <https://doi.org/10.1080/00222895.1995.9941714>
- 14 Hatfield, B.D., Landers, D.M., & Ray, W.J. (1987). Cardiovascular-CNS interactions during a self-
15 paced, intentional attentive state: Elite marksmanship performance. *Psychophysiology*, 24(5), 542-
16 549. <https://doi.org/10.1111/j.1469-8986.1987.tb00335.x>
- 17 Hassmén, P., & Koivula, N. (2001). Cardiac deceleration in elite golfers as modified by noise and
18 anxiety during putting. *Perceptual & Motor Skills*, 92(3), 947-957.
19 <https://doi.org/10.2466/pms.2001.92.3c.947>
- 20 Henry, F.M., & Rogers, D.E. (1960). Increased response latency for complicated movements and a
21 “memory drum” theory of neuromotor reaction. *Research Quarterly. American Association Health,*
22 *Physical Education & Recreation*, 31(3), 448-458. <https://doi.org/10.1080/10671188.1960.10762052>
- 23 Hoffman, M.D., & Street, G.M. (1992). Characterization of the heart rate response during biathlon.
24 *International Journal of Sports Medicine*, 13(5), 390-394. <https://doi.org/10.1055/s-2007-1021286>

- 1 Howard, J. L., Obrist, P. A., Gaebelein, C. J., & Galosy, R. A. (1974). Multiple somatic measures and
2 heart rate during classical aversive conditioning in the cat. *Journal of Comparative and Physiological*
3 *Psychology*, 87(2), 228-236. <https://doi.org/10.1037/h0036875>
- 4 Jennings, J.R., & van der Molen, M.W. (2005). Preparation for speeded action as a
5 psychophysiological concept. *Psychological Bulletin*, 131(3), 434-459. <https://doi.org/10.1037/0033->
6 2909.131.3.434
- 7 Konttinen, N., Lyytinen, H., & Viitasalo, J. (1998). Preparatory heart rate patterns in competitive
8 rifle shooting. *Journal of Sports Sciences*, 16, 235-242. <https://doi.org/10.1080/026404198366759>
- 9 Lacey, B. C., & Lacey, J. L. (1970). Some autonomic-central nervous system interrelationships. In P.
10 Black (Ed.), *Physiological correlates of emotion* (pp. 205–227). Academic Press.
- 11 Lacey, B. C., & Lacey, J. L. (1974). Studies of heart rate and other bodily processes in sensorimotor
12 behaviour. In P. A. Obrist, A. H. Black, J. Brener, & L. V. DiCara (Eds.), *Cardiovascular*
13 *psychophysiology* (pp. 538–564). Aldine.
- 14 Lacey, B.C., & Lacey, J.L. (1980). Cognitive modulation of time dependent primary bradycardia.
15 *Psychophysiology*, 17, 209–221. <https://doi.org/10.1111/j.1469-8986.1980.tb00137.x>
- 16 Land, M. F. (2009). Vision, eye movements, and natural behavior. *Visual Neuroscience*, 26(1), 51-62.
17 <https://doi.org/10.1017/S0952523808080899>
- 18 Maxwell, J.P., Masters, R.S.W., & Eves, F.F. (2003). The role of working memory in motor learning
19 and performance. *Consciousness & Cognition*, 12(3), 376-402. <https://doi.org/10.1016/S1053->
20 8100(03)00005-9
- 21 Moore, L. J., Vine, S. J., Cooke, A., Ring, C., & Wilson, M. R. (2012). Quiet eye training expedites
22 motor learning and aids performance under heightened anxiety: The roles of response
23 programming and external attention. *Psychophysiology*, 49(7), 1005-1015.
24 <https://doi.org/10.1111/j.1469-8986.2012.01379.x>

- 1 Neumann, D. L., & Thomas, P. R. (2009). The relationship between skill level and patterns in
2 cardiac and respiratory activity during golf putting. *International Journal of Psychophysiology*, 72(3),
3 276-282. <https://doi.org/10.1016/j.ijpsycho.2009.01.001>
- 4 Neumann, D. L., & Thomas, P. R. (2011). Cardiac and respiratory activity and golf putting
5 performance under attentional focus instructions. *Psychology of Sport & Exercise*, 12(4), 451-459.
6 <https://doi.org/10.1016/j.psychsport.2011.02.002>
- 7 Obrist, P. A. (1968). Heart rate and somatic-motor coupling during classical aversive conditioning
8 in humans. *Journal of Experimental Psychology*, 77(2), 180-193. <https://doi.org/10.1037/h0025814>
- 9 Obrist, P.A., Howard, J.L., Lawler, J.E., Galosy, R.A., Meyers, K.A., & Gaebelein, C.J. (1974). The
10 cardiac-somatic interaction. In: P.A. Obrist, A.H. Black, J. Brener, & L.V. DiCara (Eds.),
11 *Cardiovascular psychophysiology* (pp. 136-162). Aldine.
- 12 Obrist, P. A., Webb, R. A., & Sutterer, J. R. (1969). Heart rate and somatic changes during aversive
13 conditioning and a simple reaction time task. *Psychophysiology*, 5(6), 696-723.
14 <http://psycnet.apa.org/doi/10.1111/j.1469-8986.1969.tb02872.x>
- 15 Obrist, P. A., Webb, R. A., Sutterer, J. R., & Howard, J. L. (1970). Cardiac deceleration and
16 reaction time: An evaluation of two hypotheses. *Psychophysiology*, 6(6), 695-706.
17 <https://doi.org/10.1111/j.1469-8986.1970.tb02257.x>
- 18 Radlo, S. J., Steinberg, G. M., Singer, R. N., Barba, D. A., & Melnikov, A. (2002). The influence of an
19 attentional focus strategy on alpha brain wave activity, heart rate and dart-throwing
20 performance. *International Journal of Sport Psychology*, 33(2), 205-217.
- 21 Requin, J., Brener, J., & Ring, C. (1991). Preparation for action. In: J.R. Jennings & M.G.H. Coles
22 (Eds.) *Handbook of cognitive psychophysiology* (357-448). Wiley.
- 23 Schweizer, G., Furley, P., Rost, N., & Barth, K. (2020). Reliable measurement in sport psychology:
24 The case of performance outcome measures. *Psychology of Sport and Exercise*, 48, 101663.
25 <https://doi.org/10.1016/j.psychsport.2020.101663>
- 26 Siegel, S., & Castellan, J.N. (1956). *Nonparametric studies for the behavioral sciences*. McGraw-Hill.

- 1 Smith, A.M., Malo, S.A., Laskowski, E.R., Sabick, M., Cooney, W.P., Finnie, S.B., Crews, D. J.,
2 Eischen, J.J., Hay, I.D., Detling, N.J. (2000). A multidisciplinary study of the 'yips' phenomenon in
3 golf. *Sports Medicine*, 30(6), 423-437. <https://doi.org/10.2165/00007256-200030060-00004>
- 4 Stern, R. M. (1976). Reaction time and heart rate between the GET SET and GO of simulated
5 races. *Psychophysiology*, 13(2), 149-154. <https://doi.org/10.1111/j.1469-8986.1976.tb00091.x>
- 6 Stinear, C. M., Coxon, J. P., Fleming, M. K., Lim, V. K., Prapavessis, H., & Byblow, W. D. (2006).
7 The yips in golf: Multimodal evidence for two subtypes. *Medicine & Science in Sports &*
8 *Exercise*, 38(11), 1980-1989. <https://doi.org/10.1249/01.mss.0000233792.93540.10>
- 9 Tremayne, P., & Barry, R.J. (2001). Elite pistol shooters: Physiological patterning of best vs. worst
10 shots. *International Journal of Psychophysiology*, 41(1), 19-29. [https://doi.org/10.1016/S0167-](https://doi.org/10.1016/S0167-8760(00)00175-6)
11 [8760\(00\)00175-6](https://doi.org/10.1016/S0167-8760(00)00175-6)
- 12 Vasey, M. W., & Thayer, J. F. (1987). The continuing problem of false positives in repeated
13 measures ANOVA in psychophysiology: A multivariate solution. *Psychophysiology*, 24(4), 479-
14 486. <https://doi.org/10.1111/j.1469-8986.1987.tb00324.x>
- 15 Walters-Symons, R., Wilson, M., Klostermann, A., & Vine, S. (2018). Examining the response
16 programming function of the Quiet Eye: Do tougher shots need a quieter eye? *Cognitive*
17 *Processing*, 19(1), 47-52. <https://doi.org/10.1007/s10339-017-0841-6>
- 18 Williams, E. J. (1949). Experimental designs balanced for the estimation of residual effects of
19 treatments. *Australian Journal of Chemistry*, 2(2), 149-168. <https://doi.org/10.1071/CH9490149>
- 20 Woodman, T., & Davis, P. A. (2008). The role of repression in the incidence of ironic errors. *The*
21 *Sport Psychologist*, 22, 183-198. <https://doi.org/10.1123/tsp.22.2.183>
- 22 Wulf, G. (2007). *Attention and motor skill learning*. Human Kinetics.

Figure 1. Mean heart rate change (bpm) in the 10 s prior to and 5 s following putter-ball impact as a function of (A) distance from the hole, (B) size of the hole, and (C) surface contour.




Table 1. Effects of distance from hole on perception, performance, kinematics and physiology.

Measure	Distance								F(3, 37)	η_p^2
	0.5 m		1 m		2 m		3 m			
	M	SD	M	SD	M	SD	M	SD		
Difficulty (1-7)	1.61	0.86	2.79 ^a	1.36	2.74 ^a	1.17	4.68 ^{a,b,c}	1.38	61.63 ^{***}	.83
Number of Holed Putts (0-9)	8.98	0.16	6.90 ^a	1.89	6.55 ^a	1.88	3.78 ^{a,b,c}	1.98	95.26 ^{***}	.89
Radial Error (cm)	0.04	0.26	7.36 ^a	6.54	9.30 ^a	7.96	28.14 ^{a,b,c}	18.72	39.22 ^{***}	.76
X-axis Acceleration (m·s ⁻²)	0.28	0.07	0.34 ^a	0.09	0.50 ^{a,b}	0.14	0.62 ^{a,b,c}	0.18	77.86 ^{***}	.86
Y-axis Acceleration (m·s ⁻²)	0.46	0.34	0.48 ^a	0.36	0.58 ^{a,b}	0.49	0.67 ^{a,b,c}	0.50	21.67 ^{***}	.64
Z-axis Acceleration (m·s ⁻²)	2.34	0.53	2.66 ^a	0.61	3.22 ^{a,b}	0.76	3.65 ^{a,b,c}	0.75	84.44 ^{***}	.87
Jerk (m·s ⁻³)	2.35	0.54	2.65 ^a	0.60	3.18 ^{a,b}	0.71	3.60 ^{a,b,c}	0.73	85.24 ^{***}	.87
Smoothness	61.53	10.54	63.99 ^a	10.94	67.74 ^{a,b}	12.56	68.21 ^{a,b,c}	10.30	13.72 ^{***}	.53
Change in Flexor Carpi Radialis EMG (μV)	7.45	8.68	8.49	11.03	8.72	10.10	9.24	11.34	1.38	.26
Change in Biceps Brachii EMG (μV)	1.38	6.04	1.54	7.79	0.59	6.11	0.71	6.62	1.72	.12
Heart Rate (bpm)	84.97	12.39	85.10	12.40	85.95	12.50	85.47	12.74	1.18	.09
Change in Heart Rate (bpm)	-12.96	7.26	-14.60	8.57	-14.25	8.39	-11.77 ^{b,c}	6.85	5.17 ^{**}	.30
Rate of Heart Rate Deceleration (bpm)	-194.37	108.97	-134.81 ^a	79.10	-122.10 ^a	71.88	-78.47 ^{a,b,c}	45.64	35.10 ^{***}	.74
Heart Rate at Impact (bpm)	74.56	11.48	73.69	10.30	75.09	12.65	76.06 ^{a,b}	11.92	4.79 ^{**}	.28

Note: Superscripted letters ^a, ^b and ^c indicate significant difference from 0.5 m, 1 m, and 2 m (control) conditions, respectively. * $p < .05$, ** $p < .01$, *** $p < .001$.

Table 2. Effects of size of the hole on perception, performance, kinematics and physiology.

Measure	Size						F(2, 38)	η_p^2
	100%		75%		50%			
	M	SD	M	SD	M	SD		
Difficulty (1-7)	2.74	1.17	4.18 ^a	1.45	5.00 ^{a,b}	1.20	49.87 ^{***}	.72
Number of Holed Putts (0-9)	6.55	1.88	5.65 ^a	2.20	3.73 ^{a,b}	2.24	23.10 ^{***}	.55
Radial Error (cm)	9.30	7.96	12.10 ^a	10.06	17.55 ^{a,b}	12.10	11.33 ^{***}	.37
X-axis Acceleration (m·s ⁻²)	0.50	0.14	0.48 ^a	0.14	0.45 ^{a,b}	0.13	15.04 ^{***}	.42
Y-axis Acceleration (m·s ⁻²)	0.58	0.49	0.59	0.53	0.57	0.51	0.47	.02
Z-axis Acceleration (m·s ⁻²)	3.22	0.76	3.07 ^a	0.71	3.00 ^a	0.67	8.84 ^{***}	.32
Jerk (m·s ⁻³)	3.18	0.71	3.03 ^a	0.67	2.95 ^{a,b}	0.63	10.06 ^{***}	.35
Smoothness	67.74	12.56	69.03	12.40	67.59	11.82	2.99	.14
Change in Flexor Carpi Radialis EMG (μV)	8.72	10.10	8.27	10.17	8.09	10.92	0.64	.03
Change in Biceps Brachii EMG (μV)	0.59	6.11	1.69	8.99	0.87	7.66	0.53	.03
Heart Rate (bpm)	85.95	12.50	84.84	12.14	84.89	11.98	2.43	.11
Change in Heart Rate (bpm)	-14.25	8.39	-13.45	7.22	-12.38	7.43	2.50	.12
Rate of Heart Rate Deceleration (bpm)	-122.10	71.88	-115.25	61.93	-92.89 ^{a,b}	55.69	11.19 ^{***}	.37
Heart Rate at Impact (bpm)	75.09	12.65	74.87	11.57	75.18	11.96	0.19	.01

Note: Superscripted letters ^a and ^b indicate significant difference from 100% (control) and 75% conditions, respectively. * $p < .05$, ** $p < .01$, *** $p < .001$.

Table 3. Effects of surface contour on perception, performance, kinematics and physiology.

Measure	Contour						F(2, 38)	η_p^2
	Straight		Right-to-Left		Left-to-Right			
	M	SD	M	SD	M	SD		
Difficulty (1-7)	2.74	1.17	4.32 ^a	1.43	4.50 ^a	1.51	26.53 ^{***}	.58
Number of Holed Putts (0-9)	6.55	1.88	6.28	2.40	5.13 ^{a,b}	2.13	6.44 ^{**}	.25
Radial Error (cm)	9.30	7.96	12.65	12.91	24.62 ^{a,b}	16.33	16.80 ^{***}	.47
X-axis Acceleration (m·s ⁻²)	0.50	0.14	0.56 ^a	0.16	0.58 ^{a,b}	0.17	11.19 ^{***}	.37
Y-axis Acceleration (m·s ⁻²)	0.58	0.49	0.61	0.54	0.66 ^{a,b}	0.58	7.02 ^{**}	.27
Z-axis Acceleration (m·s ⁻²)	3.22	0.76	3.51 ^a	0.75	3.50 ^a	0.71	22.27 ^{***}	.54
Jerk (m·s ⁻³)	3.18	0.71	3.47 ^a	0.72	3.46 ^a	0.69	24.97 ^{***}	.57
Smoothness	67.74	12.56	65.46 ^a	10.95	64.35 ^a	11.15	8.07 ^{***}	.30
Change in Flexor Carpi Radialis EMG (μ V)	8.72	10.10	8.33	10.09	9.00	10.84	0.51	.03
Change in Biceps Brachii EMG (μ V)	0.59	6.11	0.47	5.77	0.77	6.14	0.41	.02
Heart rate (bpm)	85.95	12.50	83.89 ^a	12.18	84.08 ^a	11.62	7.03 ^{**}	.27
Change in Heart Rate (bpm)	-14.25	8.39	-12.80	6.32	-13.07	6.65	1.50	.07
Rate of Heart Rate Deceleration (bpm)	-122.10	71.88	-109.69	54.17	-120.67	61.39	2.30	.11
Heart Rate at Impact (bpm)	75.09	12.65	74.47	11.50	73.60	11.22	2.63	.12

Note: Superscripted letters ^a and ^b indicate significant difference from straight (control) and right-to-left slope conditions, respectively. * $p < .05$, ** $p < .01$, *** $p < .001$.

Highlights

- Heart rate slowing before movement reflects attentional and/or motor processes
- We confirm bradycardia during preparation for action
- Its timing and extent depended on task difficulty
- Data favor *intake-rejection hypothesis* rather than *general inhibition hypothesis*

Journal Pre-proof

Declaration of Interest

We declare no conflicts of interest.

Journal Pre-proof