

Mapping local variation in household overcrowding across Africa from 2000 to 2018: a modelling study



Michael G Chipeta, Emmanuelle P A Kumaran, Annie J Browne, Bahar H Kashef Hamadani, Georgina Haines-Woodhouse, Benn Sartorius, Robert C Reiner Jr, Christiane Dolecek, Simon I Hay, Catrin E Moore



Summary

Background Household overcrowding is a serious public health threat associated with high morbidity and mortality. Rapid population growth and urbanisation contribute to overcrowding and poor sanitation in low-income and middle-income countries, and are risk factors for the spread of infectious diseases, including COVID-19, and antimicrobial resistance. Many countries do not have adequate surveillance capacity to monitor household overcrowding. Geostatistical models are therefore useful tools for estimating household overcrowding. In this study, we aimed to estimate household overcrowding in Africa between 2000 and 2018 by combining available household survey data, population censuses, and other country-specific household surveys within a geostatistical framework.

Methods We used data from household surveys and population censuses to generate a Bayesian geostatistical model of household overcrowding in Africa for the 19-year period between 2000 and 2018. Additional sociodemographic and health-related covariates informed the model, which covered 54 African countries.

Findings We analysed 287 surveys and population censuses, covering 78 695 991 households. Spatial and temporal variability arose in household overcrowding estimates over time. In 2018, the highest overcrowding estimates were observed in the Horn of Africa region (median proportion 62% [IQR 57–63]); the lowest regional median proportion was estimated for the north of Africa region (16% [14–19]). Overall, 474·4 million (95% uncertainty interval [UI] 250·1 million–740·7 million) people were estimated to be living in overcrowded conditions in Africa in 2018, a 62·7% increase from the estimated 291·5 million (180·8 million–417·3 million) people who lived in overcrowded conditions in the year 2000. 48·5% (229·9 million) of people living in overcrowded conditions came from six African countries (Nigeria, Ethiopia, Democratic Republic of the Congo, Sudan, Uganda, and Kenya), with a combined population of 538·3 million people.

Interpretation This study incorporated survey and population censuses data and used geostatistical modelling to estimate continent-wide overcrowding over a 19-year period. Our analysis identified countries and areas with high numbers of people living in overcrowded conditions, thereby providing a benchmark for policy planning and the implementation of interventions such as in infectious disease control.

Funding UK Department of Health and Social Care, Wellcome Trust, Bill & Melinda Gates Foundation.

Copyright © 2022 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

Household overcrowding is associated with higher morbidity and mortality, and with poorer sociodemographic conditions in urban areas (eg, as shown in Beirut) than in households that are not overcrowded.^{1–4} Moreover, it is known that transmission of infectious diseases occurs at a higher rate when people are in close contact with others.^{5–9} Although it is difficult to prove that transmission is determined solely by overcrowded conditions, household overcrowding has been implicated in several diseases, such as tuberculosis¹⁰ and acute respiratory infections, including pneumonia.^{11,12} Infectious diseases that are associated with increased transmission in overcrowded households include SARS-CoV-2, influenza, tuberculosis, meningococcal disease, norovirus, and drug-resistant infections (ie, antimicrobial resistance [AMR]).^{8,11–18}

In addition to viral spread, household overcrowding, together with several social factors including poverty and

level of education, are associated with an increased carriage risk and increased spread of drug-resistant bacteria in high-income countries and low-income and middle-income countries (LMICs).¹³ Household overcrowding is associated with a higher incidence of tuberculosis^{18–20} and a higher carriage risk of resistant *Staphylococcus aureus*.²¹ Urbanisation in LMICs contributes to overcrowding and poor sanitation, which might also be linked to the increase and spread of AMR.^{22–25}

The world population is projected to increase to 9·7 billion people by 2050, with most living in LMICs, primarily in Africa, where countries in Africa comprise eight of the top ten countries that have an annual increase in population of between 2·1% and 3·6%. This population increase—without suitable growth in housing stock—will, in turn, increase overcrowding over time.^{26,27} Housing is a fundamental human right, as defined by the UN, which is essential to human

Lancet Planet Health 2022;
6: e670–81

Big Data Institute, Li Ka Shing Centre for Health Information and Discovery (M G Chipeta PhD, E P A Kumaran MSc, A J Browne PhD, G Haines-Woodhouse MRes, C E Moore PhD) and Centre for Tropical Medicine and Global Health (B H K Hamadani MPH, B Sartorius PhD, C Dolecek PhD), University of Oxford, Oxford, UK; African Institute for Development Policy, Lilongwe, Malawi (M G Chipeta); Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand (B H K Hamadani, B Sartorius, C Dolecek); Institute for Health Metrics and Evaluation (B Sartorius, R C Reiner Jr PhD, S I Hay DSc) and Department of Health Metrics Sciences, School of Medicine (R C Reiner Jr, S I Hay), University of Washington, Seattle, WA, USA; Centre for Neonatal and Paediatric Infection, St George's, University of London, London, UK (C E Moore)

Correspondence to:
Dr Catrin E Moore, Centre for Neonatal and Paediatric Infection, St George's, University of London, London, UK
camoore@sgul.ac.uk

Research in context

Evidence before this study

We conducted a literature search on household overcrowding in Africa, searching PubMed from January, 2000, to November, 2020. We searched for studies using the terms ("housing" OR "housing conditions" OR "crowding" OR "household overcrowding") AND ("infectious diseases") AND ("Africa" OR "low-income" OR "middle-income") in the title, with no language restrictions. We found several studies published since 2010 that addressed household overcrowding as a risk factor for infectious diseases and antimicrobial resistance, but there were only few studies, and these were for localised settings (countries and sub-national locations). Similarly, we identified several studies addressing the status of housing conditions in specific low-income and middle-income settings. We found no systematic assessments of household overcrowding across the African continent.

Added value of this study

In our study, we assessed continent-wide household overcrowding in Africa between the years 2000 and 2018. We estimated the proportions and absolute counts of people living in overcrowded conditions resulting from a rapid growth in population and urbanisation. We used data from countries and islands in Africa from 2000 to 2018 using a geospatial

modelling technique that incorporated multiple data sources, including surveys and population censuses. The high resolution maps and data for overcrowding estimates generated in our study could be a useful covariate for predicting several infectious diseases, providing information to inform interventions, including for SARS-CoV-2. To the best of our knowledge, household overcrowding has not been assessed at the continent-wide level over such a long duration. A shortage of information on housing conditions impedes interventions and development planning and implementation, as it prevents a full assessment of the threats presented by household overcrowding in low-income and middle-income countries.

Implications of all the available evidence

African countries are experiencing substantial and rapid population and urban growth. These changes can be viewed by policy makers as an opportunity for leveraging public health improvements that also affirm wider human development goals, for example, for effective policies for urban growth that support economic development and the eradication of poverty. The estimates provided in our study could help to direct investments towards the provision of adequate, safe, and affordable housing, and basic services for everyone (Sustainable Development Goal 11).

security, nutrition, and health. Housing is also an important social determinant of health; living in poor housing conditions has a profound negative effect on health and wellbeing.^{2,3} WHO defines healthy housing as the physical structure of a dwelling that supports, inter alia, human wellbeing, adequate sanitation, illumination, and sufficient space.^{4,28–30} Adequate space is necessary for maintaining clean indoor air, reducing disease transmission risk and noise pollution, and for meeting the occupant's privacy needs.^{31,32} Furthermore, good and adequate housing enables access to basic services, inclusive growth (ie, socioeconomic growth that is distributed fairly across society and creates opportunities for all), and sustainable development.^{31,32} This point is underscored by UN Sustainable Development Goal (SDG) 11, which aims to "ensure access for all to adequate, safe and affordable housing and basic services and upgrade slums by 2030".³

Many different metrics of household overcrowding have been suggested and applied.^{1,33,34} An overarching definition is any situation in which the number of individuals occupying a dwelling exceeds the capacity of the dwelling space (appendix p 2). This definition can be measured in terms of rooms, bedrooms, or floor area where the shortage of space can result in adverse physical and mental health outcomes.³⁵ A standard definition of household crowding, as used by the UN and WHO, is any situation in which more than two people occupy a sleeping space in a dwelling.⁴

When applied with other covariates, household overcrowding patterns can accurately predict and parameterise transmission models of infectious disease.^{36–38} Moreover, household overcrowding patterns can inform the design of targeted interventions for infectious diseases and poor living conditions, which are useful for policy making at all levels. However, there is currently a lack of detailed, fine-scale information for the patterns and levels of overcrowding that policy makers can act upon.

Current understanding of overcrowding in households across Africa is based on national data. By combining household survey data, population censuses, and other country-specific household surveys within a geostatistical framework, we aimed to provide household overcrowding estimates at finer spatial resolution (5×5 km). Household surveys contain a wealth of information and have previously been used as input data for statistical models (ie, geospatial models) to estimate health outcomes and indicators,^{39–44} especially in LMICs, where health registries are either incomplete or non-existent. Our spatiotemporal analysis focuses on household size and the number of sleeping rooms or spaces.

Methods

Overview

Geolocated data for the prevalence of household overcrowding from household and census datasets were synthesised by MGC and EPAK, then a two-stage Bayesian model-based geostatistical (MBG) framework was applied

See Online for appendix

to produce fine-scale-resolution (5×5 km) estimates of household overcrowding proportions between 2000 and 2018 across Africa. Our analysis covered 54 countries from both mainland Africa and islands for which survey data were available (ie, Cabo Verde, Comoros, Madagascar, and São Tomé & Príncipe). Analytical and model validation steps are described later in this Article and in the appendix (pp 31–32).

Data

Household overcrowding data were extracted from household surveys by authors MGC and EPAK, including Demographic Health Surveys (DHSs), Multiple Indicator Cluster Surveys, Integrated Public Use Microdata Series population censuses, and other country-specific surveys (appendix p 5). These surveys are regularly conducted in LMICs, use similar formats, and provide internationally comparable data collected over many years. We extracted data for the number of rooms or spaces for sleeping in each household and the number of people who slept in the house the night before the survey for the period between 2000 and 2018. These data were linked to the finest spatial resolution or location available: GPS cluster coordinates (latitudes and longitudes) or the smallest identifiable administrative area level, either level one (which were often states) or administrative level two areas (which were often districts). Administrative districts were resampled to point locations using population-weighted *k*-means clustering⁴¹ (appendix p 27).

Point data and resampled administrative level data were combined and used as input data for the MBG model. We created a binary outcome indicator, with household overcrowding defined as the ratio of individuals to sleeping rooms higher than two (appendix p 2). In total, we assembled 386 surveys with information on household crowding conditions. Of these, 287 surveys had extractable information covering 78 695 991 households from 54 African countries between 2000 and 2018.

Analysis and model validation

We fitted a two-stage Bayesian MBG model to estimate the proportion of household overcrowding in any given 5×5 km pixel and year to generate predictions for household overcrowding across the African continent at a high resolution. Our chosen methods provided the best out-of-sample predictive performance at the expense of an inferential understanding of the drivers of household overcrowding.

To leverage strength from locations and observations to the entire spatial and temporal domain, we selected socioeconomic and environmental covariates (appendix pp 28–30) and fit a stacked ensemble model to capture possible non-linear effects and complex interactions between covariates (informed by plausibility and importance in the model) and household overcrowding (appendix p 30).⁴⁵ To improve computational stability, account for differences in regional overcrowding

patterns, and to allow modelling and assessment of the effects of covariates, the model was fitted separately for each region. For each region, three sub-models were fitted to the dataset using the covariate data as explanatory predictors (appendix pp 28–29): generalised additive models, boosted regression trees, and lasso regression. Each sub-model was fitted using five-fold cross-validation to avoid overfitting, and the out-of-sample predictions from across the five holdouts were compiled into a single comprehensive set of predictions per model. In the second stage, the out-of-sample sub-model predictions were fed into the full geostatistical model as the explanatory covariates when conducting the model fit.

We generated pixel-level 95% uncertainty intervals (UIs) from 1000 draws that were created from the posterior distributions of modelled parameters. We then aggregated pixel-level estimates from the 1000 candidate maps to two sub-national administrative levels (states and districts), as well as national levels. We present results at pixel, district, state, and national level resolutions with the associated UIs. All final model map outputs were shaded grey for each map where the total population density was fewer than ten individuals per 1×1 km pixel. The national-level resolution maps provide a between-country comparison of overcrowding, and we assessed the within-country overcrowding variation by calculating the relative deviation in household overcrowding. This estimate was calculated by subtracting the national estimate from each district's estimate and dividing by the national estimate. A national level summary was produced, with comparisons for mean relative deviation for each country between 2000 and 2018 (appendix pp 28–32).

We estimated absolute population counts of people living in overcrowded conditions at pixel, district, state, and national levels on the basis of location-year specific population count raster data extracted from Gridded Population of the World (v4).⁴⁶ The estimated absolute population counts (including UIs) for people living in overcrowded environments were derived by multiplying the proportions of modelled household overcrowding by the population count data for a particular location-year at the draw level, and then by calculating a mean value and 95% CIs. This study complied with the Guidelines for Accurate and Transparent Health Estimates Reporting (appendix pp 38–39).⁴⁷

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Results

We observed spatiotemporal heterogeneity in household overcrowding estimates, both between and within countries in Africa, between 2000 and 2018. In 2018, the highest prevalence of overcrowding was observed in countries within the Horn of Africa, with a slight decrease




Figure 1: Box and whisker plots of household overcrowding comparisons across Africa by region
 Median range and IQR of household overcrowding in Africa by modelling regions for the year 2000 (shown in red) and 2018 (shown in blue).

observed over the 19-year period (regional median proportion 66% [UI 63–70] in 2000, and 62% [57–63] in 2018). The highest proportions were in Somalia (66% [57–75]), south Sudan (65% [56–74]), and Sudan (63% [53–71]). Although the ratio of overcrowded households in the central sub-Saharan Africa region (out-of-sample R^2 0.81, appendix pp 35–37) varied widely, it was much lower than in the Horn of Africa region, with a median of 27% [19–35]. The highest proportions were estimated for Zambia (57% [47–66]), Angola (43% [36–49]), and Democratic Republic of the Congo (42% [36–48]). The lowest estimated proportion in central sub-Saharan Africa was for Gabon (19% [14–26%]; figures 1, 2).

Moderate to high spatial variations were observed in the eastern sub-Saharan Africa region and the southern sub-Saharan Africa region (out-of-sample R^2 0.58, appendix pp 35–37); the largest differences in household overcrowding in a small spacial area were seen in Madagascar (62%, [54–70]), Kenya (41% [33–49]), and South Africa (15% [12–20]). The eastern sub-Saharan Africa region had a median proportion of 34% (29–39). In the north of Africa region, the highest overcrowding proportion was estimated in Libya at 63% (49–74), while overcrowding was much lower in Egypt (15% [7–27]) and Tunisia (22% [9–42]; figure 2). The regional median proportion doubled in the north of Africa region (out-of-sample R^2 0.78, appendix pp 35–37; 8% [6–9] in 2000 and 16% [14–19] in 2018; figure 1). The highest overcrowding proportions in the western sub-Saharan Africa region were estimated for Mauritania (60% [45–73]) and Chad (49% [39–59]). The western sub-Saharan Africa region had an estimated median proportion of 37% (32–43; figure 1).

Several countries had high overall household overcrowding proportions at the national level in 2018, and the Horn of Africa region consistently had high proportions throughout the study period (figures 1–3). A small number of countries had a reduction in overcrowding proportions (eg, Namibia had a reduction of up to 28.8%; table 1). Conversely, some countries had an increase in overcrowding proportions (ranging from 11.4% in Chad to 280% in Algeria; table 2). Most countries in the western, eastern, southern, and central sub-Saharan Africa regions remained relatively unchanged across the study period compared with other studied regions. Sub-nationally, at pixel level (5 × 5 km), household overcrowding had decreased in several countries (figure 3) although overcrowding proportions in these countries remain high.




Figure 2: The proportion of overcrowded households in low-income and middle-income countries within Africa, 2018
 Modelled estimates are shown by national-level aggregation (A), state (level 1) administrative divisions (B), district (level 2) administrative divisions (C), and 5 × 5 km pixels (D). Pixels (1 × 1 km resolution) with a total population density fewer than ten individuals per 1 × 1 km pixel are shown in grey.

	Household overcrowding estimates, 2000 (%)	Household overcrowding estimates, 2018 (%)	Percentage change (95% CI)
Central sub-Saharan Africa			
Angola	56%	43%	-23.2% (-46.9 to -21.0)
Namibia	28%	20%	-28.8% (-34.8 to -24.2)
Western sub-Saharan Africa			
Niger	54%	48%	-11.1% (-14.0 to -6.3)
Ghana	50%	38%	-24.0% (-28.6 to -19.0)
Horn of Africa			
Eritrea	68%	59%	-13.2% (-24.6 to -5.1)
Djibouti	54%	45%	-16.7% (-28.6 to -9.6)
Ethiopia	72%	55%	-23.6% (-27.3 to -19.5)

Modelled estimates at the national level with a relative reduction between 2000 and 2018 and 95% CI of the percentage change.

Table 1: Countries in Africa that had the largest decrease ($\geq 10\%$ change) in estimated household overcrowding between 2000 and 2018

	Household overcrowding estimates, 2000 (%)	Household overcrowding estimates, 2018 (%)	Percentage change (95% CI)
North of Africa			
Algeria	5%	19%	280.0% (200.0-291.0)
Tunisia	11%	22%	100% (50.0-120.0)
Eastern sub-Saharan Africa			
Rwanda	19%	30%	57.9% (42.9-63.0)
Burundi	28%	38%	35.7% (28.0-42.1)
Western sub-Saharan Africa			
Cameroon	25%	30%	20.0% (15.8-30.0)
Central African Republic	27%	32%	18.5% (10.0-22.0)
Chad	44%	49%	11.4% (8.3-13.0)

Modelled estimates at the national level with a relative increase between 2000 and 2018 and 95% CI of the percentage change.

Table 2: Countries in Africa that had the largest increase ($\geq 10\%$ change) in estimated household overcrowding between 2000 and 2018

Within-country variations were also observed. Notably, huge within-country variations (mean relative deviation of districts from the country value) were observed within the central, eastern, and southern sub-Saharan African regions. Namibia (35%) and Mozambique (29%) recorded the highest within-country differences in 2018, followed by Kenya (26%). Egypt (26%) and Nigeria (22%) measured the most significant variations in the north of Africa and western sub-Saharan Africa.

The countries that exhibited the least relative deviations from the national mean in 2018 included Sudan, Somalia, and Cabo Verde (all between 3% and 4%; figure 4). In Egypt, the mean relative deviation of household overcrowding decreased the most between 2000 and 2018, from 53% to 26%. Significant reductions were evident in Sierra Leone (25% to 7%), Central African Republic (18% to 4%), Algeria (17% to 4%), and Rwanda (18% to 8%). Household overcrowding inequality increased over the study period in many countries in sub-Saharan Africa; as shown by the rising mean relative deviation in Comoros (18% to 34%), Republic of the Congo (6% to 14%), Senegal (7% to 11%), and Malawi (12% to 16%; figure 4).

The absolute population counts living in overcrowded conditions increased over the 19-year study period, along with population increases for Africa in general. In 2018, a total of 474.4 million (95% UI 250.1-740.7 million) people lived in overcrowded conditions in Africa; an increase from 291.5 million (180.8 million-417.3 million) in 2000. 14 countries accounted for approximately 70% of the 2018 total population living in overcrowded conditions (Nigeria, Ethiopia, Democratic Republic of the Congo, Sudan, Uganda, Kenya, Tanzania, Madagascar, Egypt, Angola, Ghana, Mozambique, Cote d'Ivoire, and Niger; figure 5). Of these, more than 20 million people lived in




Figure 3: The change in proportion of overcrowded households within Africa from 2000 to 2018
 Pixels (1 × 1 km resolution) with a total population density fewer than ten individuals per 1 × 1 km pixel are shown in grey.

overcrowded housing conditions in each of the following six countries in sub-Saharan Africa in 2018: Nigeria, Ethiopia, Democratic Republic of the Congo, Sudan, Uganda, and Kenya (table 3). These six countries have a combined population of 538.3 million, which account for 48.5% of the total population (229.9 million people) who are living in overcrowded conditions in Africa.




Figure 4: Within-country variation in household overcrowding in 2000 and 2018

(A) Bars show the range in household overcrowding within each country. Grey bars represent estimates for the year 2000, and coloured bars represent estimates for 2018. Black dots represent the mean proportions for household overcrowding for each country. (B) Bars show the mean relative deviation of household overcrowding in administrative level 2 (districts) from the national level household overcrowding estimates. Grey bars represent estimates for the year 2000, and coloured bars represent estimates for 2018. The 2018 colours are based on the country's region, and countries are ordered (along the x-axis) on the basis of mean overcrowding proportions in the year 2018 (low to high). Countries are labelled using International Organization for Standardization (ISO) codes. AGO=Angola. BEN=Benin. BDI=Burundi. BFA=Burkina Faso. BWA=Botswana. CAF=Central African Republic. CIV=Cote d'Ivoire. CMR=Cameroun. COD=Congo (the Democratic Republic of the). COG=Congo. COM=Comoros. CPV=Cabo Verde. DJI=Djibouti. DZA=Algeria. EGY=Egypt. ERI=Eritrea. ESH=Western Sahara. ETH=Ethiopia. GAB=Gabon. GHA=Ghana. GIN=Guinea. GMB=Gambia. GNB=Guinea Bissau. GNQ=Equatorial Guinea. KEN=Kenya. LBR=Liberia. LBY=Libya. LSO=Lesotho. MAR=Morocco. MDG=Madagascar. MLI=Mali. MOZ=Mozambique. MRT=Mauritania. MWI=Malawi. NAM=Namibia. NER=Niger. NGA=Nigeria. RWA=Rwanda. STP=Sao Tome and Principe. TGO=Togo. SDN=Sudan. SEN=Senegal. SLE=Sierra Leone. SOM=Somalia. SSD=South Sudan. SWZ=Eswatini. TCD=Chad. TUN=Tunisia. TZA=Tanzania, the United Republic of. UGA=Uganda. ZAF=South Africa. ZMB=Zambia. ZWE=Zimbabwe.

For more on ISO codes see <https://www.iban.com/country-codes>

The five sub-national locations with the highest number of people living in overcrowded conditions in 2018 were: north and south Nigeria; southwest Democratic Republic of the Congo; north, southeast, and central Ethiopia; north Madagascar; east Sudan; and northeast Kenya.

Pixel-level data (5×5 km) from 2018 shows that the north and south Nigeria, central and north Ethiopia, central and south Uganda, central and southern parts of Madagascar, and central and southern parts of Malawi had the highest population of overcrowded people (figure 5).

Discussion

The objectives of this modelling study were to quantify the changes in household overcrowding across Africa to a fine spatial resolution over a 19-year period (2000 to 2018) using available country-specific household survey and census data. Given the housing crisis and population growth in Africa, together with the impacts of climate change (eg, droughts and flooding) and the rising spread of infectious diseases, we wanted to examine levels of household overcrowding and assess whether these data should become part of policy for local governments going forward. In our analysis, we found that household overcrowding was stable or decreasing over time in most regions in Africa, but that there was an increase in household overcrowding in north Africa based on the results in the included surveys. Although these surveys can differ between countries, the same questions are repeated over many years, indicating that these data are robust. High proportions of household overcrowding were localised in urban areas, with heterogeneity observed across and within countries. Our findings highlight important spatiotemporal disparities in household overcrowding and provided estimates and UIs for locations where data were sparse.

Housing is explicitly outlined in target 11 of the UN SDGs^{3,48} and is implicitly an important component of other SDGs. Policy makers view reductions in household overcrowding as a key prerequisite for inclusive growth, alongside efforts to reduce poverty, unemployment, and inequality. Estimates of household overcrowding modelled at 5×5 km resolution have potentially far-reaching applications in strategies and policies intended to ensure the achievement of the SDGs. Our study highlights countries and areas that could be targeted with interventions (eg, household interventions or specific water, sanitation, and hygiene interventions) to combat infectious disease, including the ongoing SARS-CoV-2 pandemic across Africa.

In our analysis, a marked increase in household overcrowding proportions was observed in north Africa, particularly in Algeria, which had a 280% increase over the 19-year study period. One of the reasons for the increase might be increasing urbanisation, with 78% of the population living in urban areas,⁴⁹ together with a lack of planning and rapid city growth, which has led to extremely high housing deficits. Between 2000 and 2015, the annual urbanisation growth rate average was 2·8% in Algeria, with a housing deficit backlog of 1·2 million households.⁵⁰

Nine of the 54 countries analysed in our study had considerable reductions in household overcrowding proportions (between 9% and 29%). The largest decrease in household overcrowding proportions was observed in Ethiopia (table 1). Until 2005, Ethiopia was one of the least urbanised countries on the African continent. This position has since changed rapidly. Ethiopia now has one of the fastest growing economies, with an average annual




Figure 5: Population counts of people living in overcrowded conditions in Africa, 2018

Modelled estimates are shown by national level aggregation (A), state (level 1) administrative divisions (B), district (level 2) administrative divisions (C), and 5×5 km pixels (D). Pixels (1×1 km resolution) with a total population density fewer than ten individuals per 1×1 km pixel are shown in grey.

growth of 10·5% (measured between 2004 and 2018). In 2018, the urban population of Ethiopia was estimated to account for 21·2% of its 112 million people, and its urbanisation rate stood at 4·9%. From 2006 onward, the Ethiopian Government invested heavily in social housing projects, attracting foreign and local building contractors, promoting integration, and reducing household overcrowding. Ethiopia's Government integrated housing development programme condominium scheme is the most successful local housing programme,⁵¹ at least 400 000 condominiums have been built against a target of 580 000 units over a 5 year period.^{51,52} Despite this progress, a backlog of approximately 1·2 million housing units exists, with a projected demand of 655 800 housing units needed between 2015 and 2025.⁵¹

	Absolute counts			Proportions (% UI)
	Population in 2018	Population living in crowded conditions in urban areas	Population living in crowded conditions	
Nigeria	202647467	33964499	65316345 (33184819-104960938)	32.2% (16.4-51.8)
Ethiopia	98423665	11264486	53640411 (32469598-73608502)	54.5% (33.0-74.8)
Democratic Republic of the Congo	106226798	20304631	44140503 (24744843-65560635)	41.6% (23.3-61.7)
Sudan	40415598	8884130	25383229 (14378490-34298159)	62.8% (35.6-84.9)
Uganda	39362904	5473563	21052164 (11665350-29875815)	53.5% (29.6-75.9)
Kenya	51227095	5720113	20428974 (12160791-29547393)	39.9% (23.7-57.7)
United Republic of Tanzania	52806810	6739765	18215581 (10659356-26990652)	34.5% (20.2-51.1)
Madagascar	25374824	6013238	15418558 (9654250-20385703)	60.8% (38.1-80.3)
Egypt	90886136	5722960	13309209 (2530504-36212272)	14.6% (2.8-39.8)
Angola	30278009	8540180	12746537 (7501606-18430408)	42.1% (24.8-60.9)
Ghana	30660398	6635358	11640979 (6253018-17850364)	38.0% (20.4-58.2)
Mozambique	28208179	4388076	11547569 (7308287-16114246)	41.0% (25.9-57.1)
Cote d'Ivoire	24927552	5513889	10811548 (5164149-17047356)	43.4% (20.7-68.4)
Niger	21557506	1752480	10308704 (5267144-15526592)	47.8% (24.4-72.0)
Zambia	17295807	4413724	9808276 (5386556-13791360)	56.7% (31.1-79.7)
Morocco	34451860	6073526	9489884 (3370343-18238191)	27.6% (9.8-52.9)
South Sudan	14388976	2349261	9397044 (5587479-12377392)	65.3% (38.8-86.0)
South Africa	57664805	6033202	9004779 (4166378-16148805)	15.6% (7.2-28.0)
Mali	21153774	3749992	8522710 (4053278-13631197)	40.3% (19.2-64.4)
Algeria	41298619	5759785	7890116 (1059259-22990741)	19.1% (2.6-55.7)
Cameroon	26217798	4384206	7828940 (3222513-14086387)	29.9% (12.3-53.7)
Chad	15236208	1706482	7419488 (3894663-10988639)	48.7% (25.6-72.1)
Somalia	10302581	3193124	6793880 (4082442-8902759)	66.0% (39.6-86.4)
Malawi	17278582	1214981	6749896 (4213903-9557598)	39.1% (24.4-55.3)
Burkina Faso	21124265	2032739	6557222 (3440139-10426965)	31.0% (16.3-49.4)
Senegal	14315438	2648738	5405588 (2650556-8616317)	37.8% (18.5-60.2)
Zimbabwe	14001902	1800123	4737167 (2492124-7467873)	33.8% (17.8-53.3)
Benin	11598872	2072793	4318319 (2103366-6938784)	37.2% (18.1-59.8)
Libya	6489464	3132790	4016397 (2241637-5443987)	61.9% (34.5-83.9)
Burundi	10645769	561779	4012704 (2207582-6072299)	37.7% (20.7-57.0)
Guinea	11225360	1516844	3889343 (1575704-6852329)	34.7% (14.0-61.0)
Rwanda	12744898	684120	3800669 (2014509-6054304)	29.8% (15.8-47.5)
Togo	7813533	1259639	2929392 (1556752-4527726)	37.5% (19.9-58.0)
Tunisia	10971254	1690079	2414398 (400031-6366743)	22.0% (3.7-58.0)
Eritrea	4046296	1503944	2387213 (1337935-3309249)	59.0% (33.1-81.8)
Mauritania	4062871	1304511	2288615 (1244084-3179340)	56.3% (30.6-78.3)
Sierra Leone	6330537	947786	2204153 (964819-3730522)	34.8% (15.2-58.9)
Liberia	4174870	965649	1821979 (940839-2768940)	43.6% (22.5-66.3)
Central African Republic	5088636	701933	1632402 (639560-2948472)	32.1% (12.6-57.9)
Congo	3654201	695841	994059 (494755-1650515)	27.2% (13.5-45.2)
Lesotho	1831475	175015	564565 (287490-919886)	30.8% (15.7-50.2)
Gambia	2000268	325558	551794 (221268-999225)	27.6% (11.1-50.0)
Botswana	2302885	387356	530625 (232917-951986)	23.0% (10.1-41.3)
Gabon	2640378	439256	504892 (225794-914299)	19.1% (8.6-34.6)
Namibia	2353714	258068	469214 (224395-814837)	19.9% (9.5-34.6)
Djibouti	1046353	350906	444185 (212648-698895)	42.5% (20.3-66.8)
Swaziland	1077903	71357	237855 (110093-421245)	22.1% (10.2-39.1)
Equatorial Guinea	1160002	164146	224857 (93731-422955)	19.4% (8.1-36.5)
Cape Verde	475345	150916	221935 (90719-358781)	46.7% (19.1-75.5)
Comoros	716948	53039	182892 (81141-316636)	25.5% (11.3-44.2)

(Table 3 continues on next page)

	Absolute counts			Proportions (% UI)
	Population in 2018	Population living in crowded conditions in urban areas	Population living in crowded conditions	
(Continued from previous page)				
Guinea-Bissau	1628720	70486	156635 (48717–356027)	9.6% (2.9–21.9)
Sao Tome and Principe	174326	37754	51019 (25581–83653)	29.3% (14.7–47.9)
Western Sahara	448814	5	12 (5–21)	0
Absolute population counts and proportions at the national level, with 95% UI in absolute terms. UI=uncertainty interval.				
Table 3: Absolute population counts and proportions of people living in overcrowded conditions in 2018				

Other exemplar countries include Angola, Namibia, and Zambia. In Angola, the government has established the housing promotion fund (HPF) to promote access to affordable housing. The HPF is responsible for 70% of state-built housing, and the government has encouraged private developers to contribute to the national housing market for mass-scale housing production.⁵¹ A deficit of 1.7 million housing units exists in Angola. Similarly, although Zambian and Namibian national governments have invested considerably in housing, the housing deficits remain high. Overall, there is a deficit of 68.8 million housing units for people living in overcrowded conditions in Africa.

Improvements in housing conditions in Africa were observed in dwellings (based on sufficient living areas, improved water and sanitation, and durable construction), rising from 11% in 2000 to 23% in 2015.⁵³ However, high levels of household overcrowding persist, with Nigeria, Ethiopia, Democratic Republic of the Congo, Sudan, Uganda, Kenya, and Tanzania showing the highest absolute population living in overcrowded conditions in 2018. No material changes in household overcrowding proportions were observed throughout eastern, western, and central sub-Saharan Africa over the 19-year study period; these areas remain overcrowded (above a median of 27%).

Africa is the world's least urbanised continent, and has 11.3% of the world's urban population. The sub-Saharan African region is the least urbanised region; however, the region's cities are rapidly expanding.⁵⁴ The African population has been growing, on average, 2.5% per year between 2000 and 2018. Along with this population growth and urbanisation, the absolute population living in overcrowded conditions has increased drastically, with small increases in housing. Additionally, most migrants from rural areas are uneducated or unskilled and end up in the informal sector in Africa, accounting for 93% of all new jobs and 61% of all urban employment.⁵⁵ Because of the low and intermittent incomes generated by the informal sector, most migrants become slum dwellers or seek housing with slum landlords. Many governments in Africa have directly provided housing to meet the needs of growing urban populations,^{51,56} but these programmes are expensive, difficult for the urban poor to afford, and have yet to result in considerable increases in affordable

housing. As a result, the population living in overcrowded conditions increased by 62.7%, with most of these populations living in urban slums.

Urban growth rates are high on the African continent. Projections predict that Africa's cities will be home to an additional 950 million people by 2050.⁵⁰ Most of this growth will likely occur in small-sized and medium-sized towns and cities, which often struggle to attract the required infrastructure investment.⁴⁹ Without substantial investment in housing facilities and services, overcrowding and slum proliferation could increase exponentially around cities. Without essential infrastructure, large populations live in unhealthy, polluted households in congested and poor conditions with poor sanitation, insecure living conditions, and little access to utilities such as electricity and water (Frostd J J, unpublished).

Overcrowded conditions provide the perfect environment for the transmission of epidemics such as SARS-CoV-2 and other infectious diseases. Household overcrowding is positively correlated with the spread of SARS-CoV-2.^{5,8} Globally, governments have implemented non-pharmaceutical interventions to reduce the spread of SARS-CoV-2, which have included partial or total lockdowns of entire regions or countries. These measures have meant that people spend more time at home, many of whom, especially in Africa, are in overcrowded conditions.^{57,58} Living in overcrowded housing conditions makes it harder to self-isolate and shield from SARS-CoV-2. There is a correlation between transmissibility and infection outcomes with SARS-CoV-2 for those living in overcrowded spaces.^{5,8,59–62} In England, it is estimated that there were at least 70% more cases of SARS-CoV-2 in households that were overcrowded compared with non-overcrowded households.⁷ The unavoidable proximity and use of shared facilities that are inevitable with overcrowding are ideal for the spread of an airborne microbe and can increase the speed and spread of pandemics.

Our model is useful for understanding spatiotemporal differences in household overcrowding, and could also be useful for examining AMR and the spread of SARS-CoV-2 in order to shape policy across Africa. Our high-resolution overcrowding estimates could be used to improve forecasting of infectious diseases, potentially providing valuable information to assist with disease

control interventions. For example, our method could be used immediately to model household overcrowding with SARS-CoV-2 and other infectious diseases, such as *Mycobacterium tuberculosis*, as well as AMR, in addition to other research in the area.^{13,18,23,59,63–66} In a systematic review,⁶⁷ overcrowded conditions were a strong enabler of AMR in Chilean hospitals between 2008 and 2017. Our model, therefore, can be used to determine areas of high overcrowding and high transmission, and help to target interventions to reduce the transmission of infectious diseases.⁶⁸

The main limitations of our study were that we provide estimates in locations without empirical data, and that our data were derived from census data and other country-specific surveys. While these are currently the only sources of representative and comparable data, they contain multiple potential biases, such as recall and reporting bias, interviewer effects on responses, and refusal bias.^{69–72} In addition, there were data gaps in some locations (appendix p 4) that affected the robustness of our estimates, and increased uncertainty. Our resampling method increased the uncertainty intervals in our analysis. Given that our data maximised predictive performance without providing evidence of causality, additional groundwork is needed to provide this information. There are also possibly new covariates that could have an effect on our outcomes. The ideal next steps would be to conduct additional work to examine the truth of the DHS surveys in small areas, which would enhance the data already available, as well as to conduct new surveys where no data currently exists.

Additionally, our analysis used proxy information to infer urbanicity. As such, further analyses that delineate urban and rural splits would be ideal. There is an urgent need to establish and promote household surveillance programmes to accurately estimate overcrowding conditions across Africa. Household overcrowding definitions need to be harmonised (appendix p 2). We applied the definition of overcrowding as having more than two people per sleeping room. Household overcrowding is a global concern, especially in countries with humanitarian crises. The modelling of household overcrowding needs to be expanded beyond Africa. Despite these limitations, to the best of our knowledge, our study provides the first Africa-wide estimates for household overcrowding using geospatial tools, and highlights the need for interventions to increase housing infrastructure across Africa.

Our data show that 38% of the 1.26 billion people in Africa live in overcrowded households. We also provide the first detailed baseline metrics for overcrowding that build on existing measurements of housing in Africa. These results were limited to urban areas that are not currently standardised at the sub-national scale and were derived from simplistic extrapolations from survey data. Using survey data, we also provide estimates of household overcrowding at fine spatial resolutions across the

entire African continent. We found that the number of people living in overcrowded conditions increased from 291.5 million in 2000 to 474.4 in 2018, representing a 62.7% increase. 14 countries accounted for nearly 70% of overcrowding in 2018 and nearly half of people living in overcrowded conditions resided in six countries. We also describe how the proportions and patterns of household overcrowding differ between and within countries and how these data have changed over the 19-year study period. The increases were likely to have been driven by booming urbanisation and population growth in Africa. Our work should be leveraged as a major opportunity to improve public health in Africa alongside broader human development goals. Understanding where a lack of adequate living space exists is also essential for monitoring the transmission of infectious diseases, such as SARS-CoV-2, as well as the spread of AMR, and could provide priority areas for benchmarking and introducing interventions for achieving SDG 11 in Africa.

Contributors

MGC and CEM were responsible for conceptualising the study. MGC, EPAK, BHKH, and GH-W were responsible for extracting data for household overcrowding, and MGC fitted household overcrowding models with input from AJB, BS, RCR, and SIH. MGC prepared the first draft of the manuscript. CD, BS, SIH, and CEM provided a guiding role throughout the analysis and throughout the process of drafting the manuscript. All authors discussed and reviewed the results. All authors reviewed and revised the draft manuscript, and all authors approved the final version for publication. CEM and MGC had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Declaration of interests

All authors declare no competing interests.

Data sharing

A detailed table of data sources is provided in the appendix (pp 5–27). All code used for these analyses is publicly available online at <http://ghdx.healthdata.org/>.

Acknowledgments

This work was funded by a grant from the Fleming Fund of the UK Department of Health of Social care, the Wellcome Trust (209142/Z/17/Z), and the Bill & Melinda Gates Foundation (OPP1176062). We would like to thank Barney McManigal and Joseph Frostad for proofreading the manuscript, Kirsten Wiens for providing guidance on the relative deviations calculations, and Lucas Earl for producing the map figures for household overcrowding. We also gratefully acknowledge the work of the DHS Program (<https://dhsprogram.com/>), the UNICEF programme (<https://mics.unicef.org/>), and IPUMS (<https://www.ipums.org>).

References

- Melki IS, Beydoun HA, Khogali M, Tamim H, Yunis KA. Household crowding index: a correlate of socioeconomic status and inter-pregnancy spacing in an urban setting. *J Epidemiol Community Health* 2004; **58**: 476–80.
- UN. Progress towards the Sustainable Development Goals. Report of the Secretary-General, 2016. <https://digitallibrary.un.org/record/833184> (accessed July 28, 2021).
- UN. The United Nations Conference on Housing and Sustainable Urban Development Habitat III, 2016. <https://www.un.org/en/conferences/habitat/quito2016> (accessed July 28, 2021).
- WHO. WHO Housing and Health Guidelines. 2018. <https://www.who.int/publications/i/item/9789241550376> (accessed Aug 2, 2021).
- Rader B, Scarpino SV, Nande A, et al. Crowding and the shape of COVID-19 epidemics. *Nat Med* 2020; **26**: 1829–34.

- 6 Nishiura H, Oshitani H, Kobayashi T, et al. Closed environments facilitate secondary transmission of coronavirus disease 2019. *Trans Japan Acad* 2020; **75**: 27.
- 7 Adlakha D, Sallis JF. Why urban density is good for health—even during a pandemic. 2020. <https://theconversation.com/why-urban-density-is-good-for-health-even-during-a-pandemic-142108> (accessed July 10, 2020).
- 8 Zimmermann R, Sarma N, Thieme-Thörel D, et al. COVID-19 outbreaks in settings with precarious housing conditions in Germany: challenges and lessons learned. *Front Public Health* 2021; **9**: 708694.
- 9 Mossong J, Hens N, Jit M, et al. Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med* 2008; **5**: e74.
- 10 Harling G, Castro MC. A spatial analysis of social and economic determinants of tuberculosis in Brazil. *Health Place* 2014; **25**: 56–67.
- 11 Fonseca Lima EJ, Mello MJ, Albuquerque MF, et al. Risk factors for community-acquired pneumonia in children under five years of age in the post-pneumococcal conjugate vaccine era in Brazil: a case control study. *BMC Pediatr* 2016; **16**: 157.
- 12 Kristensen IA, Olsen J. Determinants of acute respiratory infections in Soweto—a population-based birth cohort. *S Afr Med J* 2006; **96**: 633–40.
- 13 Alivizda V, Mariano V, Ahmad R, et al. Investigating the impact of poverty on colonization and infection with drug-resistant organisms in humans: a systematic review. *Infect Dis Poverty* 2018; **7**: 76.
- 14 Baker RE, Mahmud AS, Miller IF, et al. Infectious disease in an era of global change. *Nat Rev Microbiol* 2021; **20**: 193–205.
- 15 Jaine R, Baker M, Venugopal K. Epidemiology of acute rheumatic fever in New Zealand 1996–2005. *J Paediatr Child Health* 2008; **44**: 564–71.
- 16 Jaine R, Baker M, Venugopal K. Acute rheumatic fever associated with household crowding in a developed country. *Pediatr Infect Dis J* 2011; **30**: 315–19.
- 17 Norheim G, Sadarangani M, Omar O, et al. Association between population prevalence of smoking and incidence of meningococcal disease in Norway, Sweden, Denmark and the Netherlands between 1975 and 2009: a population-based time series analysis. *BMJ Open* 2014; **4**: e003312.
- 18 Duarte R, Lönnroth K, Carvalho C, et al. Tuberculosis, social determinants and co-morbidities (including HIV). *Pulmonology* 2018; **24**: 115–19.
- 19 Honorio F, et al. Social determinants in household contacts and index cases of tuberculosis in Lima, Peru. *J Med Dent Sci Res* 2020; **7**: 21–29.
- 20 Baker M, Das D, Venugopal K, Howden-Chapman P. Tuberculosis associated with household crowding in a developed country. *J Epidemiol Community Health* 2008; **62**: 715–21.
- 21 Lestari ES, Duerink DO, Hadi U, et al. Determinants of carriage of resistant *Staphylococcus aureus* among *S. aureus* carriers in the Indonesian population inside and outside hospitals. *Trop Med Int Health* 2010; **15**: 1235–43.
- 22 Allcock S, Young EH, Holmes M, et al. Antimicrobial resistance in human populations: challenges and opportunities. *Glob Health Epidemiol Genom* 2017; **2**: e4.
- 23 Nellums LB, Friedland JS, Hargreaves S. Antimicrobial resistance among migrants in Europe—Authors' reply. *Lancet Infect Dis* 2018; **18**: 945.
- 24 Tosas Auguste O, Betley JR, Stabler RA, et al. Evidence for community transmission of community-associated but not health-care-associated methicillin-resistant *Staphylococcus Aureus* strains linked to social and material deprivation: spatial analysis of cross-sectional data. *PLoS Med* 2016; **13**: e1001944.
- 25 Connolly MA, Gayer M, Ryan MJ, Salama P, Spiegel P, Heymann DL. Communicable diseases in complex emergencies: impact and challenges. *Lancet* 2004; **364**: 1974–83.
- 26 Vollset SE, Goren E, Yuan CW, et al. Fertility, mortality, migration, and population scenarios for 195 countries and territories from 2017 to 2100: a forecasting analysis for the Global Burden of Disease Study. *Lancet* 2020; **396**: 1285–306.
- 27 Pison G. The population of the world (2011). *Popul Soc (Paris)* 2011; **480**: 2–8.
- 28 Pepin C, Muckle G, Moisan C, Forget-Dubois N, Riva M. Household overcrowding and psychological distress among Nunavik Inuit adolescents: a longitudinal study. *Int J Circumpolar Health* 2018; **77**: 1541395.
- 29 Nkosi V, Haman T, Naicker N, Mathee A. Overcrowding and health in two impoverished suburbs of Johannesburg, South Africa. *BMC Public Health* 2019; **19**: 1358.
- 30 Waters A-M. Do housing conditions impact on health inequalities between Australia's rich and poor? 2001. <https://www.ahuri.edu.au/research/final-reports/4> (accessed Aug 4, 2021).
- 31 Braubach M. Key challenges of housing and health from WHO perspective. *Int J Public Health* 2011; **56**: 579–80.
- 32 Kubba S. Indoor environmental quality. In: Handbook of Green Building Design and Construction. 2nd edn. Boca Raton, FL, USA: Butterworth-Heinemann, 2017: 353–412.
- 33 Blake KS, Kellerson RL, Simic A. Measuring overcrowding in housing. 2007. https://www.huduser.gov/publications/pdf/measuring_overcrowding_in_hsg.pdf (accessed Aug 10, 2021).
- 34 Ferraro CF, Trotter CL, Nascimento MC, et al. Household crowding, social mixing patterns and respiratory symptoms in seven countries of the African meningitis belt. *PLoS One* 2014; **9**: e101129.
- 35 Evans GW. The built environment and mental health. *J Urban Health* 2003; **80**: 536–55.
- 36 Edmunds WJ, Kafatos G, Wallinga J, Mossong JR. Mixing patterns and the spread of close-contact infectious diseases. *Emerg Themes Epidemiol* 2006; **3**: 10.
- 37 Wallinga J, Teunis P, Kretzschmar M. Using data on social contacts to estimate age-specific transmission parameters for respiratory-spread infectious agents. *Am J Epidemiol* 2006; **164**: 936–44.
- 38 Ogunjimi B, Hens N, Goeyvaerts N, Aerts M, Van Damme P, Beutels P. Using empirical social contact data to model person to person infectious disease transmission: an illustration for varicella. *Math Biosci* 2009; **218**: 80–87.
- 39 Graetz N, Friedman J, Osgood-Zimmerman A, et al. Mapping local variation in educational attainment across Africa. *Nature* 2018; **555**: 48–53.
- 40 Osgood-Zimmerman A, Milliar AI, Stubbs RW, et al. Mapping child growth failure in Africa between 2000 and 2015. *Nature* 2018; **555**: 41–47.
- 41 Golding N, Burstein R, Longbottom J, et al. Mapping under-5 and neonatal mortality in Africa, 2000–15: a baseline analysis for the Sustainable Development Goals. *Lancet* 2017; **390**: 2171–82.
- 42 Wiens KE, Lindstedt PA, Blacker BF, et al. Mapping geographical inequalities in oral rehydration therapy coverage in low-income and middle-income countries, 2000–17. *Lancet Glob Health* 2020; **8**: e1038–60.
- 43 Utazi CE, Thorley J, Alegana VA, et al. High resolution age-structured mapping of childhood vaccination coverage in low and middle income countries. *Vaccine* 2018; **36**: 1583–91.
- 44 Utazi CE, Thorley J, Alegana VA, et al. Mapping vaccination coverage to explore the effects of delivery mechanisms and inform vaccination strategies. *Nat Commun* 2019; **10**: 1633.
- 45 Bhatt S, Cameron E, Flaxman SR, Weiss DJ, Smith DL, Gething PW. Improved prediction accuracy for disease risk mapping using Gaussian process stacked generalization. *J R Soc Interface* 2017; **14**: 14.
- 46 University Center for international Earth Science information Network, Columbia University. Gridded Population of the World, Version 4 (GPWv4): Population Count Adjusted to Match 2015 Revision of UN WPP Country Totals. NASA Socioeconomic Data and Applications Center. SEDAC, 2016. <https://sedac.ciesin.columbia.edu/data/collection/gpw-v4> (accessed Aug 20, 2021).
- 47 Stevens GA, Alkema L, Black RE, et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *Lancet* 2016; **388**: e19–23.
- 48 UN-HABITAT. Sustainable development goals: monitoring human settlements indicators. 2020. <https://unhabitat.org/about-us/sustainable-development-goals> (accessed Oct 14, 2021).
- 49 Organisation for Economic Co-operation and Development. Africa's urbanisation dynamics 2020: Africapolis, Mapping a new urban geography. 2020. <https://www.oecd.org/development/africa-urbanisation-dynamics-2020-b6bccb81-en.htm> (accessed March 9, 2021).
- 50 Bah EHM, Faye I, Geh ZF. Housing market dynamics in Africa. London, UK: Palgrave Macmillan, 2018.
- 51 Centre for Affordable Housing Finance in Africa. 2020 yearbook: housing finance in Africa, 2020. <https://housingfinanceafrica.org/documents/2020-housing-finance-in-africa-yearbook-11th-edition/> (accessed Sept 1, 2021).

- 52 Kaysay D. Ethiopia: can mortgage banking help address Ethiopia's rapidly growing demand for housing. 2020. <https://allafrica.com/stories/202007250109.html> (accessed Sept 1, 2021).
- 53 Tusting LS, Gething PW, Gibson HS, et al. Housing and child health in sub-Saharan Africa: a cross-sectional analysis. *PLoS Med* 2020; **17**: e1003055.
- 54 Bank TW. Stocktaking of the housing sector in sub-Saharan Africa: challenges and opportunities. 2015. <https://openknowledge.worldbank.org/handle/10986/23358> (accessed Oct 14, 2021).
- 55 UNICEF. Analysis of multiple deprivations in secondary cities in sub-Saharan Africa. UNICEF and UN HABITAT. 2020. <https://www.unicef.org/esa/reports/analysis-multiple-deprivations-secondary-cities-sub-saharan-africa> (accessed Oct 21, 2021).
- 56 UN-HABITAT. Affordable land and housing in Africa. 2011. <https://unhabitat.org/sites/default/files/download-manager-files/Affordable%20Land%20and%20Housing%20in%20Africa.pdf> (accessed Oct 14, 2021).
- 57 Tinson A, Clair A. Better housing is crucial for our health and the COVID-19 recovery. 2020. <https://www.health.org.uk/publications/long-reads/better-housing-is-crucial-for-our-health-and-the-covid-19-recovery> (accessed Aug 2, 2021).
- 58 Thorstensen-Woll C, Buck D, Naylor C. Homes, health and COVID-19: how poor-quality homes have contributed to the pandemic. 2020. https://www.smf.co.uk/commentary_podcasts/homes-health-and-covid-19-how-poor-housing-adds-to-the-hardship-of-the-coronavirus-crisis/ (accessed Aug 2, 2021).
- 59 Ahmad K, Erqou S, Shah N, et al. Association of poor housing conditions with COVID-19 incidence and mortality across US counties. *PLoS One* 2020; **15**: e0241327.
- 60 Abrams EM, Szeffler SJ. COVID-19 and the impact of social determinants of health. *Lancet Respir Med* 2020; **8**: 659–61.
- 61 Morawska L, Tang JW, Bahnfleth W, et al. How can airborne transmission of COVID-19 indoors be minimised? *Environ Int* 2020; **142**: 105832.
- 62 Somsen GA, van Rijn C, Kooij S, Bem RA, Bonn D. Small droplet aerosols in poorly ventilated spaces and SARS-CoV-2 transmission. *Lancet Respir Med* 2020; **8**: 658–59.
- 63 Kenway P, Butt JA. A review of the report by Public Health England into disparities in risks and outcomes of COVID-19 between ethnic groups and by level of deprivation. 2020. <https://www.gov.uk/government/publications/covid-19-review-of-disparities-in-risks-and-outcomes> (accessed Aug 2, 2021).
- 64 Chen Q, Toorop MMA, de Boer MGJ, Rosendaal FR, Lijfering WM. Why crowding matters in the time of COVID-19 pandemic? — a lesson from the carnival effect on the 2017/2018 influenza epidemic in the Netherlands. *BMC Public Health* 2020; **20**: 1516.
- 65 von Seidlein L, Alabaster G, Deen J, Knudsen J. Crowding has consequences: prevention and management of COVID-19 in informal urban settlements. *Build Environ* 2021; **188**: 107472.
- 66 Kirenga BJ, Ssengooba W, Muwonge C, et al. Tuberculosis risk factors among tuberculosis patients in Kampala, Uganda: implications for tuberculosis control. *BMC Public Health* 2015; **15**: 13.
- 67 Alle K, García P, Labarca J, Munita JM, Rendic M, Undurraga EA. Socioeconomic factors associated with antimicrobial resistance of *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Escherichia coli* in Chilean hospitals (2008–2017). *Rev Panam Salud Publica* 2020; **44**: e30.
- 68 Tusting LS, Gething PW, Gibson HS, et al. Housing and child health in sub-Saharan Africa: a cross-sectional analysis. *PLoS Med* 2020; **17**: e1003055.
- 69 Shea RS. Potential bias and selectivity in analyses of children born in the past five years using DHS data. Rockville, Maryland. 2014. <https://dhsprogram.com/publications/publication-mmr14-methodological-reports.cfm> (accessed Oct 14, 2021).
- 70 Leone T, Sochas L, Coast E. Depends who's asking: interviewer effects in demographic and health surveys abortion data. *Demography* 2021; **58**: 31–50.
- 71 Adegboye OA, Fujii T, Leung DH. Refusal bias in HIV data from the Demographic and Health Surveys: evaluation, critique and recommendations. *Stat Methods Med Res* 2020; **29**: 811–26.
- 72 Finaret AB, Hutchinson M. Missingness of height data from the demographic and health surveys in Africa between 1991 and 2016 was not random but is unlikely to have major implications for biases in estimating stunting prevalence or the determinants of child height. *J Nutr* 2018; **148**: 781–89.