

Antimagic Labeling of Some Degree Splitting Graphs

Chirag Barasara*
Palak Prajapati†

Abstract

A graph with q edges is called *antimagic* if its edges can be labeled with $1, 2, 3, \dots, q$ without repetition such that the sums of the labels of the edges incident to each vertex are distinct. As Wang et al. [2012], proved that not all graphs are antimagic, it is interesting to investigate antimagic labeling of graph families. In this paper we discussed antimagic labeling of the larger graphs obtained using degree splitting operation on some known antimagic graphs. As discussed in Krishnaa [2016], antimagic labeling has many applications, our results will be used to expand the network on larger graphs.

Keywords: Antimagic labeling, Antimagic graph, Graph operation, Degree splitting graph.

2020 AMS subject classifications: 05C78, 05C76. ¹

*Department of Mathematics, Hemchandracharya North Gujarat University, Patan - 384265, Gujarat, INDIA; chirag.barasara@gmail.com.

†Research Scholar, Department of Mathematics, Hemchandracharya North Gujarat University, Patan - 384265, Gujarat, INDIA; palakprajapati733@gmail.com.

¹Received on June 20, 2023. Accepted on August 16, 2023. Published on September 3, 2023. DOI: 10.23755/rm.v41i0.1253. ISSN: 1592-7415. eISSN: 2282-8214. ©The Authors. This paper is published under the CC-BY licence agreement.

1 Introduction

We begin with simple, finite, connected and undirected graph $G = (V(G), E(G))$. For all standard terminologies and notations we follow Clark and Holton [1969]. We will give brief summary of definitions which are useful for the present investigations.

Definition 1.1. A graph labeling is an assignment of integers to the vertices or edges or both subject to certain condition(s). If the domain of the mapping is the set of vertices (or edges) then the labeling is called a vertex labeling (or edge labeling).

For an extensive survey on graph labeling and bibliographic references we refer to Gallian [2022].

Hartsfield and Ringel [1990] have introduced antimagic labeling as follows.

Definition 1.2. A graph with q edges is called antimagic if its edges can be labeled with $1, 2, 3, \dots, q$ without repetition such that the sums of the labels of the edges incident to each vertex are distinct.

They have also discussed antimagic labeling for paths, cycles, wheels and complete graph. Wang [2005] has shown that the toroidal grids $C_{n_1} \times C_{n_2} \times \dots \times C_{n_k}$ are antimagic. Cheng [2008] has proved that all Cartesian products of two or more regular graphs are antimagic.

Vaidya and Vyas [2012, 2013] have discuss antimagic labeling of some path and cycle related graphs as well as graph obtained using switching of a vertex. Alon et al. [2004] have obtained some conditions on degree of a vertices for graph to be antimagic.

Krishnaa [2016] has illustrated some applications of antimagic labeling.

Definition 1.3. Let $G = (V(G), E(G))$ be a graph with $V = S_1 \cup S_2 \cup S_3 \cup \dots \cup S_t \cup T$, where each S_i is a set of vertices having same degree (at least two vertices) and $T = V \setminus \cup_{i=1}^t S_i$. The degree splitting graph of G denoted by $DS(G)$ is obtained from the graph G by adding vertices $w_1, w_2, w_3, \dots, w_t$ and joining w_i to each vertex of S_i for $1 \leq i \leq t$.

Definition 1.4. The Cartesian product of graphs G_1 and G_2 denoted by $G_1 \square G_2$ is the graph with vertex set $V(G_1) \times V(G_2) = \{(u, v) / u \in V(G_1) \text{ and } v \in V(G_2)\}$ and (u, v) is adjacent to (u', v') if and only if either $u = u'$ and $vv' \in E(G_2)$ or $v = v'$ and $uu' \in E(G_1)$.

Definition 1.5. The ladder graph L_n is defined as $L_n = P_n \square K_2$.

Antimagic Labeling of Some Degree Splitting Graphs

Definition 1.6. Let G_1 and G_2 be two graphs with no vertex in common, We define the join of G_1 and G_2 , denoted by $G_1 + G_2$, to be the graph with vertex set $V(G_1 + G_2) = V(G_1) \cup V(G_2)$ and edge set $E(G_1 + G_2) = E(G_1) \cup E(G_2) \cup J$, where $J = \{x_1x_2 : x_1 \in V(G_1), x_2 \in V(G_2)\}$.

Definition 1.7. The fan graph F_n is defined as $F_n = P_n + K_1$.

Definition 1.8. Let G and H be two graphs. The corona product of G and H , denoted by $G \odot H$, is obtained by taking one copy of G and $|V(G)|$ copies of H , and by joining each vertex of the i^{th} copies of H to the i^{th} vertex of G , for $1, 2, 3, \dots, |V(G)|$.

Definition 1.9. Let C_n be a cycle of n vertices. The crown graph is defined as $C_n \odot K_1$.

Definition 1.10. Let P_n be a path of n vertices. The comb graph is defined as $P_n \odot K_1$.

Definition 1.11. Let P_n be a path of n vertices. The double comb graph is defined as $P_n \odot 2K_1$.

In this paper, we have discuss antimagic labeling of degree splitting graphs of path, ladder graph, fan graph, crown graph, comb graph and double comb graph.

2 Main Results

Theorem 2.1. The graph $DS(P_n)$ is an antimagic graph.

Proof. Let $v_1, v_2, v_3, \dots, v_n$ be the vertices of P_n with $d(v_1) = d(v_n) = 1$. To construct $DS(P_n)$, add vertices w_1 and w_2 such that w_1 is adjacent to each vertex of degree 1 and w_2 is adjacent to each vertex of degree 2 in P_n . Then $|V(DS(P_n))| = n + 2$ and $|E(DS(P_n))| = 2n - 1$ for $n \geq 4$.

We define $f : E(DS(P_n)) \rightarrow \{1, 2, \dots, 2n - 1\}$, as per following three cases.

Case - 1 For $n = 2$ and 3 .

$DS(P_2)$ and $DS(P_3)$ are also known as cycle C_3 and cycle C_4 respectively. The antimagic labeling of C_3 and C_4 have been investigated by Hartsfield and Ringel [1990].

Case - 2 For $n = 4$.




Figure 1: $DS(P_4)$ and its antimagic labeling.

Case - 3 For $n > 4$.

$$f(v_1w_1) = 2(n - 1),$$

$$f(v_nw_1) = 2n - 1,$$

$$f(v_iv_{i+1}) = i; \quad \text{For } 1 \leq i \leq n - 1,$$

$$f(v_iw_2) = 2n - i - 1; \quad \text{For } 2 \leq i \leq n - 1.$$

Above define edge labeling function will generate distinct vertex labels for all the vertices of $DS(P_n)$. Thus, f is an antimagic labeling.

Hence, the graph $DS(P_n)$ is an antimagic graph. □

Illustration 2.1. The graph $DS(P_7)$ and its antimagic labeling is shown in Figure 2.




Figure 2: $DS(P_7)$ and its antimagic labeling.

Antimagic Labeling of Some Degree Splitting Graphs

Theorem 2.2. *The graph $DS(L_n)$ is an antimagic graph.*

Proof. Let $v_1, v_2, v_3, \dots, v_n$ be the vertices of first path and $v_{n+1}, v_{n+2}, v_{n+3}, \dots, v_{2n}$ be the vertices of second path in ladder L_n . To construct $DS(L_n)$, add vertices w_1 and w_2 such that w_1 is adjacent to each vertex of degree 2 and w_2 is adjacent to each vertex of degree 3 in ladder L_n . Then $|V(DS(L_n))| = 2n + 2$ and $|E(DS(L_n))| = 5n - 2$ for $n \geq 3$.

We define $f : E(DS(L_n)) \rightarrow \{1, 2, \dots, 5n - 2\}$ as per following two cases.

Case - 1 For $n = 2$.

$DS(L_2)$ is also known as wheel W_4 . The antimagic labeling of W_4 has been investigated by Hartsfield and Ringel [1990].

Case - 2 For $n \geq 3$.

$$\begin{aligned} f(v_i v_{i+1}) &= i; & \text{For } 1 \leq i \leq n-1, \\ f(v_i v_{n+i}) &= 2n - i; & \text{For } 1 \leq i \leq n, \\ f(v_{n+i} v_{n+1+i}) &= 2n + i - 1; & \text{For } 1 \leq i \leq n-1, \\ f(v_i w_2) &= 3(n-1) + i; & \text{For } 2 \leq i \leq n-1, \\ f(v_{n+i} w_2) &= 4n - 5 + i; & \text{For } 2 \leq i \leq n-1, \\ f(v_1 w_1) &= 5(n-1) + 2, \\ f(v_n w_1) &= 5(n-1) + 1, \\ f(v_{n+1} w_1) &= 5(n-1) + 3, \\ f(v_{2n} w_1) &= 5(n-1). \end{aligned}$$

Above define edge labeling function will generate distinct vertex labels for all the vertices of $DS(L_n)$. Thus, f is an antimagic labeling.

Hence, the graph $DS(L_n)$ is an antimagic graph. □

Illustration 2.2. *The graph $DS(L_6)$ and its antimagic labeling is shown in Figure 3.*




Figure 3: $DS(L_6)$ and its antimagic labeling.

Theorem 2.3. *The graph $DS(F_n)$ is an antimagic graph.*

Proof. Let v be a apex vertex and $v_1, v_2, v_3, \dots, v_n$ be the vertices corresponding to path P_n in graph $F_n = P_n + K_1$. To construct $DS(F_n)$, add vertices w_1 and w_2 such that w_1 is adjacent to each vertex of degree 2 and w_2 is adjacent to each vertex of degree 3 in fan F_n . Then $|V(DS(F_n))| = n + 3$ and $|E(DS(F_n))| = 3n - 1$ for $n \geq 4$.

We define $f : E(DS(F_n)) \rightarrow \{1, 2, \dots, 3n - 1\}$, as per following three cases.

Case - 1 For $n = 2$.

$DS(F_2)$ is also known as wheel W_3 . The antimagic labeling of W_3 has been investigated by Hartsfield and Ringel [1990].

Case - 2 For $n = 3$.

The graph $DS(F_3)$ is an antimagic graph by Alon et al. [2004].

Case - 3 For $n \geq 4$.

$$\begin{aligned} f(v_i v_{i+1}) &= i; & \text{For } 1 \leq i \leq n - 1, \\ f(v_i w_2) &= 2n - i - 1; & \text{For } 2 \leq i \leq n - 1, \\ f(v v_i) &= 2n + i - 1; & \text{For } 1 \leq i \leq n, \\ f(v_1 w_1) &= 2(n - 1), \\ f(v_n w_1) &= 2n - 1. \end{aligned}$$

Above define edge labeling function will generate distinct vertex labels for all the vertices of $DS(F_n)$. Thus, f is an antimagic labeling.

Hence, the graph $DS(F_n)$ is an antimagic graph. \square

Antimagic Labeling of Some Degree Splitting Graphs

Illustration 2.3. *The graph $DS(F_7)$ and its antimagic labeling is shown in Figure 4.*




Figure 4: $DS(F_7)$ and its antimagic labeling

Theorem 2.4. *The graph $DS(C_n \odot K_1)$ is an antimagic graph.*

Proof. Let $v_1, v_2, v_3, \dots, v_n$ be the vertices corresponding to cycle C_n and v'_i be the pendant vertices attached to v_i for each $i = 1, 2, \dots, n$ respectively in graph $C_n \odot K_1$. To construct $DS(C_n \odot K_1)$, add vertices w_1 and w_2 such that w_1 is adjacent to each vertex of degree 3 and w_2 is adjacent to each vertex of degree 1 in $C_n \odot K_1$. Then $|V(DS(C_n \odot K_1))| = 2n + 2$ and $|E(DS(C_n \odot K_1))| = 4n$. We define $f : E(DS(C_n \odot K_1)) \rightarrow \{1, 2, \dots, 4n\}$ as follows.

$$\begin{aligned}
 f(v_i v_{i+1}) &= 2n - i + 1; & \text{For } 1 \leq i \leq n - 1, \\
 f(v_1 v_n) &= n + 1, \\
 f(v_i v'_i) &= i; & \text{For } 1 \leq i \leq n, \\
 f(v_1 w_1) &= 2n + 1, \\
 f(v_i w_1) &= 3n - i + 2; & \text{For } 2 \leq i \leq n, \\
 f(v'_i w_2) &= 3n + i; & \text{For } 1 \leq i \leq n.
 \end{aligned}$$

Above define edge labeling function will generate distinct vertex labels for all the vertices of $DS(C_n \odot K_1)$. Thus, f is an antimagic labeling.

Hence, the graph $DS(C_n \odot K_1)$ is an antimagic graph. □

Illustration 2.4. *The graph $DS(C_6 \odot K_1)$ and its antimagic labeling is shown in Figure 5.*




Figure 5: $DS(C_6 \odot K_1)$ and its antimagic labeling.

Theorem 2.5. *The graph $DS(P_n \odot K_1)$ is an antimagic graph.*

Proof. $v_1, v_2, v_3, \dots, v_n$ be the vertices corresponding to path P_n and v'_i be the pendant vertices attached to v_i for each $i = 1, 2, \dots, n$ respectively in graph $P_n \odot K_1$. To construct $DS(P_n \odot K_1)$, add vertices w_1, w_2 and w_3 such that w_1 is adjacent to each vertex of degree 1, w_2 is adjacent to each vertex of degree 2 and w_3 is adjacent to each vertex of degree 3 in $P_n \odot K_1$. Then $|V(DS(P_n \odot K_1))| = 2n + 3$ and $|E(DS(P_n \odot K_1))| = 4n - 1$ for $n \geq 4$.

We define $f : E(DS(P_n \odot K_1)) \rightarrow \{1, 2, \dots, 4n - 1\}$ as per following two cases.

Case - 1 For $n = 2$ and 3.




Figure 6: $DS(P_2 \odot K_1)$ and $DS(P_3 \odot K_1)$ and their antimagic labeling.

Antimagic Labeling of Some Degree Splitting Graphs

Case - 2 For $n \geq 4$.

$$\begin{aligned} f(v_i v'_i) &= i; & \text{For } 1 \leq i \leq n, \\ f(v_i v_{i+1}) &= 2n - i; & \text{For } 1 \leq i \leq n - 1, \\ f(v_i w_3) &= 4n - i - 1; & \text{For } 2 \leq i \leq n - 1, \\ f(v'_i w_1) &= 2n + i - 1; & \text{For } 1 \leq i \leq n, \\ f(v_1 w_2) &= 2(2n - 1), \\ f(v_n w_2) &= 4n - 1. \end{aligned}$$

Above define edge labeling function will generate distinct vertex labels for all the vertices of $DS(P_n \odot K_1)$. Thus, f is an antimagic labeling.

Hence, the graph $DS(P_n \odot K_1)$ is an antimagic graph. □

Illustration 2.5. The graph $DS(P_8 \odot K_1)$ and its antimagic labeling is shown in Figure 7.




Figure 7: $DS(P_8 \odot K_1)$ and its antimagic labeling.

Theorem 2.6. The graph $DS(P_n \odot 2K_1)$ is an antimagic graph.

Proof. $v_1, v_2, v_3, \dots, v_n$ be the vertices corresponding to path P_n and v'_i and v''_i be the pendant vertices attached to v_i for each $i = 1, 2, \dots, n$ respectively in graph

$P_n \odot 2K_1$. To construct $DS(P_n \odot 2K_1)$, add vertices w_1, w_2 and w_3 such that w_1 is adjacent to each vertex of degree 1, w_2 is adjacent to each vertex of degree 3 and w_3 is adjacent to each vertex of degree 4 in $P_n \odot 2K_1$. Then $|V(DS(P_n \odot 2K_1))| = 3n + 3$ and $|E(DS(P_n \odot 2K_1))| = 6n - 1$ for $n \geq 4$.

We define $f : E(DS(P_n \odot 2K_1)) \rightarrow \{1, 2, \dots, 6n - 1\}$ as per following two cases.

Case - 1 For $n = 2$ and 3.




Figure 8: $DS(P_2 \odot 2K_1)$ and $DS(P_3 \odot 2K_1)$ and their antimagic labeling.

Case - 2 For $n \geq 4$.

- $f(v_i v_{i+1}) = i;$ For $1 \leq i \leq n - 1,$
- $f(v_i v'_i) = n - 1 + i;$ For $1 \leq i \leq n,$
- $f(v_i v''_i) = 2n - 1 + i;$ For $1 \leq i \leq n,$
- $f(v_i w_3) = 6n - 1 - i;$ For $2 \leq i \leq n - 1,$
- $f(v'_i w_1) = 3n - 1 + i;$ For $1 \leq i \leq n,$
- $f(v''_i w_1) = 4n - 1 + i;$ For $1 \leq i \leq n,$
- $f(v_i w_2) = 2(3n - 1),$
- $f(v_n w_2) = 6n - 1.$

Above define edge labeling function will generate distinct vertex labels for all the vertices of $DS(P_n \odot 2K_1)$. Thus, f is an antimagic labeling.

Hence, the graph $DS(P_n \odot 2K_1)$ is an antimagic graph. □

Illustration 2.6. The graph $DS(P_5 \odot 2K_1)$ and its antimagic labeling is shown in Figure 9.




Figure 9: $DS(P_5 \odot 2K_1)$ and its antimagic labeling.

3 Conclusions

Hartsfield and Ringel [1990] has proved that path is an antimagic graph. Cheng [2007] has derived the antimagic labeling of ladder graph. While Sridharan and Umarani [2012] have discussed antimagic labeling of fan graph and crown graph. In this paper, we have discussed antimagic labeling of larger graphs obtained from Path, ladder graph, fan graph and crown graph using the degree splitting operation. Also antimagic labeling of degree splitting graphs of comb graph and double comb graph have been obtained. To investigated antimagic labeling for different graph families as well as in the context of various graph operations is an open area of research.

Acknowledgements

The authors are highly thankful to anonymous referees for kind comments and constructive suggestions.

References

- N. Alon, G. Kaplan, A. Lev, Y. Roditty, and R. Yuster. Dense graphs are antimagic. *Journal of Graph Theory*, 47:297–309, 2004.
- Y. Cheng. Lattice grids and prisms are antimagic. *Theoretical Computer Science*, 374:66–73, 2007.
- Y. Cheng. A new class of antimagic cartesian product graphs. *Discrete Math.*, 308(24):6441–6448, 2008.
- J. Clark and D. Holton. *A first look at Graph Theory*. World Scientific Publishing Co. Pvt. Ltd., 1969.
- J. Gallian. A dynamic survey of graph labeling. *The Electronics Journal of Combinatorics*, 25(#DS6), 2022.
- N. Hartsfield and G. Ringel. *Pearls in Graph Theory*. Academic Press, Boston, 1990.
- A. Krishnaa. Some applications of labelled graphs. *International Journal of Mathematics Trends and Technology*, 37(3):209–213, 2016.
- N. Sridharan and R. Umarani. Antimagic labeling of graphs. *International Journal of Engineering Science, Advanced Computing and Bio-Technology*, 3(1): 23–41, 2012.
- S. Vaidya and N. Vyas. Antimagic labeling in the context of switching of a vertex. *Annal of pure and Applied Mathematics*, 2(1):33–39, 2012.
- S. Vaidya and N. Vyas. Antimagic labeling of some path and cycle related graphs. *Annal of pure and Applied Mathematics*, 3(2):119–128, 2013.
- T. Wang. Toroidal grids are anti-magic. *In International Computing and Combinatorics Conference*, pages 671–679, 2005.
- T. Wang, M. Lui, and D. Li. Some classes of disconnected antimagic graphs and their joins. *Whhan Univ. J. Nat. Sci.*, 17(3):195–199, 2012.