








OPEN ACCESS

Short report

Alzheimer's disease marker phospho-tau181 is not elevated in the first year after moderate-to-severe TBI

Neil Graham ^{1,2}, Karl Zimmerman,^{1,2} Amanda J Heslegrave ³, Ashvini Keshavan,⁴ Federico Moro,^{5,6} Samia Abed-Maillard,⁷ Adriano Bernini,⁸ Vincent Dunet,⁹ Elena Garbero,¹⁰ Giovanni Nattino,¹⁰ Arturo Chierogato,¹¹ Enrico Fainardi,¹² Camelia Baciu,¹¹ Primoz Gradisek,¹³ Sandra Magnoni,¹⁴ Mauro Oddo,^{7,15} Guido Bertolini,¹⁰ Jonathan M Schott ^{3,16}, Henrik Zetterberg ^{3,17}, David Sharp ^{1,2}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/jnnp-2023-331854>).

For numbered affiliations see end of article.

Correspondence to

Dr David Sharp, Department of Brain Sciences, Imperial College London, London, SW7 2BX, UK; david.sharp@imperial.ac.uk

Received 17 May 2023

Accepted 19 September 2023

ABSTRACT

Background Traumatic brain injury (TBI) is associated with the tauopathies Alzheimer's disease and chronic traumatic encephalopathy. Advanced immunoassays show significant elevations in plasma total tau (t-tau) early post-TBI, but concentrations subsequently normalise rapidly. Tau phosphorylated at serine-181 (p-tau181) is a well-validated Alzheimer's disease marker that could potentially seed progressive neurodegeneration. We tested whether post-traumatic p-tau181 concentrations are elevated and relate to progressive brain atrophy.

Methods Plasma p-tau181 and other post-traumatic biomarkers, including total-tau (t-tau), neurofilament light (NfL), ubiquitin carboxy-terminal hydrolase L1 (UCH-L1) and glial fibrillary acidic protein (GFAP), were assessed after moderate-to-severe TBI in the BIO-AX-TBI cohort (first sample mean 2.7 days, second sample within 10 days, then 6 weeks, 6 months and 12 months, n=42). Brain atrophy rates were assessed in aligned serial MRI (n=40). Concentrations were compared patients with and without Alzheimer's disease, with healthy controls. **Results** Plasma p-tau181 concentrations were significantly raised in patients with Alzheimer's disease but not after TBI, where concentrations were non-elevated, and remained stable over one year. P-tau181 after TBI was not predictive of brain atrophy rates in either grey or white matter. In contrast, substantial trauma-associated elevations in t-tau, NfL, GFAP and UCH-L1 were seen, with concentrations of NfL and t-tau predictive of brain atrophy rates.

Conclusions Plasma p-tau181 is not significantly elevated during the first year after moderate-to-severe TBI and levels do not relate to neuroimaging measures of neurodegeneration.

traumatic brain injury-related neurodegeneration (TReND) (online supplemental ref).

A key question is how acute traumatic elevations of τ relate to progressive neurodegeneration. This is potentially mechanistically important as abnormal τ may cause neurodegeneration through prion-like proteopathic seeding. Total τ (t-tau) is increased more than one hundred-fold after moderate-to-severe TBI in brain extracellular fluid, and plasma t-tau predicts neurodegeneration. However, in contrast to the axonal degeneration marker neurofilament light (NfL) and astroglial marker glial fibrillary acidic protein (GFAP), t-tau rapidly normalises.² Advanced biomarker assays now allow investigation of links between TBI and dementia. In AD, τ phosphorylated at threonine 181 (p-tau₁₈₁) correlates with neuropathology,³ and is being incorporated clinically. P-tau₁₈₁ has not been investigated after moderate-to-severe TBI.

We assessed plasma P-tau₁₈₁ in a subset of the BIO-AX-TBI cohort of moderate-to-severe TBI (defined using the Mayo classification, see online supplemental file 1), healthy controls and AD patients, comparing blood biomarker concentrations to MRI measures of neurodegeneration in the TBI group. We have previously described the cohort and characterised trends of NfL, t-tau, neuronal marker ubiquitin C-terminal hydrolase L1 (UCH-L1) and GFAP. Hence, we were able to directly compare these markers with P-tau₁₈₁.² We hypothesised that: (1) p-tau₁₈₁ would increase early post-TBI, (2) remain elevated at 1 year and (3) predict brain atrophy.

METHODS

See online supplemental file 1.

RESULTS

P-tau₁₈₁ was quantified over 1 year in 42 patients after moderate-to-severe TBI, aged 48.7 years (mean, SD 15.6) with 76.2% male (online supplemental table 1). The lowest Glasgow Coma Scale was 3–8 in 26.8%, 9–13 in 39.0% and 14–15 in 34.1% (unknown in n=1). Diffuse injury was present on CT in 4.8%, with contusions/intraparenchymal haemorrhage in 47.6%, subdural

INTRODUCTION

Traumatic brain injury (TBI) is common occurrence and an environmental risk factor for dementia. A range of pathologies are described postinjury including the tauopathies Alzheimer's disease (AD) and chronic traumatic encephalopathy (CTE),¹ which form part of the broader complex constellation of postinjury pathologies and has been termed



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

To cite: Graham N, Zimmerman K, Heslegrave AJ, et al. *J Neurol Neurosurg Psychiatry* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jnnp-2023-331854




Figure 1 Longitudinal fluid biomarker trajectories after moderate-to-severe TBI, healthy volunteers and Alzheimer's disease. Fluid biomarkers in healthy volunteers, longitudinally in patients following TBI, people with AD and non-AD healthy controls (non-AD controls). Boxplots show median and quartiles (hinges), with whiskers extending upto 1.5 times the IQR. Individual data points are shown and connected by lines indicating within-subject trajectories. (A) shows p-tau₁₈₁; (B) shows total τ (t-tau), neurofilament light (NfL), ubiquitin c-terminal hydrolase L1 (UCH-L1) and glial fibrillar acidic protein (GFAP). AD, Alzheimer's disease; TBI, traumatic brain injury.

haematoma in 57.1%, and extradural haematoma in 14.3%. This group had less severe injuries than the broader BIO-AX-TBI cohort previously reported, indicated by lower peak 10-day injury biomarker concentrations (all $p < 0.05$, except NfL).² A group of male healthy controls underwent aligned p-tau₁₈₁ assessment (mean 45.9 years, SD 3.2). Thirty-one patients with AD were also assessed (63.7 years, SD 6.5, 48.4% male) and 14 non-AD controls (mean 60.6 years, SD 6.5, 71.4% male). Age and sex were included as confounders.

Median plasma p-tau₁₈₁ at 2.7 days (SD 2.8) post-TBI was 1.5 pg/mL (IQR 1.7) and did not differ significantly from healthy

volunteers (median 1.5 pg/mL, IQR 1.9) (see figure 1; online supplemental table 2). P-tau₁₈₁ concentrations were stable in the year after injury: day 6.3 (SD 2.4) 1.5 pg/mL (IQR 1.0); day 30.0 (SD 12.1) 1.3 pg/mL (IQR 0.8); 6 months 1.4 pg/mL (IQR 0.7) and 1 year 1.5 pg/mL (IQR 0.5). There was no significant longitudinal change. In contrast, reductions were seen in t-tau, GFAP, NfL and UCH-L1 as previously reported (all $p < 0.001$).²

There was no significant correlation of p-tau₁₈₁ measured subacutely and grey matter atrophy measured at 6 months, nor between concentrations within 6 months and white matter atrophy at 6–12 months. As previously shown in the wider

cohort,² hyperacute t-tau predicted grey matter atrophy ($p=0.046$; adjusted $R^2=0.12$), and subacute plasma NfL predicted white matter atrophy ($p=0.004$, $R^2=0.20$).² P-tau₁₈₁ was significantly raised in AD (2.0 times (1.5–2.7) higher, $p<0.001$) versus non-AD controls.

DISCUSSION

Plasma p-tau₁₈₁ was not elevated after moderate-to-severe TBI and concentrations did not vary significantly over 1 year. This contrasts with our previous findings in other markers t-tau, NfL, GFAP and UCH-L1, which showed substantial elevations.² Unlike plasma t-tau and NfL, we found no correlation of p-tau₁₈₁ at any time point and brain atrophy, a measure of neurodegeneration.

P-tau₁₈₁ is a neuropathologically validated in vivo marker of amyloid-induced τ phosphorylation in AD.³ Given the presence of amyloid and τ pathologies post-TBI, there is interest in whether TBI may trigger neurodegeneration through mechanisms similar to AD, and whether this is reflected in AD-specific blood biomarkers. AD patients showed substantial p-tau₁₈₁ elevations, demonstrating the assay's sensitivity, but this was not seen post-TBI. The lack of p-tau₁₈₁ elevation at 1-year contrasts with neurodegeneration marker NfL and astroglial marker GFAP, both of which remained chronically elevated, as previously reported.² It is possible that phosphorylated τ accumulates as a late consequence of TBI and we would not have identified this in our 1-year follow-up period. Post-traumatic neurodegeneration is likely to be dynamic over time and a specific temporal pattern of plasma p-tau isoform changes, as seen in AD where plasma p-tau₂₃₁ precedes p-tau₁₈₁ positivity, with both markers well correlated with τ PET.⁴ Very long-term follow-up incorporating comprehensive longitudinal fluid biomarker assessment, brain volumetry and molecular imaging would likely be highly informative.

It is possible that other blood markers may be more specific to post-traumatic neurodegeneration. For example, post-mortem work suggests that p-tau₂₀₂ may have greatest specificity for CTE. This has yet to be assessed clinically as there is not currently a reliable assay to do so at scale. In addition, brain derived τ shows promise as a marker, correlating more closely with CSF τ and neurofibrillary tangle burden in AD better than t-tau (online supplemental ref).

There are several potential limitations. Relatively few patients were sampled <24 hours after injury, hence we may have missed an early peak in p-tau₁₈₁ as previously seen using a different assay-type quantifying p-tau₂₃₁.⁵ Second, the use of p-tau₁₈₁ controls analysed separately from TBI patients may introduce bias: however, we feel this is unlikely due to good assay performance, large numbers, and lack of longitudinal injury-associated change. Last, TBI and young controls were not well sex-matched, though this was included in statistical models.

In conclusion, plasma p-tau₁₈₁ was not increased over 1 year after moderate-to-severe TBI and was not associated with neurodegeneration. P-tau₁₈₁ dynamics were not only distinct from t-tau but differed from other biomarkers NfL, GFAP and UCH-L1. This suggests that p-tau₁₈₁ does not contribute to progressive neurodegeneration commonly seen after TBI, at least in the first year.

Author affiliations

¹Brain Sciences, Imperial College London, London, UK

²UK Dementia Research Institute Centre for Care Research and Technology, Imperial College London, London, UK

³UK Dementia Research Institute, University College London, London, UK

⁴Dementia Research Centre, UCL Queen Square Institute of Neurology, London, UK

⁵Laboratory of Acute Brain Injury and Neuroprotection, Department of Acute Brain and Cardiovascular Injury, Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Milan, Italy

⁶Dipartimento di Anestesia e Rianimazione, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

⁷Neuroscience Critical Care Research Group, Department of Intensive Care Medicine, CHUV Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland

⁸Department of Clinical Neurosciences, CHUV Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland

⁹Department of Medical Radiology, CHUV Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland

¹⁰Laboratory of Clinical Epidemiology, Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Ranica, Italy

¹¹Terapia Intensiva ad indirizzo Neurologico & Neurochirurgico, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy

¹²Department of Experimental and Clinical Sciences, Careggi University Hospital and University of Florence, Florence, Italy

¹³Clinical Department of Anaesthesiology and Intensive Therapy, University Medical Center, Ljubljana, Slovenia

¹⁴Department of Anesthesia and Intensive Care, Santa Chiara Hospital, Trento, Italy

¹⁵Directorate for Innovation and Clinical Research, CHUV Lausanne University Hospital, Lausanne, Switzerland

¹⁶Dementia Research Centre and Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, London, UK

¹⁷Department of Psychiatry and Neurochemistry, The Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

Twitter Neil Graham @nsngraham, Jonathan M Schott @jmschott and David Sharp @neurosharp

Contributors DS, HZ, NG, KZ, SA-M, MO, PG and GB were involved in the design and conception of the study. All authors were involved in data collection, collation and quality control. The first draft was prepared by NG, and DS revised initial iterations. All authors reviewed and approved the final version of the manuscript.

Funding Core funds for the BIO-AX-TBI project were from the ERA-NET NEURON Cofund (MR/R004528/1), a part of the European Research Projects on External Insults to the Nervous System call, within the Horizon 2020 funding framework, (PIs: DS, HZ, MO, GB and SM). Other sources included: Alzheimer's Research UK Clinical Research Fellowship and National Institute for Health Research (UK) Academic Clinical Lectureship (NG), UK Dementia Research Institute (DRI) Care Research and Technology Centre (NG, DS and KZ), National Institute of Health Research (NIHR) Professorship (NIHR-RP-011-048) (DS), the NIHR Clinical Research Facility and Biomedical Research Centre (BRC) at Imperial College Healthcare NHS Trust (NG, DS), Medical Research Council Clinician Scientist Fellowship (DS). Plasma p-tau₁₈₁ analysis at UCL was funded by an Alzheimer's Research UK Network Pump Priming grant (AK) AK and JMS acknowledge the support of the National Institute for Health Research University College London Hospitals Biomedical Research Centre, the Weston Brain Institute (UB170045) and the Alzheimer's Association. JMS acknowledges the support of the Wolfson Foundation, ARUK (ARUK-PG2017-1946), Brain Research UK (UCC14191), Medical Research Council, British Heart Foundation and European Union's Horizon 2020 research and innovation programme (Grant 666992). HZ is a Wallenberg Scholar supported by grants from the Swedish Research Council (#2022-01018 and #2019-02397), the European Union's Horizon Europe research and innovation programme under grant agreement No 101053962, Swedish State Support for Clinical Research (#ALFGBG-71320), the Alzheimer Drug Discovery Foundation (ADDF), USA (#201809-2016862), the AD Strategic Fund and the Alzheimer's Association (#ADSF-21-831376-C, #ADSF-21-831381-C and #ADSF-21-831377-C), the Bluefield Project, the Olav Thon Foundation, the Erling-Persson Family Foundation, Stiftelsen för Gamla Tjänarinnor, Hjärnfonden, Sweden (#FO2022-0270), the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860197 (MIRIAD), the European Union Joint Programme—Neurodegenerative Disease Research (JPND2021-00694) and the UK Dementia Research Institute at UCL (UKDRI-1003). MO is supported by the Swiss National Science Foundation (nos. 31NE30_173675 and 32003B_188501).

Competing interests HZ has served at scientific advisory boards and/or as a consultant for Abbvie, Acumen, Alector, Alzinova, ALZPath, Annexon, Apellis, Artery Therapeutics, AZTherapies, CogRx, Denali, Eisai, Nervgen, Novo Nordisk, Optoceutics, Passage Bio, Pinteon Therapeutics, Prothena, Red Abbey Labs, reMYND, Roche, Samumed, Siemens Healthineers, Triplet Therapeutics and Wave, has given lectures in symposia sponsored by Cellectric, Fujirebio, Alzecure, Biogen and Roche, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program (outside submitted work). MO received research support and speaker fees from Neuroptics, USA, unrelated to the present work, and is member of the Scientific Advisory Board of Neuroptics. DS serves on the

concussion advisory board of the UK Rugby Football Union, and undertakes clinical private practice including medicolegal assessments.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by COAT study (TBI patients): Camberwell and St Giles REC; ref 17/LO/2066 CSF study (AD patients 12/3044): Queen Square Research Ethics Committee; ref 12_LO_1504ADVANCE study (controls): Ministry of Defence Research Ethics Committee (MODREC); ref 20220405-2126MODREC22. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given,

and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Neil Graham <http://orcid.org/0000-0002-0183-3368>
Amanda J Heslegrave <http://orcid.org/0000-0002-7290-6405>
Jonathan M Schott <http://orcid.org/0000-0003-2059-024X>
Henrik Zetterberg <http://orcid.org/0000-0003-3930-4354>
David Sharp <http://orcid.org/0000-0003-4995-2240>

REFERENCES

- Livingston G, Huntley J, Sommerlad A, *et al*. Dementia prevention, intervention, and care: 2020 report of the lancet commission. *Lancet* 2020;396:413–46.
- Graham NSN, Zimmerman KA, Moro F, *et al*. Axonal marker neurofilament light predicts long-term outcomes and progressive neurodegeneration after traumatic brain injury. *Sci Transl Med* 2021;13:eabg9922.
- Lantero Rodriguez J, Karikari TK, Suárez-Calvet M, *et al*. Plasma P-Tau181 accurately predicts Alzheimer's disease pathology at least 8 years prior to post-mortem and improves the clinical characterisation of cognitive decline. *Acta Neuropathol* 2020;140:267–78.
- Tissot C, Therriault J, Kunach P, *et al*. Comparing Tau status determined via plasma pTau181, pTau231 and [¹⁸F]MK6240 tau-PET. *EBioMedicine* 2022;76:103837.
- Rubenstein R, Chang B, Yue JK, *et al*. Comparing plasma Phospho Tau, total Tau, and Phospho Tau-total Tau ratio as acute and chronic traumatic brain injury biomarkers. *JAMA Neurol* 2017;74:1063–72.

1 **Alzheimer's disease marker phospho-tau181 is not elevated in the first year**
2 **after moderate-severe TBI**

3

4 Neil S.N. Graham^{1,2,3}, Karl A. Zimmerman^{1,2,3}, Amanda Heslegrave^{4,5}, Ashvini Keshavan⁶,
5 Federico Moro^{7,8}, Samia Abed Maillard⁹, Adriano Bernini¹⁰, Vincent Dunet¹¹, Elena
6 Garbero¹², Giovanni Nattino¹², Arturo Chierogato¹³, Enrico Fainardi¹⁴, Camelia Baciu¹³,
7 Primoz Gradisek¹⁵, Sandra Magnoni¹⁶, Mauro Oddo^{9,17}, Guido Bertolini¹², Jonathan M
8 Schott^{4,5,6}, Henrik Zetterberg^{4,5,18,19,20,21}, David J. Sharp^{*1,2,3}

9

10

SUPPLEMENTARY METHODS

11

12 The BIO-AX-TBI cohort [1] assessed axonal injury over one year in patients after moderate-
13 severe TBI using advanced biomarkers, relating these to clinical outcomes. Patients
14 presenting to major trauma centres at participating study sites were eligible for inclusion if
15 they met the Mayo classification criteria for moderate-severe TBI.[2] These include: acute
16 abnormality on CT (eg. subarachnoid haemorrhage, subdural haematoma, extradural
17 haematoma, intraparenchymal bleed/contusion), injury penetrating the dura, post
18 traumatic amnesia duration > 24 hours, loss of consciousness > 30 minutes, worst Glasgow
19 Coma Scale < 13 or death due to TBI. The presence of any one of these features is sufficient
20 for the patient to meet criteria for moderate-severe injury. Blood was sampled twice within
21 ten days post-injury (first sample mean 2.7 days post-TBI [standard deviation 2.8], second
22 sample 6.3 [2.4] days), between ten days and six weeks (26.0 [2.4] days since injury), at six
23 months (189.0 [16.0] days) and twelve months (371.0 [55.2] days). Plasma was sampled
24 using standard procedures and frozen at -80C. P-tau₁₈₁ was quantified using a single-plex P-
25 tau181 V2.0-Advantage kit on a Simoa HD-X (Quanterix, Billerica, MA). UCH-L1, GFAP, total
26 tau and NfL concentrations were previously measured using multiplexed kits (Neurology 4-
27 Plex B, Quanterix, Billerica, MA).[3] A 1:4 dilution was used on-board the instrument.
28 Samples were run in duplicate and average values reported. The intraplate coefficient of
29 variation (CV) for p-tau₁₈₁ was 8.4% with an interplate CV of 5.9%.

30 Due to limitations in the Ptau₁₈₁ assay availability and the large number of longitudinal
31 samples in BIO-AX-TBI cohort, we elected analyse a subset of the group where there were
32 samples spanning an entire year, including: an acute blood sample within days 1-10, as well
33 as a chronic sample at six and twelve months post-injury. This equated to n=42 individuals.

1 Healthy control data, comprising p-tau₁₈₁ concentrations from the ADVANCE cohort (50
2 male service personnel with no history of major trauma) were used for comparison.[4, 5]
3 The intraplate CV was 8.3% and interplate CV was 8.6%. Data from a separate, pre-existing
4 group of age-matched healthy controls in BIO-AX-TBI provided norms for previously tested
5 biomarkers UCH-L1, GFAP, t-tau, NfL, using Neurology 4-plex B kits on a Simoa HD-1.

6 People with Alzheimer's disease (AD) and non-AD controls were assessed at UCL (Study
7 12/3044). Patients had a clinical diagnosis of Alzheimer's type mild cognitive impairment or
8 dementia with CSF showing amyloid pathology (defined by amyloid beta 42:40 ratio \leq
9 0.065). Age-matched controls had subjective cognitive impairment or primary mood
10 disorder and all were amyloid-negative. Analysis was performed with a single-plex P-tau181
11 kit (intraplate CV 1.16%).

12 The Marshall classification defined acute imaging pathologies. MRI was acquired in 40 of the
13 TBI patients. Pairwise volumetric T1-weighted MRI was analysed as previously [3] using
14 SPM12 (UCL), producing grey/white matter atrophy rates spanning subacute (10 day-6
15 weeks to 6 months) and early-chronic (6 months to 12 months) periods.

16 Statistical analyses were conducted using R (v4.1.1). Group differences were assessed using
17 linear regression; longitudinal biomarker assessment within TBI patients was performed
18 using linear mixed effects modelling with subject as a random effect. Due to their non-
19 normal distribution log-transformed biomarker concentrations were used; age and sex were
20 included as covariates. The relevant research ethics committee approvals were granted for
21 the investigation (UK REC ref: 17/LO/2066, Camberwell and St Giles Ethics Committee).

22
23
24

1
2
3
4
5
6
7
8
9

SUPPLEMENTARY RESULTS

Supplementary Table 1. Demographics and Plasma Biomarker Concentrations

Demographics and plasma concentrations of biomarkers reported for patients and controls. Values for patients after TBI are for the first acute testing timepoint only (ie. the first sample taken within ten days of injury). IQR: interquartile range; Nfl: neurofilament light; UCH-L1: ubiquitin c-terminal hydrolase-L1; p-tau₁₈₁:tau phosphorylated at serine 181, T-tau: total tau; GFAP: glial fibrillary acidic protein

		Traumatic brain injury	Ptau181 healthy controls	Other biomarker healthy controls	Alzheimer's disease patients	Non-AD controls
<i>Demographics</i>	n	42	50	128	31	14
	Age (mean (SD))	48.7 (15.7)	45.9 (3.20)	42.1 (16.8)	63.7 (6.48)	60.5 (6.50)
	Sex = Male (%)	32 (76.2)	50 (100.0)	78 (60.9)	15 (48.4)	10 (71.4)
<i>Plasma biomarker concentration (pg/ml), (median [IQR])</i>	P-tau ₁₈₁	1.48 [1.13, 2.85]	1.51 [1.25, 2.11]	—	3.60 [2.66, 4.64]	1.41 [1.25, 1.93]
	T-tau	3.50 [1.66, 6.57]	—	1.20 [0.88, 1.85]	—	—
	Nfl	51.3 [22.1, 93.8]	—	5.91 [4.12, 8.72]	—	—
	UCH-L1	37.6 [16.1, 84.5]	—	4.69 [4.69, 10.88]	—	—
	GFAP	3857 [589, 18942]	—	57 [40, 78]	—	—

10
11

1 **Supplementary Table 2. Longitudinal fluid biomarker concentrations following TBI**

2
3 IQR: interquartile range; NfL: neurofilament light; UCH-L1: ubiquitin c-terminal hydrolase-L1; p-tau₁₈₁: tau phosphorylated at serine 181; T-
4 tau: total tau; GFAP: glial fibrillary acidic protein.
5

	Timepoint following moderate-severe TBI				
	Day 0-10, first sample	Day 0-10, second sample	Subacute, day 10-6 weeks	6 month	12 month
P-tau₁₈₁					
n	37	30	42	42	42
Median	1.5	1.5	1.3	1.4	1.5
IQR	1.7	1	0.8	0.7	0.5
T-tau					
n	36	30	42	40	21
Median	3.5	2.1	2.7	1.3	1.3
IQR	4.9	2.8	2.3	1	1.1
NfL					
n	36	30	42	40	21
Median	51.3	124.7	328.4	24.9	12.2
IQR	71.7	152.6	445.6	30.9	6.5
UCH-L1					
n	36	30	42	37	18
Median	37.6	19.6	18	4.7	4.7
IQR	68.5	21.1	15.1	6.8	6.8
GFAP					
n	36	30	42	40	21
Median	3857	534.5	213.1	71.1	83.8
IQR	18352.7	1338.5	264.2	76	62.7

6
7

REFERENCES

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

Additional references (main manuscript)

- Smith, D.H., et al., *Collaborative Neuropathology Network Characterizing Outcomes of TBI (CONNECT-TBI)*. Acta Neuropathol Commun, 2021. **9**(1): p. 32
- Gonzalez-Ortiz, F., et al., *Brain-derived tau: a novel blood-based biomarker for Alzheimer's disease-type neurodegeneration*. Brain, 2023. **146**(3): p. 1152-1165

Supplementary references

1. Graham, N.S., et al., *Multicentre longitudinal study of fluid and neuroimaging BIOMarkers of Axonal injury after traumatic brain injury: the BIO-AX-TBI study protocol*. BMJ Open, 2020. **10**(11): p. e042093.
2. Malec, J.F., et al., *The mayo classification system for traumatic brain injury severity*. J Neurotrauma, 2007. **24**(9): p. 1417-24.
3. Graham, N.S.N., et al., *Axonal marker neurofilament light predicts long-term outcomes and progressive neurodegeneration after traumatic brain injury*. Science Translational Medicine, 2021. **13**(613): p. eabg9922.
4. Bennett, A.N., et al., *Study protocol for a prospective, longitudinal cohort study investigating the medical and psychosocial outcomes of UK combat casualties from the Afghanistan war: the ADVANCE Study*. BMJ Open, 2020. **10**(10): p. e037850.
5. Graham, N.S.N., et al., *ADVANCE-TBI study protocol: traumatic brain injury outcomes in UK military personnel serving in Afghanistan between 2003 and 2014 – a longitudinal cohort study*. BMJ Open, 2023. **13**(3): p. e069243.
6. Smith, D.H., et al., *Collaborative Neuropathology Network Characterizing Outcomes of TBI (CONNECT-TBI)*. Acta Neuropathol Commun, 2021. **9**(1): p. 32.