EXCISION OF A CHRONIC INFLAMMATORY MASS USING DIODE LASERS: A CASE REPORT

Nauman Rauf Khan<sup>1</sup>, Amna Nauman Khan<sup>2</sup>, Hira Butt<sup>3</sup>, Syeda Maryam Fatima<sup>4</sup>, Fatimah Imran<sup>5</sup>, Laiba Amer<sup>6</sup>

#### Correspondence

<sup>3</sup>Hira Butt, Demonstrator, Department of Oral Pathology, College of Dentistry, Sharif Medical and Dental College, Lahore ⊠: hira.ah.butt@gmail.com <sup>1</sup>Professor, Oral Pathology Department, College of Dentistry, Sharif Medical and Dental College, Lahore <sup>2</sup>Professor, Community and Preventive Dentistry, College of Dentistry, Sharif Medical and Dental College, Lahore <sup>4</sup>General Dental Practitioner, College of Dentistry. Sharif Medical and Dental College, Lahore <sup>5</sup>House officer, International Islamic Dental Hospital, Islamabad <sup>6</sup>General Dental Practitioner, College of Dentistry, Sharif Medical and Dental College, Lahore

How to cite this article

Khan NR, Khan AN, Butt H, Fatima SM, Imran F, Amer L. Excision of A Chronic Inflammatory Mass Using Diode Lasers: A Case Report. J Gandhara Med Dent Sci. 2023;10(2): 102-105 https://doi.org/10.37762/jgmds.10-2.419

## INTRODUCTION

Photobiomodulation (PBM), also known as low level laser therapy (LLLT) or cold laser therapy uses light energy to elicit biological responses from the cell and help regulate cell function.<sup>1</sup> This treatment procedure that nowadays is called photobiomodulation therapy (PBM) has also had more than 60 other names in scientific literature most common being -low-level laser therapy (LLLT) The reasons to prefer the term -PBM over - LLLT are twofold.<sup>2</sup> Firstly PBM does not imply that a laser is necessary to achieve the therapeutic benefits and secondly PBM implies that the therapeutic effects could in some instances be due to inhibition effects in combination with usual stimulation effects at cellular level.<sup>3</sup> According to the mechanism of action suggested by various studies LLLT affects the mitochondria of the cell, primarily Cytochrome c oxidase in the electron transfer chain(ETC) and porphyrins on the cell membrane.<sup>4,5</sup> Light photons absorbed by these receptors have been proposed to do three things : stimulation of ATP synthesis by activating ETC; transient stimulation of reactive oxygen species, increasing the conversion of ADP to ATP; and a temporary release of nitric oxide(NO) from cytochrome c oxidase, resulting in an increase in cellular respiration.<sup>6</sup> PBM utilizes nonionizing forms of

# <u>ABSTRACT</u>

Photobiomodulation (PBM), also known as low level laser therapy (LLLT) or cold laser therapy uses light energy to elicit biological responses from the cell and help regulate cell function. From a clinical point of view PBM offers dental practitioners a non-invasive treatment modality that can be used as an adjunct to the traditional therapies or as a therapeutic tool on its own. This is a case of surgical excision of a fluctuant mass using a scalpel and diode laser 980nm wavelength.

KEYWORDS:DiodeLaser,Photobiomodulation,SurgicalExcision,FluctuantMass,BoneAugmentation

light sources, such as light emitting diodes (LEDs), lasers, and broadband light to produce desired reactions in various tissues.<sup>7</sup> The results that need to be achieved by PBM depend upon the type of laser as well as the wavelength used.<sup>8</sup> The term LASER is an acronym for - light amplification by stmulated emission of radiation.<sup>9</sup> It pumps electrons of a - laser gain medium E1 to an excited state E2 using light, electricity or a chemical reaction as a source of energy.<sup>10</sup> Once majority of the electrons are in an excited state (population inversion) an incoming photon E photon will lead to stimulated outburst of new photons.<sup>10</sup> Hence, the light will be amplified. Mirrors placed at either end of the laser cavity allow the light to bounce back and forth, leading to significant amplification.<sup>10</sup> One of these mirrors is only partially reflective to allow the laser beam to escape from the cavity.<sup>10</sup>

# CASE

A 60-year old woman presented to the clinic with complaint of pain in the upper right 6 and 7 along with mobility and bleeding. Medical history revealed her to be have controlled Diabetes. The patient had undergone dental procedures before including multiple root canal treatments, crowns and scaling. Clinical examination revealed a fluctuant mass covering the upper right 6 and partially 7 as shown in figure 1.



Figure 1: Fluctuant Mass Covering the Upper Right 6 And 7

The Right upper 6 was extracted. Curettage was done and soft tissue and fluctuant mass was removed using a scalpel and diode laser 980nm wavelength as shown in figure 2.



Figure 2: Removal of the Fluctuant Mass and Associated Tooth

Excavation was done using surgical curettes and 4.0x 10mm Oostem implant tissue level was placed at 50N as shown in figure 3.



**Figure 3: Implant Placement** 

Curettage of upper right 7 and 8 was done using Gracey curettes and ultrasonic scaler. Periodontal therapy pocket treatment was done using diode laser to initiate bio stimulation and reduce mobility. Bone augmentation was done using sure Oss biosynthetic bone and membrane was placed as scaffold. Allograft Bone SureOss (FDBA) FDBA (freeze-dried bone allograft) 100% allograft cortical bone was done as shown in figure 4



Figure 4: Bone Augmentation

Silk 02 stitches were placed and removed after 7 days as shown in figure 5.



**Figure 5: Placement of Stitches** 

The patient was called for follow-up after 10 days and radiographic evaluation was done as shown in figure 6.



Figure 6: Post-Operative Radiograph

# DISCUSSION

PBM works on the principles of Arndt Schultz law, which is the basic principle of Neurosomatic therapy. It states that weak stimulus excites physiological activity, a moderate stimulus favours it, and a strong stimulus arrests it.<sup>11,12</sup> In order to achieve desired results a threshold for moderate to heavy stimuli should be calculated beyond which tissue damage may occur.13 Although the therapeutic window of these lasers is broad, it is essential to calculate the amount of dose (i.e., the energy density) required to achieve desired results before any therapeutic application.<sup>14</sup> From a clinical point of view PBM offers dental practitioners a non-invasive treatment modality that can be used as an adjunct to the traditional therapies or as a therapeutic tool on its own.<sup>15</sup> Scope of this technology includes dental analgesia, treatment of dentine hypersensitivity, healing of soft tissue lesions, reduction of pain and swelling after surgical procedures, better integration of implants into bone and faster movement of teeth during orthodontic procedures.<sup>16</sup>

# CONCLUSION

Photobiomodulation although an evolving technology has provided health practitioners with an opportunity to aid in patients wound healing, modulate the inflammatory response and regenerate nerves. It can be used effectively in dental specialties to better manage treatments that are often deemed painful by patients, without prescribing pharmaceuticals that often have a number of side effects. Apart from the health benefits that can be achieved by lasers. It is also highly acceptable among the patients, promising patient compliance required for the success of any treatment. In this the chronic inflammatory mass was excised with minimal invasion.

### **CONFLICT OF INTEREST:** None

### FUNDING SOURCES: None

## REFERENCES

- Dompe C, Moncrieff L, Matys J, Grzech-Leśniak K, Kocherova I, Bryja A, Bruska M, Dominiak M, Mozdziak P, Skiba TH, Shibli JA. Photobiomodulation—underlying mechanism and clinical applications. Journal of clinical medicine. 2020 Jun;9(6):1724.
- Anders JJ, Lanzafame RJ, Arany PR. Low-level light/laser therapy versus photobiomodulation therapy. Photomedicine and laser surgery. 2015 Apr 1;33(4):183-4.
- Mosca RC, Ong AA, Albasha O, Bass K, Arany P. Photobiomodulation therapy for wound care: a potent, noninvasive, photoceutical approach. Advances in skin & wound care. 2019 Apr 1;32(4):157-67.
- Thunshelle C, Hamblin MR. Transcranial low-level laser (light) therapy for brain injury. Photomedicine and laser surgery. 2016 Dec 1;34(12):587-98.
- 5. Karu TI. Low-power laser therapy. Biomedical photonics handbook. 2003 Feb;48:1-25.
- Asan MF, Babu GS, Castelino RL, Rao K, Pandita V. Applications of Photobiomodulation Therapy in Oral Medicine—A Review. European Journal of Therapeutics. 2021 Jun 1;27(2):177-82.
- Leal-Junior EC, Lopes-Martins RÁ, Bjordal JM. Clinical and scientific recommendations for the use of photobiomodulation therapy in exercise performance enhancement and post-exercise recovery: current evidence and future directions. Brazilian journal of physical therapy. 2019 Jan 1;23(1):71-5.
- Akçay H, Etoz OA, Guray E. The effects of led-mediatedphotobiomodulation therapy on newly formed bone in distraction osteogenesis. Ann Clin Anal Med. 2020;11(4):262-7.
- Kulkarni S, George R, Love R, Ranjitkar S. Effectiveness of photobiomodulation in reducing pain and producing dental analgesia: A systematic review. Lasers in Medical Science. 2022 Sep;37(7):3011-9.
- 10. Samaneh R, Ali Y, Mostafa J, Mahmud NA, Zohre R. Laser therapy for wound healing: A review of current techniques and

mechanisms of action. Biosci, Biotech Res Asia. 2015;12:217-3.

- Dantas EM, Amorim FK, Nóbrega FJ, Dantas PM, Vasconcelos RG, Queiroz LM. Clinical efficacy of fluoride varnish and lowlevel laser radiation in treating dentin hypersensitivity. Brazilian dental journal. 2016 Jan;27:79-82.
- Antognazza MR, Martino N, Ghezzi D, Feyen P, Colombo E, Endeman D, Benfenati F, Lanzani G. Shedding light on living cells. Advanced materials. 2015 Dec;27(46):7662-9.
- Klima K, Puła J, Synowiec A, Kliszcz A, Lepiarczyk A. Biomass yield and calorific value of Multiflora Rose (Rosa multiflora Thunb.) irradiated with laser beams and estimation of CO2 equivalent emission during the extensive cultivation. Journal of Biobased Materials and Bioenergy. 2019 Jun 1;13(3):424-7.
- Lopes-Martins RA, Marcos RL, Leal-Junior EC, Bjordal JM. Low-level laser therapy and world association for laser therapy dosage recommendations in musculoskeletal disorders and injuries. Photomedicine and Laser Surgery. 2018 Sep 1;36(9):457-9.

- Romagnoli E, Cafaro A. PBM. Theoretical and applied concepts of adjunctive use of LLLT/PBM within clinical dentistry. InLasers in Dentistry—Current Concepts 2017 (pp. 131-160). Springer, Cham.
- Luke AM, Mathew S, Altawash MM, Madan BM. Lasers: A review with their applications in oral medicine. Journal of lasers in medical sciences. 2019;10(4):324.

### CONTRIBUTORS

- 1. Nauman Rauf Khan Concept & Design; Data Acquisition; Drafting Manuscript; Critical Revision; Supervision; Final Approval
- 2. Anna Nauman Khan Concept & Design; Data Acquisition;
- Drafting Manuscript; Supervision; Final Approval 3. **Hira Butt** – Drafting Manuscript; Critical Revision; Supervision
- 4. Syeda Maryam Fatima Drafting Manuscript
- 5. Fatima Imran Drafting Manuscript
- 6. Laiba Amer Drafting Manuscript

April-June 2023

EVENCESE: JGMDS publishes its articles under a Creative Commons Attribution Non-Commercial Share-Alike license (CC-BY-NC-SA 4.0). COPYRIGHTS: Authors retain the rights without any restrictions to freely download, print, share and disseminate the article for any lawful purpose. It includes scholarlynetworks such as Research Gate, Google Scholar, LinkedIn, Academia.edu, Twitter, and other academic or professional networking sites.