



## First Physico-chemical and Traditional Characterisation of Lake Tseny, Northwest Madagascar

*Félicien Herbert Randrianandrianina, Ph.D student*

Doctoral School on Biodiversity and Tropical Environments,  
University of Toliara, Madagascar

*Daudet Andriafidison, Ph.D*

Institutes of Higher Education in Morondava-Menabe,  
University of Toliara, Madagascar

*Hanta Julie Razafimanahaka, Executive director*

Madagasikara Voakajy Association, Madagascar

*Jean Robertin Rasoloariniaina, Ph.D*

Institutes of Higher Education in Antsirabe-Vakinankaratra,  
University of Antananarivo, Madagascar

*Félicitée Rejo-Fienena, Titular Professor*

Doctoral School on Biodiversity and Tropical Environments,  
University of Toliara, Madagascar

Doi: [10.19044/esipreprint.8.2023.p431](https://doi.org/10.19044/esipreprint.8.2023.p431)

Approved: 24 August 2023

Posted: 28 August 2023

Copyright 2023 Author(s)

Under Creative Commons CC-BY 4.0

OPEN ACCESS

*Cite As:*

Randrianandrianina F.H., Andriafidison D., Razafimanahaka H.J., Rasoloariniaina J.R. & Rejo-Fienena F.(2023). *First Physico-chemical and Traditional Characterisation of Lake Tseny, Northwest Madagascar*. ESI Preprints.

<https://doi.org/10.19044/esipreprint.8.2023.p431>

### Abstract

This study was designed to assess the biophysical structure, traditional practices and water quality of Lake Tseny in northwest Madagascar. The lake was divided into five zones: the central zone and the littoral zones (east, north, west and south). Transects and direct observation were used to estimate submerged tree and vegetation cover. Information on local traditions was collected through interviews with key people and visits to sacred sites. Water quality was assessed during the rainy and dry season, 2017-2018. Samples were taken twice a day, in the early morning and evening, at randomly selected points at intervals of 400m. Over 75% of the lakeshore was covered by aquatic vegetation. Most of the submerged trunks

were recorded along the western area. Five sacred sites associated with eight forms of traditional taboos to reduce the potential source of water pollution were recorded in and around the lake. The lake is bowl shaped. The water was warm, slightly alkaline, well oxygenated and less mineralised. Between the rainy and dry seasons, we found a seasonal difference in water quality. Results showed spatial variation in water quality between zones. A comparison of the mean values with water quality standards showed that Lake Tseny has good water conditions for domestic purposes and suitable habitats for aquatic fauna throughout the year. These preliminary results provided reference data on the lake's characteristics. Due to the increasing human activities in the area, regular monitoring of water quality and physical characteristics should be carried out.

---

**Keyword:** Lake Tseny, Madagascar, Biophysical structure, Traditional practice, Physico-chemical parameters, Water quality

## Introduction

Human life and the survival of aquatic species depend heavily on water sources, including lakes and rivers. Water sources provide many services, such as use for domestic and agricultural purposes, refuge and natural habitat for aquatic species, and recreational opportunities. Many factors continue to threaten freshwater ecological integrity (Munir et al., 2016). In recent years, increasing human population, food demand, land conversion and fertiliser use have led to accelerated degradation of many freshwater resources (Jayakumar et al., 2009; Mushini et al., 2012). Various physicochemical parameters have been used to determine surface water quality for drinking purposes (Joshi et al., 2009; Kumar & Sinha, 2010), to analyse water pollution (Zinsou et al., 2016), and for bio-ecological and chemical relationships (Mbalassa et al., 2014; Pârvulescu et al., 2011).


In Madagascar, hydrobiological studies have assessed the physico-chemical parameters of different water sources as indicators of water body status, water quality, and the origin of the variability of water minerals (Rabemanana et al., 2005; Rasolofomanana, 2009; Rasolofonirina et al., 2018) and interspecific relationships (Rasoloariniaina et al., 2016). Lake Tseny, located in the northwestern part of the island, is a wetland of considerable importance for both human well-being and biodiversity conservation (Andriafidison et al., 2011). The lake is home to seven native and endemic fish species, one of which is classified as critically endangered *Paretroplus menarambo*. This species was thought to be extinct in the wild (Loiselle & De Rham, 2003), but was recently confirmed to occur in Lake Tseny (Andriafidison et al., 2011). The lake also harbours Madagascar's big-headed tortoise *Erymnochelys madagascariensis* and the Madagascar's fish

eagle *Haliaeetus vociferoides*. Both are critically endangered species. Lake Tseny is an important source of income and food from fishing activities and water for domestic use (e.g., drinking water, washing, bathing). The study area is known for its intense agricultural activity and overuse of chemical pesticides. It is one of the main producers of black-eyed peas in Madagascar. Due to the high demand, producers are increasing their efforts in terms of area under cultivation and the use of pesticides. No scientific study of water quality in the area has been carried out to date. Baseline and reference data on the lake's characteristics are needed to plan for biodiversity conservation and sustainable human use. This study aims to assess: (1) traditional beliefs and practices in and around the lake, (2) physical characteristics, and (3) water quality for aquatic species and human use.

## Materials and Methods

### *Study site*


Lake Tseny is located in the Sofia region, Port-Bergé district, northwest Madagascar (Figure 1). Lake Tseny covers an area of 641 hectares. The undulated water surface depends on the wind direction and the seasons. From May to October, during the dry season, the easterly 'varatraza' winds blow from east to west, pushing the water in the same direction. But in the rainy season, from November to April, the westerly 'talio' winds push it in the opposite direction (from west to east). The average annual temperature is 26.4°C. The average recorded rainfall varies from 1098 mm to 2053 mm.



**Figure 1.** Study site

### **Data collection**

*In situ* water sampling was carried out in September and December 2017 and in April and September 2018. The representative sample points were randomly selected using the transect method (CCME, 2011). However, we selected each point at 400 meters intervals using GPS by canoe. Fifty-eight (58) points were recorded and measured (Figure 2). Seasonal measurements were made at the same points. Based on local knowledges, the lake is divided into five fishing areas or zones: North, South, West, East, and Central zones (Figure 2). The boundaries of the areas were mainly determined base on the cardinal direction and indigenous considerations about the fishing places, passed from generation to generation in informal way.



**Figure 2.** Lake zones and sampling point locations

Sampling was carried out twice a day: between 05:00 - 08:00 in the morning and 15:00 - 18:00 in the afternoon. In this study, physico-chemical parameters such as water surface temperature, pH, electrical conductivity (EC) and total dissolved solids (TDS) were measured using a portable multi-parameter meter kit Apera instruments SX823-B. Water depth was measured using a depth mate portable water sounder. Dissolved oxygen (DO) and oxygen saturation (O<sub>2</sub>) were measured using an YSI's EcoSense DO200A handheld DO meter. Ammonium, nitrate and nitrite analyses were also carried out using a API Freshwater Master Test Kit. Sampling tubes were properly cleaned with water before the next analysis. In addition, environmental conditions (temperature and relative humidity) were recorded using special data loggers placed in the peninsula village in the middle of the lake.

The description of the vegetation cover in the littoral zones was carried out by measuring the macrophyte covering the shoreline. Measurement locations were randomly selected using the transect method

(CCME, 2011) during the selection of water sampling locations. However, the shoreline locations close to (approximately 20m) and perpendicular to the sampling locations were selected. Percent coverage of vegetation was estimated within 20 m horizontal line in the littoral zone using direct observation. Macrophyte percent coverage was determined according to the CCME (2011) categories:

- Absent 0%
- Sparse 1-30%
- Moderate 30-60%
- Dense 60-100%.

The pattern groups of vegetations all through the lake was also recorded using direct observation. A submerged woods assessment was carried out in October 2017, during the low water level by recording the location, number, and diameter of sunken woods using GPS and a canoe.

Key informant interviews were conducted to collect data on sacred sites and associated local taboos. Key informants include fokontany 'smallest administrative subdivision' or village leaders and elders, as well as community members who are familiar with the lake and its surroundings, taboos and traditional culture.


### ***Data processing and statistical analysis***

A normality test shown nonnormal distribution of data. Nonparametric statistical test variables was used. The Kruskal-Wallis test was used to compare variations in physico-chemical parameters between lake zones (north, south, east, west and central) and the Mann-Whitney test for seasonal differences (dry and wet) was performed in R 3.3.2. Spearman's correlation was used to determine the relationships between the physico-chemical parameters.

## **Results**

### ***Biophysical characteristics of the Lake Tseny***

The shape of Lake Tseny shows many bays, three tributary inlets (Ambalambato in the north, Begidroa in the east and Ambarijeby in the south) and one main outlet (Ambinany in the north). The Anjombony and Ambavany Ankazobe rivers are the two main tributaries of the lake. The tributaries flow into the lake mainly during the flood season and transport a large amount of organic and inorganic material. More than 75% of the littoral zone is covered by aquatic vegetation, mainly *Phragmites* sp. (POACEAE). The lake shows a spatial distribution of pattern groups of aquatic vegetation (*Phragmites* sp.) (Figure 3).



**Figure 3.** Spatial distribution of aquatic vegetation, submerged tree and island


### *Submerged trees*

A total of 115 submerged wood items were recorded at 81 sites. The topmost branches protrude from the water surface (Figure 4).



**Figure 4.** Emerged uppermost branches

The results show a spatial variation in the distribution of submerged wood within the lake. However, 82 submerged trunks were recorded at 53 sites on the western shore, followed by 22 submerged trunks at 19 sites in the central zone. In addition, 10 pieces of submerged wood were recorded at 8 sites in the north and only 1 piece in the south. No submerged wood was recorded on the east bank. The diameter of submerged tree trunks ranged from 0.1 to 0.6 m, with a mean of  $0.24 \pm 0.09$  m. A significant difference in wood diameter was found between lake zones (Kruskal-Wallis test:  $df = 3$ ,  $p < 0.05$ ). The largest submerged wood was found in the central zone and the smallest in the northern shore (Figure 5).



**Figure 5.** Variation in emerged uppermost tree trunks between lake zones

### *Taboos and traditional practices*

Five sacred sites were surveyed in and around Lake Tseny (Figure 3). These sites have high spiritual value and are the site of important traditional ceremonies. Eight main taboos were reported: 1) no pig or pork in and around the lake, 2) no menstruating women passing through the lake, 3) no defecating or urinating anywhere on the lake, 4) no motorised canoe, 5) no washing dishes in the lake, 6) no fishing on Tuesday, 7) no passing the lake while drunk, 8) be physically clean before fishing.

Key informants reported that these taboos are enforced in and at the villages around the lake. The penalty for breaking the taboo in the lake is the sacrifice of a zebu. Breaking the taboo is reported to have a supernatural sanction, such as low fishing catch and/or death by drowning.


### *Physico-chemical water characteristics*

The data (means, std.Err, min and max values) for water physico-chemical parameters are presented in Tables 1.


**Table 1.** Basic statistics summary for the general physico-chemical parameters (for all data together)

Water parameters	Mean	Std. Err	Min	Max
Depth (m), N= 232	4.04	0.09	1.4	7.4
Temperature (°C), N= 231	28.65	0.15	25.30	35.60
pH, N= 232	8.00	0.02	7.05	8.54
Electrical Conductivity: EC (µS/cm), N= 231	96.03	0.84	71.70	122.40
Total Dissolved Solid: TDS (mg/l), N= 232	68.27	0.60	51.10	84.90
Dissolved Oxygen: DO (mg/l), N= 232	7.62	0.06	4.63	9.60
Oxygen saturation: O <sub>2</sub> (%), N= 232	98.33	0.77	59.90	120.40
Nitrate: NO <sub>3</sub> <sup>-</sup> (mg/l), N= 208	0.09	0.05	0.00	5.00
Nitrite: NO <sub>2</sub> <sup>-</sup> (mg/l), N= 232	0.01	0.01	0.00	1.00
Ammonium: NH <sub>4</sub> <sup>-</sup> (mg/l), N= 232	0.23	0.01	0.00	1.00

The results from the statistical variation test in physico-chemical parameters are shown in figures 6 and 7.



**Figure 6.** Variation in physico-chemical parameters (p =Kruskal Wallis value test) between lake zones (centre, east, north, south and west)



**Figure 7.** Variation in physico-chemical parameters (p =Mann Whitney value test) between seasons (Dry and wet)

The lake showed very significant variation in water depth between the lake zones and seasons. However, the lake is bowl-shaped, significantly deepest in the central zone and shallowest in the littoral zone, especially in the eastern zone (Kruskal-Wallis test:  $\chi^2 = 31.48$ ,  $df = 4$ ,  $p < 0.0001$ ). The water was significantly deeper in the wet season than in the dry season (Mann-Whitney test:  $z = -4.16$ ,  $p < 0.0001$ ). The water level in the wet season (April) was 1m higher than in the dry season (November-December).

Water surface temperature was warm but not significantly differed between lake zones (Kruskal-Wallis test:  $\chi^2 = 4.61$ ,  $df = 4$ ,  $p > 0.05$ ), indicating uniform conditions for temperature. But, there was a significant difference between seasons (Mann-Whitney test:  $z = -5.73$ ,  $p < 0.0001$ ), with the wet season being warmer than the dry season.



The water was slightly alkaline. The pH value varied significantly between the zones of the lake (Kruskal-Wallis test:  $\chi^2 = 27.24$ ,  $df = 4$ ,  $p < 0.0001$ ). The highest pH was recorded in the central zone and the lowest in the littoral zone. There was also a significant difference in water pH between seasons (Mann-Whitney test:  $z = -5.57$ ,  $p < 0.0001$ ). However, it was higher in the dry season than in the wet season.

The results showed no spatial difference in conductivity between the lake zones (Kruskal-Wallis test:  $\chi^2 = 3.12$ ,  $df = 4$ ,  $p > 0.05$ ). Thus, water conductivity is relatively constant across the lake areas. A strong significant difference in conductivity between seasons was found (Mann-Whitney test:  $z = -11.39$ ,  $p < 0.0001$ ). Higher conductivity was recorded during the dry season than during the wet season.

There was no significant difference in TDS between the lake zones (Kruskal-Wallis test:  $\chi^2 = 3.35$ ,  $df = 4$ ,  $p > 0.05$ ). A similarity in water TDS was found throughout the lake. Our results showed highly significant variations in TDS between seasons (Mann-Whitney test:  $z = -1.39$ ,  $p < 0.0001$ ), and higher water TDS was observed in the dry season than in the wet season.

Dissolved oxygen (DO) concentration showed a spatial significant difference between the lake zones (Kruskal-Wallis test:  $\chi^2 = 23.64$ ,  $df = 4$ ,  $p < 0.0001$ ). However, DO was highest in the central lake and lowest in the littoral. There was also a significant variation in DO between seasons (Mann-Whitney test:  $z = -6.75$ ,  $p < 0.0001$ ), and it was higher in the dry season than in the wet season. Oxygen saturation ( $O_2$ ) is function of DO and showed significant variation between lake zones and seasons.

Our results showed no spatial difference in nitrogen components (nitrate, nitrite, and ammonium) between lake areas (Kruskal-Wallis test:  $\chi^2 < 7.15$  for all components,  $df = 4$ ,  $p > 0.05$ ) nor between seasons ( $z < 0.7$  for all components,  $p > 0.05$ ).

### ***Correlation between water physico-chemical parameters***


The results from Spearman's  $r$  correlation showed the correlation between physico-chemical parameters (Table 2).

**Table 2.** Correlation matrix between physico-chemical parameters of Lake Tseny: \*p < 0.0001. D: Depth (m); T°: Temperature (°C); EC: Electrical conductivity (µS/cm); TDS: Total Dissolved Solid (mg/l); DO: Dissolved oxygen (mg/l); O<sub>2</sub>: Oxygen Saturation (%); NO<sub>3</sub><sup>-</sup>: Nitrate (mg/l); NO<sub>2</sub><sup>-</sup>: Nitrite (mg/l); NH<sub>4</sub><sup>-</sup>: Ammonium (mg/l).

	D	T°	pH	CE	TSD	OD	O <sub>2</sub>	NO <sub>3</sub> <sup>-</sup>	NO <sub>2</sub> <sup>-</sup>
T°	0.018								
pH	0.021	0.001							
EC	0.074	0.002	0.034						
TDS	0.07	0.002	0.038	*0.997					
DO	0.007	0.066	*0.703	0.033	0.037				
O <sub>2</sub>	0.027	0.007	*0.743	0.025	0.029	*0.849			
NO <sub>3</sub> <sup>-</sup>	0.001	0.024	0.002	0.028	0.029	0.006	0.003		
NO <sub>2</sub> <sup>-</sup>	0.003	0.08	0.016	0.03	0.03	0.000	0.016	0.364	
NH <sub>4</sub> <sup>-</sup>	0.001	0.02	0.007	0.006	0.006	0.031	0.016	0.111	0.093

### *Fluctuations of ambient conditions*

Our results from the temperature and humidity data loggers showed the fluctuation of air conditions during the study period. The average temperature was 27.35 °C, with a minimum of 24.23 °C in July and a maximum of 30.16 °C in November (Figure 8). Relative humidity values varied throughout the year. The average relative humidity was 63.2%, with a minimum of 53.0% recorded in January. It reached a maximum of 73.4% in March



**Figure 8.** Temperature and humidity fluctuations during the present study

## **Discussion**

### *Physical habitat of lake Tseny*

We found that over 75% of the littoral zones was covered by aquatic vegetation and that there was a large amount of groups of aquatic vegetation throughout the lake. The most submerged tree trunks were recorded along the western lake. Our findings corroborate those of Andriafidison et al. (2011), who reported that the western area of the lake contained a

permanently submerged woods around which *P. menarambo* (a native and critically endangered fish species) was frequently netted. It was reported that the physico-chemical conditions of the western lake, especially around the submerged tree, influenced the habitat used by *P. menarambo* (Randrianandrianina unpublished). De Rham & Nourissat (2003) reported that the *Paretroplus* species are substratum spawners and require a solid surface to lay their eggs. In addition, Angermeier & Karr (1984) documented that many fish are sensitive to the effects of woody debris on habitat and food availability and are more closely related to the benefits of camouflage. The physical characteristics of the lake represent and provide essential habitats for aquatic species including fish.

### ***Importance of local traditions in Lake Tseny***

Lake Tseny is governed by various forms of taboos and traditional prohibitions. These prohibitions regulate human activities to minimise direct and indirect negative impacts on water quality by limiting potential anthropogenic pollution and to reduce pressure on fisheries resources by limiting the number of fishing days. In Madagascar, there has been a growing interest in integrating local traditions into the modern conservation and management plans (e.g. Lingard et al., 2003). The traditional customs in and around lake Tseny were more weakly respected than reported by Cinner (2007) and Rabearivony et al. (2008) (Randrianandrinina unpublished). Ambario village traditional authority asserted that to establish a lake uses and socio-cultural harmonies, the rite of purification of water should be accomplished, while the ceremony celebrates at the sacred site requires a zebu sacrifice (Randrianandrianina unpublished). To maintain the good water quality for aquatic life and human well-being through healthy aquatic ecosystem services, management implication should focus on development of the strategic plan taking into consideration the sociocultural aspect, involving the traditional authority.

### ***Water physico-chemical characteristics***

Due to rainfall, a significant difference in water depth was found between the seasons. The peak rainfall was in March (see Figure 8). The difference in water level in March may be greater than 1m the level in November-December due to the peak of precipitation. However, in terms of water depth, the research was biased by the inaccessibility of the study site in March due to the flood season when many rivers merged and overflowed, making access impossible.

Tables 3 show that all physicochemical parameters meet water quality standards for potability and suitable to sustain aquatic species.

**Table 3.** Standard limit values of drinking water and suitable for aquatic life

Physicochemical parameters (Present study)	Limit for drinking water		Limit suitable for aquatic life
	Acceptable limit	Restricted limit	
Temperature 25.30 to 35.60°C (28.65±0.15 °C)	25 to 30°C (WHO, 1984)	-	20°C à 32°C (Boyd, 1998; Lowe-McConnel, 1987)
pH 7.05 to 8.54 (8.00±0.02)	6.5 to 8.5 (WHO, 2011; WaterAid, 2004)	4.5 - 10 (WaterAid, 2004)	6.5 and 8.2 (Murdoch et al., 2001)
Conductivity 71.70 to 122.40 µS/cm (96.03±0.84 µS/cm)	2 000 µS/cm (WaterAid, 2004)	3 400 (WaterAid, 2004)	
Total dissolved solid 51.10 to 84.90 mg/l (68.27±.060 mg/l)	Less than 300 mg/l (OMS)	1 000 mg/l (Madagascar Code Eau, 1999)	400 mg/l (Boyd & Tucker, 1998)
Dissolved oxygen 4.63 to 9.60 mg/l (7.62±0.06 mg/l)	Minimal of 5 mg/l (Bhanja & Ajoy 2000)	-	-
Ammonium 0 to 1 mg/l (0.23±0.01 mg/l)	17 mg/l (EPA, 2013)	-	-
Nitrate 0 to 5 mg/l (0.09±0.05 mg/l)	50 mg/l (WHO, 2011; WaterAid, 2004)	100 (WaterAid, 2004)	-
Nitrite 0 to 1 mg/l (0.01±0.01 mg/l)	0,1 mg/l (WaterAid, 2004)	3 (WaterAid, 2004)	-

The lake temperature was similar to the mean air temperature at the study site (data from the data logger in this study) and was related to the difference in air temperature between seasons, which was slightly higher in the rainy period (28.58 °C) than in the dry period (28.07 °C), indicating that the water temperature is related to the ambient temperature of the study area and period (Tfeil et al., 2018). The comparison with Lake Kinkony, the northwestern lake, showed similar values varied between 29.8°C and 34.7°C (Rasoloariniaina, 2010). The constant surface temperature throughout the lake is in agreement with the results of Charrada (1992) who showed that the water surface temperature was homogeneous throughout the lake.

The alkalinity range of the lake may be due to the photosynthesis of high-density vegetation, a process that absorbs carbon dioxide (CO<sub>2</sub>) from the water and produces oxygen (Pedersen et al., 2013). As the plants remove CO<sub>2</sub>, the water becomes more alkaline and the pH rises. The pH of Lake Tseny was similar to that of Lake Kinkony ranged from 7.6 to 8.19 (Rasoloariniaina, 2010). Furthermore, the seasonal variation in pH, lower in the hot-wet period could be explained by the higher temperature, which accelerates chemical reactions and increases metabolic activities of organisms (Chapman et al., 1996; Usharani et al., 2010), consuming the dissolved oxygen in the water and releasing CO<sub>2</sub>, and decreasing the pH level. The spatial difference in the pH level, which was lower in the littoral zones, implied that the decomposition process was transforming high organic matter (plant debris from a large amount of aquatic vegetation on the

shoreline) and lowering the pH level. Higher organic matter levels are associated with lower pH levels (Taylor & Middleton, 2004).

A lower EC value could be due to the fact that the natural sediments are poor in organic matter and less affected by mineralisation processes (Delmas, 1980). Northwestern Lake Kinkony with EC values between 169 and 298  $\mu\text{S}/\text{cm}$  (Rasoloariniaina, 2010) showed higher values than Lake Tseny. Agarwal & Kannan (1996) reported that the deterioration of water quality was mainly due to the concentration of TDS. However, water with high TDS is generally of poorer drinking quality. Lower EC and TDS in the rainy season could be due to the high inflow of water during rainfall, which contributes to water dilution.

The high value of the DO content may be due to both the photosynthesis of the high density of vegetation, such a process absorbs carbon dioxide ( $\text{CO}_2$ ) from the water and produces oxygen (Pedersen et al., 2013), and the wind blowing increases the exchange of atmospheric oxygen with the water surface (Kateb et al., 2018). The DO content was higher than northwestern Kinkony Lake with values ranging from 2.89 to 5.38 mg/l (Rasoloariniaina, 2010). Spatial differences, the higher oxygen concentration in the centre of the lake could be explained by the fact that the centre zone is well aerated because such an area is connected to an open water body and the wind blowing creates turbulences and increases the exchange of atmospheric air with the water surface. Seasonal variation in oxygen concentrations that lower during the rainy season suggests that the lake was more polluted during the flood season when the Anjombony and Ambavany Ankazobe rivers flowed into the lake and run-off from agricultural land could carry chemical pesticides and organic matter into the lake. During the wet-hot season, the decomposition of organic matter reduces the solubility of oxygen in the water while the temperature increases (Lowe-McConnel, 1987).

The overall trend in the evolution of pH levels and oxygen contents, less basic and lower oxygen during the rainy period and in the littoral zones corroborates the positive relationship between both parameters, results from the present study.

### **Conclusion and recommendations**

First knowledge of the characteristics of Lake Tseny is important because it provides a baseline and reference data for a Lake Tseny wetland. The physical structure of the lake showed the spatial variation in habitat conditions favourable to sustain aquatic fauna. The sacred sites associated with prohibitions indicated the importance of traditional cultures in and around the lake. The physico-chemical parameters shown spatial and seasonal variations in water quality. Overall, the physico-chemical parameters indicated good water quality for domestic use and suitable

conditions for the life of aquatic species, including fish. The observed variations could change to unfavourable conditions due to the increase of agricultural activities in the surrounding areas and forest degradation in the study area, as well as the impact of climate change on the water system. This needs to be taken into account to ensure the conservation of critical and indigenous species and the well-being of people. In the short and long term, regular monitoring of physico-chemical parameters should be carried out to detect any changes in the variability of the parameters and to decide on appropriate management measures. With regard to human health, the presence or absence of microbial contamination should be verified. Monitoring water quality through the structure of phytoplankton and macroinvertebrate communities, which are among the best biological indicators of water quality. Further vertical studies at different times of the year are required to provide information on the water column. In terms of physical characteristics, the process of artificialisation of the lakeshore should be monitored to assess the rate of degradation of the river.

### **Acknowledgments**

We would like to thank the Fondation Tany Meva and ACA (American Cichlid Association) for funding the pinstriped damba conservation project. We would also like to acknowledge the Ministry of Agriculture, Livestock, and Fisheries for providing the research permit. We are also grateful to Idea wild for providing the equipment for water sampling. Special thanks go to Madagasikara Voakajy Association which led the lake Tseny project activities. We appreciate all contributions from our local assistants for their help during the field work and data collection.

### **References:**

1. Agarwal, G., & Kannan, G. (1996). Degradation of river to diffused activities and appropriate approach for management - A case study of river Mandakini. *J. IAEM.*, 23, 113–121.
2. Andriafidison, D., Jenkins, R., Loiselle, P., McCaskie, T., Rakotoarivelo, A., Rahalambomanana, J., Ravelomanana, T., Raminosoa, R., & Saunders, A. (2011). Preliminary fish survey of Lac Tseny in north- western Madagascar. *Madagascar Conservation & Development*, 6(2), 2–7.
3. Angermeier, P., & Karr, R. (1984). Relationships between woody debris and fish habitat in a small warmwater stream. *American Fisheries Society*, 113, 716–726.
4. Bhanja, K., & Ajoy, K. (2000). Studies on the water quality index of river Sanamachhakandana at Keonjhar Garh, Orissa, India. *Poll Res*, 19, 377–385.

5. Boyd, C. (1998). Water Quality for Pond Aquaculture. *Research and Development, ICAAE. Auburn University, USA, 43.*
6. Boyd, C., & Tucker, C. (1998). Pond aquaculture and water quality management. *Kluwer Academic Pub, London, 44–48.*
7. CCME. (2011). Manuel des protocoles d'échantillonnage pour l'analyse de la qualité de l'eau au Canada. *Le Conseil Canadien Des Ministres de l'Environnement, 211.*
8. Chapman, D., Meybeck, M., Friedrich, D., & Thomas, R. (1996). Rivers. Water quality assessments: a guide to the use of biota, sediments and water in environment monitoring. *Chapman Edition, 2nd Ed. E & FN Spon, London, 59–126.*
9. Charrada, B. (1992). Le Lac de Tunis après les aménagements. Paramètres physico-chimiques de l'eau et relation avec la croissance des macroalgues. *Marine Life, 1(1), 29–44.*
10. Cinner, J. E. (2007). The role of taboos in conserving coastal resources in Madagascar. *SPC Traditional Marine Resource Management and Knowledge Information Bulletin, 22, 15–23.*
11. De Rham, P., & Nourissat, J. (2003). Les Cichlidés Endémiques de Madagascar. *Publication Association France Cichlid, Solliès-Pont, France.*
12. Delmas, D. (1980). Minéralisation de la matière organique et échanges ioniques à l'interface eau-sédiment de l'étang de Berre, Méditerranée. *Oceanologica Acta, 3(3), 347–356.*
13. EPA. (2013). Aquatic life ambient water quality criteria for ammonia freshwater. *US Environmental Protection Agency.*
14. Jayakumar, P., Jothivel, N., Thimmappa, A., & Paul, V. (2009). Physico-chemical characterization of a lentic water body from Tamil Nadu with special reference to its pollution status. *The Ecoscan, 3(1–2), 59–64.*
15. Joshi, D., Kumar, A., & Agrawal, N. (2009). Studies on Physicochemical parameters to assess the water quality of river Ganga for drinking purpose in Haridwar district. *RASAYAN J. Chem., 2(1), 195–203.*
16. Kateb, A., Rüggeberg, A., Stalder, C., Neururer, C., & Spezzaferrri, S. (2018). High-resolution monitoring of water temperature and oxygen concentration in Lake Murten (Switzerland). *Swiss Journal of Geosciences, 5, 501–510.*
17. Kumar, N., & Sinha, D. (2010). Drinking water quality management through correlation studies among various physico-chemical parameters: A case study. *International Journal of Environmental Sciences, 1(2), 253–259.*
18. Lingard, M., Raharison, N., Rabakonandrianina, E., Rakotoarisoa, J.,

- & Elmqvist, T. (2003). The Role of Local Taboos in Conservation and Management of Species: The Radiated Tortoise in Southern Madagascar. *Conservation and Society*, 2(1), 223–246.
19. Loiselle, P., & De Rham, P. (2003). Paretroplus, Damba. In: *The Natural History of Madagascar*. S. M. Goodman and J. P. Benstead (Eds.), The University of Chicago Press. 877–878.
20. Lowe-McConnel, R. (1987). Ecological studies in fish communities. *Cambridge University Press*.
21. Madagascar Code Eau. (1999). Loi N°98–029. *Journal Officiel de La République de Madagascar*, 2557–2275.
22. Mbalassa, M., Bagalwa, J., Nshombo, M., & Kateyo, M. (2014). Assessment of physicochemical parameters in relation with fish ecology in Ishasha River and Lake Edward, Albertine Rift Valley, East Africa. *International Journal of Current Microbiology and Applied Sciences*, 3(6), 230–244.
23. Munir, T., Hussain, M., & Naseem, S. (2016). Water Pollution-A Menace of Freshwater Biodiversity: A Review Water Pollution-A Menace of Freshwater Biodiversity: A Review. *Journal of Entomology and Zoology*, 4(4), 578–580.
24. Murdoch, T., Cheo, M., & O’Laughlin, K. (2001). *Stream keeper’s Field Guide: Watershed Inventory and Stream Monitoring Methods*. Adopt-A-Stream Foundation, Everett, WA. 297.
25. Mushini, R., Dhilleswara, R., & Bethapudi, S. (2012). Assessment of quality of drinking water at Srikurmam in Srikakulam district, Andhra Pradesh, India. *International Research Journal of Environment Sciences*, 1(2), 13–20.
26. Pârvolescu, L., Pacioglu, O., & Hamchevici, C. (2011). The assessment of the habitat and water quality requirements of the stone crayfish (*Austropotamobius torrentium*) and noble crayfish (*Astacus astacus*) species in the rivers from the Anina Mountains (SW Romania). *Knowl. Managt. Aquatic Ecosyst*, 403(03), 1–12.
27. Pedersen, O., Colmer, T., & Sand-Jensen, K. (2013). Underwater photosynthesis of submerged plants - Recent advances and methods. *Frontiers in Plant Science*, 4, 1–19.
28. Rabearivony, J., Fanameha, E., Mampandra, J., & Thorstrom, R. (2008). Taboos and social contracts: Tools for ecosystem management – lessons from the Manambolomaty Lakes RAMSAR site, western Madagascar. *Madagascar Conservation & Development*, 3(1), 7–16.
29. Rabemanana, V., Violette, S., Marsily, G. de, Robain, H., Deffontaines, B., Andrieux, P., Bensimon, M., & Parriaux, A. (2005). Origin of the high variability of water mineral content in the bedrock



- aquifers of Southern Madagascar. *Journal of Hydrology*, 310, 143–156.
30. Rasoloariniaina, J. (2010). *Evaluation de stock et structure du peuplement ichthyologique des lacs Kinkony, Maliolio et Tsiambara dans la région de Boeny-Madagascar* (Issue March). Memoire de DEA. Option : Biologie, Ecologie et Conservation Animales, Faculté des Sciences, Université d'Antananarivo.
  31. Rasoloariniaina, J., Ganzhorn, J., Riemann, J., & Raminosoa, N. (2016). Water quality and biotic interaction of two cavefish species : *Typhleotris madagascariensis* Petit, 1933 and *Typhleotris mararybe* Sparks & Chakrabarty, 2012, in the Mahafaly Plateau groundwater system , Madagascar. *Subterranean Biology*, 18, 1–16.
  32. Rasolofomanana, L. (2009). *Characterization of Ranomafana Lake water quality - Antsirabe Madagascar*. Master's Thesis. Study program/specialization: Offshore engineering/Environmental control, Faculty of Science and Technology, University of Stavanger.
  33. Rasolofonirina, M., Ramaroson, V., & Ravelomanantsoa, S. (2018). Assessment of Nitrate Occurrence in the Shallow Groundwater of Merimandroso Area, Analamanga Region, Madagascar Using Multivariate Analysis. *American Journal of Water Resources*, 6(1), 39–47.
  34. Taylor, J., & Middleton, A. (2004). Compariosn of litter decomposition in a natural versus coal-Slurry pond reclaimed as a wetland. *Land Degrad. Develop.*, 15, 439–446.
  35. Tfeil, H., Mahfoudh, M., Mhamed, B., Aliyen, A., Yarba, L., & Hmeyada, A. (2018). Caractérisation Physico-Chimique Des Eaux De Surface Et Étude De La Diversité Ichthyologique De Quelques Zones Humides Continentales En Mauritanie Baba Aïnina Moulay Mhamed. *European Scientific Journal*, 14(6), 83–101.
  36. Usharani, K., Umarani, K., Ayyasamy, K., Lakshmanaperumalsamy, P., & Andshanthi, K. (2010). Physico-chemical and bacteriological characteristics of Noyyal River and ground water quality of Perur, India. *J Appl Sci Environ Manag*, 14, 29–35.
  37. WaterAid. (2004). Politique de la qualité de l'eau. *Wateraid Madagascar*, 3.
  38. WHO. (1984). *Guidelines for Drinking Water Quality. Health Criteria and other Supporting Information. 1*, Geneva, Switz.
  39. WHO. (2011). *Guidelines for drinking-water quality. 4th Ed*, World Health Organization.
  40. Zinsou, L., Attingli, H., Gnohossou, P., Adandedja, D., & Laleye, P. (2016). Caractéristiques physico-chimiques et pollution de l'eau du delta de l'Oueme au Benin. *J. Appl. Biosci.*, 97, 9163–9173