

University of Mississippi

eGrove

Faculty and Student Publications

Pharmacy, School of

1-1-2022

Cryptococcosis in a patient with multiple myeloma receiving pomalidomide: a case report and literature review

Daniel B. Chastain
University of Georgia

Sahand Golpayegany
University of Georgia

Andrés F. Henao-Martínez
University of Colorado Anschutz Medical Campus

Brittany T. Jackson
The Mount Sinai Hospital

Follow this and additional works at: https://egrove.olemiss.edu/pharmacy_facpubs



Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Chastain, D. B., Golpayegany, S., Henao-Martínez, A. F., Jackson, B. T., Stoudenmire, L. L., Bell, K., Stover, K. R., & Franco-Paredes, C. (2022). Cryptococcosis in a patient with multiple myeloma receiving pomalidomide: A case report and literature review. *Therapeutic Advances in Infectious Disease*, 9, 204993612211126. <https://doi.org/10.1177/20499361221112639>

This Article is brought to you for free and open access by the Pharmacy, School of at eGrove. It has been accepted for inclusion in Faculty and Student Publications by an authorized administrator of eGrove. For more information, please contact egrove@olemiss.edu.

Cryptococcosis in a patient with multiple myeloma receiving pomalidomide: a case report and literature review

Daniel B. Chastain¹, Sahand Golpayegany, Andrés F. Henao-Martínez²,
Brittany T. Jackson, Laura Leigh Stoudenmire, Kaye Bell, Kayla R. Stover³
and Carlos Franco-Paredes⁴

Abstract: While overall survival with multiple myeloma (MM) has improved, patients suffer from overwhelming tumor burden, MM-associated comorbidities, and frequent relapses requiring administration of salvage therapies. As a result, this vicious cycle is often characterized by cumulative immunodeficiency stemming from a combination of disease- and treatment-related factors leading to neutropenia, T-cell deficiency, and hypogammaglobulinemia. Infectious etiologies differ based on the duration of MM and treatment-related factors, such as number of previous treatments and cumulative dose of corticosteroids. Herein, we present the case of a patient who was receiving pomalidomide without concomitant corticosteroids for MM and was later found to have cryptococcosis, as well as findings from a literature review. Most cases of cryptococcosis are reported in patients with late-stage MM, as well as those receiving novel anti-myeloma agents, such as pomalidomide, in combination with corticosteroids or following transplantation. However, it is likely cryptococcosis may be underdiagnosed in this population. Due to the cumulative immunodeficiency present in patients with MM, clinicians must be suspicious of cryptococcosis at any stage of MM.

Keywords: *Cryptococcus*, malignancy, multiple myeloma, pomalidomide, thalidomide

Received: 25 February 2022; revised manuscript accepted: 23 June 2022.

Introduction

Cryptococcus spp. is an encapsulated yeast and opportunistic pathogen that has historically been associated with T-cell immunodeficiency due to untreated HIV infection.^{1,2} However, as the number of immunodeficient hosts increases, the incidence of cryptococcosis continues to grow in patients with malignancies, organ transplantation, autoimmune diseases, primary immunodeficiency syndromes, chronic comorbidities such as diabetes mellitus and sarcoidosis, and in those receiving immunosuppressive therapies.^{1–5}

Indeed, multiple myeloma (MM) is associated with impaired immunologic function resulting in increased susceptibility to infections.⁶ Specifically,

immunodeficiency observed in patients with MM is often multifactorial due to mucosal barrier defects, respiratory dysfunction, hypogammaglobulinemia, disrupted lymphocyte function, and treatment-related immunodeficiency.⁶ Although most patients with MM respond to initial therapies, many cases progress as conventional treatment options are not curative. Patients who relapse or develop refractory disease suffer from shorter periods of remission and more treatment-related adverse events.

Pomalidomide, a more potent and less toxic derivative of thalidomide, was introduced for patients with MM refractory to lenalidomide and bortezomib. Although infectious diseases have

Ther Adv Infect Dis

2022, Vol. 9: 1–10

DOI: 10.1177/
20499361221112639

© The Author(s), 2022.
Article reuse guidelines:
sagepub.com/journals-
permissions

Correspondence to:

Daniel B. Chastain
Clinical Associate
Professor, Department of
Clinical & Administrative
Pharmacy, College of
Pharmacy, University of
Georgia, 1000 Jefferson
Street, Albany, GA 31701,
USA.
daniel.chastain@uga.edu

Sahand Golpayegany
Department of Clinical &
Administrative Pharmacy,
College of Pharmacy,
University of Georgia,
Albany, GA, USA

Andrés F. Henao-Martínez
Division of Infectious
Diseases, University of
Colorado, Aurora, CO, USA

Brittany T. Jackson
Department of Pharmacy,
The Mount Sinai Hospital,
New York, NY, USA

Laura Leigh Stoudenmire
Department of Pharmacy,
Phoebe Putney Memorial
Hospital, Albany, GA, USA

Kaye Bell
Department of
Microbiology, Phoebe
Putney Memorial Hospital,
Albany, GA, USA

Kayla R. Stover
Department of Pharmacy
Practice, School of
Pharmacy, University of
Mississippi, Jackson, MS,
USA

Carlos Franco-Paredes
Division of Infectious
Diseases, University of
Colorado, Aurora, CO, USA
Hospital Infantil de México,
México City, México

been reported in patients receiving pomalidomide, most are viral or bacterial and are often observed when pomalidomide is administered in combination with dexamethasone.^{7,8} Herein, we present the case of a patient who was receiving pomalidomide without concomitant corticosteroids for MM and was later found to have cryptococcosis. In addition, we present findings from a literature review detailing reports of cryptococcosis in patients with MM.

Case description

A gentleman in his early 70s was admitted due to right arm and right leg weakness and paresthesia for 2 weeks. He was previously able to walk independently but was unable to bear weight on his right leg for the last 2 weeks. Other than right-sided weakness, his physical and neurological examination were unremarkable whereby he specifically denied dysarthria, facial paralysis, chest pain, fever, chills, shortness of breath, nausea, vomiting, or trauma in relation to his leg.

He had a history of MM, hypertension, atrial fibrillation, heart failure with reduced ejection fraction (EF 30% approximately 1 year earlier), automatic implantable cardioverter-defibrillator (AICD) placement, and chronic kidney disease (CKD) stage IV most likely due to a combination of MM and hypertension. He was diagnosed with stage III light chain myeloma in 2012. Notably, he did not undergo stem cell transplantation and was managed with maintenance therapy. Due to disease progression and intolerability with previous regimens, he was receiving palliative pomalidomide without corticosteroids, his fifth-line therapy, for more than 2 years with normocellular bone marrow and no evidence of residual myeloma at the time of admission (additional details of MM treatment history are provided in Table 1).

Chemistry and hematology laboratories revealed hyperkalemia [5.5 mEq/L (reference range, 3.5–5.0 mEq/L)] and transaminitis [aspartate aminotransferase (AST) 68 U/L (reference range, 10–40 U/L), alanine aminotransferase (ALT) 81 U/L (reference range, 10–40 U/L), alkaline phosphatase 79 U/L (reference range, 30–120 U/L)], but were otherwise consistent with his baseline (Table 2). He tested positive for SARS-CoV-2 *via* nasopharyngeal swab for which he underwent chest radiograph (CXR) noting ill-defined bilateral perihilar and lower lobe

infiltrates, slightly greater on the left than on the right. Computed tomography (CT) of the head without intravenous contrast revealed extensive white matter disease likely related to small vessel occlusive disease. Magnetic resonance imaging (MRI) of the brain was unable to be performed due to incompatibility with his AICD. Chest CT without intravenous contrast revealed non-specific patchy ground glass opacities throughout upper and lower lobes bilaterally and in the right middle lobe. Serologic testing was negative for hepatitis B virus (HBV) surface antigen and core antibody, as well as hepatitis C virus (HCV) antibody.

Upon admission, the neurologic complaints were attributed to a subacute cerebrovascular accident for which neurology recommended conservative management. In addition, pomalidomide was held pending COVID-19 resolution. He was considered not to be a candidate for remdesivir due to his CKD but was started on 6 mg of oral dexamethasone once daily (Figure 1). In addition, ceftriaxone and azithromycin were administered for 7 days due to concerns for a bacterial pneumonia. His oxygen demand continued to increase over the following 5 days requiring transfer to the intensive care unit (ICU) for receipt of high-flow nasal cannula (HFNC) and blood and urine cultures were obtained. Over the next 3 days, he remained on HFNC in no distress. Hematology/oncology was consulted due to his ongoing lymphopenia and recommended initiation of cefepime and micafungin for as long as he remained on dexamethasone. On day 4 of his ICU admission, his urine culture was noted to be sterile, but Gram stain of his blood cultures revealed yeast. This prompted an infectious diseases consultation resulting in continuation of micafungin and repeat blood cultures ordered for the following morning. Shortly thereafter, he experienced ventricular tachycardia and continued to deteriorate, ultimately expiring 24 hours later. Post-mortem examination was not performed.

Yeasts, identified from both sets of blood cultures, grew on Sabouraud Dextrose Agar (Emmons modification) and were identified as *Cryptococcus neoformans* *via* Thermo Scientific™ RapID™ Yeast Plus System (RapID) (Thermo Fisher Scientific, Lenexa, Kansas, U.S.) post-mortem. Due to limited capacity in the facility's microbiology laboratory, molecular identification and antifungal susceptibility testing were not performed.

Table 1. Reports of cryptococcosis in patients with multiple myeloma.

Case	Age (years)	Sex	Comorbidities	MM history prior to diagnosis of cryptococcosis	Primary site(s) of cryptococcosis	Initial management
Current case	70s	M	Atrial fibrillation, HFrEF, renal dysfunction, COVID-19	<ul style="list-style-type: none"> Stage III MM diagnosed 8 years earlier Initially treated with bortezomib, thalidomide, and dexamethasone but discontinued Subsequently received bortezomib, cyclophosphamide, and dexamethasone, but discontinued due to progression Then treated with carfilzomib, but was discontinued due to intolerance Received pomalidomide and dexamethasone with good response, but dexamethasone was discontinued for undocumented reasons Stable on palliative pomalidomide for more than 2 years with normocellular bone marrow and no evidence of residual myeloma 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from blood cultures 	<ul style="list-style-type: none"> <i>C. neoformans</i> identified post-mortem
Karnad et al. ⁹	26	M	Newly diagnosed HIV at initial diagnosis of MM	<ul style="list-style-type: none"> MM diagnosed 16 months prior following presentation with 4 × 4 cm painless lump on R parieto-occipital aspect of skull with 5 × 5 cm irregular lytic bone lesion underlying the mass Previous treatments for MM limited to localized radiation therapy 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from CSF culture 	<ul style="list-style-type: none"> Clinical improvement following treatment with amphotericin B Started vincristine, adriamycin, and dexamethasone for rapid nature of MM but expired 4 weeks later due to sustained hypotension
Mendpara et al. ¹⁰	42	F	HIV non-reactive, otherwise NR	<ul style="list-style-type: none"> Stage III-B MM (IgGκ) initially treated with vincristine, doxorubicin, and dexamethasone with good response Underwent SCT without complications for 4 months 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from CSF culture CrAg detected in CSF 	<ul style="list-style-type: none"> Clinical improvement following treatment with amphotericin B and flucytosine
Fickweiler et al. ¹¹	61	F	NR	<ul style="list-style-type: none"> Initially treated with doxorubicin and dexamethasone followed by high-dose melphalan and SCT Relapsed and started on cyclophosphamide and thalidomide, but discontinued due to pancytopenia Initiated dexamethasone and thalidomide, but discontinued due to polyneuropathy Restarted melphalan followed by re-infusion of SCT 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from CSF and blood cultures with identification of a cerebellar lesion with mass effect on the fourth ventricle causing hydrocephalus 	<ul style="list-style-type: none"> Clinical improvement following treatment with amphotericin B and flucytosine
De Oliveira et al. ¹²	58	M	NR	<ul style="list-style-type: none"> MM diagnosed 4 years earlier followed by BMT 1 year later Relapsed and started on cyclophosphamide and dexamethasone 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from an ulcerated plaque on the nasal dorsum CSF and blood cultures sterile 	<ul style="list-style-type: none"> Clinical improvement following treatment with oral fluconazole
Cerrati et al. ¹³	60s	M	NR	<ul style="list-style-type: none"> MM diagnosed several years earlier followed by radiation therapy and SCT Stable on lenalidomide and dexamethasone 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from an ulcerated plaque on the nasal dorsum 2 weeks later <i>C. neoformans</i> isolated from CSF culture 	<ul style="list-style-type: none"> Initially started on fluconazole but changed to amphotericin B plus flucytosine due to dissemination resulting in clinical improvement

(Continued)

Table 1. (Continued)

Case	Age (years)	Sex	Comorbidities	MM history prior to diagnosis of cryptococcosis	Primary site(s) of cryptococcosis	Initial management
Suner and Mathis ¹⁴	77	F	NR	<ul style="list-style-type: none"> Stage III MM (IgA-λ) diagnosed 8 years earlier After multiple relapses, started on corticosteroids, cyclophosphamide, and pomalidomide 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from CSF culture 	<ul style="list-style-type: none"> Started on amphotericin B plus flucytosine but expired 10 days later
Ferraro et al. ¹⁵	64	M	NR	<ul style="list-style-type: none"> MM diagnosed 12 years earlier Stable on bortezomib, lenalidomide, and dexamethasone 	<ul style="list-style-type: none"> Biopsy of 1.7 × 1.0 × 1.3 cm soft-tissue mass in the L sphenoid sinus revealed <i>Cryptococcus</i> spp. CrAg detected in serum, but not in CSF 	<ul style="list-style-type: none"> Clinical improvement following treatment with amphotericin B and flucytosine
Bowcock et al. ¹⁶	75	F	HIV non-reactive, otherwise NR	<ul style="list-style-type: none"> Bortezomib, cyclophosphamide, and dexamethasone as fourth line treatment 	<ul style="list-style-type: none"> CrAg detected in CSF and serum 	<ul style="list-style-type: none"> Clinical improvement following treatment with amphotericin B and flucytosine, but MM relapsed approximately 10 weeks later and expired shortly thereafter
Bowcock et al. ¹⁶	79	M	NR	<ul style="list-style-type: none"> Pomalidomide and dexamethasone as sixth line treatment 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from blood cultures 	<ul style="list-style-type: none"> Clinical improvement following treatment with amphotericin B and flucytosine, but expired 5 weeks later due to progression of MM
Sato et al. ¹⁷	62	F	Renal dysfunction	<ul style="list-style-type: none"> Stage III MM (IgG-λ) initially treated with bortezomib and dexamethasone but discontinued due to diarrhea Subsequently started on daratumumab, lenalidomide, and dexamethasone 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from CSF and blood cultures 	<ul style="list-style-type: none"> Started on amphotericin B plus flucytosine but expired approximately 2 weeks later
Stepman et al. ¹⁸	70	M	Diabetes mellitus, hypertension, renal dysfunction	<ul style="list-style-type: none"> MM diagnosed several years earlier followed by BMT Subsequently started on ixazomib, pomalidomide, and prednisone 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from CSF culture 	<ul style="list-style-type: none"> Clinical improvement following treatment with amphotericin B and flucytosine
Sassine et al. ¹⁹	77	M	NR	<ul style="list-style-type: none"> Stable on lenalidomide 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from intraoperative culture obtained from R femur CrAg detected in serum, but not CSF CSF and blood cultures sterile 	<ul style="list-style-type: none"> Clinical improvement following treatment with amphotericin B
Raheem et al. ²⁰	60	M	Hepatic cirrhosis	<ul style="list-style-type: none"> Initially treated with cyclophosphamide, bortezomib, and dexamethasone for 4 cycles, but changed to pomalidomide and dexamethasone 	<ul style="list-style-type: none"> <i>C. neoformans</i> isolated from CSF culture CrAg detected in CSF 	<ul style="list-style-type: none"> Clinical improvement following a loading dose of fluconazole, then amphotericin B and flucytosine

BMT, bone marrow transplant; CrAg, cryptococcal antigen; CSF, cerebrospinal fluid; F, female; HFREF, heart failure with reduced ejection fraction; L, left; M, male; MM, multiple myeloma; NR, not reported; R, right; SCT, stem cell transplantation.

Table 2. Laboratory values prior to and during hospitalization.

Laboratory parameter	Approximately 2 months prior to hospital admission	Approximately 1 month prior to hospital admission	Day of hospital admission	Day 3 of hospitalization	Day 6 of hospitalization	Day 9 of hospitalization	Day 12 of hospitalization
Serum creatinine, mg/dl	3.67	3.67	3.89	3.33	4.15	3.1	3.26
Blood urea nitrogen, mg/dl	36	41	56	75	104	99	113
White blood cell count, K/ μ l	2.4	3.0	2	2.6	1.6	4.0	13.7
Neutrophil count, K/ μ l	1.3	1.5	1.5	1.8	1.1	3.3	12.9
Lymphocyte count, K/ μ l	0.8	1.0	0.3	0.3	0	0.2	0.3

Baseline laboratory data are presented in relation to the day of hospitalization and include the following: BUN, blood urea nitrogen (range: 8–20 mg/dl); lymphocyte count (range: 1–4 K/ μ l); neutrophil count (range: 2.5–8 K/ μ l); SCR, serum creatinine (range: 0.70–1.30 mg/dl); WBC, white blood cell (range: 3.5–10 K/ μ l).


**Figure 1.** Timeline of hospitalization.

Figure depicts daily (D) timeline of hospitalization.

*Although blood cultures were obtained on D6 and D11, yeasts were not identified on Gram stain until 4 days later in both cases. The yeasts were then plated to Sabouraud Dextrose Agar [Emmons modification] and were eventually identified as *C. neoformans* via Thermo Scientific™ RapID™ Yeast Plus System [RapID] (Thermo Fisher Scientific, Lenexa, Kansas, U.S.). *C. neoformans* was identified after the patient expired.

Additional workup for cryptococcosis [e.g. cerebrospinal fluid (CSF) examination, cryptococcal antigen (CrAg) lateral flow assay (LFA)] was not performed due to low suspicion of *C. neoformans* in this patient.

Discussion

Patients with MM are at an increased risk for infectious diseases due to impaired quantity and function of humoral and cell-mediated immunity.⁶ In

addition, the degree of immunodeficiency observed in these patients varies from the time of initial MM diagnosis. As expected, infectious etiologies differ based on the duration of MM and treatment-related factors, such as the number of previous treatments and cumulative dose of corticosteroids.^{6,21} Compared with bacterial and viral infections, which occur during the first year and again between years 4 and 9 after MM diagnosis, invasive fungal infections occur at much later stages of MM and are associated with cumulative




Figure 2. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram.

immunodeficiency.^{21,22} Invasive aspergillosis is the most common invasive fungal infection among patients with MM due to the administration of high-dose corticosteroids and neutropenia.^{22–24} As much of the focus on invasive fungal infections in patients with MM remains on invasive aspergillosis, limited data are available about cryptococcosis in MM.

A PubMed literature search performed on 21 December 2021 using ‘cryptococcus’ or ‘cryptococcosis’ and ‘multiple myeloma’ revealed 37 results (Figure 2).²⁵ Of those providing patient

level details, 12 reports accounting for 13 patients were included (Table 1).^{9–20} Median age was 63 years (range, 26–76 years) and 69% ($n = 9$) were men. Medical history was reported for few patients, but included one patient with HIV;⁹ one with renal dysfunction;¹⁷ one with diabetes mellitus, hypertension, and diabetic nephropathy;¹⁸ and one with hepatic cirrhosis,²⁰ in addition to our case with atrial fibrillation, heart failure with reduced EF, and renal dysfunction. Two patients with no reported medical history underwent HIV testing which was non-reactive.^{16,10}

Duration of MM was reported in few cases, but was diagnosed a median of 8 years (range, 1.33–12 years) prior to diagnosis of cryptococcosis.^{9,12,14,15} Thirty-six percent ($n=5$) of patients had previously undergone transplantation for MM prior to diagnosis of cryptococcosis.^{10,11–13,18} While 86% ($n=12$) of patients were receiving anti-MM therapies, 43% ($n=6$) had received more than one previous treatment line with one patient receiving their fourth-line treatment,¹⁶ our patient receiving his fifth-line treatment, and another receiving their sixth-line treatment¹⁶ at the time of cryptococcosis was diagnosed. The most common anti-MM therapies included corticosteroids (67%, $n=8/12$), lenalidomide (33%, $n=4/12$), pomalidomide (33%, $n=4/12$), cyclophosphamide (25%, $n=3/12$), and bortezomib (17%, $n=2/12$). Notably, 75% ($n=6/8$) of patients receiving lenalidomide or pomalidomide were receiving concomitant corticosteroids, while our patient being treated with pomalidomide and one other receiving lenalidomide¹⁹ were not receiving concomitant corticosteroids.

Disseminated cryptococcosis was present in 93% ($n=13$) of patients, of which *Cryptococcus* spp. was detected in the CSF of 69% ($n=9/13$) of patients *via* microbiologic identification in 89% ($n=8/9$). Notably, one patient was found to have a cerebellar lesion with mass effect on the fourth ventricle causing hydrocephalus presumed to be a parenchymal cryptococcoma.¹¹ In addition, two patients presented with cutaneous cryptococcosis involving the nasal dorsum,^{12,13} one patient had osseous cryptococcosis of the right femur,¹⁹ while one patient had cryptococcal sinusitis involving the left sphenoid sinus.¹⁵ *C. neoformans* was identified as the causative organism in all cases, except for one where the diagnosis of cryptococcosis was established based on detection of cryptococcal antigenemia.¹⁶

Amphotericin B and flucytosine were most commonly used in the initial management of cryptococcosis [86% ($n=12$) and 71% ($n=10$), respectively]. Fluconazole monotherapy was administered to one patient with cutaneous cryptococcosis as CSF and blood cultures were sterile.¹² Clinical improvement was observed in 79% ($n=11$) of patients following initiation of antifungal therapy. However, 43% ($n=6$) of patients expired,^{9,14,16,17} which was most often attributed to progression of MM.^{9,16} Our patient had been

previously started on micafungin due to administration of dexamethasone for COVID-19, but *C. neoformans* was not identified until after he expired.

While invasive fungal infections have previously been associated with a prolonged duration of MM and receipt of concomitant corticosteroids resulting in a cumulative immunodeficiency, this might not be generalizable to cryptococcosis. In two of the excluded case reports,^{26,27} diagnosis of cryptococcosis prompted searches for an underlying cause of immunodeficiency leading to subsequent diagnosis of MM, which suggests that cryptococcosis might not be restricted to late-stage MM. Furthermore, 43% ($n=6/14$) of all patients included in this review were not receiving concomitant corticosteroids for MM prior to diagnosis of cryptococcosis. While our patient was receiving dexamethasone as part of the medical management of COVID-19, he had only received 5 days of therapy prior to growth of *C. neoformans*. Although unclear, even short courses of corticosteroids could have triggered an opportunistic fungal reactivation.

Notably, cases of *C. neoformans* in patients with COVID-19 have been reported.²⁸ The precise mechanism remains unknown, but likely involves SARS-CoV-2 infection, immunomodulatory therapy, or a combination of both. We suspect our patient had disseminated cryptococcosis due to MM-associated immunodeficiency prior to diagnosis of COVID-19 or receipt of dexamethasone. Whether the immunomodulatory activity of pomalidomide contributed to our patient's risk for cryptococcosis is unknown.

Pomalidomide, a derivative of thalidomide, inhibits tumor necrosis factor (TNF)- α and is associated with increased anti-MM activity compared with thalidomide.^{29,30} In a mouse model study of *C. neoformans*, administration of neutralizing TNF- α antiserum decreased recruitment of CD4 cells, neutrophils, and macrophages by 65%, 84%, and greater than 98%, respectively.³¹ Furthermore, mice treated with neutralizing TNF- α antiserum were unable to eradicate *C. neoformans* from their lungs or prevent cerebral or splenic dissemination. Eliminating TNF- α leads to impaired activation of CD4 cells, which are largely responsible for the cell-mediated immune response against *C. neoformans*.^{32,33} Alternatively,

thalidomide and thalidomide analogues promote T-cell proliferation, primarily the CD8+ subset, and increase production of interleukin (IL)-2 and interferon (INF)- γ .^{34,35} Previous data suggest no increased risk of infection with thalidomide and thalidomide analogues unless combined with corticosteroids.^{35–37} However, downregulation of host immune response due to receipt of corticosteroids and COVID-19-associated lymphopenia may have also contributed to reactivation of previously latent cryptococcosis.^{38,39}

Although an optimal antifungal treatment regimen has yet to be identified in MM with cryptococcosis, most patients should receive combination therapy with amphotericin B and flucytosine until the extent of disease is determined (e.g. meningoencephalitis, disseminated, or severe pulmonary disease), based on efficacy data derived from persons living with HIV and solid organ transplant recipients.^{40,41} In resource-limited settings, high-dose fluconazole with or without amphotericin B might be most efficacious due to lack of availability and prohibitive costs associated with flucytosine. Induction therapy should be continued for 4 weeks or longer depending on the extent of disease. Patients should then be transitioned to higher doses of fluconazole for a minimum of 8 weeks as consolidation therapy followed by a lower dose for at least 1 year or perhaps lifelong as the underlying immunodeficiency in patients with MM is unlikely to resolve.

Conclusion

Most cases of cryptococcosis are reported in patients with late-stage MM, as well as those receiving novel anti-myeloma agents, such as pomalidomide, in combination with corticosteroids or following transplantation. However, it is likely cryptococcosis may be underdiagnosed in this population. Due to the cumulative immunodeficiency present in patients with MM, clinicians must be suspicious of cryptococcosis at any stage of MM.

Declarations

Ethics approval and consent to participate

Our report did not require ethical board approval as it described the treatment of a single patient which does not meet the federal definition of

human subjects research. This anonymized case was documented in the context of routine care and the information presented was anonymized in accordance with the Declaration of Helsinki.

Consent for publication

Consent was unable to be obtained as the patient is deceased and relatives were not contactable as contact information was no longer available at the time this case report was developed. As such, patient details have been anonymized as much as possible.

Author contributions

Daniel B. Chastain: Conceptualization; Data curation; Visualization; Writing – original draft; Writing – review & editing.

Sahand Golpayegany: Writing – original draft; Writing – review & editing.

Andrés F. Henao-Martínez: Conceptualization; Supervision; Writing – review & editing.

Brittany T. Jackson: Writing – review & editing.

Laura Leigh Stoudenmire: Writing – review & editing.

Kaye Bell: Writing – review & editing.

Kayla R. Stover: Writing – review & editing.

Carlos Franco-Paredes: Conceptualization; Supervision; Writing – review & editing.

Acknowledgements

None.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

Competing interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Availability of data and materials

All data are available as part of this article.

ORCID iDs

Daniel B. Chastain  <https://orcid.org/0000-0002-4018-0195>

Andrés F. Henao-Martínez  <https://orcid.org/0000-0001-7363-8652>

Kayla R. Stover  <https://orcid.org/0000-0002-8635-0137>

Carlos Franco-Paredes  <https://orcid.org/0000-0001-8757-643X>

References

- Elsegeiny W, Marr KA and Williamson PR. Immunology of cryptococcal infections: developing a rational approach to patient therapy. *Front Immunol* 2018; 9: 651.
- Pappas PG, Perfect JR, Cloud GA, *et al.* Cryptococcosis in human immunodeficiency virus-negative patients in the era of effective azole therapy. *Clin Infect Dis* 2001; 33: 690–699.
- Archuleta S, Gharamti AA, Sillau S, *et al.* Increased mortality associated with uncontrolled diabetes mellitus in patients with pulmonary cryptococcosis: a single US cohort study. *Ther Adv Infect Dis* 2021; 8: 20499361211004367.
- George IA, Santos CAQ, Olsen MA, *et al.* Epidemiology of cryptococcosis and cryptococcal meningitis in a large retrospective cohort of patients after solid organ transplantation. *Open Forum Infect Dis* 2017; 4: ofx004.
- Lin YY, Shiao S and Fang CT. Risk factors for invasive *Cryptococcus neoformans* diseases: a case-control study. *PLoS ONE* 2015; 10: e0119090.
- Pratt G, Goodyear O and Moss P. Immunodeficiency and immunotherapy in multiple myeloma. *Br J Haematol* 2007; 138: 563–579.
- Lacy MQ and McCurdy AR. Pomalidomide. *Blood* 2013; 122: 2305–2309.
- Dimopoulos MA, Leleu X, Palumbo A, *et al.* Expert panel consensus statement on the optimal use of pomalidomide in relapsed and refractory multiple myeloma. *Leukemia* 2014; 28: 1573–1585.
- Karnad AB, Martin AW, Koh HK, *et al.* Nonsecretory multiple myeloma in a 26-year-old man with acquired immunodeficiency syndrome, presenting with multiple extramedullary plasmacytomas and osteolytic bone disease. *Am J Hematol* 1989; 32: 305–310.
- Mendpara SD, Ustun C, Kallab AM, *et al.* Cryptococcal meningitis following autologous stem cell transplantation in a patient with multiple myeloma. *Bone Marrow Transplant* 2002; 30: 259–260.
- Fickweiler W, Aries MJ, Enting RH, *et al.* Cryptococcal cerebellitis after chemotherapy and autologous stem cell re-infusion in a patient with multiple myeloma. *J Neurol* 2009; 256: 145–146.
- De Oliveira R, Barbosa L, Jeunon T, *et al.* Ulcerated plaque in the nasal dorsum in a patient with multiple myeloma. *Clin Exp Dermatol* 2013; 38: 678–680.
- Cerrati EW and Myssiorek D. External nasal lesion in a middle-aged man. *JAMA Otolaryngol Head Neck Surg* 2014; 140: 569–570.
- Suner L and Mathis S. Disseminated cryptococcosis in bone marrow. *Blood* 2014; 123: 3070.
- Ferraro RA, Ivanidze J, Margolskee E, *et al.* Invasive granulomatous cryptococcal sinusitis in an adult with multiple myeloma. *Clin Imaging* 2017; 41: 65–68.
- Bowcock SJ, Yip K, Majumder K, *et al.* Cryptococcosis in late stage multiple myeloma: consider it. *Br J Haematol* 2017; 176: 687.
- Sato S, Kambe E and Tamai Y. Disseminated cryptococcosis in a patient with multiple myeloma treated with daratumumab, lenalidomide, and dexamethasone. *Int Med (Tokyo, Japan)* 2019; 58: 843–847.
- Stepman G, Sanapati J, Dabb K, *et al.* Cryptococcal meningoencephalitis 10 years after bone marrow transplant in a patient with multiple myeloma. *Cureus* 2020; 12: e12094.
- Sassine J and Kontoyiannis DP. A 77-year-old man with multiple myeloma and a lytic bone lesion. *Am J Med* 2021; 134: 860–862.
- Raheem A, Rathish B, Charles D, *et al.* Pneumococcal bacteremia and cryptococcal meningitis dual infection in a patient with multiple myeloma. *Cureus* 2021; 13: e15089.
- Teh BW, Harrison SJ, Worth LJ, *et al.* Risks, severity and timing of infections in patients with multiple myeloma: a longitudinal cohort study in the era of immunomodulatory drug therapy. *Br J Haematol* 2015; 171: 100–108.
- Teh BW, Teng JC, Urbancic K, *et al.* Invasive fungal infections in patients with multiple myeloma: a multi-center study in the era of novel myeloma therapies. *Haematologica* 2015; 100: e28–e31.
- Tisi MC, Hohaus S, Cuccaro A, *et al.* Invasive fungal infections in chronic lymphoproliferative

- disorders: a monocentric retrospective study. *Haematologica* 2017; 102: e108–e111.
24. Lortholary O, Ascioğlu S, Moreau P, *et al.* Invasive aspergillosis as an opportunistic infection in nonallografted patients with multiple myeloma: a European Organization for Research and Treatment of Cancer/ Invasive Fungal Infections Cooperative Group and the Intergroupe Français du Myélome. *Clin Infect Dis* 2000; 30: 41–46.
 25. Page MJ, McKenzie JE, Bossuyt PM, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Bmj* 2021; 372: n71.
 26. Mtibaa L, Beji I, Ayoub M, *et al.* Neuromeningeal cryptococcosis revealing IgA- λ multiple myeloma. *Med Mycol Case Rep* 2021; 32: 30–33.
 27. Sayhi S, Bouzidi S, Beji I, *et al.* Neuro-meningeal cryptococcal infection revealing a multiple myeloma. *Pan Afr Med J* 2020; 36: 324.
 28. Chastain DB, Henao-Martínez AF, Dykes AC, *et al.* Missed opportunities to identify cryptococcosis in COVID-19 patients: a case report and literature review. *Ther Adv Infect Dis* 2022; 9: 20499361211066363.
 29. Hoy SM. Pomalidomide: a review in relapsed and refractory multiple myeloma. *Drugs* 2017; 77: 1897–1908.
 30. Ruchelman AL, Man H-W, Zhang W, *et al.* Isosteric analogs of lenalidomide and pomalidomide: synthesis and biological activity. *Bioorgan Med Chem Lett* 2013; 23: 360–365.
 31. Huffnagle GB, Toews GB, Burdick MD, *et al.* Afferent phase production of TNF- α is required for the development of protective T cell immunity to *Cryptococcus neoformans*. *J Immunol* 1996; 157: 4529–4536.
 32. Bauman SK, Huffnagle GB and Murphy JW. Effects of tumor necrosis factor alpha on dendritic cell accumulation in lymph nodes draining the immunization site and the impact on the anticryptococcal cell-mediated immune response. *Infect Immun* 2003; 71: 68–74.
 33. Tsiodras S, Samonis G, Boumpas DT, *et al.* Fungal infections complicating tumor necrosis factor alpha blockade therapy. *Mayo Clin Proc* 2008; 83: 181–194.
 34. Haslett PA, Corral LG, Albert M, *et al.* Thalidomide costimulates primary human T lymphocytes, preferentially inducing proliferation, cytokine production, and cytotoxic responses in the CD8+ subset. *J Exp Med* 1998; 187: 1885–1892.
 35. D’Souza C, Prince HM and Neeson PJ. Understanding the role of T-cells in the antimyeloma effect of immunomodulatory drugs. *Front Immunol* 2021; 12: 632399.
 36. Quach H, Ritchie D, Stewart AK, *et al.* Mechanism of action of immunomodulatory drugs (IMiDS) in multiple myeloma. *Leukemia* 2010; 24: 22–32.
 37. Nucci M and Anaissie E. Infections in patients with multiple myeloma in the era of high-dose therapy and novel agents. *Clinical Infectious Diseases* 2009; 49: 1211–1225.
 38. Goldman DL, Lee SC, Mednick AJ, *et al.* Persistent *Cryptococcus neoformans* pulmonary infection in the rat is associated with intracellular parasitism, decreased inducible nitric oxide synthase expression, and altered antibody responsiveness to cryptococcal polysaccharide. *Infect Immun* 2000; 68: 832–838.
 39. Ristow LC and Davis JM. The granuloma in cryptococcal disease. *PLoS Pathog* 2021; 17: e1009342.
 40. Henao-Martínez AF, Chastain DB and Franco-Paredes C. Treatment of cryptococcosis in non-HIV immunocompromised patients. *Curr Opin Infect Dis* 2018; 31: 278–285.
 41. Perfect JR, Dismukes WE, Dromer F, *et al.* Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2010; 50: 291–322.