

Chapman University

Chapman University Digital Commons

Biology, Chemistry, and Environmental Sciences
Faculty Articles and Research

Science and Technology Faculty Articles and
Research

9-4-2023

A Comparison of Methods for Excluding Light from Stems to Evaluate Stem Photosynthesis

Nadia A. Valverdi

Camilla Acosta

Gabriella R. Dauber

Gregory R. Goldsmith

Eleinis Ávila-Lovera

Follow this and additional works at: https://digitalcommons.chapman.edu/sees_articles

A Comparison of Methods for Excluding Light from Stems to Evaluate Stem Photosynthesis

Comments

This article was originally published in *Applications in Plant Sciences*, volume 11, in 2023. <https://doi.org/10.1002/aps3.11542>

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Copyright

The authors

PROTOCOL NOTE

A comparison of methods for excluding light from stems to evaluate stem photosynthesis

Nadia A. Valverdi^{1,2}  | Camilla Acosta¹ | Gabriella R. Dauber¹ |
 Gregory R. Goldsmith¹  | Eleinis Ávila-Lovera^{1,3} 

¹Schmid College of Science and Technology, Chapman University, Orange, California, USA

²Estación Experimental Agropecuaria Catamarca, Instituto Nacional de Tecnología Agropecuaria Catamarca – La Rioja, Catamarca, Argentina

³Smithsonian Tropical Research Institute, Balboa, Ancón, Panama City, Panama

Correspondence

Nadia A. Valverdi, Estación Experimental Agropecuaria Catamarca, Instituto Nacional de Tecnología Agropecuaria Catamarca – La Rioja, Ruta 33 Km 4, 4705 Sumalao, Catamarca, Argentina.
 Email: nadiavalverdi@hotmail.com

Abstract

Premise: A comparison of methods using different materials to exclude light from stems to prevent stem CO₂ exchange (i.e., photosynthesis), without affecting stem conductance to water vapor, surface temperature, and relative humidity, was conducted on stems of avocado trees in California.

Methods and Results: The experiment featured three materials: aluminum foil, paper-based wrap, and mineral-based paint. We examined stem CO₂ exchange with and without the light exclusion treatments. We also examined stem surface temperature, relative humidity, and photosynthetic active radiation (PAR) under the cover materials. All materials reduced PAR and stem CO₂ exchange. However, aluminum foil reduced stem surface temperature and increased relative humidity.

Conclusions: Methods used to study stem CO₂ exchange through light exclusion have historically relied on methods that may induce experimental artifacts. Among the methods tested here, mineral-based paint effectively reduced PAR without affecting stem surface temperature and relative humidity around the stem.

KEYWORDS

stem CO₂ exchange, stem light exclusion, stem surface temperature, stem water vapor conductance

Resumen

Premisa: Una comparación de diferentes métodos utilizando distintos materiales para bloquear la luz de los tallos y así reducir el intercambio de CO₂ (fotosíntesis) sin afectar la conductancia del tallo al vapor de agua, su temperatura superficial y la humedad relativa fue llevado a cabo en tallos de árboles de aguacate en California.

Metodología y resultados: El experimento se llevó a cabo utilizando tres materiales: papel de aluminio, papel para envoltura y pintura a base de minerales. Se examinó el intercambio de CO₂ de los tallos con y sin los materiales de bloqueo de la luz. También se examinó la temperatura de la superficie del tallo, la humedad relativa y la radiación fotosintéticamente activa (PAR por sus siglas en inglés) debajo de los materiales usados para bloquear la luz. Todos los materiales redujeron PAR y el intercambio de CO₂ del tallo. Sin embargo, el papel aluminio redujo también la temperatura de la superficie del tallo y aumento la humedad relativa.

Conclusiones: Los métodos utilizados para estudiar el intercambio de CO₂ de los tallos con el ambiente a través del bloqueo de la luz han sido métodos que pueden generar alteraciones no deseadas. Entre los métodos evaluados aquí, la pintura de base mineral fue efectiva reduciendo PAR sin alterar la temperatura superficial del tallo ni la humedad relativa alrededor de este.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Applications in Plant Sciences* published by Wiley Periodicals LLC on behalf of Botanical Society of America.

Over the past three decades, there has been growing interest in understanding the impact of stem photosynthesis on whole plant carbon gain and water transport (Ávila et al., 2014; Liu et al., 2019; Berry et al., 2021; Tomasella et al., 2021). To study this, researchers have used various strategies to block light from reaching the stem surface and reduce stem gas exchange, i.e., photosynthesis. Methods used in previous studies include nets, aluminum foil, or a combination of both (Table 1). The most commonly used method is loosely wrapping aluminum foil around the stems because of its affordability, ease of application (although time consuming), and high efficacy in blocking light (de Roo et al., 2020a). However, aluminum foil may introduce unwanted experimental artifacts by altering stem surface temperature and relative humidity. Given that photosynthesis is a chemical process that can be heavily influenced by environmental factors like temperature, radiation, and relative humidity, and that stem photosynthesis rates are generally lower than those in leaves (Valverdi et al., 2023), the impact of light exclusion methods on these physical factors needs to be taken into consideration.

Avocado trees have young and mature green stems; this characteristic sets them apart from other fruit trees and may play an essential role in their water and carbon balances through stem photosynthesis (Esteban et al., 2010). In this study, we compared new and existing methods for excluding light from stems using three different materials including aluminum foil, paper wrap, and mineral-based paint. Our

aim was to effectively halt stem gas exchange without significantly altering the surface temperature and relative humidity around the stem or affecting stem conductance to water vapor.

METHODS AND RESULTS

This experiment was carried out on field-grown avocado trees of the Gem cultivar ($n = 5$) in a common garden at the South Coast Extension and Research Center (SCERC) of the University of California in Irvine, California, USA. To investigate the effects of different light restriction methods on stem photosynthesis, we selected four sun-exposed secondary lateral branches on each tree. The branches were at a height midway to the tree canopy and one to two years old to ensure they were photosynthetically active.

We applied four light restriction methods to the branches: the first branch was coated with three layers of green mineral-based paint (White Wash Plant Guard; IV Organics, Los Angeles, California, USA); a second branch was covered with aluminum foil (Reynolds Wrap; Alcoa, Pittsburg, Pennsylvania, USA); a third branch was covered using a light brown, breathable paper wrap (model no. 350, 6 m long \times 10 cm high; Bond Manufacturing, Antioch, California, USA) secured with adhesive paper tape (Roll Products, St. Marys, Kansas, USA) (Appendix 1, Figure A1); and a fourth branch was left uncovered as a control.

TABLE 1 Methods used for stem light exclusion in different tree species.

| Light exclusion method used | Primary research question | Plant species | References |
|-----------------------------|--------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------|-----------------------------|
| Aluminum foil | What is the contribution of stem photosynthesis to growth in this plant species? | <i>Cytisus scoparius</i> | Bossard and Rejmanek (1992) |
| Light-reducing gauze | The importance of stem-internal carbon re-fixation. | <i>Populus tremula</i> and <i>Fagus sylvatica</i> | Wittmann et al. (2001) |
| Light-reducing net | Light-modulation of cortical CO ₂ -refixation. | <i>Betula pendula</i> | Wittmann et al. (2005) |
| Cloth and aluminum foil | Contribution of woody tissue photosynthesis to trunk growth and bud development. | <i>Prunus ilicifolia</i> , <i>Umbellularia californica</i> , and <i>Arctostaphylos manzanita</i> | Saveyn et al. (2010) |
| Aluminum foil | The role of corticular photosynthesis in wood production in smooth-barked branches. | <i>Eucalyptus miniata</i> | Cernusak and Hutley (2011) |
| Aluminum foil | The role of branch photosynthesis in tree functioning. | <i>Rhizophora apiculata</i> , <i>Ceriops australis</i> , and <i>Avicennia marina</i> | Schmitz et al. (2012) |
| Aluminum foil | The role of woody tissue photosynthesis in tree functioning. | <i>Populus nigra</i> 'Monviso' | Bloemen et al. (2013) |
| Aluminum foil | The role of woody tissue photosynthesis in tree functioning under drought. | <i>Populus nigra</i> 'Monviso' | Bloemen et al. (2016) |
| Aluminum foil | The role of woody tissue photosynthesis in stem carbon cycling along a gradient of water availability. | <i>Populus tremula</i> | de Roo et al. (2020b) |
| Aluminum foil | Woody tissue photosynthesis under elevated atmospheric CO ₂ concentration. | <i>Populus tremula</i> | de Roo et al. (2020a) |
| Shading net | The role of depletion of non-structural carbohydrates and xylem vulnerability to embolism. | <i>Populus nigra</i> | Tomasella et al. (2021) |

For gas exchange measurements, we used a portion of the branch with a diameter of approximately 0.5–0.8 cm and a length of 3 cm, which could fit in the gas exchange chamber (3 × 3 cm leaf chamber; LI-COR Biosciences, Lincoln, Nebraska, USA). We performed gas exchange measurements after one and two weeks of covering the branches (i.e., at weeks 2 and 3 of the experiment) to evaluate the effects of light blockage on stem gas exchange over time.

Gas exchange measurements were taken between 1100 hours and 1400 hours using an infrared gas analyzer (6800, LI-COR Biosciences) with a 3 × 3 cm leaf chamber in which the gaskets were lined with rubber foam and Terostat adhesive (Teroson; Henkel Corporation, California, USA), allowing for a tight seal around the stem with no significant leaks (Ávila-Lovera et al., 2017). Measurements were performed at 410 $\mu\text{mol}\cdot\text{mol}^{-1}$ of CO_2 , 1500 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ of PAR, 400 $\mu\text{mol}\cdot\text{s}^{-1}$ of flow rate, 10,000 RPM fan speed, 50% relative humidity, and 25°C temperature. Because the stems did not cover the whole leaf chamber, gas exchange values were recalculated using the stem surface area, which was calculated using the formula of the area of a cylinder without the top and bottom circles (i.e., $2\pi \cdot r \cdot l$, where r = the radius of the stem used and l = the length). Results showed that all cover materials (paint, aluminum, and paper) effectively reduced the stem CO_2 exchange compared to the control ($P = 0.04$) (Figure 1A). The paper wrap demonstrated the greatest absolute reduction in CO_2 exchange rates relative to the control. On the other hand, there were no significant differences between treatments for bark conductance (g_{bark}), although the paper and aluminum treatments tended to increase g_{bark} when compared to the control (Figure 1B).

To measure stem surface temperature, we used a copper-constantan thermocouple (Extech model 421509 with Type K Thermocouple; Extech Instruments, Nashua, New Hampshire, USA), while relative humidity was measured with a precision psychrometer (RH390; Extech Instruments). Photosynthetic active radiation (PAR) was measured with a light meter (Li-250A, LI-COR Biosciences) at 1200 hours on the same days that gas exchange measurements were made. To measure PAR, we secured each of the different cover materials (paper wrap, aluminum foil, or a transparent polyethylene terephthalate [PET] plastic sheet with three layers of paint) in separate cardboard frames. The frames were held parallel to the ground under a clear sky above the light meter sensor (Appendix 1, Figure A2). Our results showed that all cover materials effectively reduced PAR to nearly zero ($P < 0.001$) (Figure 2A). We found that the covers had a significant effect on stem surface temperature ($P = 0.02$), with aluminum foil having the lowest value. Relative humidity was not significantly affected by the covers ($P = 0.5$). In general, the aluminum foil cover had the lowest stem temperature and the highest relative humidity (Figure 2B, C).




FIGURE 1 Stem CO_2 exchange (A) and bark conductance to water vapor (B) for avocado stems covered with mineral-based paint, aluminum foil, and paper wrap as light exclusion treatments, as well as a control ($n = 5$, points represent mean \pm SE).

It is challenging to provide an accurate cost comparison of these materials due to fluctuating prices and differences in the amount of branch material to be covered. For the new mineral-based paint method, we estimate from a different experiment that we used ca. 30 mL of paint per layer per potted tree (2 m tall). The cost of this paint was \$32.50 USD for 473 mL (i.e., 1 U.S. pint), making it the most expensive method. However, the paint creates very little waste compared to aluminum foil and paper wrap, which are difficult to adjust due to the different branch sizes and bifurcations. Additionally, applying the paint is relatively fast (e.g., it dries quickly between layers), especially in comparison to the paper. It is worth noting that the paint is water-soluble, so it will wash off when it becomes wet from rain/irrigation. While the paint is durable, we have previously observed cracking on trees in the field after 4–6 weeks (personal observation).




FIGURE 2 Photosynthetically active radiation (PAR) (A), stem surface temperature (B), and relative humidity (C) for avocado stems covered with mineral-based paint, aluminum foil, and paper wrap as light exclusion treatments, as well as a control ($n = 5$, points represent mean \pm SE).

Statistical analysis

To test the differences among treatments, we used a linear mixed model with time (data from week 1 and week 2) as a random effect using the package lmerTest in R version 3.4.0 (R Core Team, 2017).

CONCLUSIONS

In this study, we compared different methods for creating a light exclusion environment to assess stem CO_2 exchange. In addition to the commonly used aluminum foil (see Table 1), we proposed new materials, such as using paper wrap and mineral-based paint on avocado trees. All of these cover methods effectively blocked light from reaching the

stem and altered stem CO_2 exchange without significantly affecting bark conductance to water vapor.

However, we observed that the use of aluminum foil resulted in reduced stem surface temperature (Figure 2), leading to mold formation (personal observation). The paper wrap was found to increase stem surface temperature, but we did not observe mold formation using this material. The paper wrap, however, was the most time-consuming method as it required tape to hold the paper in place, and the material was not very malleable. Overall, we found that the paint cover was the easiest and least time-consuming method to use, and it also had the least impact on stem temperature and relative humidity while effectively blocking light and reducing stem CO_2 exchange. This method is especially useful for experiments carried out in dry climates such as Southern and Central California, where the irrigation system does not wet the tree stems.

AUTHOR CONTRIBUTIONS

N.A.V., C.A., G.R.D., G.R.G. and E.A.L. made substantial contributions to conception and design, acquisition of data, analysis, and interpretation of data; were involved in drafting the manuscript or revising it critically for important intellectual content; and approved the final version of the manuscript. All authors have agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

ACKNOWLEDGMENTS

The authors thank the staff of the South Coast Extension and Research Center (SCREC) for their assistance in the field, and Mary Lu Arpaia and Eric Focht (University of California, Riverside) for their feedback while developing this project. This project was funded by the U.S. Department of Agriculture National Institute of Food and Agriculture (USDA-NIFA; award #2020-67014-30915 to E.A.L. and G.R.G.).

DATA AVAILABILITY STATEMENT

All supporting data are available within the published article as Supporting Information (Appendix S1).

ORCID

Nadia A. Valverdi  <http://orcid.org/0000-0002-6016-7764>

Gregory R. Goldsmith  <http://orcid.org/0000-0003-3567-8949>

Eleinis Ávila-Lovera  <http://orcid.org/0000-0003-3529-3600>

REFERENCES

- Ávila, E., A. Herrera, and W. Tezara. 2014. Contribution of stem CO_2 fixation to whole-plant carbon balance in nonsucculent species. *Photosynthetica* 52(1): 3–15. <https://doi.org/10.1007/s11099-014-0004-2>
- Ávila-Lovera, E., A. J. Zerpa, and L. S. Santiago. 2017. Stem photosynthesis and hydraulics are coordinated in desert plant species. *New Phytologist* 216(4): 1119–1129. <https://doi.org/10.1111/nph.14737>

- Berry, Z. C., E. Ávila-Lovera, M. E. de Guzman, K. O'Keefe, and N. C. Emery. 2021. Beneath the bark: Assessing woody stem water and carbon fluxes and its prevalence across climates and the woody plant phylogeny. *Frontiers in Forests and Global Change* 4: 675299. <https://doi.org/10.3389/ffgc.2021.675299>
- Bloemen, J., L. Overlaet-Michiels, and K. Steppe. 2013. Understanding plant responses to drought: How important is woody tissue photosynthesis? Proc. 9th International Workshop on Sap Flow. *Acta Horticulturae* 991: 149–156.
- Bloemen, J., L. L. Vergeynst, L. Overlaet-Michiels, and K. Steppe. 2016. How important is woody tissue photosynthesis in poplar during drought stress? *Trees - Structure and Function* 30(1): 63–72. <https://doi.org/10.1007/s00468-014-1132-9>
- Bossard, C. C., and M. Rejmanek. 1992. Why have green stems? *Functional Ecology* 6(2): 197–205.
- Cernusak, L. A., and L. B. Hutley. 2011. Stable isotopes reveal the contribution of cortical photosynthesis to growth in branches of *Eucalyptus miniata*. *Plant Physiology* 155(1): 515–523. <https://doi.org/10.1104/pp.110.163337>
- de Roo, L., F. Lauriks, R. L. Salomón, J. Oleksyn, and K. Steppe. 2020a. Woody tissue photosynthesis increases radial stem growth of young poplar trees under ambient atmospheric CO₂ but its contribution ceases under elevated CO₂. *Tree Physiology* 40(11): 1572–1582. <https://doi.org/10.1093/treephys/tpaa085>
- de Roo, L., R. L. Salomón, and K. Steppe. 2020b. Woody tissue photosynthesis reduces stem CO₂ efflux by half and remains unaffected by drought stress in young *Populus tremula* trees. *Plant Cell and Environment* 43(4): 981–991. <https://doi.org/10.1111/pce.13711>
- Esteban, R., B. Olascoaga, J. M. Becerril, and J. I. García-Plazaola. 2010. Insights into carotenoid dynamics in non-foliar photosynthetic tissues of avocado. *Physiologia Plantarum* 140(1): 69–78. <https://doi.org/10.1111/j.1399-3054.2010.01385.x>
- Liu, J., L. Gu, Y. Yu, P. Huang, Z. Wu, Q. Zhang, Y. Qian, et al. 2019. Cortical photosynthesis drives bark water uptake to refill embolized vessels in dehydrated branches of *Salix matsudana*. *Plant, Cell and Environment* 42(9): 2584–2596. <https://doi.org/10.1111/pce.13578>
- Saveyn, A., K. Steppe, N. Ubierna, and T. E. Dawson. 2010. Woody tissue photosynthesis and its contribution to trunk growth and bud development in young plants. *Plant, Cell and Environment* 33(11): 1949–1958. <https://doi.org/10.1111/j.1365-3040.2010.02197.x>
- Schmitz, N., J. J. G. Egerton, C. E. Lovelock, and M. C. Ball. 2012. Light-dependent maintenance of hydraulic function in mangrove branches: Do xylary chloroplasts play a role in embolism repair? *New Phytologist* 195(1): 40–46. <https://doi.org/10.1111/j.1469-8137.2012.04187.x>
- Tomasella, M., V. Casolo, S. Natale, F. Petruzzellis, W. Kofler, B. Beikircher, S. Mayr, and A. Nardini. 2021. Shade-induced reduction of stem nonstructural carbohydrates increases xylem vulnerability to embolism and impedes hydraulic recovery in *Populus nigra*. *New Phytologist* 231(1): 108–121. <https://doi.org/10.1111/nph.17384>
- Valverdi, N. A., P. Guzman-Delgado, C. Acosta, G. R. Dauber, G. R. Goldsmith, and E. Avila-Lovera. 2023. Is avocado green stem photosynthesis related to plant hydraulics? *Acta Horticulturae* 1372: 129–136. <https://doi.org/10.17660/ActaHortic.2023.1372.17>
- Wittmann, C., G. Aschan, and H. Pfan. 2001. Leaf and twig photosynthesis of young beech (*Fagus sylvatica*) and aspen (*Populus tremula*) trees grown under different light regimes. *Basic and Applied Ecology* 2(2): 145–154. <https://doi.org/10.1078/1439-1791-00047>
- Wittmann, C., H. Pfan, and F. Pietrini. 2005. Light-modulation of cortical CO₂-refixation in young birch stems (*Betula pendula* Roth.). *Phyton* 45(3): 195–212.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Appendix S1. Stem photosynthesis (A_{stem}), stomata conductance (gs), air temperature (T), relative humidity (HR), and photosynthetically active radiation (PAR) for three light exclusion treatments (paint, aluminum, paper) and an untreated control on avocado plants ($n = 5$).

How to cite this article: Valverdi, N. A., C. Acosta, G. R. Dauber, G. R. Goldsmith, and E. Ávila-Lovera. 2023. A comparison of methods for excluding light from stems to evaluate stem photosynthesis. *Applications in Plant Sciences* 11: e11542. <https://doi.org/10.1002/aps3.11542>

Appendix 1. Protocol for blocking the light from stems in avocado trees.

Materials

- Aluminum foil wrap (Reynolds Wrap; Alcoa, Pittsburg, Pennsylvania, USA)
- Paper wrap (model no. 350, 6 m long × 10 cm high; Bond Manufacturing, Antioch, California, USA)
- Mineral-based paint (White Wash Plant Guard; IV Organics, Los Angeles, California, USA)
- Paper tape (Roll Products, St. Marys, Kansas, USA)
- Scissors
- Paintbrush
- Water
- Terostat adhesive (Teroson; Henkel Corporation, California, USA)

Procedure

1. Select four sun-exposed branches in the tree.
2. For the first stem, apply two or three layers of mineral-based paint to the entire stem using a paintbrush, leaving the leaves and petioles uncovered.
3. For the second branch, cover all stem sections with aluminum foil, leaving only the leaves and petioles uncovered.
4. For the third branch, cover as much of the stem as possible with paper wrap, leaving only the leaves and petioles uncovered.
5. Leave the fourth branch uncovered as a control.
6. On all four branches, select a piece of stem between 5 and 8 mm in diameter to measure stem gas exchange and wrap two pieces of Terostat adhesive approximately 3 cm apart to ensure the stem pieces will fit into the gas exchange analyzer chamber.
7. Cut a small piece of aluminum foil or paper wrap, or apply paint to cover the piece of stem between the two Terostat pieces, except for the uncovered control (Figure A1).
8. One to two weeks after covering the branches, measure stem gas exchange as explained in the Methods section.
9. For PAR measurements, make two frames using cardboard or any available material to hold the paper wrap and a polyethylene sheet covered with mineral



FIGURE A1 Three light exclusion methods used to reduce stem CO_2 exchange in avocado stems, including (A) branch covered with mineral-based paint, (B) aluminum foil-wrapped branch, (C) paper-wrapped branch, and (D) control branch.




FIGURE A2 Examples of cardboard frames containing different cover materials to hold above the light meter and measure photosynthetic active radiation (PAR) in the field under a clear sky: (A) paper-based wrap and (B) transparent PET covered with mineral-based paint.

- paint. Place the PAR sensor on a flat surface perpendicular to the floor and place the frame with the light exclusion material above it (Figure A2).
10. For temperature and relative humidity measurements, place the sensor's head under a loose covering material

without uncovering the branch, touching the stem surface.

Note: Make sure the trees are well irrigated and monitor the covered branches for any signs of stress or damage.